

TYPES AND  $\lambda$ -CALCULUS

# Problem Sheet 7

- \*\* 1. Suppose closed term  $M$  has a normal form  $\lambda x.xx$ . Prove that all reducts of  $M$  are untypable, i.e.  $M \triangleright^* N$  implies  $N$  untypable.

Solution —————

Assume  $M$  has normal form  $\lambda x.xx$  and suppose  $M \triangleright^* N$ . Then we have  $M \triangleright^* \lambda x.xx$  and  $M \triangleright^* N$  so, by Confluence,  $N$  and  $\lambda x.xx$  have a common reduct. But  $\lambda x.xx$  is a normal form, so we must have that  $N \triangleright^* \lambda x.xx$ . We claim that  $N$  is therefore untypable. To see why, suppose that  $N$  were typable, then by Subject Reduction, so is  $\lambda x.xx$ . However, we know that this is impossible.

- \*\* 2.

- (a) Prove the following result by induction on  $M$ :

Let  $B$  be a type and  $N$  a term. For all terms  $M$ , types  $A$  and environments  $\Gamma$ : if  $\Gamma, x:B \vdash M : A$  and  $\Gamma \vdash N : B$  then  $\Gamma \vdash M[N/x] : A$ .

- (b) Prove the missing case in the proof of Lemma 12.2 from the notes: if  $\Gamma \vdash (\lambda x.M)N : A$  then  $\Gamma \vdash M[N/x] : A$ .

Solution —————

- (a) Suppose  $B$  is a type and  $N$  a term. The rest of the proof is by induction on  $M$ .

- When  $M$  is a variable  $y$ , we proceed as follows. Let  $A$  be a type and  $\Gamma$  an environment. Suppose (i)  $\Gamma, x:B \vdash y : A$  and (ii)  $\Gamma \vdash N : B$ . Now we analyse cases on  $x = y$ ?

- If  $x = y$  then, by definition of substitution,  $M[N/x] = N$  and, by inversion on (i), we obtain that  $B = A$ . Then our goal is to show  $\Gamma \vdash N : A$ , which is exactly (ii).
- Otherwise, by definition of substitution,  $M[N/x] = M$ . Our goal is to show  $\Gamma \vdash M : A$ . This follows immediately from (i) by Weakening (see the previous problem sheet).
- When  $M$  is a constant  $c$ , we proceed as follows. Let  $A$  be a type and  $\Gamma$  an environment. Suppose (i)  $\Gamma, x:B \vdash c : A$  and (ii)  $\Gamma \vdash N : B$ . By definition,  $c[N/x] = c$  and so our goal is to show  $\Gamma \vdash c : A$ , which follows immediately from (i) by Weakening.
- When  $M$  is of the form  $PQ$ , we assume the induction hypotheses:
  - (IH1) For all  $A', \Gamma'$ : if  $\Gamma', x:B \vdash P : A'$  and  $\Gamma' \vdash N : B$  then  $\Gamma' \vdash P[N/x] : A'$
  - (IH2) For all  $A', \Gamma'$ : if  $\Gamma', x:B \vdash Q : A'$  and  $\Gamma' \vdash N : B$  then  $\Gamma' \vdash Q[N/x] : A'$

Let  $A$  be a type and  $\Gamma$  an environment, then suppose (i)  $\Gamma, x:B \vdash PQ : A$  and (ii)  $\Gamma \vdash N : B$ . It follows from inversion on (i) that there is a type  $C$  such that:

- (a)  $\Gamma, x:B \vdash P : C \rightarrow A$
- (b)  $\Gamma, x:B \vdash Q : C$

Hence, we can obtain from (IH1), with  $A' := C \rightarrow A$  and  $\Gamma' := \Gamma$  that  $\Gamma \vdash P[N/x] : C \rightarrow A$ . From (IH2), with  $A' := C$  and  $\Gamma' = \Gamma$ , we obtain  $\Gamma \vdash Q[N/x] : C$ . Putting these together with (TApp) we obtain  $\Gamma \vdash P[N/x]Q[N/x] : A$ , and by definition of substitution, this is just  $\Gamma \vdash (PQ)[N/x] : A$ , which was our goal.

- When  $M$  is of the form  $\lambda y. P$ , we assume the induction hypothesis:
  - (IH) For all  $A'$  and  $\Gamma'$ , if  $\Gamma', x:B \vdash P : A'$  and  $\Gamma' \vdash N : B$  then  $\Gamma' \vdash P[N/x] : A'$ .

We may also assume, by the variable convention, that  $y$  does not occur freely in  $N$ , is not a subject in  $\Gamma$  and is distinct from  $x$  (otherwise, we rename  $y$  in the abstraction). Let  $A$  be a type and  $\Gamma$  an environment and suppose (i)  $\Gamma, x:B \vdash \lambda y. P : A$  and (ii)  $\Gamma \vdash N : B$ . It follows from (i) by inversion that there are types  $A_1$  and  $A_2$  such that  $A = A_1 \rightarrow A_2$  and (\*)  $\Gamma, x:B, y : A_1 \vdash P : A_2$ . Then it follows from this by (IH), with  $A' = A_2$  and  $\Gamma' = \Gamma \cup \{y : A_1\}$ , that  $\Gamma, y:A_1 \vdash P[N/x] : A_2$ . From this we can immediately infer  $\Gamma \vdash \lambda y. P[N/x] : A_1 \rightarrow A_2$  using (TAbs). By definition of substitution (recall we assumed  $y \neq x$  and  $y \notin \text{FV}(N)$ ) and the identity of  $A$ , this is just our goal  $\Gamma \vdash (\lambda y. P)[N/x] : A$ .

- (b) Now suppose  $\Gamma \vdash (\lambda x. M)N : A$ . By inversion, it follows that there is some type  $B$  such that  $\Gamma \vdash \lambda x. M : B \rightarrow A$  and  $\Gamma \vdash N : B$ . By inversion

on the former, we deduce that  $\Gamma, x : B \vdash M : A$ . Then the result follows from the previous part.

\*\* 3. Find pure terms that inhabit the following types (no need for justification):

- (a)  $(a \rightarrow a \rightarrow b) \rightarrow a \rightarrow b$
- (b)  $(a \rightarrow b) \rightarrow (a \rightarrow c) \rightarrow (b \rightarrow c \rightarrow d) \rightarrow a \rightarrow d$
- (c)  $((a \rightarrow a \rightarrow b) \rightarrow a \rightarrow b) \rightarrow c \rightarrow c$

Solution

---

- (a)  $\lambda x y. x y y$
- (b)  $\lambda w x y z. y (x z) (y z)$
- (c)  $\lambda x. x (\lambda y z. y z z)$

\*\* 4.

- (a) Find a *pure* term that inhabits the type  $((a \rightarrow b) \rightarrow b) \rightarrow a \rightarrow b$ .
- (b) Give the corresponding proof of the corresponding formula.

Solution

---

- (a)  $\lambda x y. x (\lambda z. z y)$
- (b) Suppose  $(a \Rightarrow b) \Rightarrow b \Rightarrow b$  (x) and assume  $a$  (y). We claim that  $(a \Rightarrow b) \Rightarrow b$ , the proof is as follows: suppose  $a \Rightarrow b$  (z), then applying this to assumption (y) we get  $b$ . Returning to our original proof, from this and (x) we obtain  $b$ , as required.

\*\*\* 5. The reason that we don't study full PCF in connection with the Curry-Howard correspondence is the presence of fix.

- (a) Use fix to show that every type is inhabited by some *PCF term* (not necessarily pure).
- (b) What is the consequence for the Curry-Howard correspondence extended to full PCF?

Solution

---

- (a) Every type  $A$  is inhabited by the term  $\text{fix } (\lambda x. x)$ . We can construct the following derivation.

$$\frac{\frac{}{\vdash \text{fix} : (A \rightarrow A) \rightarrow A} \text{ (TFix)} \quad \frac{\frac{}{x : A \vdash x : A} \text{ (TVar)} \quad \frac{}{\vdash \lambda x. x : A \rightarrow A} \text{ (TAbs)}}{\vdash \text{fix } (\lambda x. x) : A} \text{ (TApp)}$$

- (b) Hence, by the Curry-Howard correspondence, this would lead to a proof system in which every formula was provable – i.e. an inconsistent logic!

\*\*\* 6. The following property is called Subject Invariance:

if  $M \approx N$  and  $\Gamma \vdash M : A$  then  $\Gamma \vdash N : A$

Is this property true for our type system? Either prove it or give a counterexample.

Solution

---

It is not true in our system. To see why, take e.g.  $\lambda y. y \approx (\lambda x. (\lambda y. y))(\lambda x. xx)$ . We have  $\vdash \lambda y. y : a \rightarrow a$  so the hypotheses of the implication are satisfied, but  $(\lambda x. (\lambda y. y))(\lambda x. xx)$  is untypable because it includes  $\lambda x. xx$  as a subterm.