

Report Title: Example Report

Group 3
Department / Course

November 16, 2025

Contents

1	Executive Summary	2
2	Data Description	3
2.1	Key Variables	4
3	Bayesian regression model (BRM)	4
3.1	Reason for BRM	4
3.2	How BRM works	4
3.3	Model Selection	5
4	Key Variables	5
5	Priors	5
5.1	Testing Model	5
6	Calculate V_d:	5
7	Further research:	5
7.1	REPORT	5
8	Exclusive summary	5

9 Overview of Datasets	5
9.1 Variable Description	5
10 Model selection regarding key variable ‘beta60’	6
10.1 Advantage of current method	6
10.2 New model motivation	6
11 Bayesian model	6
11.1 Model fitting example	6
12 Testing another assumption(condition):	6
12.1 variable explanation	6

1 Executive Summary

In the UK it is a criminal offence to drive a motor vehicle with a blood or breath alcohol concentration above the prescribed limit. When a person is arrested for driving under the influence of alcohol it is not usually possible to perform an accurate test of the level of alcohol in the blood or breath immediately. Breath tests can be used as an initial screening tool at the scene, but these are not sufficiently accurate for prosecution. Instead, people are taken to a police station or hospital, where the test can be carried out using proper laboratory protocols. As the body clears alcohol from the blood through time this means that if the individual was over the limit, the measured blood alcohol concentration (BAC) will be lower at the time of measurement than it was when the person was driving a motor vehicle. To deal with this situation, If the BAC after time t (hours) is measured as C_t (g/kg), the BAC at time 0 is estimated as $C_0 = C_t - \beta t$, where β (g/kg/h) is BAC elimination rate.

The key point is how to find a precise β to estimate C_0 . Forensic scientists currently 2.5% percentile of β distribution constructing from samples as the estimated β value for every individuals. This method is obviously not rigorous enough for the courts, since:

- The courts will be forced to make decisions under estimated β if we only give a single estimation of β , since the calculated C_0 is either over or under the legal limit.
- Differences between individuals are ignored, for example age and sex, which may affect β .
- β value at 2.5% is over conservative and most uncertainties are hiding.

In this report, we will introduce a Bayesian regression model with considering the heterogeneity between individuals, the model gives a posterior distribution of β by using both samples and prior

knowledge. Then we randomly simulate 4000 β values from posterior distribution and find out the probability of the person's BAC over limit while driving.

In real world, β estimation is quite difficult, it depends on the individuals' liver condition, drinking habits, genetic, diet, etc. As we don't have corresponding data, in our regression model of β is mainly dominated by gender, but we still find some useful variables:

-Gender: Female's BAC elimination rate is larger than male on average, it is explained by liver weight represents a greater fraction of lean body mass in the female gender[2].

-Drinking time after BAC peak: If the person keeps drinking after BAC reaches peak, the measured β will be smaller since β is measured from start at BAC peak time till the end.

When it is too late to use a blood or breath test and the only information available is eyewitness testimony of the quantity of alcohol consumed. We have to use Widmark's equation:

$$C_t = \frac{A}{Weight \times V_d} - \beta t$$

where A is Amount of Alcohol Consumed (g), V_d is the volume of distribution that needs to be found. Forensic scientists use the same method again to estimate V_d separately with β . But V_d and β are not independent, so we build a joint Bayesian regression model, which can simulate them together to solve with the correlation. Then again after simulation, each C_t is calculated by each pair of β and V_d , so expert witness can still find a probability of C_t exceeding the limit.

To make it easier, we write the method in a function so other expert witness can also get the results by easily plugging new person's data.

2 Data Description

It is very important to normalize all variables like weight and height. Since β is a small value, the coefficients of large value variables are pretty small, near 0, which will make β estimation worse. Also when we normalize numerical variables, we can use $N(0, 1)$ to be the priors of coefficients.

Then we prefer to β to $\log(\beta)$ since the original β value has a heavy tail. After simulation we can take exponential of the results.

Since β is a constant rate, measured when the BAC curve reaches peak and starts to decrease, so variables like AAC and Maximum BAC will not be used, since they are correlated with the value of BAC, not BAC elimination rate.

From the dataset, most data comes from age smaller than 30 (85/100), so even though age is a sufficient variable for β the model can't really recognize it. Research shows that age doesn't really matter unless the subject has age-related liver disease, but because of moral and ethical, we don't find much research of alcohol test on elderly person with liver disease.

2.1 Key Variables

3 Bayesian regression model (BRM)

3.1 Reason for BRM

Comparing with fitting linear regression model for β_i , which can be expressed as:

$$\hat{\beta}_i = \hat{\gamma}_1 \text{female}_i + \hat{\gamma}_2 \text{male}_i + \hat{\gamma}_3 \text{weight}_i + \hat{\gamma}_4 \text{height}_i + \sigma^2$$

and giving a exact value of $\hat{\beta}$, BRM can model uncertainty from each coefficient γ , since β from each individuals may affect by variables in different level. We provide priors and sample likelihoods for each variables, BRM will apply bayesian rule to each variables and output simulation results of coefficients specifically for each individual:

$$\hat{\beta}_i = \hat{\gamma}_{1,i} \text{female}_i + \hat{\gamma}_{2,i} \text{male}_i + \hat{\gamma}_{3,i} \text{weight}_i + \hat{\gamma}_{4,i} \text{height}_i + \sigma_i^2.$$

Since we have simulation results of $\hat{\beta}$, we can offer a probability for the courts.

3.2 How BRM works

We use *brm* package in R to do this.

The principle theorem behind BRM is Bayes' Rule:

$$p(\theta \mid \text{data}) = \frac{p(\text{data} \mid \theta) p(\theta)}{p(\text{data})}.$$

The θ in the equation is the coefficients of γ_j . For most real models, we can't compute this posterior $p(\theta \mid \text{data})$ analytically. Markov Chain Monte Carlo (MCMC) helps us draw samples from it, which we can then summarize. BRM uses Hamiltonian Monte Carlo (One of MCMC), which is better than standard MCMC (like Metropolis-Hastings) since it is faster and cost less.

After setting priors for γ_j , we need specify a family for β as the the likelihood of the observed data. In our sample, it is better to use the most common normal distribution likelihood, it is normal and can express β values good without restrict samples' shape.

Also for the MCMC simulation method, we set 4 chains, each chains 2000 draws with 500 burn-in draws. Since MCMC is a random walk method through parameter space. One chain is a single run of the MCMC algorithm that generates a sequence of samples, so we need more chains to check the convergence and ensure that all chains convergent to same mode.

For 500 burn-in draws, since MCMC starts from an initial value which often far from the true posterior region, so the chains need some steps to walk to the correct trace. As Figure 1 showed, it is an example of the MCMC process of ‘sexfemale’ when using 4 chains, 50 draws with 30 warm-ups. Clearly that if we keep the first 30 draws, the results will be affected.

3.3 Model Selection

4 Key Variables

5 Priors

5.1 Testing Model

6 Calculate V_d :

in the Widmark’s equation assumption, beta and V_d are independent, so we need to check whether $\text{Cov}(V_d, \text{beta}) = 0$

7 Further research:

casual effect: biased data selection, (e.g. inner correlations between sex and height)

7.1 REPORT

8 Exclusive summary

background Goal - 1, 2, 3 main data + elimination rate explanation method - model we used result

9 Overview of Datasets

9.1 Variable Description

- how many observations do we have? where we get the resource, reference? explain each variables in table ## Preliminary analysis
- initial formula: explain beta60 + - motivation to improve?

variable correlations? Why? hypothesis needed? why 2.5 quantiles?

10 Model selection regarding key variable ‘beta60’

10.1 Advantage of current method

simpler model with only one parameter(beta), easier to calculate, suitable for no dataset situations

10.2 New model motivation

Current model disadvantage: 1. beta is fixed for all individuals, we need to consider the heterogeneity in individuals. 2. Deviated aim: Police Scotland aims at assessing the probabilities that a person is over the limit, while the parameter beta only concentrates on the range between 2.5% and 97.5%.

11 Bayesian model

intro New model (Bayesian) ## Model Selection and Evaluation - why Bayesian? Any advantage, any improvement? More accurate? ... ##explanation ###why prior setted? ## how result reflect?

11.1 Model fitting example

How our new model works on the 70 year-old madam?

12 Testing another assumption(condition):

12.1 variable explanation

explain all variables we used explain needed references

##Testing whether formula reasonable? whether beta60 relate to Vt? how related? hows the result? how new model fit?

#Limitation our model limit data collected reason - cannot reflect the real world model reason
other reason: variables

#N

#Reference Stan Development Team. (n.d.). Prior choice recommendations. GitHub. <https://github.com/stan-dev/stan/wiki/Prior-Choice-Recommendations>

===== »»»> bb3d1a6f16b0f748d3d6a2b63e99aaba40d20726 CTP2E1 -
 beta 1 60
<https://www.sciencedirect.com/science/article/pii/S0379073810000770?via%3Dihub>

P.Y. Kwo, V.A. Ramchandani, S. O'Connor, D. Amann, L.G. Carr, K. Sandrasegaran, K.K. Kopecky, T.K. Li Gender differences in alcohol metabolism: relationship to liver volume and effect of adjusting for body mass *Gastroenterology*, 115 (1998), pp. 1552-1557