Working Paper 342/ Computer Manual 17

THE 1981 CENSUS SMALL AREA STATISTICS FOR WEST YORKSHIRE WARDS: HOW TO ACCESS, EXTRACT AND ANALYSE.

Mark Birkin and Philip Rees

School of Geography University of Leeds LEEDS LS2 9JT UK.

July 1982

CONTENTS

Acknowledgements

Abstract

List of Tables

List of Figures

- 1. INTRODUCTION
 - 1.1 The 1981 Census and its publication
 - 1.2 Plans for use of the Census 1981 SAS by the higher education community
 - 1.3 Local use at the University of Leeds
 - 1.4 Aims of this paper
- 2. THE CENSUS 1981 SAS DATA
 - 2.1 Small areas
 - 2.2 The topics covered
- 3. CENSUS DATA FOR WEST YORKSHIRE WARDS AT THE UNIVERSITY OF LEEDS
 - 3.1 West Yorkshire wards
 - 3.2 The computer files for West Yorkshire wards
 - 3.3 Accessing the census data
- 4. A GENERAL DATA EXTRACTION AND TABULATION PROGRAM
 - 4.1 Introduction
 - 4.2 The format of the data
 - 4.3 The main features of the program
 - 4.4 Instructions for using the program
- 5. TECHNIQUES IN THE EXTRACTION AND FURTHER PROCESSING OF SPECIFIC DATA SETS
 - 5.1 Extraction
 - 5.2 Statistical analysis
 - 5.3 Mapping

6. THE ETHNIC GEOGRAPHY OF LEEDS

- 6.1 Introduction
- 6.2 An ordering and classification of the ethnic groups
- 6.3 The spatial patterns
- 6.4 Interrelationships between the ethnic group distributions
- 6.5 A composite picture of the ethnic geography of Leeds, 1981
- 6.6 Caveats and further analyses

7. CONCLUSIONS

REFERENCES

ACKNOWLEDGEMENTS

The authors would like to thank Martin Clarke and Alan Wilson for their good offices in obtaining the Small Area Statistics for West Yorkshire wards. The Office of Population and Censuses and Surveys deserve our gratitude for producing and supplying the results of the 1981 Census so rapidly. Our thanks are due to John Duke of the University Computing Service for help in overcoming some system difficulties in developing the software.

ABSTRACT

The Small Area Statistics (SAS) from the 1981 Census are now available for analysis. This paper describes a system for accessing, extracting and analysing a subset of the SAS data: the 100% tables for West Yorkshire wards. The data files, extraction and analysis programs described should help researchers prepare for the availability towards the end of 1982 of enumeration district statistics at University regional computing centres as processed by the major package SASPAC.

Since the data are of interest in themselves the paper concludes with a detailed analysis of the ethnic geography of Leeds metropolitan district in April 1981.

LIST OF TABLES

- 1. Ward names in West Yorkshire, Census 1981.
- 2. A list of Census 1981 data and associated files held on the Geography program and census data library permanent disk.
- 3. Ward statistics for persons born in the Caribbean (New Commonwealth countries), Leeds, Census 1981.
- 4. Country of birth statistics for persons resident in Headingley ward, Leeds, Census 1981.
- 5. Key indicators of ethnic residential segregation, Leeds 1981.
- 6. Indices of dissimilarity between pairs of ethnic groups.
- 7. Selected ethnic characteristics of Leeds wards, Census 1981.

LIST OF FIGURES

- 1. Ward boundaries in West Yorkshire, Census 1981
- 2. A listing of a sample memo file
- 3. An example of a census data extraction system
- 4. The SAS data for Aireborough ward, Leeds, Census 1981
- 5. Country of birth page from the coding book for the Census 1981 SAS (OPCS, 1982c)
- Selected records from the SAS for Aireborough ward, Leeds,
 Census 1981
- 7. A listing of the TABULATE program
- 8. Instructions for preparing the data file for input to the TABULATE program
- 9. A sample instructions file for the TABULATE program
- 10. A sample tabulation produced by the TABULATE program
- 11. A listing of the raw data file associated with the sample tabulation
- 12. A system of extracting and analysing ethnic variables
- 13. A listing of the DERIVE program
- 14. A specimen run using the GIMMS package
- 15. Built-up areas and electoral ward boundaries, Census 1981, Leeds metropolitan district
- 16. The proportional distribution of ethnic groups in Leeds, 1981:
 - (1) England born; Scotland to Rest of UK born (Family I)
- 17. The proportional distribution of ethnic groups in Leeds, 1981:
 - (2) Mediterranean to Far East born (Family II), East Africa born (Family III)
- 18. The proportional distribution of ethnic groups in Leeds, 1981:
 - (3) India to Rest of Africa born (Family III); Caribbean to Bangladesh born (Family IV)
- 19. Zonal percentage distribution of ethnic groups in Leeds, 1981:
 - (1) England born; Scotland to Rest of UK born (Family I)

- 20. Zonal percentage distribution of ethnic groups in Leeds, 1981:(2) Mediterranean to Far East born (Family II); East Africa born (Family III)
- 21. Zonal percentage distributions of ethnic groups in Leeds, 1981:(3) India to Rest of Africa born (Family III); Caribbean to Bangladesh born (Family IV)
- 22. Degrees of concentration (location quotients) for ethnic groups in Leeds, 1981: (1) England born; Scotland to Rest of UK born (Family I)
- 23. Degrees of concentration (location quotients) of ethnic groups in Leeds, 1981: (2) Mediterranean to Far East born (Family II); East Africa born (Family III)
- 24. Degrees of concentration (location quotients) of ethnic groups in Leeds, 1981: (3) India to Rest of Africa born (Family III); Caribbean to Bangladesh born (Family IV)
- 25. A scatterplot of Leeds wards in 'ethnic space'

1. INTRODUCTION

1.1 The 1981 Census and its publication

On the 5th April, 1981 a census of the population of England and Wales was carried out by the Office of Population Censuses and Surveys. Such censuses are administered every ten years in the United Kingdom. The 1981 Census provides a comprehensive picture of the state of the British population in terms of the characteristics of individuals, of the households they are members of and of the housing spaces they occupy.

The results of the 1981 Census are currently in the course of publication. Preliminary reports (OPCS, 1981a, 1981b) on the counts of people in administrative areas have been published, and selected tabulations (OPCS, 1982a) for those areas have been produced in the form of OPCS County Monitors. A few County Reports (OPCS, 1982b) are now available and many more are in the pipeline. Within the counties Small Area Statistics (SAS) are being published for enumeration districts, electoral wards and local government districts in a standard set of tables. Publication consists of production on printed paper, on microfilm or on magnetic tape. All this is happening at amazing and unprecedented speed within 18 months of Census day.

1.2 Plans for use of the Census 1981 SAS by the higher education community

Far reaching and ambitious steps have been taken to make the Census 1981 SAS available to the academic community. The Social Science Research Council (SSRC) has purchased a national set which is to be accessed either via the SSRC Data Archive at the University of Essex or via five regional computing centres at the Universities of London, Bristol, Manchester, Newcastle and Edinburgh. The SSRC has also bought software to make possible the easy use of this vast data set in the form of a package called SASPAC prepared at the University of Durham and at the Program Library Unit, University of Edinburgh under the direction of Professor David Rhind. Towards the end of 1982 the SAS data should be mounted on special disk packs at the regional computing centres and accessible via SASPAC from all Universities via the regional computing networks.

1.3 Local use at the University of Leeds

In order to obtain experience with the SAS prior to and in preparation for the coming online of the 'national' system, a set of ward level SAS was purchased for the School of Geography through the good offices of Professor Alan Wilson and Martin Clarke using funds available from an SSRC Grant. These ward data have been stored on diskspace assigned to the Geography program and data library (GEO6LIB) for general use by members of the School (staff, postgraduates, undergraduates).

1.4 Aims of this paper

1

This paper is designed to serve as a guide to the SAS data for West Yorkshire and as an introduction to the national SAS.

The general structure of the Census 1981 SAS data is described in section two of the paper and the key documents that need to be consulted are referenced.

The particular structure of the 1981 Census SAS for West Yorkshire wards as held on the Leeds University computer is outlined in the third section of the paper.

The fourth section of the paper presents and gives instructions for a general data extraction and tabulation program (TABULATE) written by the authors. It can be used to produce tabulations for all wards of a district on a particular topic.

The fifth section details a more specific data extraction and further analysis 'system' as an example of the kinds of program users may wish to write themselves.


Finally, some preliminary results are presented for the spatial patterning of ethnic groups in the city of Leeds to provide a flavour of what can now be done.

2. THE CENSUS 1981 SAS DATA

Here reference is made to the key documents that need to be read prior to use of the data contained in the Census 1981 SAS.

2.1 Small areas

The geography of the 1981 Census in England and Wales is described in detail in Denham (1980). Very briefly the hierarchy of small areas areas and the larger areas to which they sum is as follows:



Enumeration districts (EDs) are territorial units covered on Census day by one enumerator. The average ED in England and Wales in 1971 contained circa 450 persons. The boundaries of EDs change substantially from Census to Census, though currently OPCS proposes to put together statistics for amalgamations of EDs roughly equivalent in Census 1971 and Census 1981 (Morgan and Denham, 1982; OPCS, 1981c).

Wards are defined each year in urban areas as the territorial basis of local elections with one or more Councillors being elected from each ward. Ward boundaries change marginally from year to year and can change radically between censuses as a result of local government or parliamentary constituency reorganisation. Those used in Census 1981 are those current for the May 1981 local elections. In rural areas the equivalent units are civil parishes, whose boundaries are less subject to change. The average population size of wards and civil parishes in 1971 was circa 3,000 but in urban areas they average over 20,000 persons. The Metropolitan District of Leeds at the time of the 1981 Census contained 33 wards with average population of 21,360.

2.2 The topics covered

For each small area a standard set of tables are produced in three sections

- "1. Population and household tables on a 100 per cent basis, available for areas throughout Great Britain ...
- 2. Population tables on a 10 per cent basis available throughout Great Britain ...
- 3. Tables available for areas in England or these tables and the Welsh language table available for areas in Wales only ... or tables available for areas in Scotland only ..."

(OPCS, 1980, p. 1). Details of the table layout can be derived from:

- 1. OPCS (1980) 1981 Census: Small area statistics.
 Prospectus. OPCS Monitor CEN 80/8. OPCS, London.
- OPCS (1982c) Census 1981 Small Area Statistics. Tape specifications. 100% data. OPCS, Titchfield, Hants.
- 3. OPCS (1982b) Census 1981. County Reports. Part 1. HMSO, London.

The general topics covered are:

Demographic characteristics
Economic characteristics
Housing and amenities
Household composition
Socioeconomic and place of birth characteristics.

A full discussion of the population bases, population and household definitions, the definition and classification of communal establishments, the definitions of topic variables and the Census areas is provided in OPCS (1981b).

3. CENSUS DATA FOR WEST YORKSHIRE WARDS AT THE UNIVERSITY OF LEEDS

3.1 West Yorkshire wards

Figure 1 shows how West Yorkshire Metropolitan County is divided into 5 metropolitan districts and electoral wards. Table 1 lists and names the wards of Bradford, Calderdale, Kirklees, Leeds and Wakefield Metropolitan districts.

3.2 The computer files for West Yorkshire wards

For each Metropolitan district some four data files have been placed on the Geography program and data library permanent disk on the Amdahl V7 computer at the University of Leeds. Table 2 lists the names of the twenty files.

The files with type CDATA contain the records supplied by OPCS. There are some 210 records of 80 characters per ward containing circa 3075 data items or counts. Currently only 100% count data are available in the CDATA files. Sample or 10% count data will be added when they become available.

Type BOUNDARY files contain the raw coordinate data for ward boundaries suitable for input to the GIMMS mapping system (Waugh, 1978) or for users to employ with other mapping systems. The files of type GIMMS contain the same boundary information already processed by the GIMMS package and ready for further use with that system.

The MEMO files contain brief documentation on each district's set of files. An example is shown in Figure 2.

3.3 Accessing the census data

There are two routes to accessing the Census data, depending on username status.

(1) For undergraduates (with usernames GEO2xx or GEO3xx)


You are automatically linked to the GEO6LIB library disk when you logon, with read only access.

Files behave as if they are on your 'D' disk.

eg. To obtain a listing on the screen.

TYPE LEEDS81 MEMO D

This will inform you about the data available.



Ward names in West Yorkshire, Census 1981 Table 1.

	FORD	KIRK	CLEES .	LEEDS			
B1 B2	Baildon Bingley	K1 K2	Thornhill Dewsbury East	Ll L2	Aireborough Armley		
В3	Bingley Rural	1/3	Dewsbury West	13	Barwick & Kippax		
Bli	Bolton	K4	Dalton	L4	Beeston		
B5	Bowling	K5	Almondbury	L5	Bramley		
B6	Bradford Moor	к6	Deighton	L6	Burmantofts		
B7	Clayton	K7	Newsome & Central	L7	Chapel Allerton		
B8	Craven	к8	Birkby & Fartown	18	City & Holbeck		
B9	Eccleshill	К9	Lindley	L9	Cookridge *		
Blo	Great Horton	Klo	Paddock	LIO	Garforth &		
B11	Heaton	VKII			Swillington		
B12	Idle		Batley East		Halton		
B13	Ilkley		Batley West		Harehills		
B14	Keighley North	K14			Headingley		
B15	Keighley South	K15			Horsforth		
B16	Keighley West	K16		1115	Hunslet		
B17	Little Horton	K17		1116	Kirkstall		
B18	Odsal	K18		117	Middleton		
B19	Queensbury	K19	Colne Valley West	r_{19}	Moortown		
B 20	Rombalds	K20		Ll9	Morley North		
B21	Shipley East	K21	Holme Valley North	L20	Morley South		
B22	Shipley West	K22	Holme Valley South	L21	North		
B23	Thornton		Kirkburton	122	Otley & Wharfedal		
B24	Toller	K24	Mirfield	L23	Pudsey North		
B25	Tong			124	Pudsey South		
B26	Undercliffe	WAKE	FIELD	125	Richmond Hill		
B27	University	Wl	Wakefield No. 1	126	Rothwell		
B28	Wibsey	W2	Wakefield No. 2	L27	Roundhay		
B29	Worth Valley	W3	Wakefield No. 3	T58	Seacroft		
B30	Wyke	W4	Wakefield No. 4	L29	University		
		W5	Castleford No. 5		Weetwood		
CALD	ERDALE	w6	Castleford No. 6		Wetherby		
Cl	Brighouse	W7	Castleford No. 7	L32			
C2	Calder Valley	w8	Osset No. 8	L33	Wortley		
C3	Elland	W9	Pontefract No. 9				
C4	Greetland &		Pontefract No. 10				
	Stainland		Featherstone No. 11				
C5	Hipperholm &		Hemsworth Urban No. 12		5		
	Lightcliffe	W13	Horbury No. 13		5 14		
С6	Illingworth	W14	Knottingley No. 14				
C7	Luddenden Foot	W15	Normanton No. 15				
c8	Mixenden	W16	•				
C9	Northowram & Shelf	W17	Stanley No. 17		•		
ClO	Ov end en	W18	Hemsworth Rural No. 18				
C11	Rastrick	W19	Hemsworth Rural No. 19				
C12	Ryburn	W20	Hemsworth Rural No. 20				
C13	St. John's	W21	Wakefield Rural No. 21				
	Skircoat	W22	Wakefield Rural No. 22				
C15	Soverby Bridge		51 (1)				
C16	Todmorden						
C17	Town						
C18	Warley						
					1+40		

Table 2. A list of Census 1981 data and associated files held on the Geography program and census data library disk

filename	file type	filename	filetype
For Bradford M.D.		For Leeds M.D.	
BRADFD81	CDATA	LEEDS81	CDATA
BRADFD81	BOUNDARY	LEEDS81	BOUNDARY
BRADFD81	GIMMS	LEEDS81	GIMMS
BRADFD81	MEMO	LEEDS81	MEMO
For Calderdale M.D.		For Wakefield M.D.	
CALDER81	CDATA	WAKEFD81	CDATA
CALDER81	BOUNDARY	WAKEFD81	BOUNDARY
CALDER81	GIMMS	WAKEFD81	GIMMS
CALDER81	MEMO	WAKEFD81	MEMO
For Kirklees M.D.			
KIRKLS81	CDATA		
KIRKLS81	BOUNDARY		
KIRKLS81	GIMMS		
KIRKLS81	MEMO		

General Notes

- 1. CDATA type files contain the raw census data by ward.
- 2. BOUNDARY type files contain the boundary coordinates and associated information for use with the GIMMS package.
- 3. GIMMS type files contain the BOUNDARY data in processed form. Do not attempt to list.
- 4. MEMO type files provide brief documentation on each district set of files.
- 5. The CDATA files contain the following numbers of 80 character records:

ERADFD81: 6300
CALDER81: 3780
KIRKLS81: 5040
LEEDS81: 7140
WAKEFD81: 4620

Leeds 1981 Census Data: Memo

(1) OPCS file specifications: there are 3 copies of this document in the Geography Map Library (1 copy for reference, 1 copy for loan, 1 copy for reference in 41 University Road).

) Printouts of the raw data are also held in the Geography Map

Library.

3) Maps of the ward boundaries as of Census 1981 (April 5th/6th) are also held in the Geography Map Library.

(4) Details of software that can be used to access the data can be found in:-

Birkin, M.H. and Rees, P.H. (1982) The 1981 small area statistics for West Yorkshire wards: how to access, extract and analyze. Working Paper, School of Geography, forthcoming.

(5) The following files are held on the b disk of the Geography

program and data library (GEO6LIB):

LEEDS81 CDATA Census 1981 100% population and house-holds, small area statistics for wards.

LEEDS81 BOUNDARY Raw boundary coordinate data for Leeds

wards at Census 1981.

LEEDS81 GIMMS Processed boundary coordinate data on

Leeds wards for input to the GIMMS

mapping system.

LEEDS81 MEMO Description of the Census 1981 data on

Leeds held in the Geography program and

data library.

(6) Similar sets of files for Bradford, Kirklees, Calderdale and Wakefield should shortly be available in the Geography program and data library.

Figure 2. A listing of a sample memo file

(2) For postgraduates and staff (with usernames GEO5xxx or GEO6xxx)

You must obtain a work disk using the FILES command and then issue the following instructions

LINK GEO6LIB 191 195 RR
ACCESS 195 B/A

You are then linked to the B disk of GEO6LIB.

Having obtained access to the library files, you may wish to printout your own copies for manual extraction of data. However, it would be better to use a program to extract from the census data files those counts of direct use or the statistics of those wards of interest.

Section 4 describes a program for producing ward by count tabulations and data sets that can be employed by the user to extract information from the CDATA files. This program called TABULATE is also stored on the GEO6LIB permanent disk. Figure 3 provides a picture of the activity involved in abstracting a data set from the library. The user prepares a data file of instructions to the TABULATE program and issues LINK and ACCESS commands to connect to the GEO6LIB B disk. The issuing of a TABULATE command then initiates the reading, printing and writing to disk of the census variables that the user has requested.




Figure 3. An example of a census data extraction system

4. A GENERAL PROGRAM FOR THE EXTRACTION AND TABULATION OF CENSUS DATA

4.1 Introduction

In this section and the next we illustrate the relative ease with which it is possible to extract information from the census data files and how the analysis might then proceed. We begin by looking at the format of the data file itself, and then introduce our simple but general purpose program for extraction of the data. This may be copied and adapted by a user familiar with FORTRAN77 - sub-section 4.3 will hopefully be of assistance to such users. Alternatively the program may be used as a package with the instructions provided in subsection 4.4.

4.2 The format of the data

Figure 4 illustrates the structure of the data files for a typical ward. We see from the figure that the data for the ward are headed by three label records. The first of these locates the ward within the United Kingdom as a whole, the second gives the name of the ward and the third the name of the metropolitan district within which the ward is located. The fourth record is another label relating to the SAS themselves, telling us that the first section - containing the 100% population and household data - will follow immediately. The four label records are followed by 174 records of data, and then another label identifies the end of Section 1 and the beginning of Section 3 of the SAS. After a further 31 records of data come the header labels for the next ward and the data for that ward in the same format (and so on). Thus there are a total of 210 records for each ward.

Each of the records containing data is characterised by the appearance of the number 5 in the first column followed by three spaces. There then follow 15 integer numbers in fields of five characters each, making a total of 79 characters in each record. Note that sequential numbers will only be separated (by a space) if the second is less than 5 digits long (ie. a number less than 10000). The only data records which do not share this format are the final records in the two distinct SAS sections. Record number 178 has only two data items (14 characters) and number 210 has three items (19 characters).

| 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300| | 1300 31020080WAA00SE195414 3202AIREBOROUGH 3203LEEDS 4 81 1 SAS SECTION 1 5 24704237821144912 1132 8 9 454 8 0 0 54 5 14

Figure 4. The SAS data for Aireborough ward, Leeds, Census 1981

Figure 4. Continued

The meaning of each item of data is defined by its position in the file. Thus the first data field of the first record is assigned the identifying number 1 (or V1 for 'variable number 1'). We see from Figure 4 that for Aireborough ward the value of V1 = 24704. Identifying numbers are likewise assigned to the following data items in sequence. Thus the contents of the second field on the first data record are identified as number 2, the first field of the second data record would contain variable number 16 (since there are a total of 15 variables on the first record) and the sixth field of the twelfth data record would be defined as V171 (and so forth). Notice, however, that the simple structure of the data is somewhat disturbed by the presence of the second SAS label on record 179 (see Figure 4).

The next thing we need to know are the meanings of the identifying numbers themselves. These can be found in the 'Coding Book' (OPCS, 1982c), an illustrative sheet of which is shown in Figure 5. for example, we see that total residents in a ward may be found in location 320: that is, the number may be found in the fifth field of the twenty-second data record. The next number will give the total male population of the ward and so on. As an example the twentysecond to twenty-sixth data records for Aireborough ward in Leeds have been reproduced in full in Figure 6. The fifth field of the twentysecond record contains the number 25020-- this, then, is the total number of residents in Aireborough ward at census day. across the record we see that 12053 of Aireborough's residents were males and 12967 females, 23827 were born in England and so on. reading across records we can replace the general identifiers of Figure 5 with the actual values for Aireborough ward.

Finally in this section it is important to recognise the significance of those fields in the data which contain no information at all. These blank fields relate to quantities that have not been measured in the census but are given a location to maintain the pattern of the identifying numbers. A blank location is therefore to be distinguished from a field containing a zero, which indicates that an attribute has been counted but does not occur in the ward in question.

	4 All re	sident	s	
	Countr birt		Males	Fmles
1	TOTAL		321	322
2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20	England Scotland Wales Rest of Irish Re Old Comm New Comm East A Africa Caribbe India Banglae Far Eas Mediter Remaine Pakistan Other Eur Rest of W	th th frica Rem ean desh st cr der	324 327 330 333 336 339 342 345 348 351 354 357 360 363 366 369 372 375 378	325 328 331 334 337 340 343 346 349 352 355 358 361 364 367 370 373 376 379
	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20	TOTAL PERSOI 320 323 326 329 332 335 341 344 347 350 353 362 365 368 371 374 377	NS	

Figure 5. Country of birth page from the coding book for the Census 1981 SAS (OPCS, 1989)

5	0	0	0	72	5020	12053	L2967	238211	1495	12326	366	175	191	110	54
5	56	92	47	45	145	57	88	56	23	33	152	82	70	16	*
5	11	16	9	7	12	6	6	52	29	23	8	8	0	29	14
5	15	15	9	6	4	2	2	10	5	5	136	48	88	61	36
5	25	71	31	401	9520	9296	2825	64711	0224	3672	65521	1882	7125	1815	5310

Figure 6. Selected records from the SAS for Aireborough ward, Leeds,

Census 1981

4.3 Main features of the program

- 84

We have seen in the last section how it is possible to fill out the tables of the coding book with their actual values for particular wards. It is this task which our general purpose extraction and tabulation program performs.

The text of the FORTRAN77 program TABULATE is given in full in Figure 7. The program reads in the number of wards in the district in question and their names, and then the names and code numbers of the variables to be read off. In the latter part of the program the variables are written back in a tabular form as a wards times variables array.

Following the initial comments and parameter assignments, the key part of the program in terms of accessing the census data occurs on lines 47-74. The first 'IF' block (lines 49-53) checks to see if the data item is in Section 1 or Section 3 of the SAS. From its identifying number the program then computes the record number (NREC) and the field of that record in which the data item is to be found (lines 54-62). Once NREC has been determined an external ASSEMBLER subroutine ('READ') is called. This subroutine sets up the required data file with direct access, extracts record number NREC and transfers it to a temporary file on Channel 4. Data can then be read off as required for the tabulation routine (lines 78-99).

This somewhat circuitous procedure is necessary to obtain direct access to the 'read only' files of the GEO6LIB disk. In normal circumstances direct access requires both read and write facilities.

In addition there is a facility for variables to be printed out in matrix form for further analysis (lines 100-106)

```
IF (NGOT.NE.80.OR.IFAIL.NE.0) THEN
                                                                                TAB00640
               WRITE (2, 321) NGOT, IFAIL
                                                                                TAB00650
               STOP
                                                                                TAB00660
             ELSE
                                                                                TAB00670
               REWIND 4
                                                                                TAB00680
               WRITE (4, 322) BUFF
                                                                                TAB00690
               REWIND 4
                                                                                TAB00700
               READ (4,400) (DUMMY (K), K=1,15)
                                                                                TAB00710
             ENDIF
                                                                                TAB00720
             VAL (N, IWARD) = DUMMY(J)
                                                                                TAB00730
  1001
             CONTINUE
                                                                                TAB00740
   321 FORMAT (' BAD DATA FROM CDATA FILE, NGOT= ',15, 'IFAIL= ',15)
                                                                                TAB00750
   322 FORMAT (A80)
                                                                                TAB00760
   400 FORMAT (4X, 15F5.0)
                                                                                TAB00770
                                                                               =TAB00780
       TABULATION ROUTINE FOLLOWS.
                                                                                TAB00790
C
                                                                                TAB00800
        NV=NVARBS+1
                                                                                TAB00810
        WRITE(2,310) TITLE1, TITLE2
                                                                                TAB00820
   310 FORMAT (1H1, A80/A80)
                                                                                TAB00830
        WRITE (2, 311)
                                                                                TAB00840
   311 FORMAT (1HO, 1
                                                                                TAB00850
        WRITE (2,312) TITLE
                                                                                TAB00860
  312 FORMAT (1H0, A80)
                                                                                TAB00870
        WRITE (2,313) (DOT (K), K=1, NV)
                                                                                TAB00880
  313 FORMAT(1H , '...', 11A11)
WRITE(2,314) (VNAME1(M), M=1, NVARBS)
WRITE(2,314) (VNAME2(N), N=1, NVARBS)
                                                                                TAB00890
                                                                                TAB00900
                                                                                TAB00910
  314 FORMAT(1H ,':',12X,':',10(A10,':'))
                                                                                TAB00920
        WRITE (2, 315) (COLON (K), K=1, NV)
                                                                                TAB00930
  315 FORMAT(1H ,':::',11A11)
                                                                                TAB00940
        DO 2001, J=1,NZONE
                                                                                TAB00950
          WRITE(2,320) ZLABEL(J), (VAL(I,J),I=1,NVARBS)
                                                                                TAB00960
  320
          FORMAT(1H ,': ',A10,':',10(F8.0,':'))
                                                                                TAB00970
 2001
          CONTINUE
                                                                               TAB00980
        WRITE (2,315) (COLON (K), K=1, N)
                                                                               TAB00990
=TAB01000
C
       IF NARRAY=1 HAS BEEN SPECIFIED, NAKED VARIABLES ARE
                                                                               TAB01010
C
         WRITTEN TO CHANNEL 3 FOR FURTHER PROCESSING.
                                                                               TAB01020
          IF (NARRAY.NE.1) GOTO 1050
                                                                               TAB01030
       DO 1800, N=1,NVARBS
                                                                               TAB01040
 1800
         WRITE (3, 201) (VAL (N, IWARD), IWARD=1, NZONE)
                                                                               TAB01050
       FORMAT (7(8F7.0/))
  201
                                                                               TAB01060
C
       RETURN TO READ NEXT SET OF VARIABLES.
                                                                               TAB01070
C
                                                                               TAB01080
       CO TO 1050
                                                                               TAB01090
       END
                                                                               TAB01100
```

4.4 Instructions for using the program

4.4.1 The instructions file

Prepare, using the editor, a file that contains the instructions to the TABULATE program for running the job. The instructions consist of titles, labels and parameters (for example, the number of zones) and the names and code numbers of the variables to be extracted. Figure 8 lays out the instructions for preparing this file, which should be of filetype DATA. A simple example of such a control file is shown in Figure 9.

4.4.2 Accessing the census data

Follow the instructions given in section 3.3 to link your username to the Geography program and census data library and to access the permanent disk from your workdisk.

4.4.3 The TABULATE command for running the program

A TABULATE exec has been set up which enables you to run the program by issuing a command of the general form:

TABULATE filenamel. filename2 disklabel

The TABULATE command is followed by three arguments:

- (1) 'Filenamel' is the name of the file on which you have prepared instructions to the program. It should be of filetype DATA and be present on your workdisk. The filename is used to name the file containing the tabulations which has the filetype RESULTS and the file containing the array of raw numbers with filetype ARRAY.
- (2) 'Filename2' is the name of the GEO6LIB file that contains the census data you wish to access. The filenames are listed in Table 2.
- (3) 'Disklabel' is an alphabetic character which defines the location of the GEO6LIB disk and depends on your status as a user. For undergraduate users the letter D should be specified; for postgraduates and staff the letter B.

Record	Variable	Format	Meaning					
1	titlel	e80	Two title records for the results					
2	title2	a80	file (up to 80 characters per record)					
3	nzone	free	Number of wards in the district of interest					
4 TO N	zlabel	7a10	Ward names for nzone wards. Each ward name occupies 10 characters; 7 ward names per record					
N+1	nvarbs free Number of variables in the first table. nvarbs must be less than or equal to 10.							
N+2	2 narray free Parameter governing the production of a raw numbers array.							
			narray = 0 - no array narray = 1 - array produced					
N+3	title	a80	Label of up to 80 characters for the first table					
N+4	vnamel	alO	Labell for variable selected (<10 characters)					
	vname2	alO	Label2 for variable selected (<10 characters)					
	varb	i4	Code number of variable (as in 'Coding Book'					
	•	:	oPCS, 1982e)					
	repeat ro	or nvarbs v	ariables					
N+(m+3)	vnamel	alO	As above					
	vname2	alO						
	varb	i4						
N+(m+4)			N+1 to N+(m+3) is repeated for a new table					
	<u>Or</u> a	nd of the	ntered on a separate record to signal the analysis.					

Figure 8. Instructions for preparing the data file for input to the TABULATE program

```
TABULATION OF SELECTED VARIABLES FROM 1981 CENSUS SMALL AREA
STATISTICS : DATA FOR 33 WARDS IN LEEDS.
33
                                                 B'TOFTSCHAPEL ALL
 AIREBORO
             ARMLEY
                     BARWICK
                               BEESTON
                                         BRAMLEY
     CITY COOKRIDGE GARFORTH
                               HALTON HAREHILLSHEADINGLEY HORSFORTH
  HUNSLET KIRKSTALL MIDDLETON MOORTOWN MORLEY N. MORLEY S.
                                                               NORTH
    OTLEY PUDSEY N. PUDSEY S.RICHMOND H ROTHWELL ROUNDHAY SEACROFT
UNIVERSITY WEETWOOD WETHERBY WHINMOOR WORTLEY
 6
1
TABLE 1: TOTAL RESIDENTS BY COUNTRY OF BIRTH - BRITAIN AND IRELAND.
  TOTAL
           PERSONS 320
           ENGLAND 323
          SCOTLAND
                    326
                    329
            WALES
  REST
             UK
                    332
  IRISH
          REPUBLIC 335
```

Figure 9. A sample instruction file for the TABULATE program

The command used with the sample instructions file shown in Figure 9 was

TABULATE ETHTAB LEEDS81 B

The RESULTS and ARRAY files may be printed in the usual way.

4.4.4 The TABPRINT command for running the program

An alternative TABPRINT exec with exactly the same arguments may be used if you wish to print out the tabulations directly (when there is insufficient room on your workdisk). First of all, spool the printer to an appropriate location, then issue the command:

TABPRINT filenamel filename2 disklabel

In this case no RESULTS file is produced.

4.4.5 Output from the program

The TABULATE program produces tables of census statistics in ward times variable format with user input names for the wards and labels for the variables. Figure 10 shows the table produced by running the program with the ETHTAB instructions of Figure 9. The raw numbers from this table are also written to a separate ARRAY file for further analysis, and Figure 11 shows a listing of the contents of the raw numbers file associated with the ETHTAB example.

TABULATION OF SELECTED VARIABLES FROM 1981 CENSUS SMALL AREA STATISTICS : DATA FOR 33 WARDS IN LEEDS.

TABLE 1: TOTAL RESIDENTS BY COUNTRY OF BIRTH - BRITAIN AND IRELAND.

	• • • • • • • • • • • • • • • • • • • •			10		-		÷						
:		:	TOTAL	00		•				:	REST		IRISH	:
:		:	PERSONS	:	ENGLAND		SCOTLAND	:	WALES	:	UK	:	REPUBLIC	
:		:::		:::		::		::		:::		:::		::
:	AIREBORO	:	25020.	:	23821.	٠.	366.	:	110.	:	92.	:	145.	
:	ARMLEY	:	22887.	:	21137.	:	466.	:	91.	:	116.	:	276.	:
=	BARWICK	:	21821.	:	21013.	:	306.	:	85.	:	52.	:	76.	
:	BEESTON	:	17646.	:	16259.	:	264.	:	56.	=	93.	80	300.	
:	BRAMLEY	:	21664.	:	20746.	:	268.	:	71.	:	90.	:	127.	
:	B'TOFTS	:	21916.	:	20436.	:	286.	:	57.	:	138.	:	398.	:
:	CHAPEL ALL	:	23663.	:	17822.	:	370.	:	104.	:	186.	:	473.	:
:	CITY	:	21517.	:	19207.	:	533.	:	88.	:	146.	:	437.	
:	COOKRIDGE	:	21026.	:	19503.	:	430.	:	145.	:	120.	:	156.	:
:	CARFORTH	:	24006.	:	23110.	:	295.	:	107.	:	70.	:	92.	:
:	HALITON	:	19570.	:	18755.	:	218.	;	81.	:	64.	:	143.	:
:	HAREHILLS	:	22775.	:	17980.	:	439.	:	101.		163.	:	745.	:
:	HEADINGLEY	:	16114.	:	12713.	:	353.	:	125.	:	137.	:	389.	:
:	HORSFORTH	:	21577.	:	20477.	:	347.	:	96.	:	71.	:	139.	:
:	HUNSLET	:	16009.	:	14711.	:	315.	:	49.	:	127.	:	132.	:
:	KIRKSTALL	:	19659.	:	17758.	:	366.	:	75.	:	163.	:	389.	:
:	MIDDLETON	:	19757.	:	19028.	:	250.	:	56.	:	67.	:	116.	:
:	MOORTOWN		20113.	:	18050.	:	324.	:	127.	:	93.	•	208.	:
:	MORLEY N.	:	21424.	:	20754.	:	196.	:	69.	:	51.	:	80.	:
:	MORLEY S.		22678.	:	21986:	:	214.	:	67.	:	49.	:	73.	:
:	NORTH		20277.	:	18623.	:	393.	:	145.	0.70	108.	:	157.	:
:	OTLEY		23166.	:	22099.	•	346.	:	143.	:	65.	:	119.	:
:	PUDSEY N.	:	23023.	:	21945.	•	257.	:	96.		58.	•	86.	:
:	PUDSEY S.		22227.	:	21330.	:	223.	:	84.	:	65.	:	121.	
:	RICHMOND H		22711.	:	21418.	:	290.	:	72.	:	150.	:	334.	:
:	ROTHWELL		21271.	:	20507.	:	216.	:	86.	•	61.	:	84.	:
:	ROUNDHAY	:	20304.	:	18042.	:	446.		172.	:	132.	•	295.	:
:	SEACROFT		20992.	:	19916.	:	289.	:	79.	:	146.	:	223.	:
	UNIVERSITY	:	17706.	:	14392.	:	478.	:	99.	:	207.	:	491.	:
:	WEETWOOD		18250.	:	16669.		406.	:	134.	8:8	95.	:	265.	:
:	WETHERBY		23396.	:	21802.	:	607.	:	298.	:	103.	:	146.	:
:	WHINMOOR		19279.	:	18214.	:	311.	:	71.	:	102.	:	141.	:
:	WORTLEY		23263.	:	21982.	:	327.	:	76.	1886	147.	:	206.	:
:		::		:::		:::		:::		:::		:::		::

Figure 10. A sample tabulation produced by the TABULATE program

25020.	22887.	21821.	17646.	21664.	21916.	23663	21517
ZIUZU.	24000.	19570.	22775.	16114.	21577.	16000	10550
19757.		Z14Z4.	22678.	20277.	23166.	23023	22222
	21271.	20304.	20992.	17706.	18250	23396.	70270
23263.							
23821.	21137.	21013.	16259.	20746.	20436.	17822	10207
19503.	23110.	18755.	17980.	12713.	20477.	14711	17758.
19028.	18050.	20754.	21986.	18623.		21945.	
21418.	20507.	18042.	19916.	14392.	16669.	21802	18214.
21982.							-
366.	466.	306.	264.	268.	286.	370	533.
43V.	295.	218.	439.	353.	347.	315.	
250.	324.	196.	214.	393.		257.	
290.	216.	446.	289.	478.		607.	311.
327.							
110.	91.	85.	56.	71.	57.	104	90
145.	107.	81.	101.	125.	96.	40	75
56.		69.			143.	96	9/
72.	86.	172.	79.	99.	134.	208	71
76.	1						
92.	116.	52.	93.	90-	138.	186	146.
120.	70.	64.	163.	137.	71.	127	163.
67.	93.	51.			65.	58	65.
150.	61.	132.			95.	103.	
147.	7				,,,,	103.	102.
145.	276.	76.	300.	127.	398.	473.	437.
156.	92.		745.		139.		
116.	208.	80.	73.	-	119.		
334.	84.	295.	223.	491.	265.	146.	141.
206.			-			TTU.	TATE

Figure 11. A listing of the raw data file associated with the sample tabulation

5. TECHNIQUES IN THE EXTRACTION AND FURTHER PROCESSING OF SPECIFIC DATA SETS

In this section we narrow the focus of our attention to one particular table of the SAS. We have chosen to investigate spatial patterns of place-of-birth groups of the resident population, the variables for which were listed in Figure 5.

In general terms there are three operations which we would like to perform on the data. Firstly, we would like to undertake a statistical analysis. Secondly, we should like to map the variables so derived (or possibly in some variant of their crude form). Finally, it is necessary to make various tabulations so that the information is not lost in the cartographic exercise. It will be appreciated that the aim of mapping work in this context is better seen in terms of 'clarification through simplification', rather than an attempt to represent the whole of the data in a map form which could then be used for autonomous statistical 'quarrying'. The overall process is summarised as Figure 12 around which the following sections are built.

5.1 Extraction

Our need for data from the census file in this analysis is specific to the extent that only one output file is required, to be used as an input to IOFD FORTRAN, a program written at Leeds (Rees, Brookes and Sellers, 1981) to compute the indices of dissimilarity between social groups in cities. The extraction program used is reproduced in full as Figure 13.

We use essentially the same direct access read procedure as for the TABULATE program with a temporary file on channel 4. The first data item is number 320 which, as we saw earlier, comes five fields into the twenty-second data record - the twenty-sixth in all - hence the determination of NREC in line 16. The five records of data containing the ethnic variables are all read into the temporary file (lines 18-29). Since the variable codes follow simple and well-defined patterns (see Figure 5), the 'DO' loops can be nested straightforwardly to extract the data from the temporary file (lines 32-33).




Figure 12. A system of extracting and analysing ethnic variables

```
DER00010
@PROCESS SC (READ, GTARG)
DER00020
     A FORTRAN PROGRAMME TO EXTRACT VARIABLES FROM
                                                                                DER00030
                                                                                DER00040
  TABLE 4 OF THE UNPROCESSED CENSUS DATA FILE, AND
                                                                                DER00050
   TO ARRANGE THE DATA FOR INPUT TO IOFD.
                                                                                DER00060
                                                                                DER00070
      REAL*8 ETHNIC (33, 20, 3), COLTOT (20), IX (33)
      CHARACTER *6 GROUP (20)
                                                                                DER00080
                                                                                DER00090
      CHARACTER #10 WARD(33)
      CHARACTER*80 BUFF
                                                                                DER00100
                                                                                DER00110
      CHARACTER*8 ARGS (3)
      ARGS (1)="LEEDS81
                                                                                DER00120
      ARGS (2) = CDATA
                                                                                DER00130
                                                                                DER00145
      ARGS (3) = 'B
      DO 999, I=1,33
NREC=((I-1)*210)+26
                                                                                DER00150
                                                                                DER00160
                                                                                DER00170
        MREC=NREC+4
                                                                                DER00180
        DO 998, NUM-NREC, HREC
           CALL READ (ARGS, NUM, BUFF, 80, 'F', NGOT, IFAIL)
                                                                                DER00190
             IF ((NGOT.NE.80).OR. (IFAIL.NE.0)) THEN
                                                                                DER00200
               WRITE (5, 321) NGOT, IFAIL
FORMAT ('BAD DATA FROM CDATA FILE'/
'NGOT= ', 15/
'IFAIL=', 15)
                                                                                DER00210
                                                                                DER00220
  321
                                                                                DER00230
                                                                                DER00240
     2
                                                                                DER00250
               STOP
                                                                                DER00260
                                                                                DER00270
               WRITE (4, 322) BUFF
                                                                                DER00280
  322
               FORMAT (A80)
             ENDIF
                                                                                DER00290
                                                                                DER00300
           CONTINUE
  998
                                                                                DER00310
         REWIND 4
         READ(4,101) ((ETHNIC(I,E,X),X=1,3),E=1,20)
                                                                                DER00320
                                                                                DER00330
  101
        PORMAT (24X,11F5.0/(4X,15F5.0))
                                                                                DER00340
  999
                                                                                DER00350
                                                                                DER00360
     BEGIN ROUTINE TO PRINT DATA TO CHANNEL 2, 'IOD DATA'
                                                                                DER00370
                                                                                DER00380
      WRITE (2,401)
  401 FORMAT ('LEEDS CENSUS DATA, 1981 - ETHNIC GROUPS'/
                                                                                DER00390
               'DATA FOR 33 WARDS AND 18 GROUPS'/'18'/'33')
                                                                                DER00400
                                                                                DER00410
      READ(1,501) (WARD(N),N=1,33)
READ(1,500) (GROUP(M),M=1,20)
                                                                                DER00420
                                                                                DERO0430
  500 FORMAT (10A7)
                                                                                DER00440
  501 PORMAT (6A11)
      DO 600, I=1,20
                                                                                DER00450
                                                                                 DER00460
  600 COLTOT (I) =0.0
                                                                                 DER00470
      DO 1001, J=1,20
                                                                                 DER00480
         DO 1000, I=1,33
                                                                                DER00490
 1000
           COLTOT(J) = COLTOT(J) + ETHNIC(I,J,1)
                                                                                 DER00500
 1001
           CONTINUE
       DO 1005, K=2,20
IF (K.EQ.8) GO TO 1005
                                                                                DER00510
                                                                                 DER00520
         WRITE (2,410) GROUP (K), COLTOT (K)
                                                                                 DER00530
                                                                                 DER00540
  410
        FORMAT (A10/F7.0)
                                                                                 DER0055(
 1005
        CONTINUE
                                                                                 DER00560
       DO 1010, L=1,33
                                                                                 DER0057(
         WRITE(2,420) WARD(L), ETHNIC(L,1,1)
                                                                                 DER0058(
   420
         FORMAT (A10/F6.0)
                                                                                 DER0059(
 1010
        CONTINUE
                                                                                 DER00600
   WRITE (2, 430)
430 FORMAT ('1')
                                                                                 DER0061(
                                                                                 DER0062
       DO 1050, II=1,33
         WRITE(2,440) (ETHNIC(II,IJ,1),IJ=2,7)
WRITE(2,441) (ETHNIC(II,JJ,1),JJ=9,20)
                                                                                 DER0063:
                                                                                 DER0064
                                                                                 DER0065
         FORMAT (6F7.0)
   440
                                                                                 DER0066
   441
         FORMAT (6F7.0/6F7.0)
                                                                                 DER0067
         CONTINUE
 1050
                                                                                 DER0068
       STOP
                                                                                 DER0069
```

In the second half of the program the data are sorted into exactly the right form for input into the next program in the chain. Some tables of 'primary' information may also be constructed from the raw data file using the TABULATE program described in the last section. The use of this program allows us to exert tighter control over the presentation of this basic data than is possible with IOFD.

5.2 Statistical analysis

Once the data file has been prepared as described above it is run with the IOFD program. This produces the following sets of statistics:

- Tabular output of zone by group populations. (This information can be used to check that the data has been input correctly but as noted above is not suitable for final presentation).
- Tables of indices of dissimilarity (see next section) between pairs of social groups and between each social group and the whole population.
- Tables of indices of dissimilarity between pairs of city zones and between each city zone and the whole region.
- For each group a table across zones of the following characteristics:
 - (a) the proportion of a zone's population belonging to the said group
 - (b) the proportion of the total group population locating in the said zone
 - (c) the location quotient an index of the relative concentration of a group's population in particular zones

These tables are particularly important because the information they contain is set to the file IDSRACE DATA and then used in the GIMMS package to produce cartographic representations of the three features. An example of one of these tables is shown in Table 3.

- For each zone a table across groups of the following:
 - (a) the proportion of the zone's population of particular ethnic origins
 - (b) the proportion of different group populations locating in the zone
 - (c) the location quotients for each group in that zone.

In fact what we have, therefore, are the same variables as in the group tables but in a transposed form, to highlight the characteristics of the spatial units rather than the structure of the 'sectoral' units (racial groups). An example of such a table is given in Table 4.

5.3 Mapping

The GIMMS (Geographic Information Management and Mapping System) package is a relatively new implementation at the University of Leeds and so a few comments regarding its salient features may be useful. The major source of information is the Reference Manual (Waugh, 1978). In addition a 'HELP' file on the Amdahl elucidates some of the (many!) grey areas in the manual.

Two types of information are needed for GIMMS - locational information detailing the spatial boundaries of the region under consideration, and non-locational information giving values for different variables to be mapped for the various zones. Any project is likely to have three iterative and ongoing phases according to Waugh (1978, p. 7).

(1) Locational data banks are set up. In our case this means setting up a BOUNDARY data file for Leeds wards, and from this a processed GIMMS file (see Section 3). Processed GIMMS files exist for all the West Yorkshire metropolitan districts and are available on the Geography Library (GEO6LIB) disk. Thus for users interested in analysing West Yorkshire census data this stage of the operation has been effectively short-circuited. For users interested in setting up further locational data files, instructions for digitising area boundaries using the GTCO digitiser and Apple II microcomputer with links to the Amdahl mainframe are given in Birkin (1982).

Table 4. Country of birth statistics for persons resident in Headingley ward, Leeds, Census 1981

ZONE13 :	HEADINGLEY		
GROUP	PERCENT IN GROUP OF ZONAL POPULATION	PERCENT IN ZONE OF GROUP POPULATION	LOCATION QUOTIENT
1 ENGLISH	78.89	1.98	0.86
2 SCOTS	2.19	3.15	1.36
3 WELSH	0.78	3 .7 7	1.63
4 REST OF UK	0.85	3.88	1.68
5 IRISH	2.41	5.14	2.22
6 OLD COMM	0.22	2.98	1.29
7 EAST AFRIC	1.44	12.76	5.52
8 REST AFRIC	0.63	13.00	5.62
9 CARIBBEAN	1.01	4.17	1.80
10 INDIA	3.34	11.61	5.02
11 BANGLADESH	0.05	1.25	0.54
12 FAR EAST	1.94	21.01	9.08
13 MEDIT	0.10	2.86	1.24
14 REST NEWOW	0.10	8.84	3.82
15 PAKISTAN	2.57	12.44	5.38
16 REST EEC	0.86	4.18	1.81
17 REST EUR	0.59	3.09	1.34
18 REST WORLD	2.04	8.14	3.52
TOTAL	100.00	2.31	1.00

- (2) Set up non-locational data banks. This operation has basically two elements. Firstly, the data to be stored must be obtained. In the above example the file LDSRACE DATA is created from the 'group tables' of the IOFD run. Secondly, this data must be stored in a processed form by an initial GIMMS run to create a file called here LDSRACE DATMAN. Details of this relatively simple operation are given in Chapter 5 of the Reference Manual.
- (3) The two sets of data banks are brought together and maps produced using the application modules described in Chapter 5 of the manual. An example is given in Figure 14 to demonstrate the simplicity of the procedure.

To begin with, an instructions file of type 'GIMS' is set up let us call this EXAMPLE GIMMS. To get access to the GIMMS routines the command SETUP PACKAGES is issued. It is also necessary to link to the GEO6LIB disk to use the locational data files stored there. Having done this the appropriate locational data file, in this case for Leeds, is linked to input channel 10. The non-locational data file will normally be on the user's own disk (filemode = a) - this is linked to channel 11. To process the file the command GIMMS EXAMPLE Clo39 is This produces two files - the file EXAMPLE LISTING traces the steps in the execution procedure; the EXAMPLE PLOT file contains the codes for plotting the map produced by the run. If everything has gone smoothly this may be set to the Calcomp 1039 using the PLOT EXAMPLE command.

For the ethnic groups project a total of 54 maps were produced using three measures for each of eighteen groups. Percentage of zonal population in a group was mapped using a simple grid shade; the proportion of group population in a zone using proportional squares; and location quotients using a combination of grid and bar shading. It is hoped that these maps are illustrative of the power of the GIMMS system, although they are in no sense exhaustive of its potential.

TYPE EXAMPLE GIMMS

```
*PLOTPARM GINO, 'M.H.BIRKIN - SCHOOL OF GEOGRAPHY'
*SYSPARM DIAGS=3
*TEXTPARM ALPHABET=13
*PLOTPROG
  *NEWMAP 10 9.5 FRAME
  *GIMMSFILE 10, FACTOR=0.001, PLOT=0.5,0.5, MAP=550,900
    *DATMAN 33 6 6
   *DATA FILE 11 LEEDSDAT
   *DEFINE IRISH=13
   *SET 5 (IRISH)
   *END
 *LEGEND 0.1 9.5 0.4 2 COMPACT
 *LEVELS 6
 *INTERVALS 0.0 0.25 0.5 1.0 2.0 4.0 8.0 5
 *SYMBOLISM AREA GRID 0.24 0.20 0.16 0.12
                       0.08 0.04
 *TEXT X=0.1,Y=9.0,SIZE=0.3,
   TEXT='@ZPOP. BORN IN REP. OF IRELAND'
 *MAP 5
 *END
                                    POP. BORN IN REP. OF IRELAND
```

END OF FILE R; T=0.01/0.07 17:11:52

*STOP

SETUP PACKAGES R; T=0.03/0.06 16:58:48

LINK GEO6LIB 191 195 RR R: T=0.01/0.01 16:58:58

ACCESS 195 B/A '195' REPLACES ' B (195) ' B (195) R/O R; T=0.01/0.01 16:59:04

FI 10 DISK LEEDS81 GIMMS B4 R; T=0.01/0.01 16:59:18

FI 11 DISK LDSRACE DATA A4 R; T=0.01/0.01 16:59:28

GIMMS EXAMPLE Cl039 VM/CMS Version 1.1 GIMMS 3.0 YOUR PLOTFILE IS - EXAMPLE PLOT EXECUTION BEGINS...





R; T=1.19/1.41 16:59:44

PLOT EXAMPLE PRT FILE 6407 TO GUP COPY 001 NOHOLD - YOUR PRINTER IS NOW SPOOLED TO SYSTEM ! R; T=0.04/0.12 16:59:53

> Figure 14. A specimen run using the GIMMS package

6. THE ETHNIC GEOGRAPHY OF LEEDS, 1981

6.1 Introduction

In this section of the paper, the ethnic geography of Leeds at Census 1981 is described as revealed by the Table 4 SAS statistics for persons by country of birth. There are a number of drawbacks (see section 6.6) affecting the interpretation and use of these statistics as indicators of ethnicity, and there are additional data in Table 37 of the SAS (OPCS, 1982c) and Tables 10, 11, 32 of the County statistics (OPCS, 1982b) which should be employed. Here we assume a rough equivalence between birth in a country and membership of that national ethnic group. However, the Table 4 data are sufficiently interesting in their own right to justify a careful description of the patterns of residential location, and thus to provide the reader of this technical manual filled with computing jargon with his geographic reward.

6.2 An ordering and classification of the ethnic groups

The first question we pose is 'how segregated spatially is each ethnic group?'. We can answer this question by comparing the spatial distribution of the group with that of the whole population, using the most frequently used index - the index of dissimilarity. The index of dissimilarity is defined as:

$$D_{ef} = 100 \times 0.5 \times \sum_{i=1}^{N} \begin{vmatrix} \frac{P^{i}}{e} & \frac{P^{i}}{*} \\ P_{e} & P_{f} \end{vmatrix}$$

where P_{ef} is the index of dissimilarity between group e and group f of the population; P_{e}^{i} and P_{f}^{i} are the populations of group e and f respectively in zone i; P_{e}^{*} and P_{f}^{*} are the populations of groups e and f in all zones, of which there are N; and |a| indicates the absolute value function where |a| = a and |-a| = a. The index has a range of 0 (no segregation) to 100 (maximum segregation).

The first column of Table 5 gives the index of dissimilarity between each of 18 ethnic groups and the whole population. The country of birth groups have been arranged in ascending order of the index. Comparison of the ethnic group with the whole population has the drawback than that we are comparing the group with itself, to a certain

Table 5. Key indicators of ethnic residential segregation, Leeds, 1981

Country of No. 43		ssimilarity v	ersus	Percentage	No.
Country of birth group	Whole population (1)	Rest of population (2)	English (3)	of Leeds population (4)	in group (5) -
England	2.1	26.9	C.0	92.18	642,202
FAMILY I			i	, ., .	, , , , , , , , , , , , , , , , , , ,
Scotland	12.3	12.5	14.3	1.61	11 102
Old Commonwealth	15.7	15.8	17.2	0.17	11,193
EED	15.7	15.8	17.4	0.47	1,173 3,305
Wales	16.0	16.1	17.3	0.48	3,314
Rest of the UK	17.7	17.8	19.3	0.51	3,530
FAMILY II					0,,00
NC Mediterranean	21.7	21.7	23.2	0.08	562
Irish Republic	28.2	28.5	30.1	1.09	7,563
Rest of Europe	33.5	33.7	35.4	0.44	3,069
Rest of World	34.4	34.6	36.3	0.58	4,044
NC Far East	38.8	38.9	40.8	0.21	1,493
FAMILY III				-	-, .,,
NC East Africa	43.0	43.1	45.0	0.26	1,816
NC India	43.7	44.0	45.6	0.66	4,627
NC Rest of New Comm.	44.8	44.9	45.4	0.03	177
NC Rest of Africa	45.1	45.1	46.9	0.11	779
FAMILY IV			! -		• • • •
NC Caribbean	61.1	61.4	62.9	0.56	3,889
Pakistan	62.7	63.0	64.6	0.48	
NC Bangladesh	75.6	75.7	77.1	0.09	3,337 641

Source: Output of the IOFD program using Country of birth statistics for Leeds wards, Census 1981.

Column (2): the index of residential segregation Column (3): the index of spatial assimilation

NC = New Commonwealth

EEC = European Economic Community

All statistics refer to the usually resident population. The total usually resident population of Leeds at Census 1981 was 696,714.

extent. It is better to subtract the group itself from the whole population and make a comparison between the group and the rest of the population. This is done in column (2) of Table 5. The index is conventionally labelled 'the index of residential segregation'. All indices are raised, though only that for the English changes significantly because the English born make up 92 per cent of the Leeds population. An even more critical comparison is between each group and the host population (English born) which is given in the third column of the table. We will call this last statistic 'the index of spatial assimilation'. The ordering of the non-English born groups remains constant over these three indices of dissimilarity.

The ethnic groups fall fairly simply into four classes which we have termed 'families of ethnic groups'.

The groups in Family I have indices of spatial assimilation between 14 and 20. Members of the groups are either born in other parts of the United Kingdom or in the three dominions (Australia, Canada and New Zealand) or in the other countries of the European Community bar Eire. The groups in Family II exhibit spatial assimilation indices between 23 and 41, and comprise a mixed group of European (apart from the EEC bar Eire) and American countries together with the colony of Hong Kong. Family III groups show indices of spatial assimilation between 45 and 47 and comprise four groups of New Commonwealth countries. The fourth family have indices above 60 and include Caribbean born, Pakistan born and Bangladesh born Leeds residents.

6.3 The spatial patterns

6.3.1 Proportional distribution

The index of dissimilarity gives a one dimensional picture only of the residential pattern of an ethnic group. Very different spatial patterns can be associated with the same index number. One way to map spatial distributions of populations is to represent the number in a zone by a symbol proportional to the number. Because of the wide disparity in group size (columns 4 and 5 of Table 5), it is difficult to produce comparable maps using the same symbol scale. So the proportional numbers (P_e^i/P_e^*) were mapped instead, in effect reducing all maps to the same symbol scale.

Figures 16 through 18 show the resulting maps*, arranged in ascending ofter of the index of spatial assimilation and grouped roughly in families. The Family I maps (Figure 16) show fairly even distributions, although they are clearly more spatially uneven than the English The Family II maps (Figure 17) show a degree more uneveness. All show a tendency to concentration in the more central wards. Irish born show a wide spread over the central wards, the Rest of Europe born (mainly Poles) show a concentration in the Chapeltown (the ward for traditional settlement by poorer immigrant groups) and Moortown (the ward in which these groups spread as their circumstances improved) The concentration of Far East born in Headingley and University wards undoubtedly reflects their association as faculty or students with the educational institutions located there (Leeds University, Leeds Polytechnic, various Colleges of Higher Education, a major teaching hospital).

The Family III groups (Figures 17 and 18) show increased spatial concentration in particular wards, although members of the groups are found in virtually all wards. Note that the spatial patterns of those born in East Africa and in India are very close, reflecting the close ethnic ties between the groups since most of the former are displaced Indian-origin families from Kenya and Uganda. The final three maps for Family IV groups show the most intense concentration. The Bangladesh-born are the only group to show total absence from a substantial number of wards (14 out of 33). In fact, the only other group to show such absence is the tiny 'Born in the Rest of the New Commonwealth' group (5 out of 33).


6.3.2 Percentage distributions


A more frequently used method is the shading map of the percentages that a group makes up of the zonal populations. Figures 10 through 21 show the choropleth maps of zonal percentages for the 18 country-of-birth groups. The zonal percentages are defined as:

$$\pi_{e}^{i} = 100 \left(\frac{P_{e}^{i}}{P_{\pi}^{i}} \right)$$

Three different scales with six shading categories are used: one for the English, one for the larger or more concentrated groups (born in Scotland,

^{*}Figure 15 provides the ward names and boundaries associated with these maps together with an indication of the continuously built up area.





The proport and distribution of other groups in Loods, 1981; Figure 10.




Figure 17. The proportional distribution of ethnic groups in Leeds, 1981;




Figure 18. The proportional distribution of others groups in Loods, 1981:




Figure 19. Tenal percentage distribution of othnic groups is looks, 1981;




Figure 20. Zonal percentage distribution of ethnic groups in Leeds, 1981:




Figure 21. Zonal percentage distributions of ethnic groups in Leeds, 1981:

Eire, Rest of Europe, Rest of the World, Far East, India, Caribbean and Pakistan) and one for the smaller or less concentrated groups.

6.3.3 Location quotient maps

Another aspect of the concentration in a ward or absence from a ward of group is the degree to which it is more concentrated in that ward than is the population as a whole. This can be measured by the location quotient

$$IQ_e^i = \frac{P_e^i/P_e^*}{P_e^i/P_e^*}$$

where P_*^i is the total population of zone i and P_*^* is the total population of all zones. A location quotient for a ward above l indicates that a group is more concentrated there than is the whole population.

Figures 22 through 24 depict these degrees of concentration. Compare, for example, the Welsh born map with that for persons born in Bangladesh. The Welsh born are most highly concentrated in Wetherby, in the other north suburban wards but also in Headingley and University (the most affluent parts of the city). However, in comparison with the Bangladesh born, the Welsh group are not highly concentrated. The Bangladesh born show location quotients of 10.8, 8.5, 4.6 and 2.97 for Harehills, City and Holbeck, Chapel Allerton and Beeston wards (perhaps the least affluent parts of the city) respectively whereas the Welsh born location quotient for Wetherby is only 2.68.

Clearly, a great deal more could be said about the spatial distribution of each individual group. The maps; together with a good local knowledge, are good means of generating hypotheses about the residential location of the groups concerned: the testing of those hypotheses requires careful work with ther census-derived indicators and additional variables.




Figure 22. Degrees of concentration (location quotients) for ethnic groups in Locals, 1981: 1) Furland born; Cotland to Rest of DK




Figure 23. Degrees of concentration (location quotients) of ethnic groups

in Leeds, 1981: (2) Mediterranean to Far East born




Figure 24. Degrees of concentration (location quotients) of ethnic groups
in Leeds, 1981: (3) India to Rest of Africa born (Family III);
harilland to Bandlalos form (Family IV)

6.4 Interrelationships between the ethnic group distributions

We have already commented in passing on the sameness or contrast between the maps of particular pairs of ethnic groups. Those maps can be compared in one respect by computing indices of dissimilarity between ethnic groups. If we look at the interrelationships between the 17 country of birth groups (besides the English born) we can compute $(17 \times (17-1))/2$ or 136 pairwise indices of dissimilarity.

These are arranged in blocks in Table 6 in which groups of each family are compared with each family of groups. The 'British' groups of Family I are very similar in their distribution (Table 6.1) particularly the first four members of the group, whose indices of dissimilarity are lower with fellow group members than with the English born. Those born in the Rest of the UK (mainly Northern Ireland) are a little more segregated from other Family I groups. The two lowest indices for this group are with the Scots born (14) and the Irish born (15):

Other intra-family dissimilarity indices are higher than those of Family I except for the index value (19) comparing those Born in the New Commonwealth, Mediterranean and those born in the Rest of Europe, and that (value = 12) comparing those born in the New Commonwealth, East Africa and those born in India. Comparisons between families of ethnic groups (Table 6.11) tend on average to follow the ordering of the groups by degree of spatial assimilation, though there are many exceptions in detail that show that groups born in different geographical areas but linked by cultural tradition or family ties have lower indices than the general ordering would suggest.

6.5 A composite picture of the ethnic geography of Leeds, 1981

So far we have examined the ethnic geography of Leeds from a group point of view. The IOFD program provides, as we have shown, statistics, such as a matrix of inter-ward dissimilarity indices, which are the mirror image of the group statistics we have already discussed. What are the types of ethnic areas that characterise the city of Leeds? What are the corresponding statements that can be made from a zonal (ward) point of view?

Table 6. Indices of dissimilarity between pairs of ethnic groups

6.1 Family I vs Family I

BORN IN:	Scotland	Old Common- wealth	Rest of EEC	Wales	Rest of UK
Scotland	0	12	12	13	14
Old Commonwealth		0	10	8	23
Rest of EEC			. 0	12	17
Wales				0	23
Rest of UK					0

6.2 Family I vs Family II

BORN IN:	NC Medi- terranean	Irish Republic	Rest of Europe	Rest of World	NC for East
Scotland	21	23	29	28	35
Old Commonwealth	19	31	27	27	39
Rest of EEC	16	25	22	22	34
Wales	21	31	29	27	39
Rest of UK	22	15	28	26	30

6.3 Family I vs Family III

BORN IN:	NC East	NC	Rest of	NC Rest
	Africa	India	NC	of Africa
Scotland	37	39	40	41
Old Commonwealth	39	42	40	46
Rest of EEC	34	37	37	40
Wales	41	43	41	46
Rest of UK	.32	36	39	35

6.4 Family I vs Family IV

BORN IN	NC Caribbean	Pakistan	NC Bangladesh
Scotland Old Commonwealth Rest of EEC Wales Rest of UK	59	57	72
	61	63	76
	56	59	73
	61	64	77
	53	52	68

6.5 Family II vs Family II

BORN IN:	NC Medi-	Irish	Rest of	Rest of	NC Far
	terranean	Republic	Europe	World	East
NC Mediterranean Irish Republic Rest of Europe Rest of World NC Far East	0	30 0	19 31 0	25 29 23 0	35 28 34 25

6.6 Family II vs Family III

BORN IN:	NC East Africa	NC India	Rest of NC	NC Rest of Africa
NC Mediterranean	33	37	38	37
Irish Republic	24	28	43	32
Rest of Europe	28	25	41	37
Rest of World	29	31	30	24
NC Far East	28	31	36	20

6.7 Family II vs Family IV

BORN IN:	NC Caribbean	Pakistan	NC Bangladesh
NC Mediterranean	52	55	67
Irish Republic	46	40	59
Rest of Europe	43	52	61
Rest of World	49	50	69
NC Far East	47	44	68

6.8 Family III vs Family III

BORN IN	NC East Africa	NC India	Rest of NC	NC Rest of Africa
NC East Africa	0	12	42	27
NC India		0	48	32
Rest of NC			0	39
Rest of Africa				0

6.9 Family III vs Family IV

BORN IN:	NC Caribbean	Pakistan	NC Bangladesh
NC East Africa	38	25	51
NC India	35	29	48
Rest of NC	59	61	80
NC Rest of Africa	42	141	64

6.10 Family IV vs Family IV

	NC Caribbean	Pakistan	NC Bangladesh
NC Caribbean	0	41	53
Pakistan NC Bangladesh		0	34 0

6.11 Average inter-group indices of dissimilarity

Families of ethnic groups	Family I	Family II	Family III	Family IV
English born I 'British' II 'European'	17 14	33 27 28	37 39 32	68 63 53
III 'Indian & African' IV 'West Indian & Pakistan			33	48 43

Computed from indices in Table 6.1-6.10

We could carry out a sophisticated cluster and region building analysis to answer these questions, but, for present purposes a simpler analysis suffices. First, we list in Table 7 some of the key ward characteristics. The first two columns provide a measure of the ethnic diversity of a ward by recording the number of groups with location quotients above 1 and above 2. The University ward records a location quotient above 1 for 17 out of 18 groups - only the English born are underpresented there, and 11 of those 17 groups have location quotients above 2. Neighbouring Headingley ward shows similar characteristics, again associated with having a University-Polytechnic-Colleges of Higher Education linked resident population. Almost as ethnically diverse are the wards of Harehills and Chapel At the other end of the spectrum are the intensely Allerton. 'English' wards of Barwick and Kippax, Bramley, Garforth and Swillington, Morley North, Morley South, Pudsey South and Rothwell in which the only over-represented group is the English born.

If we look at the groups (columns (3) to (5) in Table 7) with the highest percentages in a ward (apart from the English born who are in a 75% or over majority in every ward), then the list is dominated by the numerically largest groups of Scots and Irish born, displaced from first place only in Chapel Allerton by the Caribbean born, in Harehills by the Pakistani born and in Headingley by the Indian born. The picture of groups with the highest concentration in particular wards (columns (6) to (8) in Table 7) is much more diverse and the first column includes 17 of the 18 groups.

A convenient summary of the ethnic character of the different parts of the city is provided if we compute the tripartite percentage distribution of each ward's population into:

- (A) English born
- (B) Born in Family I or II (Scotland, Old Commonwealth, Rest of EEC, Wales, Rest of UK, NC Mediterranean, Eire, Rest of Europe, Rest of World, NC Far East)
- (C) Born in Family III or IV (NC East Africa, NC India, Rest of NC, NC Rest of Africa, NC Caribbean, Pakistan, NC Bangladesh)

NC East Africa IC East Africa Rest of Europe Rest of Europe Rest of World highest LQ Rest of EEC Rest of UK Caribbean Caribbean Old Comm. Caribbean Third Old Comm. (8) English English English English English Snglish Indian Irish Irish Welsh Irish Groups with highest LQ >1 NC Rest Africa Rest of Europe Mediterranean Mediterranean Rest of World Mediterranean highest LQ Rest of EEC Rest of NC Rest of NC Rest of NC Rest of UK Rest of UK Old Comm. Old Comm. Old Comm. Pakistani Pakisteni Pakistani (2) English English English Welsh Irish Welsh Scots Scots NC Rest Africa NC Rest Africa Rest of Europe Mediterranean Mediterranean Rest of World Mediterranean Rest of NC English NC Far East Rest of MC Rest of UK Bangladeshi Bangladeshi Rest of NC Rest of UK Rest of NC Highest Bengladesh Pakistani Old Comm. Caribbean Pakisteni <u>සුම</u> Old Comm. English English English English English English Indian Welsh Irish Irish Rest of Europe Rest of Europe Rest of World Rest of World Rest of World Rest of EEC Rest of UK Rest of EEC Rest of UK Rest of UK Rest of EEC Rest of UK Rest of UK Rest of UK Caribbean Pakistani Third akistani (2) Rest of Indian Indian Indian Irish Irish Irish Welsh delsh Irish Welsh rish Welsh Irish Welsh rish Group with highest Europe Rest of World Rest of World Rest of World Second Pakistani $\widehat{\Xi}$ dest of ndian ndian Irish Irish rish Irish Welsh Irish rish Frish 3cots rish Welsh Welsh rish [rish Fish /elsh rish Scots Crish Welsh Scots Scot B rish rish Scots Caribbean Pakistani First (3) Indian Irish Irish Irish Scots Scots Scots Scots Irish Scots Scots Scots Scots Scots Scots rish Scots Scots Scots Scots Scots Scots Scots Scot 8 Scots Scots Scots Scots Scots Scots Scots with LQs above No. of groups ×2 (2) Ξ Garforth & Swillington Otley & Wharfedale Barwick & Kippex Chapel Allerton City & Holbeck Richmond Hill Pudsey North Pudsey South Morley North Morley South Burmantofts Ward Aireborough Headingley University Cookridge Harchills Horsforth Kirkstall Middleton Roundhay Moortown Rothwell Wetherby Seacroft Weetwood Whinmoor Bramley Hunslet Wortley Beeston Halton Armley North 13. -63 50.00 Ξ. N. <u>8</u> တ္ဆု 29 o (−α) 0, 50 100 98

Table 7. Selected ethnic characteristics of Leeds wards, Census 1981

and plot the position of the wards on triangular coordinate paper (Figure 25). Three clusters can be distinguished on the scatter plot:

- Cluster I 'native areas' with more than 94% of their population born in England, less than 5% born in ethnic families I or II and less than 2% born in ethnic families III or IV;
- Cluster II 'moderately ethnic areas' with between 89
 and 94% of their population born in England,
 Between 5 and 10% of their population born
 ethnic families I or II and up to 5% born
 in ethnic families III or IV;
- Cluster III 'intensely ehtnic areas' with between 75% and 82% of their population born in England, between 9 and 13% born in ethnic families I and II and between 6 and 15% born in ethnic families III and IV.

The inset map shows the spatial location of these ward clusters. Cluster I wards cover the outer suburban and small town ring around the periphery of Leeds metropolitan district. Cluster II wards occupy parts of the inner city, suburbs of the city immediately beyond the inner city plus the outer suburbs on the north-northeastern fringe. Cluster III wards occupy the inner city north of the Central Business district.

6.6 Caveats and further analysis

Our analysis of the ethnic geography of Leeds in 1981 has shown how rich a source of information the decennial census is. The findings have served to underline the value of easy access to that data on disk, the value of writing specific software designed for the subject of interest, and the value of using a well written general package.

However, there is scope for improvement of the analysis in several ways.




Figure 25. A scutterplot of Leeds wards in 'ethnic space'

6.6.1 The definition of ethnicity

The country of birth place statistics undoubtedly underestimate the size of the non-English ethnic groups. Further SAS tables, County report tables and tables from the General Household Survey and Labour Force Survey can be used to improve estimates of the distribution of the population among ethnic groups. What is needed is a model for estimating from available statistics the numbers in more sharply defined ethnic groups. Such a model would have to deal with the conceptual difficulties of mixed birth for individuals and of mixed ethnicity for households.

6.6.2 The definition of spatial segregation

The index of dissimilarity which we have used to summarise the spatial pattern or interrelationships of groups needs improvement. Further indices need to be used. Lieberson (1981) has suggested re-use of an index that reflects the asymmetry of segregation between a larger and a smaller group. It would also be interesting to use our knowledge of the spatial arrangement of wards to compute a set of spatial indices such as the mean inter-group and intra-group distances: this would provide a very interpretable measure of spatial separation.

6.6.3 The scale problem

And, of course, there is the scale problem. If we measured the patterns at enumeration district scale or in terms of clusters of enumeration districts the patterns identified at ward scale are likely to be much sharper. Levels of segregation are likely to be higher (cf. Poole and Boal, 1973; Rees, 1979, p. 299-302). Inter-ethnic distance measures are also likely to be affected, but perhaps to a lesser extent.

6.6.4 Evolution over time

In order to understand the processes that have produced the spatial patterns of 1981 an analysis of their evolution over time is necessary. Severe problems of change in ward and enumeration district definition need to be overcome. It will also be necessary to build up a picture of the demographic changes experienced by the various ethnic groups. The work of King (1974, 1976) will be an invaluable guide here. It should

also be possible to follow and improve on King's work in estimating the socioeconomic characteristics of ethnic groups, and the housing they live in.

6.6.5 Models of ethnic residential location

Given understanding of the spatial and social patterns and their evolution over time, it should then be possible to test rigorously a set of hypotheses concerning the determinants of ethnic residential patterns, either through cross-sectional analysis or through simulation modelling. A major improvement on these previous studies would be the use of a much wider variety of ethnic spatial histories and the use of a more general microsimulation methodology.

6.6.6 Further issues

The analysis should conclude with a careful evaluation of the costs and benefits of the various residential patterns to group members and to society as a whole (cf. Boal, 1976), and an assessment of the influence of public policy in housing and in employment on ethnic residential location.

7. CONCLUSIONS

We have in this paper described an information system. An information system consists of data, a storage device, a description, an accessing system and an analysis system. Traditionally, data have been stored in printed books in libraries, described in the contents and introductory pages of those books; the accessing system has been the way in which people may join libraries and borrow books and the analysis system has been reading and making notes on the information contained in the books.

Using data stored in a library is an elementary activity learnt early in life. We have attempted to make the information system for 'the 1981 Census Small Area Statistics for West Yorkshire Wards' almost as easy to use. The data are numbers contained in machine readable records stored on a computer disk (GEC6LIB's B disk); the data are described in a simple coding book (OPCS, 1982c). Just as you have to gain permission to use a library so you have to acquire permission to use the computer (your username), and the accessing system is the easy-to-use TABULATE program. The analysis system we have left, as does the library, in the hands of the reader of the data, but have shown how simple spatial statistics may be computed using the GIMMS package.

We hope that the information system we have designed minimises the routine tasks in getting access to part of the 1981 Census SAS so as much time as possible can be spent in the task of thought and innovation in its analysis. We trust that in the near future the same will apply to the national information system for accessing enumeration district data at regional computing centres.

REFERENCES

OPCS = Office of Population Censuses and Surveys.

- Birkin, M.H. (1982) Digitizing maps on the GTCO digitizer and Apple II system for use with the GIMMS package. Computer Manual, forthcoming, School of Geography, University of Leeds.
- Boal, F.W. (1976) Ethnic residential segregation. In Herbert, D.T. and Johnston, R.J. (eds.) Social areas in cities, Volume 1. Spatial processes and form. Wiley, London.
- Denham, C.J. (1980) The geography of the Census 1971 and 1981. Population Trends, 19, 6-12.
- King, J.R. (1974) Immigrant fertility trends and population growth in Leeds. Environment and Planning, 6, 509-46.
- King, J.R. (1976) Immigrants in Leeds: an investigation into their socioeconomic characteristics, spatial distribution, fertility trends and population growth. PhD dissertation, School of Geography, University of Leeds.
- Lieberson, S. (1981) An asymmetrical approach to segregation. In Read, C., Robinson, V. and Smith, S. (ed.) Ethnic segregation in cities. Croom Helm, London.
- Morgan, C. and Denham, C.J. (1982) Census small area statistics (SAS): measuring change and spatial variation. *Population Trends*, 28, 12-17.
- OPCS (1980) Census 1981. OPCS Monitor. CEN 80/8. OPCS, London.
- OPCS (1981a) Census 1981. Preliminary report. England and Wales. HMSO, London. CEN 81 PR(1).
- OPCS (1981b) Census 1981. Preliminary report for towns, urban and rural population. England and Wales. HMSO, London. CEN 81 PR(2).
- OPCS (1981c) Census 1981. OPCS Monitor. CEN 81/85. OPCS, London.
- OPCS (1982a) Census 1981. OPCS County Monitors. OPCS, London. CEN 81 CM1-46.
- OPCS (1982b) Census 1981. County Reports. Part 1. HMSO, London.
- OPCS (1982c) Census 1981 Small Area Statistics. Tape specifications. 100% data. OPCS, Titchfield, Hants.
- OPCS (1982d) Census 1981. Definitions. Great Britain. CEN 81 DEF. HMSO, London.
- Poole, M.A. and Boal, F.W. (1973) Religious residential segregation in Belfast in mid-1969: a multi-level analysis. In Clark, B.D. and Gleave, M.B. (ed.) Social patterns in cities. IBG Special Publication No. 5.
- Rees, P.H. (1979) Residential patterns in American cities: 1960.
 Research Paper 189, Department of Geography, University of Chicago, Chicago.
- Rees, P.H., Brookes, S.M. and Sellers, P.J. (1981) A manual for using *IOD*, an index of dissimilarity program. Second edition. Computer Manual 5, School of Geography, University of Leeds.
- Waugh, T.C. (1978) GIMMS. Second Edition, Inter-University Research Council Series, Report No. 30. Program Library Unit, University of Edinburgh.