

EKT-816 Lecture 1

Probability Review (1)

Jesse Naidoo

University of Pretoria

PDFs, CDFs, and Quantiles

- Discrete distribution:

- mass functions: $f(x) = P(X = x)$.
- cumulative distribution functions: $F(x) = P(X \leq x)$.
- Examples: Bernoulli(p); binomial(n, p); Poisson(λ).

- continuous distributions:

- density function $f_X(x)$ such that

$$\int_{-\infty}^{\infty} f_X(x) dx = 1.$$

- CDF $F_X(x)$ is increasing and such that $\lim_{x \rightarrow -\infty} F_X(x) = 0$, $\lim_{x \rightarrow \infty} F_X(x) = 1$.
- $F'_X(x) = f_X(x)$, or

$$F_X(x) = \int_{-\infty}^x f_X(t) dt$$

- the τ -th quantile of the distribution F is the value x_τ such that

$$F(x_\tau) = \tau$$

PDFs, CDFs, and Quantiles

- Discrete distribution:
 - mass functions: $f(x) = P(X = x)$.
 - cumulative distribution functions: $F(x) = P(X \leq x)$.
 - Examples: Bernoulli(p); binomial(n, p); Poisson(λ).
- continuous distributions:

- density function $f_X(x)$ such that

$$\int_{-\infty}^{\infty} f_X(x) dx = 1.$$

- CDF $F_X(x)$ is increasing and such that $\lim_{x \rightarrow -\infty} F_X(x) = 0$, $\lim_{x \rightarrow \infty} F_X(x) = 1$.
- $F'_X(x) = f_X(x)$, or

$$F_X(x) = \int_{-\infty}^x f_X(t) dt$$

- the τ -th quantile of the distribution F is the value x_τ such that

$$F(x_\tau) = \tau$$

PDFs, CDFs, and Quantiles

- Discrete distribution:
 - mass functions: $f(x) = P(X = x)$.
 - cumulative distribution functions: $F(x) = P(X \leq x)$.
 - Examples: Bernoulli(p); binomial(n, p); Poisson(λ).
- continuous distributions:

- density function $f_X(x)$ such that

$$\int_{-\infty}^{\infty} f_X(x) dx = 1.$$

- CDF $F_X(x)$ is increasing and such that $\lim_{x \rightarrow -\infty} F_X(x) = 0$, $\lim_{x \rightarrow \infty} F_X(x) = 1$.
- $F'_X(x) = f_X(x)$, or

$$F_X(x) = \int_{-\infty}^x f_X(t) dt$$

- the τ -th quantile of the distribution F is the value x_τ such that

$$F(x_\tau) = \tau$$

PDFs, CDFs, and Quantiles

- Discrete distribution:
 - mass functions: $f(x) = P(X = x)$.
 - cumulative distribution functions: $F(x) = P(X \leq x)$.
 - Examples: Bernoulli(p); binomial(n, p); Poisson(λ).
- continuous distributions:

- density function $f_X(x)$ such that

$$\int_{-\infty}^{\infty} f_X(x) dx = 1.$$

- CDF $F_X(x)$ is increasing and such that $\lim_{x \rightarrow -\infty} F_X(x) = 0$, $\lim_{x \rightarrow \infty} F_X(x) = 1$.
- $F'_X(x) = f_X(x)$, or

$$F_X(x) = \int_{-\infty}^x f_X(t) dt$$

- the τ -th quantile of the distribution F is the value x_τ such that

$$F(x_\tau) = \tau$$

PDFs, CDFs, and Quantiles

- Discrete distribution:
 - mass functions: $f(x) = P(X = x)$.
 - cumulative distribution functions: $F(x) = P(X \leq x)$.
 - Examples: Bernoulli(p); binomial(n, p); Poisson(λ).
- continuous distributions:

- density function $f_X(x)$ such that

$$\int_{-\infty}^{\infty} f_X(x) dx = 1.$$

- CDF $F_X(x)$ is increasing and such that $\lim_{x \rightarrow -\infty} F_X(x) = 0$, $\lim_{x \rightarrow \infty} F_X(x) = 1$.
 - $F'_X(x) = f_X(x)$, or

$$F_X(x) = \int_{-\infty}^x f_X(t) dt$$

- the τ -th quantile of the distribution F is the value x_τ such that

$$F(x_\tau) = \tau$$

PDFs, CDFs, and Quantiles

- Discrete distribution:
 - mass functions: $f(x) = P(X = x)$.
 - cumulative distribution functions: $F(x) = P(X \leq x)$.
 - Examples: Bernoulli(p); binomial(n, p); Poisson(λ).
- continuous distributions:
 - density function $f_X(x)$ such that

$$\int_{-\infty}^{\infty} f_X(x) dx = 1.$$

- CDF $F_X(x)$ is increasing and such that $\lim_{x \rightarrow -\infty} F_X(x) = 0$, $\lim_{x \rightarrow \infty} F_X(x) = 1$.
- $F'_X(x) = f_X(x)$, or

$$F_X(x) = \int_{-\infty}^x f_X(t) dt$$

- the τ -th quantile of the distribution F is the value x_τ such that

$$F(x_\tau) = \tau$$

PDFs, CDFs, and Quantiles

- Discrete distribution:
 - mass functions: $f(x) = P(X = x)$.
 - cumulative distribution functions: $F(x) = P(X \leq x)$.
 - Examples: Bernoulli(p); binomial(n, p); Poisson(λ).
- continuous distributions:
 - density function $f_X(x)$ such that

$$\int_{-\infty}^{\infty} f_X(x) dx = 1.$$

- CDF $F_X(x)$ is increasing and such that $\lim_{x \rightarrow -\infty} F_X(x) = 0$, $\lim_{x \rightarrow \infty} F_X(x) = 1$.
- $F'_X(x) = f_X(x)$, or

$$F_X(x) = \int_{-\infty}^x f_X(t) dt$$

- the τ -th quantile of the distribution F is the value x_τ such that

$$F(x_\tau) = \tau$$

PDFs, CDFs, and Quantiles

- Discrete distribution:
 - mass functions: $f(x) = P(X = x)$.
 - cumulative distribution functions: $F(x) = P(X \leq x)$.
 - Examples: Bernoulli(p); binomial(n, p); Poisson(λ).
- continuous distributions:
 - density function $f_X(x)$ such that

$$\int_{-\infty}^{\infty} f_X(x) dx = 1.$$

- CDF $F_X(x)$ is increasing and such that $\lim_{x \rightarrow -\infty} F_X(x) = 0$, $\lim_{x \rightarrow \infty} F_X(x) = 1$.
- $F'_X(x) = f_X(x)$, or

$$F_X(x) = \int_{-\infty}^x f_X(t) dt$$

- the τ -th quantile of the distribution F is the value x_τ such that

$$F(x_\tau) = \tau$$

PDFs, CDFs, and Quantiles

- Discrete distribution:
 - mass functions: $f(x) = P(X = x)$.
 - cumulative distribution functions: $F(x) = P(X \leq x)$.
 - Examples: Bernoulli(p); binomial(n, p); Poisson(λ).
- continuous distributions:
 - density function $f_X(x)$ such that

$$\int_{-\infty}^{\infty} f_X(x) dx = 1.$$

- CDF $F_X(x)$ is increasing and such that $\lim_{x \rightarrow -\infty} F_X(x) = 0$, $\lim_{x \rightarrow \infty} F_X(x) = 1$.
- $F'_X(x) = f_X(x)$, or

$$F_X(x) = \int_{-\infty}^x f_X(t) dt$$

- the τ -th quantile of the distribution F is the value x_τ such that

$$F(x_\tau) = \tau$$

Moments

- the *mean* of a distribution is

$$E[X] = \int_{-\infty}^{\infty} xf_X(x)dx$$

- the *variance* of the distribution is

$$V[X] = E[(X - \mu)^2]$$

where μ is the mean of the distribution

- note, these moments may not exist!
 - but, if $V[X] < \infty$, the mean will exist (why?)
 - also notice that $V[X] = E[X^2] - E[X]^2$
 - third (centered) moment is called *skewness*
 - fourth (centered) moment is called *kurtosis*

Moments

- the *mean* of a distribution is

$$E[X] = \int_{-\infty}^{\infty} xf_X(x)dx$$

- the *variance* of the distribution is

$$V[X] = E[(X - \mu)^2]$$

where μ is the mean of the distribution

- note, these moments may not exist!
 - but, if $V[X] < \infty$, the mean will exist (why?)
 - also notice that $V[X] = E[X^2] - E[X]^2$
 - third (centered) moment is called *skewness*
 - fourth (centered) moment is called *kurtosis*

Moments

- the *mean* of a distribution is

$$E[X] = \int_{-\infty}^{\infty} xf_X(x)dx$$

- the *variance* of the distribution is

$$V[X] = E[(X - \mu)^2]$$

where μ is the mean of the distribution

- note, these moments may not exist!
 - but, if $V[X] < \infty$, the mean will exist (why?)
 - also notice that $V[X] = E[X^2] - E[X]^2$
 - third (centered) moment is called *skewness*
 - fourth (centered) moment is called *kurtosis*

Moments

- the *mean* of a distribution is

$$E[X] = \int_{-\infty}^{\infty} xf_X(x)dx$$

- the *variance* of the distribution is

$$V[X] = E[(X - \mu)^2]$$

where μ is the mean of the distribution

- note, these moments may not exist!
 - but, if $V[X] < \infty$, the mean will exist (why?)
 - also notice that $V[X] = E[X^2] - E[X]^2$
 - third (centered) moment is called *skewness*
 - fourth (centered) moment is called *kurtosis*

Moments

- the *mean* of a distribution is

$$E[X] = \int_{-\infty}^{\infty} xf_X(x)dx$$

- the *variance* of the distribution is

$$V[X] = E[(X - \mu)^2]$$

where μ is the mean of the distribution

- note, these moments may not exist!
 - but, if $V[X] < \infty$, the mean will exist (why?)
 - also notice that $V[X] = E[X^2] - E[X]^2$
 - third (centered) moment is called *skewness*
 - fourth (centered) moment is called *kurtosis*

Moments

- the *mean* of a distribution is

$$E[X] = \int_{-\infty}^{\infty} xf_X(x)dx$$

- the *variance* of the distribution is

$$V[X] = E[(X - \mu)^2]$$

where μ is the mean of the distribution

- note, these moments may not exist!
 - but, if $V[X] < \infty$, the mean will exist (why?)
 - also notice that $V[X] = E[X^2] - E[X]^2$
 - third (centered) moment is called *skewness*
 - fourth (centered) moment is called *kurtosis*

Moments

- the *mean* of a distribution is

$$E[X] = \int_{-\infty}^{\infty} xf_X(x)dx$$

- the *variance* of the distribution is

$$V[X] = E[(X - \mu)^2]$$

where μ is the mean of the distribution

- note, these moments may not exist!
 - but, if $V[X] < \infty$, the mean will exist (why?)
 - also notice that $V[X] = E[X^2] - E[X]^2$
 - third (centered) moment is called *skewness*
 - fourth (centered) moment is called *kurtosis*

Example: Pareto Distributions

- let $\alpha > 0$ be some constant
- density is

$$f_X(x) = \begin{cases} \alpha x^{-(\alpha+1)} & \text{if } x > 1 \\ 0 & \text{else} \end{cases}$$

- what is the CDF, $F_X(x)$?
- what is the mean, $E[X]$? do we have to impose any conditions to ensure the mean exists?
- what is the variance, $V[X]$?

Example: Pareto Distributions

- let $\alpha > 0$ be some constant
- density is

$$f_X(x) = \begin{cases} \alpha x^{-(\alpha+1)} & \text{if } x > 1 \\ 0 & \text{else} \end{cases}$$

- what is the CDF, $F_X(x)$?
- what is the mean, $E[X]$? do we have to impose any conditions to ensure the mean exists?
- what is the variance, $V[X]$?

Example: Pareto Distributions

- let $\alpha > 0$ be some constant
- density is

$$f_X(x) = \begin{cases} \alpha x^{-(\alpha+1)} & \text{if } x > 1 \\ 0 & \text{else} \end{cases}$$

- what is the CDF, $F_X(x)$?
- what is the mean, $E[X]$? do we have to impose any conditions to ensure the mean exists?
- what is the variance, $V[X]$?

Example: Pareto Distributions

- let $\alpha > 0$ be some constant
- density is

$$f_X(x) = \begin{cases} \alpha x^{-(\alpha+1)} & \text{if } x > 1 \\ 0 & \text{else} \end{cases}$$

- what is the CDF, $F_X(x)$?
- what is the mean, $E[X]$? do we have to impose any conditions to ensure the mean exists?
- what is the variance, $V[X]$?

Example: Pareto Distributions

- let $\alpha > 0$ be some constant
- density is

$$f_X(x) = \begin{cases} \alpha x^{-(\alpha+1)} & \text{if } x > 1 \\ 0 & \text{else} \end{cases}$$

- what is the CDF, $F_X(x)$?
- what is the mean, $E[X]$? do we have to impose any conditions to ensure the mean exists?
- what is the variance, $V[X]$?

Inverse CDF Trick

- suppose we want to generate random numbers from some distribution with CDF F
 - we can compute F and F^{-1}
 - we can generate uniformly distributed random numbers, $U \sim U(0, 1)$
- then, you can generate $X \sim F$ as follows:

$$X = F^{-1}(U)$$

- proof: let's find the CDF of such X s.
 - let x be an arbitrary number; we're going to show that $P(X \leq x) = F(x)$
 - $P(X \leq x) = P(F^{-1}(U) \leq x) = P(U \leq F(x)) = F(x)$

Inverse CDF Trick

- suppose we want to generate random numbers from some distribution with CDF F
 - we can compute F and F^{-1}
 - we can generate uniformly distributed random numbers, $U \sim U(0, 1)$
- then, you can generate $X \sim F$ as follows:

$$X = F^{-1}(U)$$

- proof: let's find the CDF of such X s.
 - let x be an arbitrary number; we're going to show that $P(X \leq x) = F(x)$
 - $P(X \leq x) = P(F^{-1}(U) \leq x) = P(U \leq F(x)) = F(x)$

Inverse CDF Trick

- suppose we want to generate random numbers from some distribution with CDF F
 - we can compute F and F^{-1}
 - we can generate uniformly distributed random numbers, $U \sim U(0, 1)$
- then, you can generate $X \sim F$ as follows:

$$X = F^{-1}(U)$$

- proof: let's find the CDF of such X s.
 - let x be an arbitrary number; we're going to show that $P(X \leq x) = F(x)$
 - $P(X \leq x) = P(F^{-1}(U) \leq x) = P(U \leq F(x)) = F(x)$

Inverse CDF Trick

- suppose we want to generate random numbers from some distribution with CDF F
 - we can compute F and F^{-1}
 - we can generate uniformly distributed random numbers, $U \sim U(0, 1)$
- then, you can generate $X \sim F$ as follows:

$$X = F^{-1}(U)$$

- proof: let's find the CDF of such X s.
 - let x be an arbitrary number; we're going to show that $P(X \leq x) = F(x)$
 - $P(X \leq x) = P(F^{-1}(U) \leq x) = P(U \leq F(x)) = F(x)$

Inverse CDF Trick

- suppose we want to generate random numbers from some distribution with CDF F
 - we can compute F and F^{-1}
 - we can generate uniformly distributed random numbers, $U \sim U(0, 1)$
- then, you can generate $X \sim F$ as follows:

$$X = F^{-1}(U)$$

- proof: let's find the CDF of such X s.
 - let x be an arbitrary number; we're going to show that $P(X \leq x) = F(x)$
 - $P(X \leq x) = P(F^{-1}(U) \leq x) = P(U \leq F(x)) = F(x)$

Inverse CDF Trick

- suppose we want to generate random numbers from some distribution with CDF F
 - we can compute F and F^{-1}
 - we can generate uniformly distributed random numbers, $U \sim U(0, 1)$
- then, you can generate $X \sim F$ as follows:

$$X = F^{-1}(U)$$

- proof: let's find the CDF of such X s.
 - let x be an arbitrary number; we're going to show that $P(X \leq x) = F(x)$
 - $P(X \leq x) = P(F^{-1}(U) \leq x) = P(U \leq F(x)) = F(x)$

Inverse CDF Trick

- suppose we want to generate random numbers from some distribution with CDF F
 - we can compute F and F^{-1}
 - we can generate uniformly distributed random numbers, $U \sim U(0, 1)$
- then, you can generate $X \sim F$ as follows:

$$X = F^{-1}(U)$$

- proof: let's find the CDF of such X s.
 - let x be an arbitrary number; we're going to show that $P(X \leq x) = F(x)$
 - $P(X \leq x) = P(F^{-1}(U) \leq x) = P(U \leq F(x)) = F(x)$

Marginal and Conditional Distributions

- take a joint density $f_{XY}(x, y)$ that integrates to 1 over \mathbb{R}^2
 - the marginal density of X is

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy$$

- analogous for marginal of Y
- the *conditional* density of Y given that $X = x$ is

$$f_{Y|X}(y|X=x) = \frac{f_{XY}(x, y)}{f_X(x)}$$

Marginal and Conditional Distributions

- take a joint density $f_{XY}(x, y)$ that integrates to 1 over \mathbb{R}^2
 - the marginal density of X is

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy$$

- analogous for marginal of Y
- the *conditional* density of Y given that $X = x$ is

$$f_{Y|X}(y|X=x) = \frac{f_{XY}(x, y)}{f_X(x)}$$

Marginal and Conditional Distributions

- take a joint density $f_{XY}(x, y)$ that integrates to 1 over \mathbb{R}^2
 - the marginal density of X is

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy$$

- analogous for marginal of Y
- the *conditional* density of Y given that $X = x$ is

$$f_{Y|X}(y|X=x) = \frac{f_{XY}(x, y)}{f_X(x)}$$

Marginal and Conditional Distributions

- take a joint density $f_{XY}(x, y)$ that integrates to 1 over \mathbb{R}^2
 - the marginal density of X is

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy$$

- analogous for marginal of Y
- the *conditional* density of Y given that $X = x$ is

$$f_{Y|X}(y|X = x) = \frac{f_{XY}(x, y)}{f_X(x)}$$

Basic Rules

- expectations are linear: $E[aX + Y] = aE[X] + E[Y]$
- $V[aX] = a^2 V[X]$
- $V[X + Y] = V[X] + V[Y] + 2 \operatorname{cov}(X, Y)$

Basic Rules

- expectations are linear: $E[aX + Y] = aE[X] + E[Y]$
- $V[aX] = a^2 V[X]$
- $V[X + Y] = V[X] + V[Y] + 2 \operatorname{cov}(X, Y)$

Basic Rules

- expectations are linear: $E[aX + Y] = aE[X] + E[Y]$
- $V[aX] = a^2 V[X]$
- $V[X + Y] = V[X] + V[Y] + 2 \operatorname{cov}(X, Y)$

Example: Zero Correlation, But Not Independent

- consider the following distribution:

$$f_{XY}(x, y) = \begin{cases} 3/4 & \text{if } x \in (-1, 1) \text{ and } y \in (0, 1 - x^2) \\ 0 & \text{otherwise} \end{cases} \quad (1)$$

- show that $\text{cov}(X, Y) = 0$
 - yet, the two are not independent!
 - to see this, compute the conditional expectation $E[Y|X]$

Example: Zero Correlation, But Not Independent

- consider the following distribution:

$$f_{XY}(x, y) = \begin{cases} 3/4 & \text{if } x \in (-1, 1) \text{ and } y \in (0, 1 - x^2) \\ 0 & \text{otherwise} \end{cases} \quad (1)$$

- show that $\text{cov}(X, Y) = 0$
 - yet, the two are not independent!
 - to see this, compute the conditional expectation $E[Y|X]$

Example: Zero Correlation, But Not Independent

- consider the following distribution:

$$f_{XY}(x, y) = \begin{cases} 3/4 & \text{if } x \in (-1, 1) \text{ and } y \in (0, 1 - x^2) \\ 0 & \text{otherwise} \end{cases} \quad (1)$$

- show that $\text{cov}(X, Y) = 0$
 - yet, the two are not independent!
 - to see this, compute the conditional expectation $E[Y|X]$

Example: Zero Correlation, But Not Independent

- consider the following distribution:

$$f_{XY}(x, y) = \begin{cases} 3/4 & \text{if } x \in (-1, 1) \text{ and } y \in (0, 1 - x^2) \\ 0 & \text{otherwise} \end{cases} \quad (1)$$

- show that $\text{cov}(X, Y) = 0$
 - yet, the two are not independent!
 - to see this, compute the conditional expectation $E[Y|X]$

Law of Iterated Expectations and Variance Decomposition

- law of iterated expectations:

$$E[E[Y|X]] = E[Y]$$

- variance decomposition:

$$V[Y] = V[E[Y|X]] + E[V[Y|X]]$$

Law of Iterated Expectations and Variance Decomposition

- law of iterated expectations:

$$E[E[Y|X]] = E[Y]$$

- variance decomposition:

$$V[Y] = V[E[Y|X]] + E[V[Y|X]]$$

Example: Censored Normal Distribution

Classical Statistical Paradigm

Modes of Convergence

Law(s) of Large Numbers

Central Limit Theorems

Desirable Properties of Estimators

References

Table of Contents

Univariate Distributions

Joint Distributions

Classical (Frequentist) Estimation