

EKT-816 Lecture 5

OLS Consistency and Inference

Jesse Naidoo

University of Pretoria

Preliminaries

- Continuous Mapping Theorem: if $X_n \rightarrow_p X_0$ and $g(\cdot)$ is continuous, then $g(X_n) \rightarrow_p g(X_0)$.
- Slutsky's Theorem:
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n + Y_n \rightarrow_d X_0 + Y$.
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n Y_n \rightarrow_d X_0 Y$.
- Delta method: if $X_n \rightarrow_d N(\mu, \Sigma)$, and $g(\cdot)$ is smoothly differentiable, then $g(X_n) \rightarrow_d N(g(\mu), \nabla g(\mu) \Sigma \nabla g(\mu)')$.
 - here, $\nabla g(x)$ is the gradient of g (recall X can be a vector)
- you can find proofs of these statements in, e.g. Appendix A of Cameron and Trivedi (2005)
 - Ch.3 of Wooldridge (2010) or Ch. 6 of Stachurski (2016) also cover basic asymptotic theory

Preliminaries

- Continuous Mapping Theorem: if $X_n \rightarrow_p X_0$ and $g(\cdot)$ is continuous, then $g(X_n) \rightarrow_p g(X_0)$.
- Slutsky's Theorem:
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n + Y_n \rightarrow_d X_0 + Y$.
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n Y_n \rightarrow_d X_0 Y$.
- Delta method: if $X_n \rightarrow_d N(\mu, \Sigma)$, and $g(\cdot)$ is smoothly differentiable, then $g(X_n) \rightarrow_d N(g(\mu), \nabla g(\mu) \Sigma \nabla g(\mu)')$.
 - here, $\nabla g(x)$ is the gradient of g (recall X can be a vector)
- you can find proofs of these statements in, e.g. Appendix A of Cameron and Trivedi (2005)
 - Ch.3 of Wooldridge (2010) or Ch. 6 of Stachurski (2016) also cover basic asymptotic theory

Preliminaries

- Continuous Mapping Theorem: if $X_n \rightarrow_p X_0$ and $g(\cdot)$ is continuous, then $g(X_n) \rightarrow_p g(X_0)$.
- Slutsky's Theorem:
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n + Y_n \rightarrow_d X_0 + Y$.
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n Y_n \rightarrow_d X_0 Y$.
- Delta method: if $X_n \rightarrow_d N(\mu, \Sigma)$, and $g(\cdot)$ is smoothly differentiable, then $g(X_n) \rightarrow_d N(g(\mu), \nabla g(\mu) \Sigma \nabla g(\mu)')$.
 - here, $\nabla g(x)$ is the gradient of g (recall X can be a vector)
- you can find proofs of these statements in, e.g. Appendix A of Cameron and Trivedi (2005)
 - Ch.3 of Wooldridge (2010) or Ch. 6 of Stachurski (2016) also cover basic asymptotic theory

Preliminaries

- Continuous Mapping Theorem: if $X_n \rightarrow_p X_0$ and $g(\cdot)$ is continuous, then $g(X_n) \rightarrow_p g(X_0)$.
- Slutsky's Theorem:
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n + Y_n \rightarrow_d X_0 + Y$.
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n Y_n \rightarrow_d X_0 Y$.
- Delta method: if $X_n \rightarrow_d N(\mu, \Sigma)$, and $g(\cdot)$ is smoothly differentiable, then $g(X_n) \rightarrow_d N(g(\mu), \nabla g(\mu) \Sigma \nabla g(\mu)')$.
 - here, $\nabla g(x)$ is the gradient of g (recall X can be a vector)
- you can find proofs of these statements in, e.g. Appendix A of Cameron and Trivedi (2005)
 - Ch.3 of Wooldridge (2010) or Ch. 6 of Stachurski (2016) also cover basic asymptotic theory

Preliminaries

- Continuous Mapping Theorem: if $X_n \rightarrow_p X_0$ and $g(\cdot)$ is continuous, then $g(X_n) \rightarrow_p g(X_0)$.
- Slutsky's Theorem:
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n + Y_n \rightarrow_d X_0 + Y$.
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n Y_n \rightarrow_d X_0 Y$.
- Delta method: if $X_n \rightarrow_d N(\mu, \Sigma)$, and $g(\cdot)$ is smoothly differentiable, then $g(X_n) \rightarrow_d N(g(\mu), \nabla g(\mu) \Sigma \nabla g(\mu)')$.
 - here, $\nabla g(x)$ is the gradient of g (recall X can be a vector)
- you can find proofs of these statements in, e.g. Appendix A of Cameron and Trivedi (2005)
 - Ch.3 of Wooldridge (2010) or Ch. 6 of Stachurski (2016) also cover basic asymptotic theory

Preliminaries

- Continuous Mapping Theorem: if $X_n \rightarrow_p X_0$ and $g(\cdot)$ is continuous, then $g(X_n) \rightarrow_p g(X_0)$.
- Slutsky's Theorem:
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n + Y_n \rightarrow_d X_0 + Y$.
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n Y_n \rightarrow_d X_0 Y$.
- Delta method: if $X_n \rightarrow_d N(\mu, \Sigma)$, and $g(\cdot)$ is smoothly differentiable, then $g(X_n) \rightarrow_d N(g(\mu), \nabla g(\mu) \Sigma \nabla g(\mu)')$.
 - here, $\nabla g(x)$ is the gradient of g (recall X can be a vector)
- you can find proofs of these statements in, e.g. Appendix A of Cameron and Trivedi (2005)
 - Ch.3 of Wooldridge (2010) or Ch. 6 of Stachurski (2016) also cover basic asymptotic theory

Preliminaries

- Continuous Mapping Theorem: if $X_n \rightarrow_p X_0$ and $g(\cdot)$ is continuous, then $g(X_n) \rightarrow_p g(X_0)$.
- Slutsky's Theorem:
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n + Y_n \rightarrow_d X_0 + Y$.
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n Y_n \rightarrow_d X_0 Y$.
- Delta method: if $X_n \rightarrow_d N(\mu, \Sigma)$, and $g(\cdot)$ is smoothly differentiable, then $g(X_n) \rightarrow_d N(g(\mu), \nabla g(\mu) \Sigma \nabla g(\mu)')$.
 - here, $\nabla g(x)$ is the gradient of g (recall X can be a vector)
- you can find proofs of these statements in, e.g. Appendix A of Cameron and Trivedi (2005)
 - Ch.3 of Wooldridge (2010) or Ch. 6 of Stachurski (2016) also cover basic asymptotic theory

Preliminaries

- Continuous Mapping Theorem: if $X_n \rightarrow_p X_0$ and $g(\cdot)$ is continuous, then $g(X_n) \rightarrow_p g(X_0)$.
- Slutsky's Theorem:
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n + Y_n \rightarrow_d X_0 + Y$.
 - if $X_n \rightarrow_p X_0$ (a constant) and $Y_n \rightarrow_d Y$ (a nondegenerate distribution), then $X_n Y_n \rightarrow_d X_0 Y$.
- Delta method: if $X_n \rightarrow_d N(\mu, \Sigma)$, and $g(\cdot)$ is smoothly differentiable, then $g(X_n) \rightarrow_d N(g(\mu), \nabla g(\mu) \Sigma \nabla g(\mu)')$.
 - here, $\nabla g(x)$ is the gradient of g (recall X can be a vector)
- you can find proofs of these statements in, e.g. Appendix A of Cameron and Trivedi (2005)
 - Ch.3 of Wooldridge (2010) or Ch. 6 of Stachurski (2016) also cover basic asymptotic theory

Asymptotic Distribution of the OLS Estimator

- to build up intuition, think of the single-regressor case:

$$y = x\beta + \varepsilon$$

with $\varepsilon \perp\!\!\!\perp x$ and $E[x] = E[\varepsilon] = 0$.

- we know $\hat{\beta} = \widehat{\text{cov}}(y, x) / \widehat{V}[x] = \sum_i y_i x_i / \sum_i x_i^2$.
- we also know $V[\hat{\beta}] = \sigma_\varepsilon^2 / V[x]$
 - in the usual picture, this corresponds to the fact that estimating β is "harder" with:
 - more noise (i.e. larger variance to ε)
 - less horizontal dispersion in x (i.e. smaller values of $V[x]$)
- now, we are going to extend this result to more complicated settings:
 - multiple regressors
 - unequal variances for ε at different values of x ("heteroskedasticity")
 - correlations between the errors of different observations (serial correlation or clustering)

Asymptotic Distribution of the OLS Estimator

- to build up intuition, think of the single-regressor case:

$$y = x\beta + \varepsilon$$

with $\varepsilon \perp\!\!\!\perp x$ and $E[x] = E[\varepsilon] = 0$.

- we know $\hat{\beta} = \widehat{\text{cov}}(y, x) / \widehat{V}[x] = \sum_i y_i x_i / \sum_i x_i^2$.
- we also know $V[\hat{\beta}] = \sigma_\varepsilon^2 / V[x]$
 - in the usual picture, this corresponds to the fact that estimating β is "harder" with:
 - more noise (i.e. larger variance in ε)
 - less horizontal dispersion in x (i.e. smaller values of $V[x]$)
- now, we are going to extend this result to more complicated settings:
 - multiple regressors
 - unequal variances for ε at different values of x ("heteroskedasticity")
 - correlations between the errors of different observations (serial correlation or clustering)

Asymptotic Distribution of the OLS Estimator

- to build up intuition, think of the single-regressor case:

$$y = x\beta + \varepsilon$$

with $\varepsilon \perp\!\!\!\perp x$ and $E[x] = E[\varepsilon] = 0$.

- we know $\hat{\beta} = \widehat{\text{cov}}(y, x) / \widehat{V}[x] = \sum_i y_i x_i / \sum_i x_i^2$.
- we also know $V[\hat{\beta}] = \sigma_\varepsilon^2 / V[x]$
 - in the usual picture, this corresponds to the fact that estimating β is “harder” with:
 - ▶ more vertical dispersion in y (i.e. larger values of σ_ε^2)
 - ▶ less horizontal dispersion in x (i.e. smaller values of $V[x]$)
- now, we are going to extend this result to more complicated settings:
 - multiple regressors
 - unequal variances for ε at different values of x (“heteroskedasticity”)
 - correlations between the errors of different observations (serial correlation or clustering)

Asymptotic Distribution of the OLS Estimator

- to build up intuition, think of the single-regressor case:

$$y = x\beta + \varepsilon$$

with $\varepsilon \perp\!\!\!\perp x$ and $E[x] = E[\varepsilon] = 0$.

- we know $\hat{\beta} = \widehat{\text{cov}}(y, x) / \widehat{V}[x] = \sum_i y_i x_i / \sum_i x_i^2$.
- we also know $V[\hat{\beta}] = \sigma_\varepsilon^2 / V[x]$
 - in the usual picture, this corresponds to the fact that estimating β is “harder” with:
 - ▶ more vertical dispersion in y (i.e. larger values of σ_ε^2)
 - ▶ less horizontal dispersion in x (i.e. smaller values of $V[x]$)
- now, we are going to extend this result to more complicated settings:
 - multiple regressors
 - unequal variances for ε at different values of x (“heteroskedasticity”)
 - correlations between the errors of different observations (serial correlation or clustering)

Asymptotic Distribution of the OLS Estimator

- to build up intuition, think of the single-regressor case:

$$y = x\beta + \varepsilon$$

with $\varepsilon \perp\!\!\!\perp x$ and $E[x] = E[\varepsilon] = 0$.

- we know $\hat{\beta} = \widehat{\text{cov}}(y, x) / \widehat{V}[x] = \sum_i y_i x_i / \sum_i x_i^2$.
- we also know $V[\hat{\beta}] = \sigma_\varepsilon^2 / V[x]$
 - in the usual picture, this corresponds to the fact that estimating β is “harder” with:
 - ▶ more vertical dispersion in y (i.e. larger values of σ_ε^2)
 - ▶ less horizontal dispersion in x (i.e. smaller values of $V[x]$)
- now, we are going to extend this result to more complicated settings:
 - multiple regressors
 - unequal variances for ε at different values of x (“heteroskedasticity”)
 - correlations between the errors of different observations (serial correlation or clustering)

Asymptotic Distribution of the OLS Estimator

- to build up intuition, think of the single-regressor case:

$$y = x\beta + \varepsilon$$

with $\varepsilon \perp\!\!\!\perp x$ and $E[x] = E[\varepsilon] = 0$.

- we know $\hat{\beta} = \widehat{\text{cov}}(y, x) / \widehat{V}[x] = \sum_i y_i x_i / \sum_i x_i^2$.
- we also know $V[\hat{\beta}] = \sigma_\varepsilon^2 / V[x]$
 - in the usual picture, this corresponds to the fact that estimating β is “harder” with:
 - ▶ more vertical dispersion in y (i.e. larger values of σ_ε^2)
 - ▶ less horizontal dispersion in x (i.e. smaller values of $V[x]$)
- now, we are going to extend this result to more complicated settings:
 - multiple regressors
 - unequal variances for ε at different values of x (“heteroskedasticity”)
 - correlations between the errors of different observations (serial correlation or clustering)

Asymptotic Distribution of the OLS Estimator

- to build up intuition, think of the single-regressor case:

$$y = x\beta + \varepsilon$$

with $\varepsilon \perp\!\!\!\perp x$ and $E[x] = E[\varepsilon] = 0$.

- we know $\hat{\beta} = \widehat{\text{cov}}(y, x) / \widehat{V}[x] = \sum_i y_i x_i / \sum_i x_i^2$.
- we also know $V[\hat{\beta}] = \sigma_\varepsilon^2 / V[x]$
 - in the usual picture, this corresponds to the fact that estimating β is “harder” with:
 - ▶ more vertical dispersion in y (i.e. larger values of σ_ε^2)
 - ▶ less horizontal dispersion in x (i.e. smaller values of $V[x]$)
- now, we are going to extend this result to more complicated settings:
 - multiple regressors
 - unequal variances for ε at different values of x (“heteroskedasticity”)
 - correlations between the errors of different observations (serial correlation or clustering)

Asymptotic Distribution of the OLS Estimator

- to build up intuition, think of the single-regressor case:

$$y = x\beta + \varepsilon$$

with $\varepsilon \perp\!\!\!\perp x$ and $E[x] = E[\varepsilon] = 0$.

- we know $\hat{\beta} = \widehat{\text{cov}}(y, x) / \widehat{V}[x] = \sum_i y_i x_i / \sum_i x_i^2$.
- we also know $V[\hat{\beta}] = \sigma_\varepsilon^2 / V[x]$
 - in the usual picture, this corresponds to the fact that estimating β is “harder” with:
 - ▶ more vertical dispersion in y (i.e. larger values of σ_ε^2)
 - ▶ less horizontal dispersion in x (i.e. smaller values of $V[x]$)
- now, we are going to extend this result to more complicated settings:
 - multiple regressors
 - unequal variances for ε at different values of x (“heteroskedasticity”)
 - correlations between the errors of different observations (serial correlation or clustering)

Asymptotic Distribution of the OLS Estimator

- to build up intuition, think of the single-regressor case:

$$y = x\beta + \varepsilon$$

with $\varepsilon \perp\!\!\!\perp x$ and $E[x] = E[\varepsilon] = 0$.

- we know $\hat{\beta} = \widehat{\text{cov}}(y, x) / \widehat{V}[x] = \sum_i y_i x_i / \sum_i x_i^2$.
- we also know $V[\hat{\beta}] = \sigma_\varepsilon^2 / V[x]$
 - in the usual picture, this corresponds to the fact that estimating β is “harder” with:
 - ▶ more vertical dispersion in y (i.e. larger values of σ_ε^2)
 - ▶ less horizontal dispersion in x (i.e. smaller values of $V[x]$)
- now, we are going to extend this result to more complicated settings:
 - multiple regressors
 - unequal variances for ε at different values of x (“heteroskedasticity”)
 - correlations between the errors of different observations (serial correlation or clustering)

Asymptotic Distribution of the OLS Estimator

- to build up intuition, think of the single-regressor case:

$$y = x\beta + \varepsilon$$

with $\varepsilon \perp\!\!\!\perp x$ and $E[x] = E[\varepsilon] = 0$.

- we know $\hat{\beta} = \widehat{\text{cov}}(y, x) / \widehat{V}[x] = \sum_i y_i x_i / \sum_i x_i^2$.
- we also know $V[\hat{\beta}] = \sigma_\varepsilon^2 / V[x]$
 - in the usual picture, this corresponds to the fact that estimating β is “harder” with:
 - ▶ more vertical dispersion in y (i.e. larger values of σ_ε^2)
 - ▶ less horizontal dispersion in x (i.e. smaller values of $V[x]$)
- now, we are going to extend this result to more complicated settings:
 - multiple regressors
 - unequal variances for ε at different values of x (“heteroskedasticity”)
 - correlations between the errors of different observations (serial correlation or clustering)

Asymptotic Distribution of the OLS Estimator

- we want to apply a central limit theorem to $\hat{\beta}$
- because $\hat{\beta} = (X'X)^{-1}X'Y = \beta + (X'X)^{-1}X'\varepsilon$, we have

$$\sqrt{N}(\hat{\beta} - \beta) = (X'X/N)^{-1}(X'\varepsilon/\sqrt{N}) \quad (1)$$

- we will maintain the assumption that $X'\varepsilon/N \rightarrow_p 0$
 - an easy sufficient condition is that ε is *mean independent* of X , i.e. $E[\varepsilon|X] = 0$
 - we don't want to go as far as assuming ε is independent of X though
 - why not? Full independence implies no heteroskedasticity or clustering
 - if $X'\varepsilon/N \rightarrow_p 0$, we get that OLS is *consistent* for β
- the simple case is one where $E[\varepsilon\varepsilon'|X] = \sigma^2 I$
 - take variances on both sides of (1) to get that

$$\sqrt{N}(\hat{\beta} - \beta) \rightarrow_d N(0, \sigma^2(X'X)^{-1})$$

- notice, this is just a multivariable generalization of $V[\hat{\beta}] = \sigma_\varepsilon^2/V[X]$
- so, to do inference on the elements of $\hat{\beta}$ (or functions of them) in practice, we'd use the *asymptotic covariance matrix* $s^2(X'X)^{-1}/N$

Asymptotic Distribution of the OLS Estimator

- we want to apply a central limit theorem to $\hat{\beta}$
- because $\hat{\beta} = (X'X)^{-1}X'Y = \beta + (X'X)^{-1}X'\varepsilon$, we have

$$\sqrt{N}(\hat{\beta} - \beta) = (X'X/N)^{-1}(X'\varepsilon/\sqrt{N}) \quad (1)$$

- we will maintain the assumption that $X'\varepsilon/N \rightarrow_p 0$
 - an easy sufficient condition is that ε is *mean independent* of X , i.e. $E[\varepsilon|X] = 0$
 - we don't want to go as far as assuming ε is independent of X though
 - why not? Full independence implies no heteroskedasticity or clustering
 - if $X'\varepsilon/N \rightarrow_p 0$, we get that OLS is *consistent* for β
- the simple case is one where $E[\varepsilon\varepsilon'|X] = \sigma^2 I$
 - take variances on both sides of (1) to get that

$$\sqrt{N}(\hat{\beta} - \beta) \rightarrow_d N(0, \sigma^2(X'X)^{-1})$$

- notice, this is just a multivariable generalization of $V[\hat{\beta}] = \sigma_e^2/V[X]$
- so, to do inference on the elements of $\hat{\beta}$ (or functions of them) in practice, we'd use the *asymptotic covariance matrix* $s^2(X'X)^{-1}/N$

Asymptotic Distribution of the OLS Estimator

- we want to apply a central limit theorem to $\hat{\beta}$
- because $\hat{\beta} = (X'X)^{-1}X'Y = \beta + (X'X)^{-1}X'\varepsilon$, we have

$$\sqrt{N}(\hat{\beta} - \beta) = (X'X/N)^{-1}(X'\varepsilon/\sqrt{N}) \quad (1)$$

- we will maintain the assumption that $X'\varepsilon/N \rightarrow_p 0$
 - an easy sufficient condition is that ε is *mean independent* of X , i.e. $E[\varepsilon|X] = 0$
 - we don't want to go as far as assuming ε is independent of X though
 - ▶ why not? Full independence implies no heteroskedasticity or clustering
 - if $X'\varepsilon/N \rightarrow_p 0$, we get that OLS is *consistent* for β
- the simple case is one where $E[\varepsilon\varepsilon'|X] = \sigma^2 I$
 - take variances on both sides of (1) to get that

$$\sqrt{N}(\hat{\beta} - \beta) \rightarrow_d N(0, \sigma^2(X'X)^{-1})$$

- notice, this is just a multivariable generalization of $V[\hat{\beta}] = \sigma_e^2/V[X]$
- so, to do inference on the elements of $\hat{\beta}$ (or functions of them) in practice, we'd use the *asymptotic covariance matrix* $s^2(X'X)^{-1}/N$

Asymptotic Distribution of the OLS Estimator

- we want to apply a central limit theorem to $\hat{\beta}$
- because $\hat{\beta} = (X'X)^{-1}X'Y = \beta + (X'X)^{-1}X'\varepsilon$, we have

$$\sqrt{N}(\hat{\beta} - \beta) = (X'X/N)^{-1}(X'\varepsilon/\sqrt{N}) \quad (1)$$

- we will maintain the assumption that $X'\varepsilon/N \rightarrow_p 0$
 - an easy sufficient condition is that ε is *mean independent* of X , i.e. $E[\varepsilon|X] = 0$
 - we don't want to go as far as assuming ε is independent of X though
 - ▶ why not? Full independence implies no heteroskedasticity or clustering
 - if $X'\varepsilon/N \rightarrow_p 0$, we get that OLS is *consistent* for β
- the simple case is one where $E[\varepsilon\varepsilon'|X] = \sigma^2 I$
 - take variances on both sides of (1) to get that

$$\sqrt{N}(\hat{\beta} - \beta) \rightarrow_d N(0, \sigma^2(X'X)^{-1})$$

- notice, this is just a multivariable generalization of $V[\hat{\beta}] = \sigma_e^2/V[X]$
- so, to do inference on the elements of $\hat{\beta}$ (or functions of them) in practice, we'd use the *asymptotic covariance matrix* $s^2(X'X)^{-1}/N$

Asymptotic Distribution of the OLS Estimator

- we want to apply a central limit theorem to $\hat{\beta}$
- because $\hat{\beta} = (X'X)^{-1}X'Y = \beta + (X'X)^{-1}X'\varepsilon$, we have

$$\sqrt{N}(\hat{\beta} - \beta) = (X'X/N)^{-1}(X'\varepsilon/\sqrt{N}) \quad (1)$$

- we will maintain the assumption that $X'\varepsilon/N \rightarrow_p 0$
 - an easy sufficient condition is that ε is *mean independent* of X , i.e. $E[\varepsilon|X] = 0$
 - we don't want to go as far as assuming ε is independent of X though
 - ▶ why not? Full independence implies no heteroskedasticity or clustering
 - if $X'\varepsilon/N \rightarrow_p 0$, we get that OLS is *consistent* for β
- the simple case is one where $E[\varepsilon\varepsilon'|X] = \sigma^2 I$
 - take variances on both sides of (1) to get that

$$\sqrt{N}(\hat{\beta} - \beta) \rightarrow_d N(0, \sigma^2(X'X)^{-1})$$

- notice, this is just a multivariable generalization of $V[\hat{\beta}] = \sigma_e^2/V[X]$
- so, to do inference on the elements of $\hat{\beta}$ (or functions of them) in practice, we'd use the *asymptotic covariance matrix* $s^2(X'X)^{-1}/N$

Asymptotic Distribution of the OLS Estimator

- we want to apply a central limit theorem to $\hat{\beta}$
- because $\hat{\beta} = (X'X)^{-1}X'Y = \beta + (X'X)^{-1}X'\varepsilon$, we have

$$\sqrt{N}(\hat{\beta} - \beta) = (X'X/N)^{-1}(X'\varepsilon/\sqrt{N}) \quad (1)$$

- we will maintain the assumption that $X'\varepsilon/N \rightarrow_p 0$
 - an easy sufficient condition is that ε is *mean independent* of X , i.e. $E[\varepsilon|X] = 0$
 - we don't want to go as far as assuming ε is independent of X though
 - ▶ why not? Full independence implies no heteroskedasticity or clustering
 - if $X'\varepsilon/N \rightarrow_p 0$, we get that OLS is *consistent* for β
- the simple case is one where $E[\varepsilon\varepsilon'|X] = \sigma^2 I$
 - take variances on both sides of (1) to get that

$$\sqrt{N}(\hat{\beta} - \beta) \rightarrow_d N(0, \sigma^2(X'X)^{-1})$$

- notice, this is just a multivariable generalization of $V[\hat{\beta}] = \sigma_e^2/V[X]$
- so, to do inference on the elements of $\hat{\beta}$ (or functions of them) in practice, we'd use the *asymptotic covariance matrix* $s^2(X'X)^{-1}/N$

Asymptotic Distribution of the OLS Estimator

- we want to apply a central limit theorem to $\hat{\beta}$
- because $\hat{\beta} = (X'X)^{-1}X'Y = \beta + (X'X)^{-1}X'\varepsilon$, we have

$$\sqrt{N}(\hat{\beta} - \beta) = (X'X/N)^{-1}(X'\varepsilon/\sqrt{N}) \quad (1)$$

- we will maintain the assumption that $X'\varepsilon/N \rightarrow_p 0$
 - an easy sufficient condition is that ε is *mean independent* of X , i.e. $E[\varepsilon|X] = 0$
 - we don't want to go as far as assuming ε is independent of X though
 - ▶ why not? Full independence implies no heteroskedasticity or clustering
 - if $X'\varepsilon/N \rightarrow_p 0$, we get that OLS is *consistent* for β
- the simple case is one where $E[\varepsilon\varepsilon'|X] = \sigma^2 I$
 - take variances on both sides of (1) to get that

$$\sqrt{N}(\hat{\beta} - \beta) \rightarrow_d N(0, \sigma^2(X'X)^{-1})$$

- notice, this is just a multivariable generalization of $V[\hat{\beta}] = \sigma_e^2/V[X]$
- so, to do inference on the elements of $\hat{\beta}$ (or functions of them) in practice, we'd use the *asymptotic covariance matrix* $s^2(X'X)^{-1}/N$

Asymptotic Distribution of the OLS Estimator

- we want to apply a central limit theorem to $\hat{\beta}$
- because $\hat{\beta} = (X'X)^{-1}X'Y = \beta + (X'X)^{-1}X'\varepsilon$, we have

$$\sqrt{N}(\hat{\beta} - \beta) = (X'X/N)^{-1}(X'\varepsilon/\sqrt{N}) \quad (1)$$

- we will maintain the assumption that $X'\varepsilon/N \rightarrow_p 0$
 - an easy sufficient condition is that ε is *mean independent* of X , i.e. $E[\varepsilon|X] = 0$
 - we don't want to go as far as assuming ε is independent of X though
 - ▶ why not? Full independence implies no heteroskedasticity or clustering
 - if $X'\varepsilon/N \rightarrow_p 0$, we get that OLS is *consistent* for β
- the simple case is one where $E[\varepsilon\varepsilon'|X] = \sigma^2 I$
 - take variances on both sides of (1) to get that

$$\sqrt{N}(\hat{\beta} - \beta) \rightarrow_d N(0, \sigma^2(X'X)^{-1})$$

- notice, this is just a multivariable generalization of $V[\hat{\beta}] = \sigma_\varepsilon^2/V[X]$
- so, to do inference on the elements of $\hat{\beta}$ (or functions of them) in practice, we'd use the *asymptotic covariance matrix* $s^2(X'X)^{-1}/N$

Asymptotic Distribution of the OLS Estimator

- we want to apply a central limit theorem to $\hat{\beta}$
- because $\hat{\beta} = (X'X)^{-1}X'Y = \beta + (X'X)^{-1}X'\varepsilon$, we have

$$\sqrt{N}(\hat{\beta} - \beta) = (X'X/N)^{-1}(X'\varepsilon/\sqrt{N}) \quad (1)$$

- we will maintain the assumption that $X'\varepsilon/N \rightarrow_p 0$
 - an easy sufficient condition is that ε is *mean independent* of X , i.e. $E[\varepsilon|X] = 0$
 - we don't want to go as far as assuming ε is independent of X though
 - ▶ why not? Full independence implies no heteroskedasticity or clustering
 - if $X'\varepsilon/N \rightarrow_p 0$, we get that OLS is *consistent* for β
- the simple case is one where $E[\varepsilon\varepsilon'|X] = \sigma^2 I$
 - take variances on both sides of (1) to get that

$$\sqrt{N}(\hat{\beta} - \beta) \rightarrow_d N(0, \sigma^2(X'X)^{-1})$$

- notice, this is just a multivariable generalization of $V[\hat{\beta}] = \sigma_\varepsilon^2/V[X]$
- so, to do inference on the elements of $\hat{\beta}$ (or functions of them) in practice, we'd use the *asymptotic covariance matrix* $s^2(X'X)^{-1}/N$

Asymptotic Distribution of the OLS Estimator

- we want to apply a central limit theorem to $\hat{\beta}$
- because $\hat{\beta} = (X'X)^{-1}X'Y = \beta + (X'X)^{-1}X'\varepsilon$, we have

$$\sqrt{N}(\hat{\beta} - \beta) = (X'X/N)^{-1}(X'\varepsilon/\sqrt{N}) \quad (1)$$

- we will maintain the assumption that $X'\varepsilon/N \rightarrow_p 0$
 - an easy sufficient condition is that ε is *mean independent* of X , i.e. $E[\varepsilon|X] = 0$
 - we don't want to go as far as assuming ε is independent of X though
 - ▶ why not? Full independence implies no heteroskedasticity or clustering
 - if $X'\varepsilon/N \rightarrow_p 0$, we get that OLS is *consistent* for β
- the simple case is one where $E[\varepsilon\varepsilon'|X] = \sigma^2 I$
 - take variances on both sides of (1) to get that

$$\sqrt{N}(\hat{\beta} - \beta) \rightarrow_d N(0, \sigma^2(X'X)^{-1})$$

- notice, this is just a multivariable generalization of $V[\hat{\beta}] = \sigma_\varepsilon^2/V[X]$
- so, to do inference on the elements of $\hat{\beta}$ (or functions of them) in practice, we'd use the *asymptotic covariance matrix* $s^2(X'X)^{-1}/N$

Asymptotic Distribution of the OLS Estimator

- we want to apply a central limit theorem to $\hat{\beta}$
- because $\hat{\beta} = (X'X)^{-1}X'Y = \beta + (X'X)^{-1}X'\varepsilon$, we have

$$\sqrt{N}(\hat{\beta} - \beta) = (X'X/N)^{-1}(X'\varepsilon/\sqrt{N}) \quad (1)$$

- we will maintain the assumption that $X'\varepsilon/N \rightarrow_p 0$
 - an easy sufficient condition is that ε is *mean independent* of X , i.e. $E[\varepsilon|X] = 0$
 - we don't want to go as far as assuming ε is independent of X though
 - ▶ why not? Full independence implies no heteroskedasticity or clustering
 - if $X'\varepsilon/N \rightarrow_p 0$, we get that OLS is *consistent* for β
- the simple case is one where $E[\varepsilon\varepsilon'|X] = \sigma^2 I$
 - take variances on both sides of (1) to get that

$$\sqrt{N}(\hat{\beta} - \beta) \rightarrow_d N(0, \sigma^2(X'X)^{-1})$$

- notice, this is just a multivariable generalization of $V[\hat{\beta}] = \sigma_\varepsilon^2/V[X]$
- so, to do inference on the elements of $\hat{\beta}$ (or functions of them) in practice, we'd use the *asymptotic covariance matrix* $s^2(X'X)^{-1}/N$

Asymptotic Distribution of the OLS Estimator

- we always get that $(X'X/N)^{-1} \rightarrow_p (E[X'X])^{-1} = M^{-1}$, by the LLN
- so, the key part of the previous argument was characterizing the limiting value of

$$\text{plim } X'\epsilon\epsilon'X/N = V, \text{ say}$$

- “best practice” in applied micro is *not* to try and explicitly model V
 - you might imagine writing down a parametric model $V(\gamma)$ and trying to estimate γ simultaneously with β
 - e.g., using the estimated OLS residuals $\hat{\epsilon}$ as a first-stage input into the estimation of γ , then using $\hat{\gamma}$ to re-estimate β
 - If you specify the model for $V(\gamma)$ correctly, this can yield efficiency gains over OLS
 - But OLS is major distribution-free tool you get for free! For a given γ , you don't need to know anything about V to get the best linear unbiased estimator of β
 - This approach is usually called “generalized least squares” (GLS)
 - instead, various data-driven approximations of V are used to get “robust” standard errors

Asymptotic Distribution of the OLS Estimator

- we always get that $(X'X/N)^{-1} \rightarrow_p (E[X'X])^{-1} = M^{-1}$, by the LLN
- so, the key part of the previous argument was characterizing the limiting value of

$$\text{plim } X' \varepsilon \varepsilon' X / N = V, \text{ say}$$

- “best practice” in applied micro is *not* to try and explicitly model V
 - you might imagine writing down a parametric model $V(\gamma)$ and trying to estimate γ simultaneously with β
 - i.e., using the estimated OLS residuals $\hat{\varepsilon}$ as a first-stage input into the estimation of γ , then using $\hat{\gamma}$ to re-estimate β
 - if you specify the model for $V(\gamma)$ correctly, this can yield efficiency gains over OLS
 - but OLS is simpler, doesn't require specifying the model for V , and is robust to misspecification
 - so, instead, various data-driven approximations of V are used to get “robust” standard errors

Asymptotic Distribution of the OLS Estimator

- we always get that $(X'X/N)^{-1} \rightarrow_p (E[X'X])^{-1} = M^{-1}$, by the LLN
- so, the key part of the previous argument was characterizing the limiting value of

$$\text{plim } X' \varepsilon \varepsilon' X / N = V, \text{ say}$$

- “best practice” in applied micro is *not* to try and explicitly model V
 - you might imagine writing down a parametric model $V(\gamma)$ and trying to estimate γ simultaneously with β
 - ▶ or, using the estimated OLS residuals $\hat{\varepsilon}$ as a first-stage input into the estimation of γ , then using $\hat{\gamma}$ to re-estimate β
 - ▶ if you specify the model for $V(\gamma)$ correctly, this can yield efficiency gains over OLS
 - ▶ BUT, a major disadvantage is that if you get the model for V wrong, you end up losing consistency of $\hat{\beta}$
 - ▶ this approach is usually called “generalized least squares” (GLS)
 - instead, various data-driven approximations of V are used to get “robust” standard errors

Asymptotic Distribution of the OLS Estimator

- we always get that $(X'X/N)^{-1} \rightarrow_p (E[X'X])^{-1} = M^{-1}$, by the LLN
- so, the key part of the previous argument was characterizing the limiting value of

$$\text{plim } X' \varepsilon \varepsilon' X / N = V, \text{ say}$$

- “best practice” in applied micro is *not* to try and explicitly model V
 - you might imagine writing down a parametric model $V(\gamma)$ and trying to estimate γ simultaneously with β
 - ▶ or, using the estimated OLS residuals $\hat{\varepsilon}$ as a first-stage input into the estimation of γ , then using $\hat{\gamma}$ to re-estimate β
 - ▶ if you specify the model for $V(\gamma)$ correctly, this can yield efficiency gains over OLS
 - ▶ BUT, a major disadvantage is that if you get the model for V wrong, you end up losing consistency of $\hat{\beta}$
 - ▶ this approach is usually called “generalized least squares” (GLS)
- instead, various data-driven approximations of V are used to get “robust” standard errors

Asymptotic Distribution of the OLS Estimator

- we always get that $(X'X/N)^{-1} \rightarrow_p (E[X'X])^{-1} = M^{-1}$, by the LLN
- so, the key part of the previous argument was characterizing the limiting value of

$$\text{plim } X' \varepsilon \varepsilon' X / N = V, \text{ say}$$

- “best practice” in applied micro is *not* to try and explicitly model V
 - you might imagine writing down a parametric model $V(\gamma)$ and trying to estimate γ simultaneously with β
 - ▶ or, using the estimated OLS residuals $\hat{\varepsilon}$ as a first-stage input into the estimation of γ , then using $\hat{\gamma}$ to re-estimate β
 - ▶ if you specify the model for $V(\gamma)$ correctly, this can yield efficiency gains over OLS
 - ▶ BUT, a major disadvantage is that if you get the model for V wrong, you end up losing consistency of $\hat{\beta}$
 - ▶ this approach is usually called “generalized least squares” (GLS)
 - instead, various data-driven approximations of V are used to get “robust” standard errors

Asymptotic Distribution of the OLS Estimator

- we always get that $(X'X/N)^{-1} \rightarrow_p (E[X'X])^{-1} = M^{-1}$, by the LLN
- so, the key part of the previous argument was characterizing the limiting value of

$$\text{plim } X' \varepsilon \varepsilon' X / N = V, \text{ say}$$

- “best practice” in applied micro is *not* to try and explicitly model V
 - you might imagine writing down a parametric model $V(\gamma)$ and trying to estimate γ simultaneously with β
 - ▶ or, using the estimated OLS residuals $\hat{\varepsilon}$ as a first-stage input into the estimation of γ , then using $\hat{\gamma}$ to re-estimate β
 - ▶ if you specify the model for $V(\gamma)$ correctly, this can yield efficiency gains over OLS
 - ▶ BUT, a major disadvantage is that if you get the model for V wrong, you end up losing consistency of $\hat{\beta}$
 - ▶ this approach is usually called “generalized least squares” (GLS)
- instead, various data-driven approximations of V are used to get “robust” standard errors

Asymptotic Distribution of the OLS Estimator

- we always get that $(X'X/N)^{-1} \rightarrow_p (E[X'X])^{-1} = M^{-1}$, by the LLN
- so, the key part of the previous argument was characterizing the limiting value of

$$\text{plim } X' \varepsilon \varepsilon' X / N = V, \text{ say}$$

- “best practice” in applied micro is *not* to try and explicitly model V
 - you might imagine writing down a parametric model $V(\gamma)$ and trying to estimate γ simultaneously with β
 - ▶ or, using the estimated OLS residuals $\hat{\varepsilon}$ as a first-stage input into the estimation of γ , then using $\hat{\gamma}$ to re-estimate β
 - ▶ if you specify the model for $V(\gamma)$ correctly, this can yield efficiency gains over OLS
 - ▶ BUT, a major disadvantage is that if you get the model for V wrong, you end up losing consistency of $\hat{\beta}$
 - ▶ this approach is usually called “generalized least squares” (GLS)
- instead, various data-driven approximations of V are used to get “robust” standard errors

Asymptotic Distribution of the OLS Estimator

- we always get that $(X'X/N)^{-1} \rightarrow_p (E[X'X])^{-1} = M^{-1}$, by the LLN
- so, the key part of the previous argument was characterizing the limiting value of

$$\text{plim } X' \varepsilon \varepsilon' X / N = V, \text{ say}$$

- “best practice” in applied micro is *not* to try and explicitly model V
 - you might imagine writing down a parametric model $V(\gamma)$ and trying to estimate γ simultaneously with β
 - ▶ or, using the estimated OLS residuals $\hat{\varepsilon}$ as a first-stage input into the estimation of γ , then using $\hat{\gamma}$ to re-estimate β
 - ▶ if you specify the model for $V(\gamma)$ correctly, this can yield efficiency gains over OLS
 - ▶ BUT, a major disadvantage is that if you get the model for V wrong, you end up losing consistency of $\hat{\beta}$
 - ▶ this approach is usually called “generalized least squares” (GLS)
- instead, various data-driven approximations of V are used to get “robust” standard errors

Asymptotic Distribution of the OLS Estimator

- we always get that $(X'X/N)^{-1} \rightarrow_p (E[X'X])^{-1} = M^{-1}$, by the LLN
- so, the key part of the previous argument was characterizing the limiting value of

$$\text{plim } X' \varepsilon \varepsilon' X / N = V, \text{ say}$$

- “best practice” in applied micro is *not* to try and explicitly model V
 - you might imagine writing down a parametric model $V(\gamma)$ and trying to estimate γ simultaneously with β
 - ▶ or, using the estimated OLS residuals $\hat{\varepsilon}$ as a first-stage input into the estimation of γ , then using $\hat{\gamma}$ to re-estimate β
 - ▶ if you specify the model for $V(\gamma)$ correctly, this can yield efficiency gains over OLS
 - ▶ BUT, a major disadvantage is that if you get the model for V wrong, you end up losing consistency of $\hat{\beta}$
 - ▶ this approach is usually called “generalized least squares” (GLS)
- instead, various data-driven approximations of V are used to get “robust” standard errors

Asymptotic Distribution of the OLS Estimator

- the goal is to obtain estimates of the precision of $\hat{\beta}$ that are approximately correct under a wide range of assumptions about the exact form of $E[\varepsilon\varepsilon'|X]$
 - after all, we know OLS is consistent (if possibly inefficient)
 - GLS may not even be consistent if we misspecify the model for the error covariances!
- if you carry out the matrix multiplication you will see that

$$X'\varepsilon\varepsilon'X/N = N^{-1} \sum_{i=1}^N \sum_{j=1}^N X_j X_i' \varepsilon_i \varepsilon_j$$

- there are different choices of “robust” standard errors
 - Newey-West, Eicker-White, HC0, HC1, etc
 - all of these amount to different choices of weights ω_{ij} in a formula like

$$\hat{V} = \sum_{i=1}^N \sum_{j=1}^N \omega_{ij} X_j X_i' \hat{\varepsilon}_i \hat{\varepsilon}_j$$

Asymptotic Distribution of the OLS Estimator

- the goal is to obtain estimates of the precision of $\hat{\beta}$ that are approximately correct under a wide range of assumptions about the exact form of $E[\varepsilon\varepsilon'|X]$
 - after all, we know OLS is consistent (if possibly inefficient)
 - GLS may not even be consistent if we misspecify the model for the error covariances!
- if you carry out the matrix multiplication you will see that

$$X'\varepsilon\varepsilon'X/N = N^{-1} \sum_{i=1}^N \sum_{j=1}^N X_j X_i' \varepsilon_i \varepsilon_j$$

- there are different choices of “robust” standard errors
 - Newey-West, Eicker-White, HC0, HC1, etc
 - all of these amount to different choices of weights ω_{ij} in a formula like

$$\hat{V} = \sum_{i=1}^N \sum_{j=1}^N \omega_{ij} X_j X_i' \hat{\varepsilon}_i \hat{\varepsilon}_j$$

Asymptotic Distribution of the OLS Estimator

- the goal is to obtain estimates of the precision of $\hat{\beta}$ that are approximately correct under a wide range of assumptions about the exact form of $E[\varepsilon\varepsilon'|X]$
 - after all, we know OLS is consistent (if possibly inefficient)
 - GLS may not even be consistent if we misspecify the model for the error covariances!
- if you carry out the matrix multiplication you will see that

$$X'\varepsilon\varepsilon'X/N = N^{-1} \sum_{i=1}^N \sum_{j=1}^N X_j X_i' \varepsilon_i \varepsilon_j$$

- there are different choices of “robust” standard errors
 - Newey-West, Eicker-White, HC0, HC1, etc
 - all of these amount to different choices of weights ω_{ij} in a formula like

$$\hat{V} = \sum_{i=1}^N \sum_{j=1}^N \omega_{ij} X_j X_i' \hat{\varepsilon}_i \hat{\varepsilon}_j$$

Asymptotic Distribution of the OLS Estimator

- the goal is to obtain estimates of the precision of $\hat{\beta}$ that are approximately correct under a wide range of assumptions about the exact form of $E[\varepsilon\varepsilon'|X]$
 - after all, we know OLS is consistent (if possibly inefficient)
 - GLS may not even be consistent if we misspecify the model for the error covariances!
- if you carry out the matrix multiplication you will see that

$$X'\varepsilon\varepsilon'X/N = N^{-1} \sum_{i=1}^N \sum_{j=1}^N X_j X_i' \varepsilon_i \varepsilon_j$$

- there are different choices of “robust” standard errors
 - Newey-West, Eicker-White, HC0, HC1, etc
 - all of these amount to different choices of weights ω_{ij} in a formula like

$$\hat{V} = \sum_{i=1}^N \sum_{j=1}^N \omega_{ij} X_j X_i' \hat{\varepsilon}_i \hat{\varepsilon}_j$$

Asymptotic Distribution of the OLS Estimator

- the goal is to obtain estimates of the precision of $\hat{\beta}$ that are approximately correct under a wide range of assumptions about the exact form of $E[\varepsilon\varepsilon'|X]$
 - after all, we know OLS is consistent (if possibly inefficient)
 - GLS may not even be consistent if we misspecify the model for the error covariances!
- if you carry out the matrix multiplication you will see that

$$X'\varepsilon\varepsilon'X/N = N^{-1} \sum_{i=1}^N \sum_{j=1}^N X_j X_i' \varepsilon_i \varepsilon_j$$

- there are different choices of “robust” standard errors
 - Newey-West, Eicker-White, HC0, HC1, etc
 - all of these amount to different choices of weights ω_{ij} in a formula like

$$\hat{V} = \sum_{i=1}^N \sum_{j=1}^N \omega_{ij} X_j X_i' \hat{\varepsilon}_i \hat{\varepsilon}_j$$

Asymptotic Distribution of the OLS Estimator

- the goal is to obtain estimates of the precision of $\hat{\beta}$ that are approximately correct under a wide range of assumptions about the exact form of $E[\varepsilon\varepsilon'|X]$
 - after all, we know OLS is consistent (if possibly inefficient)
 - GLS may not even be consistent if we misspecify the model for the error covariances!
- if you carry out the matrix multiplication you will see that

$$X'\varepsilon\varepsilon'X/N = N^{-1} \sum_{i=1}^N \sum_{j=1}^N X_j X_i' \varepsilon_i \varepsilon_j$$

- there are different choices of “robust” standard errors
 - Newey-West, Eicker-White, HC0, HC1, etc
 - all of these amount to different choices of weights ω_{ij} in a formula like

$$\hat{V} = \sum_{i=1}^N \sum_{j=1}^N \omega_{ij} X_j X_i' \hat{\varepsilon}_i \hat{\varepsilon}_j$$

Asymptotic Distribution of the OLS Estimator

- the goal is to obtain estimates of the precision of $\hat{\beta}$ that are approximately correct under a wide range of assumptions about the exact form of $E[\varepsilon\varepsilon'|X]$
 - after all, we know OLS is consistent (if possibly inefficient)
 - GLS may not even be consistent if we misspecify the model for the error covariances!
- if you carry out the matrix multiplication you will see that

$$X'\varepsilon\varepsilon'X/N = N^{-1} \sum_{i=1}^N \sum_{j=1}^N X_j X_i' \varepsilon_i \varepsilon_j$$

- there are different choices of “robust” standard errors
 - Newey-West, Eicker-White, HC0, HC1, etc
 - all of these amount to different choices of weights ω_{ij} in a formula like

$$\hat{V} = \sum_{i=1}^N \sum_{j=1}^N \omega_{ij} X_j X_i' \hat{\varepsilon}_i \hat{\varepsilon}_j$$

Asymptotic Distribution of the OLS Estimator

- for more details about this, see Ch. 4.4 of Cameron and Trivedi (2005) or Ch. 8 of Angrist and Pischke (2008)
 - Cameron and Miller (2015) is a useful reference about clustering, in particular
- next, we turn to a different question: why are we running OLS in the first place?

Asymptotic Distribution of the OLS Estimator

- for more details about this, see Ch. 4.4 of Cameron and Trivedi (2005) or Ch. 8 of Angrist and Pischke (2008)
 - Cameron and Miller (2015) is a useful reference about clustering, in particular
- next, we turn to a different question: why are we running OLS in the first place?

Asymptotic Distribution of the OLS Estimator

- for more details about this, see Ch. 4.4 of Cameron and Trivedi (2005) or Ch. 8 of Angrist and Pischke (2008)
 - Cameron and Miller (2015) is a useful reference about clustering, in particular
- next, we turn to a different question: why are we running OLS in the first place?

OLS as the Best Linear Approximation of $E[Y|X]$

- we can motivate OLS without literally believing $Y = X\beta + \varepsilon$ is the data-generating process
- instead, consider the following problems
 - $\beta^* = \arg \min_b E[(Y - Xb)^2]$, finding the best linear predictor of Y
 - $\beta^{**} = \arg \min_b E[(E[Y|X] - Xb)^2]$, finding the best linear approximation to $E[Y|X]$
- the OLS coefficient is β^* by definition, but these two problems have identical solutions
 - so, we can always think of the OLS coefficient as providing the best linear approximation to the conditional mean $E[Y|X]$, even if it is nonlinear
- of course, these facts tell us *nothing* about causality!
 - the *causal* question “what would happen to Y on average if we manipulated X by one unit” makes no sense without a model!
 - “what would happen to Y if we changed the car’s color?”
 - “what would happen to Y if we changed the car’s weight?”
 - “what would happen to Y if we changed the car’s engine?”
 - on the other hand, if you start with a causal model (say from economic theory), knowing that OLS estimates approximate $E[Y|X]$ helps you think about whether you are going to get a good estimate of the causal effect you are trying to measure

OLS as the Best Linear Approximation of $E[Y|X]$

- we can motivate OLS without literally believing $Y = X\beta + \varepsilon$ is the data-generating process
- instead, consider the following problems
 - $\beta^* = \arg \min_b E[(Y - Xb)^2]$, finding the best linear predictor of Y
 - $\beta^{**} = \arg \min_b E[(E[Y|X] - Xb)^2]$, finding the best linear approximation to $E[Y|X]$
- the OLS coefficient is β^* by definition, but these two problems have identical solutions
 - so, we can always think of the OLS coefficient as providing the best linear approximation to the conditional mean $E[Y|X]$, even if it is nonlinear
- of course, these facts tell us *nothing* about causality!
 - the *causal* question “what would happen to Y on average if we manipulated X by one unit” makes no sense without a model!
 - “what would happen to Y if we changed X ?”
 - “what would happen to Y if we changed X ?”
 - “what would happen to Y if we changed X ?”
 - on the other hand, if you start with a causal model (say from economic theory), knowing that OLS estimates approximate $E[Y|X]$ helps you think about whether you are going to get a good estimate of the causal effect you are trying to measure

OLS as the Best Linear Approximation of $E[Y|X]$

- we can motivate OLS without literally believing $Y = X\beta + \varepsilon$ is the data-generating process
- instead, consider the following problems
 - $\beta^* = \arg \min_b E[(Y - Xb)^2]$, finding the best linear predictor of Y
 - $\beta^{**} = \arg \min_b E[(E[Y|X] - Xb)^2]$, finding the best linear approximation to $E[Y|X]$
- the OLS coefficient is β^* by definition, but these two problems have identical solutions
 - so, we can always think of the OLS coefficient as providing the best linear approximation to the conditional mean $E[Y|X]$, even if it is nonlinear
- of course, these facts tell us *nothing* about causality!
 - the *causal* question “what would happen to Y on average if we manipulated X by one unit” makes no sense without a model!
 - “causal” questions produce reliable
 - “statistical” predictions can be wrong
 - on the other hand, if you start with a causal model (say from economic theory), knowing that OLS estimates approximate $E[Y|X]$ helps you think about whether you are going to get a good estimate of the causal effect you are trying to measure

OLS as the Best Linear Approximation of $E[Y|X]$

- we can motivate OLS without literally believing $Y = X\beta + \varepsilon$ is the data-generating process
- instead, consider the following problems
 - $\beta^* = \arg \min_b E[(Y - Xb)^2]$, finding the best linear predictor of Y
 - $\beta^{**} = \arg \min_b E[(E[Y|X] - Xb)^2]$, finding the best linear approximation to $E[Y|X]$
- the OLS coefficient is β^* by definition, but these two problems have identical solutions
 - so, we can always think of the OLS coefficient as providing the best linear approximation to the conditional mean $E[Y|X]$, even if it is nonlinear
- of course, these facts tell us *nothing* about causality!
 - the *causal* question “what would happen to Y on average if we manipulated X by one unit” makes no sense without a model!
 - “causal model” → “causal graph” → “causal model”
 - “causal model” → “causal graph” → “causal model”
 - on the other hand, if you start with a causal model (say from economic theory), knowing that OLS estimates approximate $E[Y|X]$ helps you think about whether you are going to get a good estimate of the causal effect you are trying to measure

OLS as the Best Linear Approximation of $E[Y|X]$

- we can motivate OLS without literally believing $Y = X\beta + \varepsilon$ is the data-generating process
- instead, consider the following problems
 - $\beta^* = \arg \min_b E[(Y - Xb)^2]$, finding the best linear predictor of Y
 - $\beta^{**} = \arg \min_b E[(E[Y|X] - Xb)^2]$, finding the best linear approximation to $E[Y|X]$
- the OLS coefficient is β^* by definition, but these two problems have identical solutions
 - so, we can always think of the OLS coefficient as providing the best linear approximation to the conditional mean $E[Y|X]$, even if it is nonlinear
- of course, these facts tell us *nothing* about causality!
 - the *causal* question “what would happen to Y on average if we manipulated X by one unit” makes no sense without a model!
 - *causal* questions are *not* predictive questions
 - *causal* questions are *not* predictive questions
 - on the other hand, if you start with a causal model (say from economic theory), knowing that OLS estimates approximate $E[Y|X]$ helps you think about whether you are going to get a good estimate of the causal effect you are trying to measure

OLS as the Best Linear Approximation of $E[Y|X]$

- we can motivate OLS without literally believing $Y = X\beta + \varepsilon$ is the data-generating process
- instead, consider the following problems
 - $\beta^* = \arg \min_b E[(Y - Xb)^2]$, finding the best linear predictor of Y
 - $\beta^{**} = \arg \min_b E[(E[Y|X] - Xb)^2]$, finding the best linear approximation to $E[Y|X]$
- the OLS coefficient is β^* by definition, but these two problems have identical solutions
 - so, we can always think of the OLS coefficient as providing the best linear approximation to the conditional mean $E[Y|X]$, even if it is nonlinear
- of course, these facts tell us *nothing* about causality!
 - the *causal* question “what would happen to Y on average if we manipulated X by one unit” makes no sense without a model!
 - on the other hand, if you start with a causal model (say from economic theory), knowing that OLS estimates approximate $E[Y|X]$ helps you think about whether you are going to get a good estimate of the causal effect you are trying to measure

OLS as the Best Linear Approximation of $E[Y|X]$

- we can motivate OLS without literally believing $Y = X\beta + \varepsilon$ is the data-generating process
- instead, consider the following problems
 - $\beta^* = \arg \min_b E[(Y - Xb)^2]$, finding the best linear predictor of Y
 - $\beta^{**} = \arg \min_b E[(E[Y|X] - Xb)^2]$, finding the best linear approximation to $E[Y|X]$
- the OLS coefficient is β^* by definition, but these two problems have identical solutions
 - so, we can always think of the OLS coefficient as providing the best linear approximation to the conditional mean $E[Y|X]$, even if it is nonlinear
- of course, these facts tell us *nothing* about causality!
 - the *causal* question “what would happen to Y on average if we manipulated X by one unit” makes no sense without a model!
 - ▶ umbrella prevalence predicts rainfall
 - ▶ ambulances predict car crashes
 - on the other hand, if you start with a causal model (say from economic theory), knowing that OLS estimates approximate $E[Y|X]$ helps you think about whether you are going to get a good estimate of the causal effect you are trying to measure

OLS as the Best Linear Approximation of $E[Y|X]$

- we can motivate OLS without literally believing $Y = X\beta + \varepsilon$ is the data-generating process
- instead, consider the following problems
 - $\beta^* = \arg \min_b E[(Y - Xb)^2]$, finding the best linear predictor of Y
 - $\beta^{**} = \arg \min_b E[(E[Y|X] - Xb)^2]$, finding the best linear approximation to $E[Y|X]$
- the OLS coefficient is β^* by definition, but these two problems have identical solutions
 - so, we can always think of the OLS coefficient as providing the best linear approximation to the conditional mean $E[Y|X]$, even if it is nonlinear
- of course, these facts tell us *nothing* about causality!
 - the *causal* question “what would happen to Y on average if we manipulated X by one unit” makes no sense without a model!
 - ▶ umbrella prevalence predicts rainfall
 - ▶ ambulances predict car crashes
 - on the other hand, if you start with a causal model (say from economic theory), knowing that OLS estimates approximate $E[Y|X]$ helps you think about whether you are going to get a good estimate of the causal effect you are trying to measure

OLS as the Best Linear Approximation of $E[Y|X]$

- we can motivate OLS without literally believing $Y = X\beta + \varepsilon$ is the data-generating process
- instead, consider the following problems
 - $\beta^* = \arg \min_b E[(Y - Xb)^2]$, finding the best linear predictor of Y
 - $\beta^{**} = \arg \min_b E[(E[Y|X] - Xb)^2]$, finding the best linear approximation to $E[Y|X]$
- the OLS coefficient is β^* by definition, but these two problems have identical solutions
 - so, we can always think of the OLS coefficient as providing the best linear approximation to the conditional mean $E[Y|X]$, even if it is nonlinear
- of course, these facts tell us *nothing* about causality!
 - the *causal* question “what would happen to Y on average if we manipulated X by one unit” makes no sense without a model!
 - ▶ umbrella prevalence predicts rainfall
 - ▶ ambulances predict car crashes
 - on the other hand, if you start with a causal model (say from economic theory), knowing that OLS estimates approximate $E[Y|X]$ helps you think about whether you are going to get a good estimate of the causal effect you are trying to measure

OLS as the Best Linear Approximation of $E[Y|X]$

- we can motivate OLS without literally believing $Y = X\beta + \varepsilon$ is the data-generating process
- instead, consider the following problems
 - $\beta^* = \arg \min_b E[(Y - Xb)^2]$, finding the best linear predictor of Y
 - $\beta^{**} = \arg \min_b E[(E[Y|X] - Xb)^2]$, finding the best linear approximation to $E[Y|X]$
- the OLS coefficient is β^* by definition, but these two problems have identical solutions
 - so, we can always think of the OLS coefficient as providing the best linear approximation to the conditional mean $E[Y|X]$, even if it is nonlinear
- of course, these facts tell us *nothing* about causality!
 - the *causal* question “what would happen to Y on average if we manipulated X by one unit” makes no sense without a model!
 - ▶ umbrella prevalence predicts rainfall
 - ▶ ambulances predict car crashes
 - on the other hand, if you start with a causal model (say from economic theory), knowing that OLS estimates approximate $E[Y|X]$ helps you think about whether you are going to get a good estimate of the causal effect you are trying to measure

OLS as the Best Linear Approximation of $E[Y|X]$

- we can motivate OLS without literally believing $Y = X\beta + \varepsilon$ is the data-generating process
- instead, consider the following problems
 - $\beta^* = \arg \min_b E[(Y - Xb)^2]$, finding the best linear predictor of Y
 - $\beta^{**} = \arg \min_b E[(E[Y|X] - Xb)^2]$, finding the best linear approximation to $E[Y|X]$
- the OLS coefficient is β^* by definition, but these two problems have identical solutions
 - so, we can always think of the OLS coefficient as providing the best linear approximation to the conditional mean $E[Y|X]$, even if it is nonlinear
- of course, these facts tell us *nothing* about causality!
 - the *causal* question “what would happen to Y on average if we manipulated X by one unit” makes no sense without a model!
 - ▶ umbrella prevalence predicts rainfall
 - ▶ ambulances predict car crashes
 - on the other hand, if you start with a causal model (say from economic theory), knowing that OLS estimates approximate $E[Y|X]$ helps you think about whether you are going to get a good estimate of the causal effect you are trying to measure

More on Causality

- consider the following setup:
 - y_i is output per acre on farm i
 - x_{i1} is an index of soil quality
 - x_{i2} is an index of weather quality
 - x_{i3} is an index of pesticide use
 - e_i is a measure of insect population density
- We know that crop yields are determined as

$$y = x_1\beta_1 + x_2\beta_2 + x_3\beta_3 + e$$

- x_1 , x_2 , and e are mutually independent with mean zero

More on Causality

- consider the following setup:
 - y_i is output per acre on farm i
 - x_{i1} is an index of soil quality
 - x_{i2} is an index of weather quality
 - x_{i3} is an index of pesticide use
 - e_i is a measure of insect population density
- We know that crop yields are determined as

$$y = x_1\beta_1 + x_2\beta_2 + x_3\beta_3 + e$$

- x_1 , x_2 , and e are mutually independent with mean zero

More on Causality

- consider the following setup:
 - y_i is output per acre on farm i
 - x_{i1} is an index of soil quality
 - x_{i2} is an index of weather quality
 - x_{i3} is an index of pesticide use
 - e_i is a measure of insect population density
- We know that crop yields are determined as

$$y = x_1\beta_1 + x_2\beta_2 + x_3\beta_3 + e$$

- x_1 , x_2 , and e are mutually independent with mean zero

More on Causality

- consider the following setup:
 - y_i is output per acre on farm i
 - x_{i1} is an index of soil quality
 - x_{i2} is an index of weather quality
 - x_{i3} is an index of pesticide use
 - e_i is a measure of insect population density
- We know that crop yields are determined as

$$y = x_1\beta_1 + x_2\beta_2 + x_3\beta_3 + e$$

- x_1 , x_2 , and e are mutually independent with mean zero

More on Causality

- consider the following setup:
 - y_i is output per acre on farm i
 - x_{i1} is an index of soil quality
 - x_{i2} is an index of weather quality
 - x_{i3} is an index of pesticide use
 - e_i is a measure of insect population density
- We know that crop yields are determined as

$$y = x_1\beta_1 + x_2\beta_2 + x_3\beta_3 + e$$

- x_1 , x_2 , and e are mutually independent with mean zero

More on Causality

- consider the following setup:
 - y_i is output per acre on farm i
 - x_{i1} is an index of soil quality
 - x_{i2} is an index of weather quality
 - x_{i3} is an index of pesticide use
 - e_i is a measure of insect population density
- We know that crop yields are determined as

$$y = x_1\beta_1 + x_2\beta_2 + x_3\beta_3 + e$$

- x_1 , x_2 , and e are mutually independent with mean zero

More on Causality

- consider the following setup:
 - y_i is output per acre on farm i
 - x_{i1} is an index of soil quality
 - x_{i2} is an index of weather quality
 - x_{i3} is an index of pesticide use
 - e_i is a measure of insect population density
- We know that crop yields are determined as

$$y = x_1\beta_1 + x_2\beta_2 + x_3\beta_3 + e$$

- x_1 , x_2 , and e are mutually independent with mean zero

More on Causality

- consider the following setup:
 - y_i is output per acre on farm i
 - x_{i1} is an index of soil quality
 - x_{i2} is an index of weather quality
 - x_{i3} is an index of pesticide use
 - e_i is a measure of insect population density
- We know that crop yields are determined as

$$y = x_1\beta_1 + x_2\beta_2 + x_3\beta_3 + e$$

- x_1 , x_2 , and e are mutually independent with mean zero

More on Causality

- consider two different assumptions about how pesticide use is determined:
 - model A: farmers ignore soil quality (or do not observe it), but they do observe the level of insect populations
 - ▶ they set $x_3 = -e/\gamma + \eta$ where η is independent of all the x variables
 - model B: farmers set $x_3 = x_1 - e/\gamma_2 + \varepsilon$, where ε is independent of the x variables
- You have access to data on y , x_1 , x_2 , and x_3
- do you want to control for pesticide use if your goal is to estimate β_1 ?
- suppose model A generates the data and you use x_3 in your regression
 - $\hat{\beta}_1$ will be consistent for β_1 (why?)
 - however, you cannot estimate β_3 consistently:

$$\text{plim } \hat{\beta}_3 = \beta_3 - \frac{\sigma_e^2/\gamma}{\sigma_e^2/\gamma^2 + \sigma_\eta^2} < \beta_3$$

More on Causality

- consider two different assumptions about how pesticide use is determined:
 - model A: farmers ignore soil quality (or do not observe it), but they do observe the level of insect populations
 - ▶ they set $x_3 = -e/\gamma + \eta$ where η is independent of all the x variables
 - model B: farmers set $x_3 = x_1 - e/\gamma_2 + \varepsilon$, where ε is independent of the x variables
- You have access to data on y , x_1 , x_2 , and x_3
- do you want to control for pesticide use if your goal is to estimate β_1 ?
- suppose model A generates the data and you use x_3 in your regression
 - $\hat{\beta}_1$ will be consistent for β_1 (why?)
 - however, you cannot estimate β_3 consistently:

$$\text{plim } \hat{\beta}_3 = \beta_3 - \frac{\sigma_e^2/\gamma}{\sigma_e^2/\gamma^2 + \sigma_\eta^2} < \beta_3$$

More on Causality

- consider two different assumptions about how pesticide use is determined:
 - model A: farmers ignore soil quality (or do not observe it), but they do observe the level of insect populations
 - ▶ they set $x_3 = -e/\gamma + \eta$ where η is independent of all the x variables
 - model B: farmers set $x_3 = x_1 - e/\gamma_2 + \varepsilon$, where ε is independent of the x variables
- You have access to data on y , x_1 , x_2 , and x_3
- do you want to control for pesticide use if your goal is to estimate β_1 ?
- suppose model A generates the data and you use x_3 in your regression
 - $\hat{\beta}_1$ will be consistent for β_1 (why?)
 - however, you cannot estimate β_3 consistently:

$$\text{plim } \hat{\beta}_3 = \beta_3 - \frac{\sigma_e^2/\gamma}{\sigma_e^2/\gamma^2 + \sigma_\eta^2} < \beta_3$$

More on Causality

- consider two different assumptions about how pesticide use is determined:
 - model A: farmers ignore soil quality (or do not observe it), but they do observe the level of insect populations
 - ▶ they set $x_3 = -e/\gamma + \eta$ where η is independent of all the x variables
 - model B: farmers set $x_3 = x_1 - e/\gamma_2 + \varepsilon$, where ε is independent of the x variables
- You have access to data on y , x_1 , x_2 , and x_3
- do you want to control for pesticide use if your goal is to estimate β_1 ?
- suppose model A generates the data and you use x_3 in your regression
 - $\hat{\beta}_1$ will be consistent for β_1 (why?)
 - however, you cannot estimate β_3 consistently:

$$\text{plim } \hat{\beta}_3 = \beta_3 - \frac{\sigma_e^2/\gamma}{\sigma_e^2/\gamma^2 + \sigma_\eta^2} < \beta_3$$

More on Causality

- consider two different assumptions about how pesticide use is determined:
 - model A: farmers ignore soil quality (or do not observe it), but they do observe the level of insect populations
 - ▶ they set $x_3 = -e/\gamma + \eta$ where η is independent of all the x variables
 - model B: farmers set $x_3 = x_1 - e/\gamma_2 + \varepsilon$, where ε is independent of the x variables
- You have access to data on y , x_1 , x_2 , and x_3
- do you want to control for pesticide use if your goal is to estimate β_1 ?
- suppose model A generates the data and you use x_3 in your regression
 - $\hat{\beta}_1$ will be consistent for β_1 (why?)
 - however, you cannot estimate β_3 consistently:

$$\text{plim } \hat{\beta}_3 = \beta_3 - \frac{\sigma_e^2/\gamma}{\sigma_e^2/\gamma^2 + \sigma_\eta^2} < \beta_3$$

More on Causality

- consider two different assumptions about how pesticide use is determined:
 - model A: farmers ignore soil quality (or do not observe it), but they do observe the level of insect populations
 - ▶ they set $x_3 = -e/\gamma + \eta$ where η is independent of all the x variables
 - model B: farmers set $x_3 = x_1 - e/\gamma_2 + \varepsilon$, where ε is independent of the x variables
- You have access to data on y , x_1 , x_2 , and x_3
- do you want to control for pesticide use if your goal is to estimate β_1 ?
- suppose model A generates the data and you use x_3 in your regression
 - $\hat{\beta}_1$ will be consistent for β_1 (why?)
 - however, you cannot estimate β_3 consistently:

$$\text{plim } \hat{\beta}_3 = \beta_3 - \frac{\sigma_e^2/\gamma}{\sigma_e^2/\gamma^2 + \sigma_\eta^2} < \beta_3$$

More on Causality

- consider two different assumptions about how pesticide use is determined:
 - model A: farmers ignore soil quality (or do not observe it), but they do observe the level of insect populations
 - ▶ they set $x_3 = -e/\gamma + \eta$ where η is independent of all the x variables
 - model B: farmers set $x_3 = x_1 - e/\gamma_2 + \varepsilon$, where ε is independent of the x variables
- You have access to data on y , x_1 , x_2 , and x_3
- do you want to control for pesticide use if your goal is to estimate β_1 ?
- suppose model A generates the data and you use x_3 in your regression
 - $\hat{\beta}_1$ will be consistent for β_1 (why?)
 - however, you cannot estimate β_3 consistently:

$$\text{plim } \hat{\beta}_3 = \beta_3 - \frac{\sigma_e^2/\gamma}{\sigma_e^2/\gamma^2 + \sigma_\eta^2} < \beta_3$$

More on Causality

- consider two different assumptions about how pesticide use is determined:
 - model A: farmers ignore soil quality (or do not observe it), but they do observe the level of insect populations
 - ▶ they set $x_3 = -e/\gamma + \eta$ where η is independent of all the x variables
 - model B: farmers set $x_3 = x_1 - e/\gamma_2 + \varepsilon$, where ε is independent of the x variables
- You have access to data on y , x_1 , x_2 , and x_3
- do you want to control for pesticide use if your goal is to estimate β_1 ?
- suppose model A generates the data and you use x_3 in your regression
 - $\hat{\beta}_1$ will be consistent for β_1 (why?)
 - however, you cannot estimate β_3 consistently:

$$\text{plim } \hat{\beta}_3 = \beta_3 - \frac{\sigma_e^2/\gamma}{\sigma_e^2/\gamma^2 + \sigma_\eta^2} < \beta_3$$

More on Causality

- consider two different assumptions about how pesticide use is determined:
 - model A: farmers ignore soil quality (or do not observe it), but they do observe the level of insect populations
 - ▶ they set $x_3 = -e/\gamma + \eta$ where η is independent of all the x variables
 - model B: farmers set $x_3 = x_1 - e/\gamma_2 + \varepsilon$, where ε is independent of the x variables
- You have access to data on y , x_1 , x_2 , and x_3
- do you want to control for pesticide use if your goal is to estimate β_1 ?
- suppose model A generates the data and you use x_3 in your regression
 - $\hat{\beta}_1$ will be consistent for β_1 (why?)
 - however, you cannot estimate β_3 consistently:

$$\text{plim } \hat{\beta}_3 = \beta_3 - \frac{\sigma_e^2/\gamma}{\sigma_e^2/\gamma^2 + \sigma_\eta^2} < \beta_3$$

References

- Angrist, Joshua D, and Jörn-Steffen Pischke. 2008. *Mostly Harmless Econometrics: An Empiricist's Companion*. Princeton University Press.
- Cameron, A Colin, and Douglas L Miller. 2015. "A Practitioner's Guide to Cluster-Robust Inference." *Journal of Human Resources* 50 (2): 317–72.
- Cameron, A Colin, and Pravin K Trivedi. 2005. *Microeconometrics: Methods and Applications*. New York: Cambridge University Press.
- Stachurski, John. 2016. *A Primer in Econometric Theory*. Cambridge, MA: MIT Press.
- Wooldridge, Jeffrey M. 2010. *Econometric Analysis of Cross Section and Panel Data*. 2nd ed. MIT Press Books. The MIT Press.

Table of Contents

Preliminaries

Asymptotic Distribution of the OLS Estimator

OLS as the Best Linear Approximation of $E[Y|X]$

More on Causality

References