Integral Techniques

Linearity

$$\int \left[af(x) \pm bg(x)\right] dx = a \int f(x) dx \pm b \int g(x) dx, \text{ for constants } a \text{ and } b$$

Power Rule

$$\int x^a dx = \frac{x^{a+1}}{a+1} + C, \quad \text{for constant } a \neq -1$$

Integration by Substitution

$$\int f(u(x)) \frac{du}{dx} \ dx = \int f(u) du$$

Integration by Parts

$$\int udv = uv - \int vdu$$

Trigonometric Substitution

	$x = a\sin\left(\theta\right)$	
$\sqrt{a^2-x^2}$	$dx = a\cos(\theta)d\theta$	
	$\sqrt{a^2 - x^2} = a\cos\left(\theta\right)$	
$\sqrt{a^2 + x^2}$	$x = a \tan (\theta)$	
	$dx = a\sec^2(\theta)d\theta$	
	$\sqrt{a^2 + x^2} = a\sec\left(\theta\right)$	
$\sqrt{x^2 - a^2}$	$x = a \sec(\theta)$	
	$dx = \sec(\theta)\tan(\theta)d\theta$	
	$\sqrt{x^2 - a^2} = a \tan(\theta)$	

Partial Fractions

$\frac{P(x)}{Q(x)}$, degree $(P(x)) < \text{degree}(Q(x))$	distinct	Q(x) = (ax+b)(cx+d)	$\frac{P(x)}{Q(x)} = \frac{A}{ax+b} + \frac{B}{cx+d}$
	repeated	$Q(x) = (ax+b)^2$	$\frac{P(x)}{Q(x)} = \frac{A}{ax+b} + \frac{B}{(ax+b)^2}$
Q(II)	irreducible	$Q(x) = ax^2 + bx + c$	$\frac{P(x)}{Q(x)} = \frac{Ax + B}{ax^2 + bx + c}$
	repeated	$Q(x) = (ax^2 + bx + c)^2$	$\frac{P(x)}{Q(x)} = \frac{Ax+B}{ax^2+bx+c} + \frac{Cx+D}{(ax^2+bx+c)^2}$
	irreducible	Q(x) = (ax + bx + c)	$Q(x) = \frac{1}{ax^2 + bx + c} + \frac{1}{(ax^2 + bx + c)^2}$
$\frac{P(x)}{Q(x)}$, degree $(P(x)) \ge \text{degree}(Q(x))$	long division		