

Black body radiation

cube with perfectly reflecting walls

standing electromagnetic waves inside cavity

modes

$$\mathbf{E}(\mathbf{x}, t) = -2E_0 \sin \frac{2\pi}{\lambda} x \cos 2\pi\nu t$$

$$\mathbf{B}(\mathbf{x}, t) = 2B_0 \cos \frac{2\pi}{\lambda} x \sin 2\pi\nu t$$



modes are characterized by vector \mathbf{n}

$$\lambda_{n_x} = \frac{2L}{n} \quad \nu_{n_x} = \frac{n_x c}{2L}$$

what is the energy density in the cube?

number of modes as function of frequency

average energy of each modes



Black body radiation

cube with perfectly reflecting walls

number of modes as function of frequency?

smallest frequency 'volume' ($\mathbf{n} = \mathbf{1}$)

$$V_{111} = \left(\frac{c}{2L}\right)^3$$

'frequency density'

$$\rho(\nu) = \left(\frac{2L}{c}\right)^3$$

total frequency 'volume' in interval $\nu + d\nu$

$$V(\nu)d\nu = \frac{1}{8}4\pi\nu^2 d\nu$$



with polarization up & down

$$V(\nu)d\nu = \pi\nu^2 d\nu$$

number of modes f in interval $\nu + d\nu$

$$f(\nu)d\nu = \pi\nu^2 d\nu \left(\frac{2L}{c}\right)^3 = \frac{8\pi L^3}{c^3} \nu^2 d\nu$$

$$\nu_{n_x} = \frac{n_x c}{2L}$$



Black body radiation

cube with perfectly reflecting walls

number of modes as function of frequency?

number of modes f in interval $\nu + d\nu$

$$f(\nu)d\nu = \pi\nu^2 d\nu \left(\frac{2L}{c}\right)^3 = \frac{8\pi L^3}{c^3} \nu^2 d\nu$$

average energy per mode?

classical mechanics: equipartition theory

$$\langle E \rangle = k_{\text{B}}T$$


energy density

classical mechanics

$$u(\nu, T) = \frac{1}{L^3} f(\nu) \langle E \rangle = \frac{8\pi\nu^2}{c^3} k_{\text{B}}T$$

ultraviolet catastrophe

$$U(T) = \int_0^\infty \frac{8\pi\nu^2}{c^3} k_{\text{B}}T d\nu \rightarrow \infty$$



Black body radiation

cube with perfectly reflecting walls

number of modes as function of frequency?

number of modes f in interval $\nu + d\nu$

$$f(\nu)d\nu = \pi\nu^2 d\nu \left(\frac{2L}{c}\right)^3 = \frac{8\pi L^3}{c^3} \nu^2 d\nu$$

average energy per mode?

Planck

$$\langle E \rangle = \frac{h\nu}{1 - \exp[h\nu/k_{\text{B}}T]}$$

Black body radiation


system with discrete equidistant energy levels

partition function

$$Z = \sum_{n=0}^{\infty} \exp[-\beta n \delta E] = \frac{1}{1 - \exp[-\beta \delta E]} \quad \beta = \frac{1}{k_B T}$$

average energy

$$\langle E \rangle = -\frac{\partial}{\partial \beta} \ln Z = \frac{\delta E \exp[-\beta \delta E]}{1 - \exp[-\beta \delta E]} = \frac{\delta E}{\exp[\beta \delta E] - 1}$$



using

$$\delta E = h\nu$$

leads to Planck's expression

$$\langle E \rangle = \frac{h\nu}{1 - \exp[h\nu/k_B T]}$$

electromagnetic radiation apparently has discrete energies

atoms in the wall (oscillatory sources of EM radiation) emit energy in discrete packages

Black body radiation

cube with perfectly reflecting walls

number of modes as function of frequency?

number of modes f in interval $\nu + d\nu$

$$f(\nu)d\nu = \pi\nu^2 d\nu \left(\frac{2L}{c}\right)^3 = \frac{8\pi L^3}{c^3} \nu^2 d\nu$$

average energy per mode?

Planck

$$\langle E \rangle = \frac{h\nu}{1 - \exp[h\nu/k_B T]}$$

energy density

walls emit radiation in discrete 'quanta' of energy $n h \nu$ with $n = 1, 2, 3, ..$

$$u(\nu, T) = \frac{1}{L^3} f(\nu) \langle E \rangle = \frac{8\pi\nu^2}{c^3} \frac{h\nu}{1 - \exp[h\nu/k_B T]}$$

