

Lecture 3

Linear Maps: Kernel, Image, Rank–Nullity

Throughout, $\mathbb{F}$ is $\mathbb{R}$ or $\mathbb{C}$, and $\mathcal{L}(V, W)$ denotes the linear maps $V \rightarrow W$. Tags indicate the flavour of each problem; a ★ marks a harder one.

Core tutorial route

Work **3.1, 3.4, 3.8, 3.9, 3.15, 3.19, 3.20, and 3.24** in that order (about 65–80 minutes before discussion). This eight-problem route checks linearity and basis-defined maps, a complete kernel/image calculation, an image proof, rank–nullity and dimension obstructions, projection structure, and the finite injective–surjective transfer.

Additional practice: 3.2, 3.5, 3.7, 3.10–3.12, 3.14, 3.17–3.18, 3.21–3.23, 3.26. **Bridge:** 3.3, 3.6, 3.13, 3.16 (calculus applications). **Extension:** 3.25 and 3.27–3.34 (quotients and the First Isomorphism Theorem).

Linear maps

Exercise 3.1 (computational) Which of the following maps $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ (or $\mathbb{R}^2 \rightarrow \mathbb{R}$) are linear?

$$T(x, y) = (x - y, 2x), \quad S(x, y) = (xy, x), \quad R(x, y) = |x| + y.$$

Exercise 3.2 (computational) Define $T \in \mathcal{L}(\mathbb{R}^2, \mathbb{R}^3)$ by $T(1, 0) = (1, 2, 3)$ and $T(0, 1) = (0, 1, -1)$. Find a formula for $T(x, y)$, and determine $\text{rank } T$ and $\dim \text{null } T$.

Exercise 3.3 (computational) On $\mathcal{P}(\mathbb{R})$ let D be differentiation and M multiplication by t . Compute the commutator $[D, M^2] = DM^2 - M^2D$ as an operator.

Exercise 3.4 (computational) Let $T \in \mathcal{L}(\mathbb{R}^2, \mathbb{R}^2)$ satisfy $T(1, 1) = (3, 1)$ and $T(1, -1) = (1, 3)$. Find a formula for $T(x, y)$.

Exercise 3.5 (computational) Which of the following maps $\mathbb{R}^3 \rightarrow \mathbb{R}^2$ are linear?

$$F(x, y, z) = (x + z, 2y), \quad G(x, y, z) = (x, yz), \quad H(x, y, z) = (x - y, 0).$$

Exercise 3.6 (computational) On $\mathcal{P}(\mathbb{R})$ let D be differentiation and M multiplication by t . Compute the commutator $[D, M^3] = DM^3 - M^3D$ as an operator.

Exercise 3.7 (proof) Fix $v_0 \in V$. Prove that the evaluation map $E: \mathcal{L}(V, W) \rightarrow W$ defined by $E(T) = Tv_0$ is linear.

Kernel and image

Exercise 3.8 (computational) Let $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be $Tx = Ax$ with $A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 2 & 1 \end{pmatrix}$. Find bases for $\text{null } T$ and $\text{range } T$, and verify the rank–nullity theorem.

Exercise 3.9 (proof) Suppose $v_1, \dots, v_n$ is a basis of V and $T \in \mathcal{L}(V, W)$. Prove that $\text{range } T = \text{span}(Tv_1, \dots, Tv_n)$.

Exercise 3.10 (computational) Let $\varphi: \mathbb{C}^3 \rightarrow \mathbb{C}$ be $\varphi(z_1, z_2, z_3) = z_1 + 2z_2 + 3z_3$. Describe $\text{null } \varphi$ and its dimension, and decide whether φ is surjective.

Exercise 3.11 (computational) Let $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be $Tx = Ax$ with $A = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 1 \\ 1 & 1 & 2 \end{pmatrix}$. Find bases for $\text{null } T$ and $\text{range } T$, and verify rank–nullity.

Exercise 3.12 (computational) Let $T: \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be $Tx = Ax$ with $A = \begin{pmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 1 & 3 \\ 1 & 1 & 3 & 4 \end{pmatrix}$. Find bases for $\text{null } T$ and $\text{range } T$.

Exercise 3.13 (computational) Let $T: \mathcal{P}_3(\mathbb{R}) \rightarrow \mathbb{R}^2$ be $Tp = (p(0), p'(0))$. Find $\text{null } T$ and $\text{rank } T$.

Exercise 3.14 (proof) Let $T \in \mathcal{L}(U, V)$ and $S \in \mathcal{L}(V, W)$. Prove that $\text{range}(ST) \subseteq \text{range } S$ and $\text{null } T \subseteq \text{null}(ST)$.

Rank–nullity

Exercise 3.15 (computational) Use the rank–nullity theorem to fill in each blank. (a) $T \in \mathcal{L}(\mathbb{R}^4, \mathbb{R}^3)$ with $\dim \text{null } T = 2$: find $\text{rank } T$. (b) $T \in \mathcal{L}(\mathbb{R}^6, \mathbb{R}^2)$ surjective: find $\dim \text{null } T$. (c) $T \in \mathcal{L}(\mathcal{P}_4(\mathbb{R}), \mathbb{R}^3)$ with $\text{rank } T = 3$: find $\dim \text{null } T$.

Exercise 3.16 (computational) Let D^2 be the second-derivative operator on $\mathcal{P}_4(\mathbb{R})$. Find $\dim \text{null}(D^2)$ and $\text{rank}(D^2)$.

Exercise 3.17 (computational) Let $T: \mathbb{R}^4 \rightarrow \mathbb{R}^4$ be $Tx = Ax$ with $A = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$. Find $\text{rank } T$ and a basis for $\text{null } T$.

Exercise 3.18 (proof) Suppose V is finite-dimensional and $T \in \mathcal{L}(V)$ satisfies $T^2 = 0$. Prove $\text{range } T \subseteq \text{null } T$, and deduce $\dim \text{range } T \leq \frac{1}{2} \dim V$.

Exercise 3.19 (proof) Suppose $P \in \mathcal{L}(V)$ satisfies $P^2 = P$ (a *projection*). Prove that $V = \text{null } P \oplus \text{range } P$.

Exercise 3.20 (★) Prove that every $T \in \mathcal{L}(\mathbb{F}^5, \mathbb{F}^2)$ has $\dim \text{null } T \geq 3$.

Exercise 3.21 (proof) Suppose V is finite-dimensional and $S, T \in \mathcal{L}(V)$. Prove that $\text{rank}(S + T) \leq \text{rank } S + \text{rank } T$.

Operators: injective versus surjective

Exercise 3.22 (computational) Is the operator $T \in \mathcal{L}(\mathbb{R}^3)$ given by $Tx = Ax$ with $A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$ invertible?

Exercise 3.23 (computational) Is the operator $T \in \mathcal{L}(\mathbb{R}^3)$ given by $Tx = Ax$ with $A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 1 & 1 & 1 \end{pmatrix}$ invertible? If not, give $\dim \text{null } T$.

Exercise 3.24 (proof) Suppose V, W are finite-dimensional with $\dim V = \dim W$, and $T \in \mathcal{L}(V, W)$ is injective. Prove that T is surjective.

Exercise 3.25 (★) Suppose V is finite-dimensional and $S, T \in \mathcal{L}(V)$ satisfy $ST = I$. Prove that $TS = I$ as well (so a one-sided inverse of an operator is automatically two-sided).

Exercise 3.26 (proof) Suppose V is finite-dimensional and $T \in \mathcal{L}(V)$ is surjective. Prove that T is injective.

Quotient spaces [self-study]

Exercise 3.27 (computational) Let $U = \text{span}((1, 1, 0)) \subseteq \mathbb{R}^3$. Describe the coset of $(2, 0, 1)$ modulo U , and determine $\dim(\mathbb{R}^3/U)$.

Exercise 3.28 (proof) Prove that the quotient-space operations in the lecture notes are well-defined: if $v + U = v' + U$ and $w + U = w' + U$, show $(v + w) + U = (v' + w') + U$; also show that $v + U = v' + U$ implies $av + U = av' + U$ for every scalar a .

Exercise 3.29 (computational) Let $T: \mathbb{R}^4 \rightarrow \mathbb{R}^2$ be $T(x_1, x_2, x_3, x_4) = (x_1 + x_2, x_3 - x_4)$. (a) Find $\text{null } T$. (b) Write the First Isomorphism Theorem for T explicitly: what is the isomorphism $\tilde{T}: \mathbb{R}^4/\text{null } T \rightarrow \text{range } T$?

Exercise 3.30 (computational) Let $U = \text{span}((1, 1, 1)) \subseteq \mathbb{R}^3$. Describe the coset of $(2, 0, 1)$ modulo U , and determine $\dim(\mathbb{R}^3/U)$.

Exercise 3.31 (computational) Let $U = \text{span}((1, 0, 1, 0), (0, 1, 0, 1)) \subseteq \mathbb{R}^4$. Do $(1, 1, 1, 1)$ and the zero vector lie in the same coset modulo U ? Find $\dim(\mathbb{R}^4/U)$.

Exercise 3.32 (computational) Let D be differentiation on $\mathcal{P}_3(\mathbb{R})$. Identify $\text{null } D$ and the isomorphism $\mathcal{P}_3(\mathbb{R})/\text{null } D \cong \text{range } D$ furnished by the First Isomorphism Theorem.

Exercise 3.33 (★) Let $T \in \mathcal{L}(V, W)$. Prove that $\tilde{T}: V/\text{null } T \rightarrow W$, defined by $\tilde{T}(v + \text{null } T) = Tv$, is well-defined, linear, and injective.

Exercise 3.34 (proof) Let U be a subspace of a finite-dimensional space V . Prove that the quotient map $\pi: V \rightarrow V/U$, $\pi(v) = v + U$, is linear and surjective with $\text{null } \pi = U$, and deduce $\dim(V/U) = \dim V - \dim U$.