

# Lecture 14

## Operators on Hilbert Space

---

Throughout,  $\mathcal{H}$  is complex,  $\mathcal{B}(\mathcal{H})$  denotes bounded operators, and an unbounded operator includes its specified dense domain.  $\text{spec}(T)$  is the spectrum;  $T^\dagger$  the adjoint; inner products follow the physics convention. Quantum questions distinguish the finite ideal-projective model from the general density-operator/POVM-instrument/channel framework. A ★ marks a harder problem.

### Core tutorial route

Work **14.1, 14.7, 14.9, 14.10, 14.17, 14.19, 14.26, and 14.27** in that order (about 90–105 minutes before discussion). This route covers norm and adjoint; point/continuous/residual comparison; normalized sequences proving both position and momentum unbounded; common-domain CCR plus the finite trace obstruction; boundary conditions and symmetric versus self-adjoint reasoning; the qualified compact theorem and expansion; and domain-aware PVM, Stone, and quantum-model interpretation.

**Additional practice:** 14.2–14.6 and 14.8. **Bridge:** 14.20–14.25 (finite ideal measurement and dynamics). **Extension:** 14.11–14.16 and 14.18 (polynomial commutators and ladder algebra). Bridge and extension tasks do not count toward core progress.

### Bounded operators, adjoints, spectrum

**Exercise 14.1** (computational) Find the operator norm and the adjoint of multiplication by  $m(x) = e^{ix}$  on  $L^2[0, 2\pi]$ .

**Exercise 14.2** (computational) The diagonal operator  $De_n = \frac{n}{n+1}e_n$  on  $\ell^2$ . Find its point spectrum and full spectrum.

**Exercise 14.3** (computational) Let  $De_n = 2^{-n}e_n$  on  $\ell^2$ , where  $e_1, e_2, \dots$  is the standard orthonormal basis ( $n \geq 1$ ). Find  $\|D\|$ , the adjoint  $D^\dagger$ , the point spectrum, and the full spectrum; is  $D$  self-adjoint? compact?

**Exercise 14.4** (computational) Find the operator norm, the adjoint, and the spectrum of multiplication by  $g(x) = 2x$  on  $L^2[0, 1]$ . Does it have eigenvalues?

**Exercise 14.5** (computational) Let  $S$  be the right shift  $S(x_1, x_2, \dots) = (0, x_1, x_2, \dots)$  on  $\ell^2$ . Compute  $S^\dagger S$  and  $SS^\dagger$ , give  $\|S\|$ , and decide whether  $S$  is an isometry and whether it is unitary.

**Exercise 14.6** (computational) For  $A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$  on  $\mathbb{C}^2$ , find the adjoint  $A^\dagger$ , classify  $A$  (self-adjoint / unitary / normal), and give  $\text{spec}(A)$ .

**Exercise 14.7** (proof) Classify  $\lambda = 0$  as point, continuous, or residual spectrum in each case:

1.  $P$  is a nonzero orthogonal projection with nontrivial kernel;
2.  $De_n = n^{-1}e_n$  on  $\ell^2$ ;
3.  $S$  is the unilateral right shift on  $\ell^2$ .

For the last two, justify the density or non-density of the range. Then explain why a bounded normal operator cannot have residual spectrum.

**Exercise 14.8** (proof) Show that a bounded diagonal operator  $De_n = d_n e_n$  on  $\ell^2$  (with  $\sup_n |d_n| < \infty$ ) has  $\|D\| = \sup_n |d_n|$ .

### Position, momentum, and the commutator

**Exercise 14.9** (proof) On  $L^2(\mathbb{R})$  prove both position and momentum are unbounded from normalized inputs.

1. For  $X\psi = x\psi$  and  $\psi_n = \mathbf{1}_{[n, n+1]}$ , prove  $\|\psi_n\| = 1$  and compute  $\|X\psi_n\|^2$ .
2. Fix a normalized  $\phi \in \mathcal{S}(\mathbb{R})$  and, for the wave number  $k \in \mathbb{R}$ , set  $\eta_k(x) = e^{ikx}\phi(x)$ . Prove  $\eta_k$  is normalized and in  $\mathcal{S}(\mathbb{R})$ , compute  $\hat{p}\eta_k$  for  $\hat{p} = -i\hbar d/dx$ , and use the reverse triangle inequality to conclude that  $\hat{p}$  is unbounded.

**Exercise 14.10** (proof) On the common invariant domain  $\mathcal{S}(\mathbb{R})$ , first explain why  $\hat{x}\hat{p}\psi$  and  $\hat{p}\hat{x}\psi$  are both defined. Then verify  $[\hat{x}, \hat{p}]\psi = i\hbar\psi$  for  $\hat{p} = -i\hbar d/dx$ . Finally prove that no finite matrices satisfy this exact CCR and state precisely what the trace obstruction does—and does not—show about quantum models.

*Domain convention for Exercises 14.11–14.16 and 14.18:* all displayed identities are equalities on  $\mathcal{S}(\mathbb{R})$ . This dense space is invariant under  $\hat{x}$ ,  $\hat{p}$ , and their polynomial products, so every composition used below is defined. These repeated calculations are extension practice rather than part of the selected core route.

**Exercise 14.11** (computational) Using  $[A, BC] = [A, B]C + B[A, C]$ , compute  $[\hat{x}, \hat{p}^2]$ .

**Exercise 14.12** (computational) Using  $[A, BC] = [A, B]C + B[A, C]$  and  $[\hat{x}, \hat{p}] = i\hbar$ , compute  $[\hat{p}, \hat{x}^2]$ .

**Exercise 14.13** (computational) Compute  $[\hat{x}, \hat{p}^3]$ .

**Exercise 14.14** (computational) Compute  $[\hat{x}^2, \hat{p}^2]$ , leaving the answer in terms of  $\hat{x}$  and  $\hat{p}$ .

**Exercise 14.15** (computational) In natural units ( $\hbar = 1$ , so  $[\hat{x}, \hat{p}] = i$ ) set  $a = \frac{1}{\sqrt{2}}(\hat{x} + i\hat{p})$  and  $a^\dagger = \frac{1}{\sqrt{2}}(\hat{x} - i\hat{p})$ . Compute  $[a, a^\dagger]$ .

**Exercise 14.16** (computational) With  $a, a^\dagger$  as above and the number operator  $N = a^\dagger a$ , use  $[a, a^\dagger] = \text{id}$  to compute  $[N, a]$ .

**Exercise 14.17** (conceptual) (*The boundary condition is part of the observable.*) Let  $T = -\frac{d^2}{dx^2}$  act on twice-differentiable functions on  $[0, 1]$  with the  $L^2$  inner product.

1. Integrating by parts twice, show that

$$\langle Tu, v \rangle - \langle u, Tv \rangle = \left[ \bar{u} v' - \overline{u'} v \right]_0^1.$$

2. Conclude that  $T$  is symmetric ( $\langle Tu, v \rangle = \langle u, Tv \rangle$ ) on the Dirichlet domain  $u(0) = u(1) = 0$ , and explain why the same operator with *no* boundary condition is not.
3. Explain why this boundary-form calculation proves symmetry but does not by itself prove self-adjointness; what further domain equality is needed?

**Exercise 14.18** (proof) Prove  $[\hat{x}, \hat{p}^n] = n i\hbar \hat{p}^{n-1}$  for all  $n \geq 1$  by induction.

## The spectral theorem and dynamics

**Exercise 14.19** (proof) State the compact self-adjoint spectral theorem without assuming separability, including the nonzero eigenvalues, their multiplicities, the kernel, and the finite-dimensional case. Then apply it to  $Te_n = n^{-1}e_n$  on  $\ell^2$ : write  $Tv$  and  $\|Tv\|^2$  for  $v = \sum_n c_n e_n$  and classify 0.

**Bridge practice: finite ideal measurement and dynamics.** Exercises 14.20–14.25 transfer the core spectral language to finite models; they do not count toward core progress.

**Exercise 14.20** (computational) For  $H = \frac{\hbar\omega}{2}\sigma_z$ , evolve  $|\psi\rangle(0) = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$  to time  $t$ .

**Exercise 14.21** (computational) Measuring  $\sigma_x$  on  $|0\rangle$ : give the outcome probabilities and  $\langle\sigma_x\rangle$ .

**Exercise 14.22** (computational) Let  $Te_n = \frac{1}{n}e_n$  on  $\ell^2$  (compact, self-adjoint) and  $v = 3e_1 + 4e_2$ . Compute  $Tv$ ,  $\|v\|^2$ ,  $\|Tv\|^2$ ,  $\langle v, Tv \rangle$ , and the Rayleigh quotient  $\langle v, Tv \rangle / \|v\|^2$ .

**Exercise 14.23** (computational) A three-level system has  $H|n\rangle = n\varepsilon|n\rangle$  for  $n = 0, 1, 2$ . Evolve  $|\psi\rangle(0) = \frac{1}{\sqrt{3}}(|0\rangle + |1\rangle + |2\rangle)$  to time  $t$ .

**Exercise 14.24** (computational) Measuring  $\sigma_z$  in  $|\psi\rangle = \frac{1}{2}(\sqrt{3}|0\rangle + |1\rangle)$ : give the outcome probabilities and  $\langle\sigma_z\rangle$ .

**Exercise 14.25** (computational) Using the spectral decomposition  $\sigma_z = |0\rangle\langle 0| - |1\rangle\langle 1|$ , compute  $e^{i\alpha\sigma_z}$  for real  $\alpha$ .

**Exercise 14.26** (proof) Let  $A$  be a bounded self-adjoint operator,  $\psi$  normalized, and  $\rho = |\psi\rangle\langle\psi|$ .

1. Prove  $\text{tr}(\rho A) = \langle\psi, A\psi\rangle$ .
2. If  $P$  is the PVM of  $A$ , state the probability of a spectral region  $\Omega$  and the ideal Lüders post-measurement state, including the nonzero-probability condition.
3. State the scope of this simple quantum dictionary and name the general replacements for states, measurements, and open dynamics.

**Exercise 14.27** (proof) Let  $A$  be a possibly unbounded self-adjoint operator with PVM  $P$  and  $\mu_\psi(\Omega) = \langle\psi, P(\Omega)\psi\rangle$ . State and explain:

1. the domain  $D(A)$ ;
2. the difference between bounded Borel  $f(A)$  and unbounded  $f(A)$ ;
3. when  $(A - zI)^{-1}$  is the bounded resolvent;
4. the continuity hypothesis in Stone's theorem and the domain of its generator derivative.