

Lecture 5

Eigenvalues and Eigenvectors

Throughout, $T \in \mathcal{L}(V)$ is an operator and $E(\lambda, T) = \text{null}(T - \lambda I)$ its eigenspace. Tags mark the flavour of each problem; a ★ marks a harder one.

Finding eigenvalues and eigenvectors

Exercise 5.1 (computational) Find the eigenvalues and a basis of each eigenspace for $A = \begin{pmatrix} 4 & -5 \\ 2 & -3 \end{pmatrix}$.

Exercise 5.2 (computational) Compute the eigenvalues of the rotation matrix $\begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix}$ over $\mathbb{C}$. For which α are they real?

Exercise 5.3 (computational) Let $D \in \mathcal{L}(\mathcal{P}_2(\mathbb{R}))$ be differentiation. Find all eigenvalues and eigenvectors of D .

Exercise 5.4 (computational) Find the eigenvalues and a basis of each eigenspace for $A = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$.

Exercise 5.5 (computational) Find the eigenvalues and a basis of each eigenspace for $A = \begin{pmatrix} 0 & 1 \\ -6 & 5 \end{pmatrix}$.

Exercise 5.6 (computational) Working over $\mathbb{C}$, find the eigenvalues and a basis of each eigenspace for $A = \begin{pmatrix} 1 & -2 \\ 1 & -1 \end{pmatrix}$.

Exercise 5.7 (proof) Suppose T is invertible and λ is an eigenvalue of T . Prove that $\lambda \neq 0$ and that $1/\lambda$ is an eigenvalue of T^{-1} , with the same eigenvectors.

Exercise 5.8 (★) Let J be the $n \times n$ real matrix with every entry equal to 1. Find all eigenvalues of J , their algebraic and geometric multiplicities, and a basis of each eigenspace.

Structure

Exercise 5.9 (computational) Let $A = \begin{pmatrix} 1 & 2 \\ 0 & 3 \end{pmatrix}$. Find the eigenvalues of $A^2 - A$.

Exercise 5.10 (computational) Verify that $A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ is an involution and find $E(1, A)$ and $E(-1, A)$.

Exercise 5.11 (computational) The matrix $A = \begin{pmatrix} 2 & 0 \\ 1 & 3 \end{pmatrix}$ is invertible. Find the eigenvalues of A^{-1} together with a corresponding eigenvector for each.

Exercise 5.12 (proof) Suppose λ is an eigenvalue of T and $p \in \mathcal{P}(\mathbb{F})$. Prove that $p(\lambda)$ is an eigenvalue of $p(T)$, with the same eigenvector.

Exercise 5.13 (proof) Prove that T is invertible if and only if 0 is not an eigenvalue of T .

Exercise 5.14 (proof) Suppose $T^2 = I$ (an involution). Prove that every eigenvalue of T is ± 1 , and that $V = E(1, T) \oplus E(-1, T)$.

Exercise 5.15 (★) Let $S, T \in \mathcal{L}(V)$. Prove that ST and TS have the same eigenvalues.

Exercise 5.16 (proof) Call T *nilpotent* if $T^k = 0$ for some $k \geq 1$. Prove that the only eigenvalue of a nilpotent operator is 0.

Upper-triangular form

Exercise 5.17 (computational) Read off the eigenvalues of $A = \begin{pmatrix} 5 & 2 & 7 \\ 0 & 5 & -1 \\ 0 & 0 & 3 \end{pmatrix}$ and state its trace and determinant. Is A invertible?

Exercise 5.18 (computational) Read off the eigenvalues of $A = \begin{pmatrix} 2 & 3 & 1 \\ 0 & -1 & 4 \\ 0 & 0 & 5 \end{pmatrix}$ and state $\text{tr } A$ and $\det A$. Is A invertible?

Exercise 5.19 (computational) Read off the eigenvalues of the lower-triangular matrix $A = \begin{pmatrix} 3 & 0 & 0 & 0 \\ 7 & 1 & 0 & 0 \\ 2 & 5 & -2 & 0 \\ 6 & 1 & 4 & 2 \end{pmatrix}$ and state $\text{tr } A$ and $\det A$.

Exercise 5.20 (computational) Read off the eigenvalues of $A = \begin{pmatrix} 4 & 1 & 7 \\ 0 & 0 & 2 \\ 0 & 0 & 3 \end{pmatrix}$ and state $\text{tr } A$ and $\det A$. Is A invertible?

Exercise 5.21 (proof) Suppose T has an upper-triangular matrix with diagonal entries $\lambda_1, \dots, \lambda_n$ with respect to a basis $v_1, \dots, v_n$. Prove that $(T - \lambda_1 I) \cdots (T - \lambda_n I) = 0$.

Exercise 5.22 (★) Prove that if $T \in \mathcal{L}(V)$ has $\dim V$ distinct eigenvalues, then T is diagonalizable (its eigenvectors form a basis).

Exercise 5.23 (computational) For $A = \begin{pmatrix} 3 & 1 \\ 0 & 3 \end{pmatrix}$, find the eigenspace $E(3, A)$ and the generalized eigenspace $G(3, A)$, and exhibit a Jordan chain.

Exercise 5.24 (proof) Let $T \in \mathcal{L}(V)$ and $c \in \mathbb{F}$. Prove that T and $T + cI$ have exactly the same eigenvectors, and that λ is an eigenvalue of T if and only if $\lambda + c$ is an eigenvalue of $T + cI$.