

Lecture 9

Adjoint; Self-Adjoint, Unitary, and Normal Operators

Inner products are in the physics convention (conjugate-linear in the first slot); the adjoint satisfies $\langle T^\dagger u, v \rangle = \langle u, Tv \rangle$. Tags mark the flavour; a ★ marks a harder one.

Adjoint

Exercise 9.1 (computational) Find the adjoint of $A = \begin{pmatrix} 2 & 1+i \\ 3 & -i \end{pmatrix}$, and state whether A is self-adjoint, unitary, or normal.

Exercise 9.2 (computational) On $\mathbb{R}^2$ with the standard inner product, find the adjoint of $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$, and say whether A is self-adjoint.

Exercise 9.3 (computational) Find the adjoint of $A = \begin{pmatrix} i & 2 \\ 1-i & 3i \end{pmatrix}$ on $\mathbb{C}^2$.

Exercise 9.4 (computational) Let $T(z_1, z_2) = (2z_1 + iz_2, z_1 - z_2)$ on $\mathbb{C}^2$. Find $T^\dagger$ as a formula.

Exercise 9.5 (proof) Prove that $(ST)^\dagger = T^\dagger S^\dagger$ for all $S, T \in \mathcal{L}(V)$.

Exercise 9.6 (proof) Prove that $\text{null } T^\dagger = (\text{range } T)^\perp$.

Exercise 9.7 (proof) Prove that for every $T \in \mathcal{L}(V)$ both $T + T^\dagger$ and $T^\dagger T$ are self-adjoint.

Self-adjoint operators

Exercise 9.8 (computational) Verify that $A = \begin{pmatrix} 3 & 2-i \\ 2+i & -1 \end{pmatrix}$ is Hermitian, and explain why its eigenvalues must be real.

Exercise 9.9 (computational) Verify that $A = \begin{pmatrix} 4 & 1 \\ 1 & 4 \end{pmatrix}$ is self-adjoint on $\mathbb{R}^2$, and find its eigenvalues and an orthonormal eigenbasis.

Exercise 9.10 (computational) Check that $A = \begin{pmatrix} 3 & i \\ -i & 3 \end{pmatrix}$ is Hermitian and find its (real) eigenvalues and corresponding eigenvectors.

Exercise 9.11 (computational) Find all $a \in \mathbb{C}$ for which $A = \begin{pmatrix} 5 & a \\ 3-2i & -1 \end{pmatrix}$ is self-adjoint.

Exercise 9.12 (proof) Prove that the eigenvalues of a self-adjoint operator are real, directly from $Tv = \lambda v$.

Exercise 9.13 (proof) Suppose T is self-adjoint and $Tu = \alpha u$, $Tv = \beta v$ with $\alpha \neq \beta$. Prove $\langle u, v \rangle = 0$.

Exercise 9.14 (proof) Suppose T is self-adjoint and $T^2 = 0$. Prove that $T = 0$.

Unitary and normal operators

Exercise 9.15 (computational) Show that $U = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & i \\ i & 1 \end{pmatrix}$ is unitary and find $|\lambda|$ for its eigenvalues.

Exercise 9.16 (proof) Prove that a unitary operator preserves the inner product: $\langle Uu, Uv \rangle = \langle u, v \rangle$.

Exercise 9.17 (computational) Let $S(w, z) = (-z, w)$ on $\mathbb{C}^2$. Find $S^\dagger$, show S is normal, and find its eigenvalues.

Exercise 9.18 (computational) Show that $Q = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$ is unitary, and state $|\lambda|$ for its eigenvalues.

Exercise 9.19 (computational) Determine whether $A = \begin{pmatrix} 2 & -3 \\ 3 & 2 \end{pmatrix}$ is normal, self-adjoint, or unitary, and find its eigenvalues.

Exercise 9.20 (computational) Classify $D = \text{diag}(i, -i, 1)$ on $\mathbb{C}^3$ as self-adjoint, unitary, and/or normal.

Exercise 9.21 (computational) Show that $A = \begin{pmatrix} 1 & i \\ i & 1 \end{pmatrix}$ is normal, and find its eigenvalues. Is it self-adjoint or unitary?

Exercise 9.22 (★) Prove that T is normal if and only if $\|Tv\| = \|T^\dagger v\|$ for all v .

Exercise 9.23 (★) Suppose A, B are self-adjoint. Prove AB is self-adjoint if and only if $AB = BA$, and that $i(AB - BA)$ is always self-adjoint.

Exercise 9.24 (proof) An operator is *skew-adjoint* if $B^\dagger = -B$. Prove that every eigenvalue of a skew-adjoint operator is purely imaginary.