

LUIGI VANVITELLI UNIVERSITY OF
CAMPANIA

DOCTORAL THESIS

Manifestation of the cluster
structure of weakly bound light
nuclei in the direct nuclear
reactions

Author:

Bakytzhan URAZBEKOV

Supervisor:

Prof. Nunzio ITACO

Co-Supervisor:

Prof. Andrey DENIKIN

Department of Mathematics and Physics



Declaration of Authorship

I, Bakytzhan URAZBEKOV, declare that this thesis titled, “Manifestation of the cluster structure of weakly bound light nuclei in the direct nuclear reactions” and the work presented in it are my own. I confirm that:

- This work was done wholly or mainly while in candidature for a research degree at this University.
- Where any part of this thesis has previously been submitted for a degree or any other qualification at this University or any other institution, this has been clearly stated.
- Where I have consulted the published work of others, this is always clearly attributed.
- Where I have quoted from the work of others, the source is always given. With the exception of such quotations, this thesis is entirely my own work.
- I have acknowledged all main sources of help.
- Where the thesis is based on work done by myself jointly with others, I have made clear exactly what was done by others and what I have contributed myself.

Signed:

Date:

*“By the love of God, the nations are created, and you love them like yourself.
Love people as brothers, like freedom, then truth and life are for you. ”*

Abay Qunanbayev

LUGI VANVITELLI UNIVERSITY OF CAMPANIA

Abstract

Department of Mathematics and Physics

Doctor of Philosophy

Manifestation of the cluster structure of weakly bound light nuclei in the direct nuclear reactions

by Bakytzhan URAZBEKOV

In this work, the following light weakly bound atomic nuclei were studied: ^6He , ^6Li and ^9Be . A three-body model $\alpha + 2\text{N}$ for ^6He and ^6Li , and $2\alpha + \text{N}$ model for ^9Be are proposed. The wave function of the three-body system was obtained within the framework of variational calculations based on the Gaussian basis. The interaction potentials for three-body systems have a Pauli projector, which excludes forbidden states. An analytical expression is obtained for the density distribution function of nuclear matter using the three-body wave function. Calculated and given comparisons of root-mean-square matter radii.

The interactions of ^6He , ^6Li and ^9Be have been studied in detail with the simplest particles, such as α , d , ^3He . On the basis of the obtained calculations on the density distribution function of nuclear matter, the folding interaction potential was obtained, and shown their comparison with optical potentials with global optical parametrizations. The resulting folding interaction potential is used to calculate the differential cross section for elastic scattering. Using the Coupled Channels approach for ^9Be the contribution of channels is studied, and deformation parameters are derived. Comparisons of theoretical cross sections with experimental data in inelastic channels for nuclear reactions $\alpha + ^9\text{Be}$, $d + ^9\text{Be}$ are given. Within the framework of the Coupled Reaction Channels method, the cross sections for nuclear reactions of single-nucleon and cluster transfers were calculated taking into account the internal structure of atomic nuclei ^6He , ^6Li and ^9Be . Features of the internal structure of these nuclei in direct nuclear reactions are manifested in their mechanism of transfer of clusters and nucleons. In this regard, several mechanisms have been proposed for the transfer of clusters and nucleons in ^6He , ^6Li and ^9Be .

Acknowledgements

I believe that if it were not for these people, then the thesis in the form in which you are reading, would not have taken place.

I am very grateful to my scientific supervisors Prof. Nunzio Itaco and Prof. Andrey Denikin, who gave a great chance to implement my scientific ideas. Their suggestions, ideas and comments always gave me goals, raised my understanding of science higher and higher.

Undoubtedly, I think it is worthwhile to thank the staff of the Department of Physics and Astronomy and the CIRCE laboratory: Prof. D'Onofrio, Prof. Gialanella, Giuseppe Porzio, Liz, and all members.

In addition, I would like to express my gratitude to Prof. Vladimir Kukulin (Moscow State University), who gave advice on a scientific career, Prof Sayabek Sakhiyev (Abay State University), who gave not only scientific guides but also about life.

I also want to thank my dear friends Riccardo Mancino and Felice Pignatelli, without whom I could not just cope with simple life questions in Caserta.

Thanks to Prof. Ian Thompson (LLNL) for explaining the input file for the FRESCO code, Prof. Alexandr Volya (MSU), for providing the values of the spectroscopic amplitudes of alpha particles for p-shell nuclei.

I also dedicate my big thanks to my wife Mervet, who has always supported me at all stages of my scientific career, and my newborn daughter Dina, who has been a source of strength and positiveness.

Contents

Declaration of Authorship	i
Abstract	iii
Acknowledgements	iv
1 Introduction	1
2 The three body model	7
2.1 The basis function	7
2.2 Transformation of the basis function	9
2.3 Overlap matrix elements	10
2.3.1 By means of the exponential function expansion	11
2.3.2 By means of the projection into the \mathbf{T} matrix	12
2.4 Normalization and correlation density of the three-body wave function	13
2.5 The density distribution function of nuclear matter	14
3 Theoretical models describing nuclear reactions	18
3.1 Description of elastic scattering	18
3.2 The coupled channels method for inelastic scattering	20
3.3 The coupled-reaction-channels method for the transfer reactions	23
3.4 Distorted Wave Born Approximation	25
3.5 Spectroscopic Amplitudes within the Shell Model	27
3.6 Interaction potentials. The double folding model	30
4 Results and discussions	33
4.1 The three-body wave function	33
4.2 The density distribution function of nuclear matter	36
4.3 The $\alpha + {}^6\text{He}$ nuclear reaction	40
4.3.1 Elastic scattering	40
4.3.2 Elastic transfer	42
4.4 The $\alpha + {}^9\text{Be}$ nuclear reactions	44

4.4.1	Elastic scattering	44
4.4.2	Inelastic scattering	46
4.4.3	One nucleon transfer reactions	46
4.5	The d+ ⁹ Be nuclear reactions	48
4.5.1	The elastic channel	48
4.5.2	Inelastic scattering	49
4.5.3	One-nucleon transfer reactions	51
4.5.4	Cluster-transfer reaction	55
5	Conclusion	61
A	Some aspects from the quantum theory of angular momenta	62
B	Parameters of the three body wave function	65
B.1	Helium-6	65
B.2	Lithium-6	72
B.3	Berillium-9	80
	Bibliography	88

List of Figures

2.1	The schemes of Jacobi coordinate sets for the three body system.	8
2.2	The (k, pq) scheme of Jacobi coordinate sets for calculation of the density distribution function of nuclear matter in the framework of the three body system.	14
2.3	The (p, qk) scheme of Jacobi coordinate sets for calculation of the density distribution function of nuclear matter in the framework of the three body system.	15
3.1	An incoming plane wave scattering off a body making a distorted wave . .	19
3.2	The arrangement of radial vectors in the DWBA on example of the $A + b \rightarrow a + B$ nuclear reaction. Here $A = a + \nu$ and $B = b + \nu$	26
3.3	Radius vectors for the double folding model.	31
4.1	The correlation function $W(x, y)$ of the three-body wave function of the ground state of ${}^6\text{He}$ in the a) 3D-Plot and b) contour-plot.	35
4.2	The same caption as in the Fig.4.1, but for ${}^6\text{Li}$	35
4.3	The same caption as in the Fig.4.1, but for ${}^9\text{Be}$	36
4.4	The density distribution function of nuclear matter $\rho(R) = \rho_{\alpha}^{(k)} + \rho_N^{(p)} + \rho_N^{(q)}$ illustrated with the densities of the ${}^6\text{He}$ constituents.	36
4.5	The same caption as in Fig. 4.4, but for ${}^6\text{Li}$	37
4.6	The density distribution function of nuclear matter $\rho(R) = \rho_N^{(k)} + \rho_{\alpha}^{(p)} + \rho_{\alpha}^{(q)}$ illustrated with the densities of the ${}^9\text{Be}$ constituents.	37
4.7	The folding potential $V^{DF}(r)$ in the framework of the three-body model. The potential is for the interaction of the α particle with the constituents of ${}^6\text{He}$	40
4.8	a) The differential cross section of the elastic scattering of ${}^6\text{He}$ by α -particles at $E_{lab} = 151\text{MeV}$ with the different potentials: the double folding potential (blue solid), the WS2 (gray dashed) [15] and WS1 (gray dotted) [15]. b) The cross sections of the elastic transfer reaction in terms of the transfer mechanisms: elastic scattering (solid gray), simultaneous transfer of $2N$ (gray dashed), sequential transfer of $2N$ (dotted) and total coherent sum (solid blue). Experimental data taken from [15].	41

4.9	Schematic representation of the elastic transfer nuclear reaction, on the left - the simultaneous transfer of two nucleons, on the right - the sequential transfer of two nucleons.	43
4.10	Folding potential of α interaction with ${}^9\text{Be}$ nucleus with allowance for the Coulomb interaction.	44
4.11	Differential cross sections of elastic scattering ${}^9\text{Be}(\alpha, \alpha){}^9\text{Be}$ for different collision energies, compared to the experimental data in [17–19] : (1) 4.5, (2) 7.25, (3) 10.00, (4) 11.25, (5) 12.62, and (6) 26.00 MeV/nucleon.	45
4.12	Differential cross sections for $E_{lab} = 16.25$ MeV/nucleon: (a) inelastic scattering ${}^9\text{Be}(\alpha, \alpha){}^9\text{Be}^*$ with excitation of: (1) $5/2^-$ states, (2) $7/2^-$ states; (b) single nucleon transfer reactions: (1) ${}^9\text{Be}(\alpha, t){}^{10}\text{B}$ and (2) ${}^9\text{Be}(\alpha, {}^3\text{He}){}^{10}\text{Be}$. Experimental data are taken from [21–23]	47
4.13	The angular distribution of elastic scattering data of d from ${}^9\text{Be}$ at laboratory energy 19.5 MeV in comparison with theoretical calculations within the OM (solid curve).	49
4.14	The cross sections of inelastic scattering ${}^9\text{Be}(d, d){}^9\text{Be}^*$ ($E_{exc}=2.43$ MeV) at laboratory energies 19.5 MeV (full circle) and 35 MeV (full triangle). Theoretical curves are described in the text.	50
4.15	The target coupling schemes in the ${}^9\text{Be}(d, p){}^{10}\text{Be}$ (upper) and the ${}^9\text{Be}(d, t){}^8\text{Be}$ (lower) nuclear reactions. The bold two-headed arrows indicate $E\lambda$ transitions. The spin re-orientation effects are indicated as back-pointing arrows.	51
4.16	Differential cross sections for the ${}^9\text{Be}(d, p){}^{10}\text{Be}^*$ reactions at 19.5 MeV leading to different final states (labelled in the figure) in ${}^{10}\text{Be}$. The experimental data are shown in comparison with theoretical results obtained within the CRC method.	52
4.17	Differential cross sections for the ${}^9\text{Be}(d, t){}^8\text{Be}^*$ reactions at 19.5 and 35 MeV leading to different final states (labelled in the figure) in ${}^8\text{Be}$. The experimental data are shown in comparison with theoretical results obtained within the CRC method.	53
4.18	The scheme illustrates the reaction mechanisms taken into account in CRC calculations of the cross sections for ${}^9\text{Be}(d, \alpha){}^7\text{Li}$ reaction.	56
4.19	Differential cross sections for the ${}^9\text{Be}(d, \alpha){}^7\text{Li}$ reactions measured at 19.5 MeV and 35 MeV energy with the ${}^7\text{Li}$ observed in the ground or low-lying excited states in the exit channels.	57

- 4.20 The scheme illustrates the energy balance of the different intermediate stages for the two-step mechanisms of ${}^9\text{Be}(d, \alpha){}^7\text{Li}$ transfer reaction. The Q -values for the different intermediate channels are shown near the corresponding lines. The numbers near the arrows correspond to the spectroscopic amplitudes of the heaviest reaction participants. For example, spectroscopic amplitude for the ${}^9\text{Be} = {}^8\text{Li} + p$ configuration is equal $\mathcal{S} = -0.947$ 58
- 4.21 Contributions of the different mechanisms to the cross section of the ${}^9\text{Be}(d, \alpha){}^7\text{Li}_{g.s.}$ reaction. See Fig. 4.18 for explanation of the curve notations. 59

List of Tables

3.1	Parameters of the M3Y-Paris [4] NN-potential used in double-folding calculations.	32
4.1	Results of the search for the minimum energy of three-body systems found by means of the variational approach: ${}^6\text{He}$, ${}^6\text{Li}$ and ${}^9\text{Be}$. The obtained minimum energy E_{min} , used set of quantum numbers $\gamma \equiv \lambda L S$, the dimension D and the probability P of each γ are listed.	34
4.2	Comparisons of the binding energies obtained by means of variational approach BE_{min} with the experimental data BE_{exp} and the rms matter radii $\langle R_m^2 \rangle^{\frac{1}{2}}$ with other sources.	38
4.3	The optical parameters of the potentials (DF, WS1, WS2) used in calculations of differential cross section of ${}^6\text{He}$ by α -particle at $E_{lab} = 151$ MeV within the optical model.	42
4.4	Parameters of imaginary part of the optical potential for calculating elastic scattering cross section $\alpha + {}^9\text{Be}$. The a_w parameter for all energies is 0.97 fm	46
4.5	Parameters of the optical potentials for calculations using the distorted waves and channel coupling methods	48
4.6	Parameterized double-folding potentials of the $d+{}^9\text{Be}$ system used in the OM, CC and DWBA calculations.	49
4.7	Spectroscopic amplitudes used in CRC calculations for the Composite = Core + Nucleon system. The one-nucleon spectroscopic amplitudes have been calculated by means of the <i>ANTOINE</i> code [31]. The alpha spectroscopic amplitudes were taken from [33, 34].	54
B.1	The three-body wave function parameters found by means of variational approach for the ground state of ${}^6\text{He}$	65
B.2	The three-body wave function parameters found by means of variational approach for the ground state of ${}^6\text{Li}$	72
B.3	The three-body wave function parameters found by means of variational approach for the ground state of ${}^9\text{Be}$	80

List of Abbreviations

OM	optical model
DF	double folding
CC	coupled channel
CRC	coupled reaction channels
DWBA	distorted wave Born approximation
SA	spectroscopic amplitude
cm	center of mass

Physical Constants

Speed of Light $c_0 = 2.997\,924\,58 \times 10^8 \text{ m s}^{-1}$ (exact)

Constant Name *Symbol* = *ConstantValue* with units

List of Symbols

Symbol	Name	Unit
ω	angular frequency	rad

to my mom

whom we had immeasurable love of mother and child...

Chapter 1

Introduction

In recent years, the study of light weakly bound nuclei has not lost interest due to the successful development of experimental technology. In particular, it should be noted a significant achievement in the field of production of secondary radioactive beams. It is known that in light nuclei nucleons tend to group into clusters, the relative motion of which mainly determines the properties and characteristics of the being explored nuclei. The cluster structure of the ground and low-lying excited states of light exotic nuclei is one of the priority areas of both experimental and theoretical nuclear physics.

Exotic states in most cases have a rarefied structure and an increased size, which are reflected in rms radii. Such peculiar properties are manifested in the second excited state of the ^{12}C 7.65 MeV nucleus ($J^p = 0^+$), which in the framework of many well-known models has a cluster structure. A similar behavior is predicted for excited states in nuclei ^{11}B and ^{13}C , possibly also having Hoyle states. In the framework of the α -condensate theory [4, 5], the radius of the Hoyle state is approximately 1.4–1.7 times the radius of the ground state, and the existence of states with a radius close to the radius of uranium nuclei is predicted in $^{12,13}\text{C}$ and ^{11}B nuclei.

In the framework of the cluster model, atomic nuclei can be formed from simple particles, such as ions: deuterium-2, tritium-3, helium-3 alpha and others. As an example, ^6He and ^6Li nuclei can be used. Based on the successful application of theoretical research within the framework of the hyperspherical harmonic model [6], as well as correlation gausoids [7], it can be deduced that the ^6He and ^6Li nuclei are well described by three-body cluster models $\alpha + n + n$ and $\alpha + n + p$, respectively. The clustering effect is manifested in many cases in unstable nuclei located on the border of the stability of the nuclear map. But there are stable atomic nuclei that clearly show the cluster structure as ^7Li , ^9Be , ^{12}C , and ^{16}O . The ^9Be nucleus is of particular interest for research, since it is a stable, but at the same time weakly bound nucleus. For example, the binding energy of one neutron $\text{Sn} (^9\text{Be}) = 1.7 \text{ MeV}$ in ^9Be is less than the binding energy

of one neutron Sn (${}^6\text{He}$) = 1.9 MeV in the unstable ${}^6\text{He}$ nucleus. Moreover, the ${}^9\text{Be}$ nucleus has a Borromian structure in which each pair combination in the triple $\alpha + \alpha + \text{n}$ structure has no bound state.

The cluster configuration of the ${}^9\text{Be}$ nucleus is also of interest in nuclear technology. In particular, this nucleus plays an important role in thermonuclear fusion. It is known that the larger the proton number in the atomic nucleus of the wall material, the more the material mixture is formed. In [8] calculations show that with a low proton number the ${}^9\text{Be}$ nucleus is the most suitable material for the wall of thermonuclear devices. Moreover, if one take into account that the ${}^9\text{Be}$ nucleus is easily decomposed by electrons and gamma particles into two fast alpha particles, then their kinetic energy contributes to the burning of fuel in the active zone.

It is widely acknowledged that ${}^8\text{Be}$ and ${}^5\text{He}$ nuclei do not exist in nature. The lifetime of these nuclei is very short - approximately 10-20 seconds. However, the formation of new elements through these unstable nuclei exists. In particular, the formation of the ${}^9\text{Be}$ nucleus based on ${}^8\text{Be}$ and ${}^5\text{He}$ nuclei is strongly suppressed. However, in [9] it was shown that under certain circumstances this synthesis is possible. It is interesting to note the properties of neutron-rich beryllium isotopes. The addition of three neutrons to ${}^9\text{Be}$ leads to the filling of the neutron p-shell. Along with other beryllium isotopes, the ${}^{12}\text{Be}$ nucleus has a sufficient lifetime (21.3 ms) for registration and a binding energy of 3.17 MeV greater than the binding energy of the stable beryllium isotope. Therefore, the ${}^{12}\text{Be}$ core is a remarkable subject of study within the framework of both the shell and cluster models. The study of the structure of radioactive atomic nuclei within the framework of the shell model is interesting because it allows one to take into account all degrees of freedom of nucleons, use realistic effective NN interactions and take into account the contribution of three-nucleon forces when describing the structure of the nucleus [10]. The model describes well the ground states p, sd shell nuclei, but not resonant excitations in the continuous spectrum. To study the state of the continuum, it is necessary to use theoretical approaches that correctly take into account the asymptotic behavior of wave functions at large distances. Beryllium isotopes exhibit an exceptional α -cluster structure — the ground and excited states form molecular structures with two α -particles bound by additional neutrons. The configuration of this kind of light nuclei is manifested not only in the structure, but also in the mechanism of interaction with other nuclei. To a large extent, this is observed in the cross sections of the channels of the fusion and transfer reactions. It is also interesting to note that the clustering effect in light nuclei

can manifest themselves during the interaction by means of exotic systems, such as $2n$, $8Be$, $5He$, etc. For example, $8Be$ and $5He$ nuclei are associated with a breakup as an intermediate channel for a stable 9Be nucleus. In this case, it is important to note the works [9, 11], which shows the path of the breakup of this nucleus. In this experimental work, it was proved that low-lying states of the 9Be nucleus up to an energy of 4 MeV have an $n + 8Be$ configuration, and high-lying excited states from 4 MeV have a $5He + \alpha$ structure. In addition to studying the cluster structure in breakup reactions, it is interesting to note the interaction of deuterons with clustered nuclei [12]. In [12] conducted experiments to study the interaction of deuterons with 9Be nuclei. In particular, the nuclear reaction $^9Be(d,a)7Li$ should be noted. For this nuclear reaction, different reaction mechanisms were evaluated: direct deuteron transfer, contribution of evaporation residues, and transfer of the $5He$ heavy cluster. The contribution of the mechanism through the compound nucleus, calculated in the framework of the statistical method, has a small contribution for this reaction at a laboratory energy of 7 MeV. It turned out that a large contribution to the cross section for the (d,a) reaction on the 9Be nucleus is mainly due to the mechanism of $5He$ cluster transfer at backward scattering angles. However, [13] concluded that the nuclear reaction $^9Be(d, a)7Li$ goes through the mechanism the compound nucleus. One of the main arguments in favor of this mechanism is that the energy of the nuclear reaction at 12 and 14 MeV lies precisely in the range of giant resonance in the excitation spectrum of the ^{11}B compound nucleus. In addition, in the work [12], theoretical calculations for the cross section (d, a) of the reaction underestimate the experimental data at the front scattering angles, which suggests the presence of two more direct mechanisms: sequential transfer of the $n + p$ system. Nevertheless, one more experiment should be carried out, and the data should be analyzed in a different energy range. It is assumed that on the basis of a new set of experiments in the framework of the proposed Project, it will be possible to answer the question which mechanism has the greatest contribution to the reaction cross section $^9Be(d,a)7Li$. In Ref. [14 - 16] theoretically studied the reactions of elastic and inelastic scattering, as well as single-nucleon transfers in the interactions of d and $3,4He$ and with the 9Be nucleus. The interaction potential of light d and $3,4He$ particles with 9Be was calculated in the framework of the folding model using the wave function of the 9Be ground state in the three-particle $2\alpha + n$ approximation. Within the framework of the coupled channel method and the distorted wave Born approximatton method, the differential cross sections of inelastic scattering and the reactions of single nucleon transfer were calculated using the folding potential.

The obtained angular distributions are in good agreement with experimental data. Thus, it was concluded that the three-cluster approximation ${}^9\text{Be} = 2\alpha + n$ constructed on the basis of the multicluster dynamic model taking into account the Pauli principle, provides an adequate description of the internal structure and properties of the ${}^9\text{Be}$ nucleus. In particular, interesting information about the properties of exotic nuclei is manifestation of their cluster structure, which can also be obtained from experiments on elastic scattering. The study of exotic states in light nuclei is a priority in the development of nuclear physics in recent decades. Exotic conditions in most cases have a rarefied structure and increased size. Of particular interest is the second excited ${}^{12}\text{C}$ state, the Hoyle state, which in many models has a cluster structure and increased dimensions [17, 18]. A similar increase is predicted for states in nuclei ${}^{11}\text{B}$ and ${}^{13}\text{C}$ [19, 20], possible analogues of the Hoyle state. As mentioned above, in the framework of the α -condensate theory, the radius of the Hoyle state is 1.4–1.7 times the radius of the ground state and the existence of states which has a radius close to the radius of the uranium nucleus are predicted in the ${}^{12}\text{C}$ and ${}^{11}\text{B}$ nuclei. Hoyle's state plays an important astrophysical role in the synthesis of ${}^{12}\text{C}$ in the Universe. The formation of elements heavier than carbon goes through this state. If it would not exist, the rate of carbon formation reaction was 7 orders of magnitude lower. This state is interesting from the hypothesis describing it as a gas of interacting alpha particles, which can be represented as a Bose condensate. Despite numerous studies, the properties of this state, lying above the threshold of decay into three alpha particles, are still poorly understood. To study the theory of nuclear reactions and its use to describe experimental data a method the Distorted Wave Born Approximation (DWBA) has been successfully applied. The distorted wave method is a good tool for describing nuclear reactions with products registered in the ground state. For transfer channels with excited states, calculations using the DWBA method in many cases give disagreement with the experimental data. In this case, the disagreement is explained by the fact that the wave function of the bound state for the output channel corresponds to a discrete eigenvalue, while the state of this excited nucleus in the experimental data has a resonance with a certain width in the order of several MeV. An adequate theoretical analysis of such problems requires other approaches and methods that require taking into account the degrees of freedom associated with the dynamics of the movement of three clusters and its influence on the mechanisms of nuclear reactions. The consideration of the continuum in the processes of nuclear reactions is used in the Continuum Discretized Coupled Channels (CDCC) approach [21]. According to the name

of the method, the essence of the method consists in discretizing the continuum and in coupling the reaction channels. As in the DWBA method, CDCC requires data which take into account the internal structure of colliding nuclei. The structure is determined by calculations of spectroscopic amplitudes, which strongly depend on the theoretical models under consideration. To date, a well-developed approach for studying the internal structure of the light nuclei of the ground state is ab initio. The ab initio method, which uses realistic NN forces, differs from the shell model in independence from model spaces. There is another method developed on the basis of the CDCC method, this is XCDCC [22]. The addition to the name “X” means taking into account the excited state of the target nucleus. Recently, a continuum discretization method was proposed based on imaginary correlated gaussian functions [23]. This approach successfully describes the displacement phases of elastic scattering of the p + system and reproduces well the real Coulomb functions. The aim of this project are: an adequate description of the properties and characteristics of ${}^9\text{Be}$ and ${}^{12}\text{C}$ nuclei using the above methods, obtaining new experimental data on the scattering of light particles by ${}^9\text{Be}$ and ${}^{12}\text{C}$ nuclei, extraction of experimental differential cross sections for elastic, inelastic, single-nucleon and cluster transmission channels. It is expected that the features of the experimental methodology, the latest equipment and modern theoretical approaches will provide an opportunity to better understand the properties and characteristics of the studied nuclei. Research on this topic is one of the rapidly developing areas of modern nuclear physics of all major scientific centers of the world. Kazakhstan scientists also conduct intensive research in these areas, and work closely with renowned scientists, major research centers, and the results are quite competitive internationally. This is evidenced by the scientific publications of Project managers in international journals with a high impact factor, and thereby show the high importance of this project. New data on the cross sections of nuclear reactions, reliable parameters of the optical potential, and spectroscopic characteristics of the excited states of ${}^5\text{He}$, ${}^8\text{Be}$, ${}^{10,11,12}\text{B}$, and ${}^{12}\text{C}$ nuclei will be useful for testing various cluster nuclear models and for carrying out model calculations of nucleosynthesis reactions for astrophysical and thermonuclear applications. The project is expected to obtain a number of new and relevant results. The new experimental technique will allow, firstly, to conduct experiments at high energies, and secondly, to experimentally measure cross sections at large scattering angles. In the first case, it is important to note that in the scientific world there are still no enough data on the scattering of ${}^3\text{He}$ and α particles by ${}^9\text{Be}$ nuclei above energies of 35 MeV. It is very important to conduct an

experiment at such energies, since they make it possible to measure the cross sections of relatively heavy reaction products, which reflect direct evidence of cluster transfer. In the second case, modern devices provide a great opportunity to extract information from nuclear reactions at the backward scattering angles. The available information from world-famous databases on the cross section for the ${}^3\text{He} + {}^9\text{Be}$ reactions for today, unfortunately, has limitations due to poor resolution of detectors. Therefore, the realization of this project, which is not only of academic interest, but also of great practical importance, is highly desirable.

Chapter 2

The three body model

The wave function for the three-body system in the current work was taken from [1]. The work [1] great progress has been made in understanding the structure and properties of the ${}^6\text{He}$, ${}^6\text{Li}$, ${}^9\text{Be}$ nuclei based on a three-particle model. It should be noted the progress achieved as a result of the study of the three-body structure, reproduced by the three-body wave function:

- structure and static characteristics of the ground state;
- low-lying excited states and geometric structure;
- processes of ${}^6\text{Li}$ interaction with hadrons, including quasi-elastic scattering (α , 2α), (p, d) and ($p, p\alpha$), high energy proton scattering, and the lightest nuclei on ${}^6\text{Li}$
- processes of absorption and scattering of π^\pm -mesons by ${}^6\text{Li}$
- electromagnetic structure;
- β -decay of ${}^6\text{He}$ and μ -capture by ${}^6\text{Li}$ and *etc.*

2.1 The basis function

The model [1] uses three pair pseudo-potentials to describe the three-body system, taking into account forbidden states by the Pauli principle:

$$\tilde{V}_{ij} = V_{ij} + \Delta_{ij}, \quad (2.1)$$

where V_{ij} is an interaction potential of (ij) subsystem and $\Delta_{ij} = \lambda \Gamma$ is a orthogonalizer with the λ constant and with Γ projector, which for forbidden f state is as follows:

$$\Gamma = \Gamma(f) = \sum_{m_f} |\phi_{fm_f}(\mathbf{x})\rangle \langle \phi_{fm_f}(\mathbf{x}')| \delta(\mathbf{y} - \mathbf{y}'). \quad (2.2)$$

The principle plays a huge role in the structure of the nucleus, which does not allow overlapping of two constituent particles. Thus, the three body pseudo-Hamiltonian including kinetic energy and pseudo potentials looks like this

$$\tilde{H} = H_0 + \sum_{i < j} \tilde{V}_{ij}. \quad (2.3)$$

As an example, in this work the $2\alpha - n$ three body model of the ${}^9\text{Be}$ nucleus is taken. The three-body wave function of the ${}^9\text{Be}$ nucleus with total spin J and spin projection M_J is represented as

$$\Psi^{JM_J} = \sum_i C_i \psi_i^{JM_J} (k, pq). \quad (2.4)$$

For simplicity the Jacobi coordinates \mathbf{x}_k and \mathbf{y}_k are down, the symbols k, p and q comply with the cluster indices (see Fig. 2.1), and the combination of indices (k, pq) corresponds to a certain choice of Jacobi coordinates \mathbf{x}_k and \mathbf{y}_k of the three-body system, where \mathbf{x}_k is a vector of the relative distance between the pair of particles pq and k , and \mathbf{y}_k is the vector of the relative distance between the center of mass of the pair pq and the particle k . The coefficients C_i in Eq. (ref totwf) are the parameters of the wave function expansion and are found as a result of solving the generalized eigenvalue problem.

The explicit form of the basis functions $\psi_i^\gamma (k, pq)$ is chosen in the form of the multiplication of the spatial and spin wave functions:

$$\psi_i^{JM_J} (k, pq) = [\phi_i^\gamma (k, pq) \times \chi^S]_{JM_J}, \quad (2.5)$$

here the index γ includes the quantum numbers $L\lambda l$. The spatial part $\phi_i^\gamma (k, pq)$ of the wave function (2.5) is constructed using the Gaussian functions:

$$\phi_i^\gamma (k, pq) = x_k^\lambda y_k^l \exp \left(-\frac{1}{2} \alpha_i^{(k)} x_k^2 - \frac{1}{2} \beta_i^{(k)} y_k^2 \right) [Y_\lambda (\hat{\mathbf{x}}_k) \times Y_l (\hat{\mathbf{y}}_k)]_{LM_L}, \quad (2.6)$$




FIGURE 2.1: The schemes of Jacobi coordinate sets for the three body system.

where L and M_L are the total orbital momentum of the system and its projection, λ, l are the orbital moments conjugated to the coordinates \mathbf{x}_k and \mathbf{y}_k respectively, $\alpha_i^{(k)}, \beta_i^{(k)}$ are the linear parameters of the three-body wave function.

2.2 Transformation of the basis function

The chosen form of the basis function (2.5) is convenient in that it can be easily transformed for use with an alternative set of Jacobi coordinates. A rotation matrix connecting different sets of the Jacobi coordinates is provided through

$$\begin{pmatrix} \mathbf{x}_k \\ \mathbf{y}_k \end{pmatrix} = \mathbf{T}^{(kq)} \begin{pmatrix} \mathbf{x}_q \\ \mathbf{y}_q \end{pmatrix}. \quad (2.7)$$

here, particularly for the (kq) transition, the $\mathbf{T}^{(kq)}$ matrix is

$$\mathbf{T}^{(kp)} = \begin{pmatrix} \mathbf{T}_{11}^{(kq)} & \mathbf{T}_{12}^{(kq)} \\ \mathbf{T}_{21}^{(kq)} & \mathbf{T}_{22}^{(kq)} \end{pmatrix} = \begin{pmatrix} -\frac{m_p}{m_p+m_q} & 1 \\ -\frac{m_q(m_k+m_p+m_q)}{(m_p+m_q)(m_k+m_q)} & -\frac{m_k}{(m_k+m_q)} \end{pmatrix} \quad (2.8)$$

There is no need to consider the transformation of spin part of the basis function. The reason is that the S spin of the $2\alpha - n$ three-body system is defined only by the vacant nucleon, and it doesn't depend on the choice of the Jacobi coordinate. That is to say, the coupling of the vacant nucleon spin with the $\alpha - \alpha$ subsystem spin gives again the spin of the nucleon due to the α particles having a spin of zero.

The transformation of the space part of the wave function from the set (k, pq) to the set (q, kp) can be expressed in the following way

$$\phi_i^{\gamma}(k, pq) = \sum_{\tilde{\gamma}} A_{\tilde{\gamma}\gamma}(\mathbf{T}^{(kq)}) \phi_i^{\tilde{\gamma}}(q, kp), \quad (2.9)$$

where the sum is over quantum numbers of the $\tilde{\gamma}$ new set, and the new basis function is

$$\begin{aligned} \phi_i^{\tilde{\gamma}}(q, kp) = & x_q^{\tilde{\lambda}} y_q^{\tilde{l}} \exp \left(-\frac{1}{2} \alpha_i^{(q)} x_q^2 - \frac{1}{2} \beta_i^{(q)} y_q^2 - \rho_i^{(q)} \mathbf{x}_q \cdot \mathbf{y}_q \right) \times \\ & \times [Y_{\tilde{\lambda}}(\hat{\mathbf{x}}_q) \times Y_{\tilde{l}}(\hat{\mathbf{y}}_q)]_{LM_L}, \end{aligned} \quad (2.10)$$

where new parameters of the wave function are given by

$$\begin{pmatrix} \alpha_i^{(q)} & \rho_i^{(q)} \\ \rho_i^{(q)} & \beta_i^{(q)} \end{pmatrix} = (\mathbf{T}^{(kq)})^T \begin{pmatrix} \alpha_i^{(k)} & \rho_i^{(k)} \\ \rho_i^{(k)} & \beta_i^{(k)} \end{pmatrix} \mathbf{T}^{(kq)}. \quad (2.11)$$

Note, that for the (k, pq) coordinate set the radial wave function (2.6) does not include the scalar product $\mathbf{x}_k \cdot \mathbf{y}_k$, which means $\rho_i^{(k)} = 0$. Using Eq. (A.9) and (A.13) it is easy to get the coupling coefficient $A_{\tilde{\gamma}\gamma}(\mathbf{T}^{(kq)})$ from Eq. (2.9), which is defined as follow

$$A_{\tilde{\gamma}\gamma}(\mathbf{T}^{(kq)}) = \sum_{\lambda_1 \lambda_2 l_1 l_2} \left(\mathbf{T}_{11}^{(kq)} \right)^{\lambda_1} \left(\mathbf{T}_{12}^{(kq)} \right)^{\lambda_2} \left(\mathbf{T}_{21}^{(kq)} \right)^{l_1} \left(\mathbf{T}_{22}^{(kq)} \right)^{l_2} \times \\ \times E_{\tilde{\lambda}l}^{\lambda_1 \lambda_2 \lambda l_1 l_2 L} \mathcal{D}(\lambda, \lambda_1, \lambda_2) \mathcal{D}(l, l_1, l_2). \quad (2.12)$$

The sum in this equation is carried out in conditions: $\lambda_1 + \lambda_2 = \lambda$, $l_1 + l_2 = l$ and $\lambda + l = \tilde{\lambda} + \tilde{l}$.

2.3 Overlap matrix elements

Within the (k, pq) scheme an overlap matrix element, in particular for the space part $\phi_i^\gamma(k, pq)$ of the basis function, is expressed by

$$\langle \phi_i^\gamma(k, pq) | \phi_j^{\gamma'}(k, pq) \rangle = \int \int d\mathbf{x}_k d\mathbf{y}_k \phi_i^\gamma(k, pq) \left(\phi_j^{\gamma'}(k, pq) \right)^*. \quad (2.13)$$

Following the properties of spherical harmonics the latter six dimensional integral gets $\delta_{\gamma\gamma'}$. Then, it is reduced to analytical expression as

$$\mathcal{I} \left(\lambda, l, \alpha_{ij}^{(k)}, \beta_{ij}^{(k)} \right) = 2^{1+\lambda+l} \frac{\Gamma \left(\frac{3}{2} + \lambda \right) \Gamma \left(\frac{3}{2} + l \right)}{\left(\alpha_{ij}^{(k)} \right)^{\frac{3}{2}+\lambda} \left(\beta_{ij}^{(k)} \right)^{\frac{3}{2}+l}}. \quad (2.14)$$

with

$$\alpha_{ij}^{(k)} = \alpha_i^{(k)} + \alpha_j^{(k)} \quad \beta_{ij}^{(k)} = \beta_i^{(k)} + \beta_j^{(k)} \quad (2.15)$$

However, to calculate overlap matrix elements for arbitrary basis functions, for example

$$\langle \phi_i^{\tilde{\gamma}}(q, kp) | \phi_j^{\tilde{\gamma}'}(q, kp) \rangle = \int \int d\mathbf{x}_k d\mathbf{y}_k \phi_i^{\tilde{\gamma}}(q, kp) \left(\phi_j^{\tilde{\gamma}'}(q, kp) \right)^*, \quad (2.16)$$

one must handle with the scalar product $\exp(-\rho \mathbf{x} \cdot \mathbf{y})$ in Eq. (2.10), in which radial and angular parts are mixed. It makes a problem in integration procedure, consequently, mathematical techniques must be applied. There are two tricks to solve this problem. First is the solution of problem is in expansion of the exponential function into the partial waves. Another one is in projection of the

factor of scalar product into the rotation matrix \mathbf{T} . Both approaches give the same results in calculation of the matrix elements.

2.3.1 By means of the exponential function expansion

The expansion of exponential function is given by

$$\exp(-\rho \mathbf{x} \cdot \mathbf{y}) = 4\pi \sum_{\kappa} \sqrt{2\kappa + 1} \epsilon(\kappa, \rho) i_{\kappa}(|\rho|xy) Y_{00}^{(\kappa\kappa)}(\hat{\mathbf{x}}, \hat{\mathbf{y}}) \quad (2.17)$$

where $i_{\kappa}(x)$ – modified spherical Bessel function of the first kind, $\epsilon(\kappa, \rho) = (-1)^{\kappa}$ for $\rho \leq 0$, otherwise it equals to 1. Once radial part is separated, defining an integral

$$\int_0^\infty \int_0^\infty dx dy x^{2\lambda+\kappa+2} y^{2l+\kappa+2} \exp(-\alpha x^2 - \beta y^2) i_{\kappa}(|\rho|xy), \quad (2.18)$$

one can get its analytical form

$$\begin{aligned} \mathcal{I}(\lambda, l, n, \alpha, \beta, |\rho|) = & \sqrt{\frac{\pi}{8}} (2l)!! \Gamma(l + n + \frac{3}{2}) |\rho|^n \beta^{-l-n-\frac{3}{2}} \times \\ & \times \sum_{\kappa=0}^l \frac{\Gamma(\kappa + \lambda + n + \frac{3}{2})}{\kappa!(l-\kappa)!\Gamma(\kappa+n+\frac{3}{2})} \left(\frac{\rho^2}{2\beta}\right)^{\kappa} \left(\frac{\alpha}{2} - \frac{\rho^2}{2\beta}\right)^{-\kappa-\lambda-n-\frac{3}{2}}. \end{aligned} \quad (2.19)$$

The angular part is integrated all over angular variables, then, it can be expressed analytically in the following way

$$\int \int d\hat{\mathbf{x}} d\hat{\mathbf{y}} Y_{00}^{(\kappa\kappa)}(\hat{\mathbf{x}}, \hat{\mathbf{y}}) Y_{L'M'L'}^{(\lambda'l')}(\hat{\mathbf{x}}, \hat{\mathbf{y}}) \left(Y_{LM_L}^{(\lambda l)}(\hat{\mathbf{x}}, \hat{\mathbf{y}})\right)^* = E_{\lambda l}^{\kappa\kappa 0\lambda' l' LL} \delta_{LL'} \quad (2.20)$$

Using the property of the 9-j symbol, in which one of the numbers is zero, $E_{\lambda l}^{\kappa\kappa 0\lambda' l' LL}$ can be reduced as

$$E_{\lambda l}^{\kappa\kappa 0\lambda' l' LL} = U(\lambda L \kappa l'; l \lambda') \frac{\mathcal{C}(\lambda', \lambda, \kappa) \mathcal{C}(l', l, \kappa)}{\sqrt{(2L+1)(2\kappa+1)}}. \quad (2.21)$$

The Jacobian matrix $\mathbf{J}^{(kq)}$ for transformation from the $\mathbf{x}_k, \mathbf{y}_k$ coordinates to the $\mathbf{x}_q, \mathbf{y}_q$ coordinates gives the $\mathbf{T}^{(kq)}$ matrix

$$\mathbf{J}^{(kq)} = \begin{pmatrix} \frac{\partial \mathbf{x}_k(\mathbf{x}_q, \mathbf{y}_q)}{\partial \mathbf{x}_q} & \frac{\partial \mathbf{x}_k(\mathbf{x}_q, \mathbf{y}_q)}{\partial \mathbf{y}_q} \\ \frac{\partial \mathbf{y}_k(\mathbf{x}_q, \mathbf{y}_q)}{\partial \mathbf{x}_q} & \frac{\partial \mathbf{y}_k(\mathbf{x}_q, \mathbf{y}_q)}{\partial \mathbf{y}_q} \end{pmatrix} = \mathbf{T}^{(kq)}. \quad (2.22)$$

Accordingly, the determinant $|\mathbf{J}^{(kq)}|$ is a determinant of the $\mathbf{T}^{(kq)}$ matrix, which equals to 1:

$$|\mathbf{J}^{(kq)}| = |\mathbf{T}^{(kq)}| = 1 \quad (2.23)$$

Therefore, the integration variables in Eq. (2.16) can be changed without any factorization.

Lastly, an expression for the overlap matrix element of the (q, kp) coordinate sets (2.16) can be determined as follow

$$\begin{aligned} \langle \phi_i^{\tilde{\gamma}}(q, kp) | \phi_j^{\tilde{\gamma}'}(q, kp) \rangle &= \int \int d\mathbf{x}_k d\mathbf{y}_k \phi_i^{\tilde{\gamma}}(q, kp) \left(\phi_j^{\tilde{\gamma}'}(q, kp) \right)^* = \\ &= 4\pi \sum_{\tilde{\gamma}\tilde{\gamma}'} A_{\gamma\tilde{\gamma}}(T^{(kq)}) A_{\gamma\tilde{\gamma}'}(T^{(kq)}) \sum_{\kappa} \sqrt{2\kappa+1} \epsilon(\kappa, \rho) E_{\lambda l}^{\kappa\kappa 0\lambda' l' LL} \times \\ &\quad \times \mathcal{I} \left(\frac{\tilde{\lambda}+\tilde{\lambda}'-\kappa}{2}, \frac{\tilde{l}+\tilde{l}'-\kappa}{2}, \kappa, \alpha_{ij}^{(q)}, \beta_{ij}^{(q)}, |\rho_{ij}^{(q)}| \right). \end{aligned} \quad (2.24)$$

with $\rho_{ij}^{(q)} = \rho_i^{(q)} + \rho_j^{(q)}$.

2.3.2 By means of the projection into the T matrix

In this approach the rotation matrix \mathbf{Q} , projecting the scalar product in the radial part of the wave function, is implemented by

$$\mathbf{Q}_i^{(kq)} = \mathbf{T}^{(kq)} \times \begin{pmatrix} 1 & -\frac{\rho_i^{(q)}}{\alpha_i^{(q)}} \\ 0 & 1 \end{pmatrix}. \quad (2.25)$$

Consequently, the radial part of the wave function can be rewritten with no scalar product term as

$$\begin{aligned} \phi_i^{\tilde{\gamma}}(q, kp) &= x_q^{\tilde{\lambda}} y_q^{\tilde{l}} \exp \left(-\frac{1}{2} \alpha_i^{(q)} x_q^2 - \frac{1}{2} \left(\beta_i^{(q)} - \frac{(\rho_i^{(q)})^2}{\alpha_i^{(q)}} \right) y_q^2 \right) \times \\ &\quad \times [Y_{\tilde{\lambda}}(\hat{\mathbf{x}}_q) \times Y_{\tilde{l}}(\hat{\mathbf{y}}_q)]_{LM_L}. \end{aligned} \quad (2.26)$$

One can get easily the expression of the overlap matrix element for the (q, kp) coordinate sets (2.16) can be determined as follow

$$\begin{aligned} \langle \phi_i^{\tilde{\gamma}}(q, kp) | \phi_j^{\tilde{\gamma}'}(q, kp) \rangle &= \int \int d\mathbf{x}_k d\mathbf{y}_k \phi_i^{\tilde{\gamma}}(q, kp) \left(\phi_j^{\tilde{\gamma}'}(q, kp) \right)^* = \\ &= \sum_{\tilde{\gamma}\tilde{\gamma}'} A_{\gamma\tilde{\gamma}}(\mathbf{Q}_{ij}^{(kq)}) A_{\gamma\tilde{\gamma}'}(\mathbf{Q}_{ij}^{(kq)}) \mathcal{I} \left(\tilde{\lambda}, \tilde{l}, \alpha_{ij}^{(q)}, \left(\beta_{ij}^{(q)} - \frac{(\rho_{ij}^{(q)})^2}{\alpha_{ij}^{(q)}} \right) \right) \delta_{\tilde{\gamma}\tilde{\gamma}'}. \end{aligned} \quad (2.27)$$

Analogously, changing the integration variables is carried out with no factorizations due to the $|\mathbf{Q}_i^{(kq)}| = 1$. Notably, rotation matrix becomes depended on the ij indexes in this case.

2.4 Normalization and correlation density of the three-body wave function

Using the overlap matrix elements the normalization of the total three-body wave function is given by

$$\begin{aligned}\mathcal{N}^{(k)} &= \langle \Psi^{JM_J} | \Psi^{JM_J} \rangle = \sum_{\gamma} \mathcal{N}_{\gamma}^{(k)}, \\ \mathcal{N}_{\gamma}^{(k)} &= \sum_{ij} C_i C_j \mathcal{I} \left(\lambda, l, \alpha_{ij}^{(k)}, \beta_{ij}^{(k)} \right).\end{aligned}\quad (2.28)$$

For the alternative set of Jacobi coordinates the overlap of the total wave function is given by

$$\mathcal{N}^{(q)} = \langle \Psi^{JM_J} | \Psi^{JM_J} \rangle = \sum_{\gamma} \sum_{\gamma'} \mathcal{N}_{\gamma\gamma'}^{(q)} \quad (2.29)$$

It should be mentioned, that in the latter expression the basis functions is not orthogonal. Therefore, according to the Eq. (2.20) the sum is limited with the condition $\delta_{LL'}$ only. Using the analytical form of overlap matrix element (2.19), $\mathcal{N}_{\gamma\gamma'}^{(q)}$ can be given by

$$\begin{aligned}\mathcal{N}_{\gamma\gamma'}^{(q)} &= 4\pi \sum_{ij} C_i C_j \sum_{\tilde{\gamma}\tilde{\gamma}'} A_{\gamma\tilde{\gamma}} \left(\mathbf{T}^{(kq)} \right) A_{\gamma'\tilde{\gamma}'} \left(\mathbf{T}^{(kq)} \right) \times \\ &\quad \times \sum_{\kappa} \sqrt{2\kappa + 1} \epsilon(\kappa, \rho_{ij}^{(k)}) E_{\lambda l}^{\kappa\kappa 0\lambda' l' LL} \times \\ &\quad \times \mathcal{I} \left(\frac{\tilde{\lambda} + \tilde{\lambda}' - \kappa}{2}, \frac{\tilde{l} + \tilde{l}' - \kappa}{2}, \kappa, \alpha_{ij}^{(q)}, \beta_{ij}^{(q)}, |\rho_{ij}^{(q)}| \right),\end{aligned}\quad (2.30)$$

or using the Eq. (2.27)

$$\begin{aligned}\mathcal{N}_{\gamma\gamma'}^{(q)} &= \sum_{ij} C_i C_j \sum_{\tilde{\gamma}\tilde{\gamma}'} A_{\gamma\tilde{\gamma}} \left(\mathbf{Q}_{ij}^{(kq)} \right) A_{\gamma'\tilde{\gamma}'} \left(\mathbf{Q}_{ij}^{(kq)} \right) \times \\ &\quad \times \mathcal{I} \left(\tilde{\lambda}, \tilde{l}, \alpha_{ij}^{(q)}, \left(\beta_{ij}^{(q)} - \frac{(\rho_{ij}^{(q)})^2}{\alpha_{ij}^{(q)}} \right) \right) \delta_{\tilde{\gamma}\tilde{\gamma}'}.\end{aligned}\quad (2.31)$$

A correlation density function of the total wave function (2.4) can be expressed in the following way

$$\begin{aligned} W(x_k, y_k) &= \sum_{\gamma} W_{\gamma}(x_k, y_k) \\ W_{\gamma}(x_k, y_k) &= \sum_{ij} C_i C_j x_k^{2+2\lambda} y_k^{2+2l} \exp \left(-\frac{1}{2} \alpha_{ij}^{(k)} x_k^2 - \frac{1}{2} \beta_{ij}^{(k)} y_k^2 \right). \end{aligned} \quad (2.32)$$

2.5 The density distribution function of nuclear matter

An operator of matter density distribution of the three body system takes the sums of all three clusters and brings it from the center of system mass (see. Fig. 2.2, 2.3)

$$\rho(R) = \sum_{i=k,p,q} \rho_i^{(i)}(R), \quad (2.33)$$

where $0i$ stands for the α particle or the nucleon inside the three-body system.

In particular, the internal structure of the nucleon is treated as point like particle, while the α cluster is taken with its internal structure as follow

$$\rho_{\alpha}(r) = \rho_0 \exp(-\gamma_0 r^2) \quad (2.34)$$

where parameters $\gamma_0 = 0.7024$, $\rho_0 = 0.4229$ [2]. So corresponding matrix elements of the density function for both the nucleon and the α cluster, which are




FIGURE 2.2: The (k, pq) scheme of Jacobi coordinate sets for calculation of the density distribution function of nuclear matter in the framework of the three body system.




FIGURE 2.3: The (p, qk) scheme of Jacobi coordinate sets for calculation of the density distribution function of nuclear matter in the framework of the three body system.

spectators in the three-body system (k, pq), can be written as

$$\begin{aligned} \rho_N^{(k)}(R) &= \sum_{\gamma} \sum_{ij} \langle \varphi_i^{\gamma}(k, pq) | \delta(\mathbf{R} - y_0^{(k)} \mathbf{y}_k) | \varphi_j^{\gamma}(k, pq) \rangle \\ \rho_{\alpha}^{(k)}(R) &= \sum_{\gamma} \sum_{ij} \langle \varphi_i^{\gamma}(k, pq) | \rho_{\alpha}(\mathbf{r}) | \varphi_j^{\gamma}(k, pq) \rangle \end{aligned} \quad (2.35)$$

Here the \mathbf{r} radius is from Eq. 2.34, and it has the relation $\mathbf{r} = y_0^{(k)} \mathbf{y}_k + \mathbf{R}$.

Using the basis function in the (k, pq) coordinate set, one can express the density function of the spectator of the three body system (p, qk) by means of the equation (2.9) for transformation into the (p, qk) set as follows:

$$\begin{aligned} \rho_N^{(q)}(R) &= \sum_{\gamma\tilde{\gamma}} \sum_{ij} A_{\gamma\tilde{\gamma}} \left(\mathbf{Q}_{ij}^{(kq)} \right) A_{\gamma\tilde{\gamma}} \left(\mathbf{Q}_{ij}^{(kq)} \right) \langle \varphi^{\tilde{\gamma}}(q, kp) | \delta(\mathbf{R} - \mathbf{y}_q) | \varphi^{\tilde{\gamma}}(q, kp) \rangle \\ \rho_{\alpha}^{(q)}(R) &= \sum_{\gamma\tilde{\gamma}} \sum_{ij} A_{\gamma\tilde{\gamma}} \left(\mathbf{Q}_{ij}^{(kq)} \right) A_{\gamma\tilde{\gamma}} \left(\mathbf{Q}_{ij}^{(kq)} \right) \langle \varphi^{\tilde{\gamma}}(q, kp) | \rho_{\alpha}(\mathbf{r}) | \varphi^{\tilde{\gamma}}(q, kp) \rangle. \end{aligned}$$

Using the properties of the spherical functions, well known expression of the partial expansion of the exponent function (2.17) and pre-defined table integrals (2.19) and (2.14) above mentioned six and eight dimensional integrals may be expressed analytically. In particular, for the spectators in the (k, qp) set the

density functions are defined like

$$\begin{aligned}\rho_k^{(N)}(R) &= \sum_{\gamma} \sum_{ij} C_i C_j R^{2+2l} \mathcal{I} \left(2 + 2\lambda, \frac{1}{2} \alpha_{ij}^{(k)} \right) \exp \left(-\frac{1}{2} \beta_{ij}^{(k)} \left(y_0^{(k)} R \right)^2 \right) \\ \rho_k^{(\alpha)}(R) &= (4\pi)^2 \rho_0 \exp(-\gamma_0 R^2) \sum_{\gamma} \sum_{ij} \mathcal{I} \left(2 + 2\lambda, \frac{1}{2} \alpha_{ij}^{(k)} \right) \times \\ &\quad \times \mathcal{I} \left(l, 0, (\beta_{ij}^{(k)} + 2\gamma_0) \left(y_0^{(k)} \right)^2, 2\gamma_0 R \right).\end{aligned}\quad (2.36)$$

The density distribution function of the nuclear matter for the N spectators in the (p, qk) three-body system can be written as follow

$$\begin{aligned}\rho_q^{(N)}(R) &= \sum_{\gamma\tilde{\gamma}} \sum_{ij} C_i C_j A_{\gamma\tilde{\gamma}}^{(N)} \left(\mathbf{Q}_{ij}^{(kq)}, R \right) A_{\gamma\tilde{\gamma}}^{(N)} \left(\mathbf{Q}_{ij}^{(kq)}, R \right) \times \\ &\quad \times \exp \left(-\frac{1}{2} \left(\beta_{ij}^{(q)} - \frac{(\rho_{ij}^{(q)})^2}{\alpha_{ij}^{(q)}} \right) \left(y_0^{(q)} R \right)^2 \right)\end{aligned}\quad (2.37)$$

The coefficient of transformation of the basis function (2.12) now depends on R and it has form as

$$\begin{aligned}A_{\tilde{\gamma}\gamma}^{(N)} \left(\mathbf{Q}_{ij}^{(kq)}, R \right) &= \sum_{\lambda_1 \lambda_2 l_1 l_2} \left(\mathbf{Q}_{11}^{(kq)} \right)^{\lambda_1} \left(\mathbf{Q}_{12}^{(kq)} \right)^{\lambda_2} \left(\mathbf{Q}_{21}^{(kq)} \right)^{l_1} \left(\mathbf{Q}_{22}^{(kq)} \right)^{l_2} \times \\ &\quad \times E_{\tilde{\lambda}\tilde{l}}^{\lambda_1 \lambda_2 \lambda l_1 l_2 L} \mathcal{D}(\lambda, \lambda_1, \lambda_2) \mathcal{D}(l, l_1, l_2) \times \\ &\quad \times R^{\lambda + l - \ell/2} \mathcal{I} \left(\ell + 2, \frac{1}{2} \alpha_{ij}^{(q)} \right)^{1/2},\end{aligned}\quad (2.38)$$

where $\ell = \lambda_1 + \lambda_2 + l_1 + l_2$, and the sum is carried out in conditions: $\lambda_1 + \lambda_2 = \lambda$, $l_1 + l_2 = l$ and $\lambda + l = \tilde{\lambda} + \tilde{l}$.

The density distribution function of the nuclear matter for the α spectator in the (p, qk) scheme has the form

$$\begin{aligned}\rho_q^{(\alpha)}(R) &= \sum_{\gamma\tilde{\gamma}} \sum_{ij} C_i C_j A_{\gamma\tilde{\gamma}}^{(\alpha)} \left(\mathbf{Q}_{ij}^{(kq)}, R \right) A_{\gamma\tilde{\gamma}}^{(\alpha)} \left(\mathbf{Q}_{ij}^{(kq)}, R \right) \times \\ &\quad \times \exp \left(-\frac{1}{2} \gamma_0 R^2 \right)\end{aligned}\quad (2.39)$$

where its transformation coefficient also depends on R , and turns to

$$\begin{aligned}
 A_{\tilde{\gamma}\gamma}^{(N)} \left(\mathbf{Q}_{ij}^{(kq)}, R \right) = & \sum_{\lambda_1 \lambda_2 l_1 l_2} \left(\mathbf{Q}_{11}^{(kq)} \right)^{\lambda_1} \left(\mathbf{Q}_{12}^{(kq)} \right)^{\lambda_2} \left(\mathbf{Q}_{21}^{(kq)} \right)^{l_1} \left(\mathbf{Q}_{22}^{(kq)} \right)^{l_2} \times \\
 & \times E_{\tilde{\lambda} \tilde{l}}^{\lambda_1 \lambda_2 \lambda l_1 l_2 l L} \mathcal{D}(\lambda, \lambda_1, \lambda_2) \mathcal{D}(l, l_1, l_2) \times \\
 & \times \mathcal{I} \left(\lambda + l - \ell/2, 0, \left(\left(\beta_{ij}^{(q)} - \frac{\left(\rho_{ij}^{(q)} \right)^2}{\alpha_{ij}^{(q)}} \right) + 2\gamma_0 \right) y_0^{(q)2}, 2\gamma_0 R \right)^{1/2} \times \\
 & \times \mathcal{I} \left(\ell + 2, \frac{1}{2} \alpha_{ij}^{(q)} \right), \tag{2.40}
 \end{aligned}$$

where $\ell = \lambda_1 + \lambda_2 + l_1 + l_2$, and the sum is carried out in conditions: $\lambda_1 + \lambda_2 = \lambda$, $l_1 + l_2 = l$ and $\lambda + l = \tilde{\lambda} + \tilde{l}$.

Chapter 3

Theoretical models describing nuclear reactions

3.1 Description of elastic scattering

Descriptions of the scattering of two nuclei are considered in this chapter when the interaction between them is a potential U which may depend on the spins of the two nuclei but not on their internal coordinates. Thus, it cannot excite the nuclei internally or cause the transfer between them. It can only change their relative motion and, perhaps, reorient spins to each other or to the orbital motion. In general, U will be complex.

In the case of the colliding the $a + A$ particles without spin the potential $U(r)$ is central, depending only on the magnitude of the channel coordinate \mathbf{r} . The corresponding Shrödinger equation may be written explicitly

$$\left(E + \frac{\hbar^2}{2\mu} \nabla^2 - U(r) \right) \chi(\mathbf{r}) = 0 \quad (3.1)$$

where μ is the reduced mass of the $a+A$ system, E is the energy in the centre-of-mass system. The $\chi(\mathbf{r})$ wave function is known as distorted waves describing elastic scattering. The expression "distorted wave" is meant to denote distortion away from the plane wave form due to the presence of the distorting potential $U(r)$ (see Fig. 3.1). Asymptotically, it has the form of an incident plane wave plus outgoing (scattered) spherical waves

$$\chi^{(+)}(\mathbf{k}, \mathbf{r}) \rightarrow e^{i\mathbf{k}\cdot\mathbf{r}} + f(\theta) \frac{1}{\mathbf{r}} e^{ikr}, \quad \mathbf{r} \rightarrow \infty \quad (3.2)$$

where $f(\theta)$ is the scattering amplitude. The (+) superscript stands for outgoing plane wave, while incoming spherical waves is the time-reverse of the $\chi^{(+)}$

$$\chi^{(-)}(\mathbf{k}, \mathbf{r}) = (\chi^{(+)}(-\mathbf{k}, \mathbf{r}))^* \quad (3.3)$$




FIGURE 3.1: An incoming plane wave scattering off a body making a distorted wave

Using the partial wave expansion

$$\chi^{(+)}(\mathbf{k}, \mathbf{r}) = \frac{4\pi}{kr} \sum_{LM} i^L \chi_L(k, r) Y_{LM}(\hat{\mathbf{r}}) \left(Y_{LM}(\hat{\mathbf{k}})\right)^*, \quad (3.4)$$

A solution for Eq. (3.1) in the absence of any interaction potential, $U(r) = 0$, is given by

$$\chi^{(+)}(\mathbf{k}, \mathbf{r}) \rightarrow e^{i\mathbf{k}\cdot\mathbf{r}}, \quad \mathbf{r} \rightarrow \infty, \quad (3.5)$$

for radial part only is written as

$$\chi_L(k, r) \rightarrow kr j_L(kr), \quad r \rightarrow \infty, \quad (3.6)$$

where $j_L(kr)$ is the usual spherical Bessel function. The $\chi_L(k, r)$ radial function satisfies the radial Shrödinger equation

$$\left(\nabla_r + k^2 - \frac{L(L+1)}{r^2} - \frac{2\mu}{\hbar^2} U(r)\right) \chi_L(k, r) = 0, \quad (3.7)$$

where $k^2 = \frac{2\mu E}{\hbar^2}$.

In the case of the interaction potential has a more sophisticated form, including both Coulomb and nuclear short-range potentials, the radial Shrödinger equation (3.7) may be rewritten as

$$\left(\nabla_r + k^2 - \frac{2\eta k}{r} - \frac{L(L+1)}{r^2} - \frac{2\mu}{\hbar^2} U(r)\right) \chi_L(k, r) = 0, \quad (3.8)$$

where η is the usual Sommerfeld parameter with the Z charge numbers

$$\eta = \frac{Z_a Z_A e^2 \mu}{\hbar^2 k}$$

At the radius r_0 , where the nuclear potential is negligible, Eq. (3.8) has the solution, which can be expressed in terms of the H_L outgoing and H_L^* incoming Coulomb functions, as follow

$$\chi_L(k, r) \rightarrow \frac{i}{2} e^{i\sigma_L} (H_L(kr_0)^* - S_L H_L(kr_0)), \quad r \rightarrow r_0 \quad (3.9)$$

where σ_L is the Coulomb phase shift, and it is given with the Γ Gamma function as follow

$$\sigma_L = \text{Arg}(\Gamma(L + 1 + i\eta))$$

In practice, the radial Eq. (3.8) is solved by numerical integration from $r \approx 0$, then, matched the value and the slope of the result onto the form (3.9) at $r = r_0$. This procedure then gives a value for the S_L scattering matrix elements. Having the S_L matrix elements, the amplitude of the elastic scattering, analogous from Eq. (3.2), for the $\chi_L(k, r)$ wave function in the Eq. (3.8) is given by

$$f(\theta) = \frac{1}{2ik} \sum_L (2L + 1) e^{2i\sigma_L} (S_L - 1) P_L(\cos(\theta)) \quad (3.10)$$

where $P_L(x)$ is the Legendre polynomials, which are solutions to the Legendre differential equation. By means of the Rutherford scattering amplitude

$$f_C(\theta) = -\frac{\eta}{2k \sin^2(\frac{1}{2}\theta)} e^{(i\eta \ln(\sin^2(\frac{1}{2}\theta)) + 2i\sigma_0)} \quad (3.11)$$

the differential cross section of the elastic scattering has the form

$$\frac{d\sigma_\alpha}{d\Omega} = |f_C(\theta) + f(\theta)|^2. \quad (3.12)$$

The expression (3.12) allows thus to provide comparison with the obtained experimental data.

3.2 The coupled channels method for inelastic scattering

The elastic scattering of the projectile a by the nucleus A has been denoted by $\alpha = a + A$. Let $\alpha' = a + A^*$ be an inelastic channel, in which only the A

nucleus has an extra excitation. In the framework of the coupled channel (CC) approach the total wave function Ψ for the system may be written as

$$\Psi = \phi_\alpha(x) \chi_\alpha(\mathbf{r}_\alpha) + \phi_{\alpha'}(x) \chi_{\alpha'}(\mathbf{r}_{\alpha'}) \quad (3.13)$$

where \mathbf{r} is the channel coordinate for the partitions α or α' , x represents the corresponding internal coordinates. The total wave function can be part of the Shrödinger equation kind of

$$H\Psi = E\Psi \quad (3.14)$$

with the Hamiltonian appropriate particular for the α partition

$$H = H_\alpha + K_\alpha + V_\alpha. \quad (3.15)$$

where $H_\alpha \equiv H_a + H_A$ is the internal Hamiltonian for the nuclei a and A , K_α is the kinetic energy operator of relative motion and V_α is the interaction potential operator. The wave functions of ground $\phi_\alpha(x)$ and excited state $\phi_{\alpha'}(x)$, being eigenfunctions of the internal Hamiltonian H_α

$$\begin{aligned} H_\alpha \phi_\alpha(x) &= \varepsilon_\alpha \phi_\alpha(x) \\ H_\alpha \phi_{\alpha'}(x) &= \varepsilon_{\alpha'} \phi_{\alpha'}(x), \end{aligned} \quad (3.16)$$

have an orthonormality property of the form

$$\int dx (\phi_\alpha(x))^* \phi_{\alpha'}(x) = \delta_{\alpha\alpha'}. \quad (3.17)$$

Using the expression of total wave function (3.13), multiplying Eq. (3.14) from the left by the ϕ_α^* function, then, by the $\phi_{\alpha'}^*$ function, the two coupled equations can be defined in the following form

$$\begin{aligned} (E - \varepsilon_\alpha - K_\alpha - \langle \alpha | V_\alpha | \alpha \rangle) \chi_\alpha(\mathbf{r}) &= \langle \alpha | V_\alpha | \alpha' \rangle \chi_{\alpha'}(\mathbf{r}) \\ (E - \varepsilon_\alpha - K_\alpha - \langle \alpha' | V_\alpha | \alpha' \rangle) \chi_{\alpha'}(\mathbf{r}) &= \langle \alpha' | V_\alpha | \alpha \rangle \chi_\alpha(\mathbf{r}) \end{aligned} \quad (3.18)$$

where $\langle \alpha | V_\alpha | \alpha \rangle$, or $\langle \alpha' | V_\alpha | \alpha \rangle$, is the matrix element of V_α . In particular, the matrix element $\langle \alpha' | V_\alpha | \alpha \rangle$ is given by

$$\begin{aligned} \langle \alpha' | V_\alpha | \alpha \rangle &= \int dx \phi_{\alpha'}'(x) V_\alpha(x, \mathbf{r}) \phi_\alpha(x) = \\ &= V_{\alpha'\alpha}(\mathbf{r}). \end{aligned} \quad (3.19)$$

An expansion of the coupling potential $V_{\alpha'\alpha}(\mathbf{r})$ into the λ multipoles can be given as follow

$$V_{\alpha'\alpha}(\mathbf{r}) = \sum_{\lambda\mu} V_{\alpha'\alpha}^{\lambda\mu}(r) Y_{\lambda\mu}(\hat{r}) \quad (3.20)$$

If the potential shape has a deformation, the nuclear potential can be constructed as

$$V_\alpha(x, \mathbf{r}) \equiv U(r - \delta(\hat{r}')) \quad (3.21)$$

where \hat{r}' denotes angular coordinates of (θ, ϕ) referred to the intrinsic reference frame. The function $\delta(\hat{r}')$ is normally expanded in multipoles

$$\delta(\hat{r}') = \sum_{\lambda} \delta_{\lambda} Y_{\lambda 0}(\hat{r}'). \quad (3.22)$$

In the collective model, the ground and excited states are characterized by their angular momenta I_i and I_f with projections M_i and M_f , respectively. For these state the relevant matrix element of the $V_{\alpha'\alpha}^{\lambda\mu}$ operator is described using the Wigner-Eckart theorem by

$$\langle I_i M_i | V_{\alpha'\alpha}^{\lambda\mu} | I_f M_f \rangle = \sqrt{2I_i + 1} \langle I_f M_f \lambda \mu | I_i M_i \rangle \langle I' | V_{\alpha'\alpha}^{\lambda} | I \rangle. \quad (3.23)$$

Using the definitions, (3.21) and (3.22), the reduced matrix element of radial multipoles from Eq. (3.23) can be rewritten as follow

$$\langle I_i | V_{\alpha'\alpha}^{\lambda\mu} | I_f \rangle = -\frac{\langle I_i | \delta_{\lambda} | I_f \rangle}{\sqrt{4\pi}} \frac{dU(r)}{dr} \quad (3.24)$$

where

$$\langle I_i | \delta_{\lambda} | I_f \rangle = \sqrt{2I_f + 1} \langle I_f M_f \lambda 0 | I_i M_i \rangle \langle \chi | \delta | \chi \rangle \delta_{M_i M_f}. \quad (3.25)$$

The matrix element $\langle \chi | \delta | \chi \rangle$ is the expectation value of the operator δ_{λ} in the internal state of the deformed nucleus. In the framework of the rotational model it can be given with the deformation length β_{λ} as follow

$$\langle \chi | \delta | \chi \rangle = R_0 \beta_{\lambda} \quad (3.26)$$

where R_0 is an average radius of the interaction potential.

The general asymptotic behaviour of the χ_{α} elastic channel can be taken from Eq. (3.2), while the inelastic channel $\chi_{\alpha'}$ may have

$$\chi_{\alpha'}(\mathbf{r}_{\alpha'}) \rightarrow f_{\alpha'}(\theta) \frac{e^{ikr_{\alpha'}}}{r}, \quad \mathbf{r}_{\alpha'} \rightarrow \infty \quad (3.27)$$

Note, that the plane wave expression doesn't present in this equation - only outgoing wave presents. A relevant differential cross section for the inelastic channel is obtained from the coefficient of the outgoing wave as follow

$$\frac{d\sigma_{\alpha'}(\theta)}{d\Omega} = \frac{k_{\alpha'}}{k_{\alpha}} |f_{\alpha'}(\theta)|^2. \quad (3.28)$$

The wave numbers k_{α} and $k_{\alpha'}$ follow from the energy conservation

$$E = \varepsilon_{\alpha} + \frac{\hbar^2 k_{\alpha}}{2\mu} = \varepsilon_{\alpha'} + \frac{\hbar^2 k_{\alpha'}}{2\mu}. \quad (3.29)$$

3.3 The coupled-reaction-channels method for the transfer reactions

Consider a model for the $a + A \rightarrow b + B$ nuclear reaction, in which entrance and exit channels are denoted as α and β respectively. The Ψ total wave function for this model may be given as

$$\Psi = \chi_{\alpha}(\mathbf{r}_{\alpha}) \phi_{\alpha}(x_{\alpha}) + \chi_{\beta}(\mathbf{r}_{\beta}) \phi_{\beta}(x_{\beta}). \quad (3.30)$$

with a model Hamiltonian H such that $(E - H)\Psi = 0$. From the projections of this equation onto the two channels,

$$\begin{aligned} \langle \chi_{\alpha} | (E - H) | \Psi \rangle &= 0 \\ \langle \chi_{\beta} | (E - H) | \Psi \rangle &= 0 \end{aligned} \quad (3.31)$$

with the two equivalent forms of H ,

$$\begin{aligned} H &= H_{\alpha} + K_{\alpha} + V_{\alpha} \\ H &= H_{\beta} + K_{\beta} + V_{\beta} \end{aligned} \quad (3.32)$$

one can get a pair of coupled equations for χ_{α} and χ_{β} :

$$\begin{aligned} [(E - \varepsilon_{\alpha}) - K_{\alpha} - \langle \alpha | V_{\alpha} | \alpha \rangle] \chi_{\alpha}(\mathbf{r}_{\alpha}) &= \langle \alpha | H - E | \beta \rangle \chi_{\beta} \\ [(E - \varepsilon_{\beta}) - K_{\beta} - \langle \beta | V_{\beta} | \beta \rangle] \chi_{\beta}(\mathbf{r}_{\beta}) &= \langle \beta | H - E | \alpha \rangle \chi_{\alpha}. \end{aligned} \quad (3.33)$$

These are *the coupled-reaction-channels* (CRC) equations. They are integro-differential equations, as may be seen more explicitly in the form

$$\begin{aligned} [(E - \varepsilon_\alpha) - K_\alpha - \langle \alpha | V_\alpha | \alpha \rangle] \chi_\alpha(\mathbf{r}_\alpha) &= \int d\mathbf{r}_\beta K_{\alpha\beta}(\mathbf{r}_\alpha, \mathbf{r}_\beta) \chi_\beta(\mathbf{r}_\beta) \\ [(E - \varepsilon_\beta) - K_\beta - \langle \beta | V_\beta | \beta \rangle] \chi_\beta(\mathbf{r}_\beta) &= \int d\mathbf{r}_\alpha K_{\beta\alpha}(\mathbf{r}_\beta, \mathbf{r}_\alpha) \chi_\alpha(\mathbf{r}_\alpha) \end{aligned} \quad (3.34)$$

where the kernels are

$$\begin{aligned} K_{\alpha\beta}(\mathbf{r}_\alpha, \mathbf{r}_\beta) &= J_{\alpha\beta} \int d\zeta_\alpha \phi_\alpha^*(x_\alpha) (H - E) \phi_\beta(x_\beta) \\ K_{\beta\alpha}(\mathbf{r}_\beta, \mathbf{r}_\alpha) &= J_{\beta\alpha} \int d\zeta_\beta \phi_\beta^*(x_\beta) (H - E) \phi_\alpha(x_\alpha). \end{aligned} \quad (3.35)$$

Here the internal coordinates have been transformed from the set x_α to the set $(\zeta_\alpha, \mathbf{r}_\beta)$, where the ζ_α are independent of \mathbf{r}_β . Also, $J_{\alpha\beta}$ is the Jacobian of this transformation. Then $J_{\beta\alpha}$ is the Jacobian for the analogous transformation from x_β to $(\zeta_\beta, \mathbf{r}_\alpha)$.

Since the off-diagonal matrix elements of V_α are small, they have little effect on the elastic scattering and may be neglected on the right side. This implies that the elastic scattering in the entrance α channel is described well by the potential $\langle \alpha | V_\alpha | \alpha \rangle$, and the potential $\langle \beta | V_\beta | \beta \rangle$ describes well the elastic scattering in the β channel. By doing this approximation, Eq. (3.33) may be rewritten as

$$\begin{aligned} [(E - \varepsilon_\alpha) - K_\alpha - \langle \alpha | V_\alpha | \alpha \rangle] \chi_\alpha(\mathbf{r}_\alpha) &\approx 0 \\ [(E - \varepsilon_\beta) - K_\beta - \langle \beta | V_\beta | \beta \rangle] \chi_\beta(\mathbf{r}_\beta) &\approx \langle \beta | H - E | \alpha \rangle \chi_\alpha \\ &\approx \langle \beta | V_\alpha | \alpha \rangle \chi_\alpha + \langle \beta | \alpha \rangle (H_\alpha - E_\alpha) \chi_\alpha \\ &\approx \langle \beta | V_\alpha | \alpha \rangle \chi_\alpha \end{aligned} \quad (3.36)$$

Here the prior interaction, that is the interaction of the α channel, is used. The non-orthogonal term $\langle \beta | \alpha \rangle (H_\alpha - E_\alpha) \chi_\alpha$ vanishes because of the χ_α is on-shell. The usual Green function techniques may then be used to solve these equations and give *the distorted wave Born approximation transition* (DWBA) amplitude. A detailed representation of the amplitude will be done in the next section.

3.4 Distorted Wave Born Approximation

Let consider the nuclear reaction of transfer ν particle from a A nucleus into a B nucleus:

$$\begin{aligned} A + b &\rightarrow a + B \\ A = a + \nu, \quad B = b + \nu \end{aligned} \tag{3.37}$$

In the case of weak coupling between intermediate channels, it is reasonable to evaluate the transition amplitude in Born Approximation.

In the rearrangement reactions there are few ways to describe the interaction between the different fragments, one for each partition. For example, if we choose to describe the scattering in terms of the nuclei of the entrance partition, the projectile-target interaction will be written as

$$V_{Ab} = V_{\nu b} + U_{ab}. \tag{3.38}$$

Here, $V_{\nu b}$ – a binding potential of the ν valence particle with the b core, and it is real potential, U_{ab} – a complex optical potential describing the scattering state of the $a + b$ system. In this representation, known as prior mode, the transition amplitude for the transfer process is given by

$$\begin{aligned} T_{prior} &= \langle \chi_{\beta}^{(-)} \phi_a \phi_B | V_{\nu b} + U_{ab} - U_{\alpha} | \chi_{\alpha}^{(+)} \phi_A \phi_b \rangle = \\ &= \int \int d\mathbf{r}_{\alpha} d\mathbf{r}_{\beta} \chi_{\beta}^{-}(\mathbf{r}_{\beta})^* I_{\beta\alpha}((\mathbf{r}_{\beta}, \mathbf{r}_{\alpha})) \chi_{\alpha}^{+}(\mathbf{r}_{\alpha}), \end{aligned} \tag{3.39}$$

where \mathbf{r}_{α} and \mathbf{r}_{β} – the radius vectors, illustrated in Fig.3.2, describing relative distance of the $b + A$ and $a + B$ systems respectively, U_{α} – the optical potential describing elastic scattering of α -channel, and $I_{\beta\alpha}$ is the kernel function having form

$$I_{\beta\alpha}((\mathbf{r}_{\beta}, \mathbf{r}_{\alpha})) = (\phi_a \phi_B | V_{\nu b} + U_{ab} - U_{\alpha} | \phi_A \phi_b). \tag{3.40}$$

Analogously, for the exit channel with $V_{aB} = V_{a\nu} + U_{ab}$ the transition amplitude turns to

$$T_{post} = \langle \chi_{\beta}^{(-)} \phi_a \phi_B | V_{a\nu} + U_{ab} - U_{\beta} | \chi_{\alpha}^{(+)} \phi_A \phi_b \rangle. \tag{3.41}$$

Here, U_{β} stands for the optical potential corresponding to β elastic channel.




FIGURE 3.2: The arrangement of radial vectors in the DWBA on example of the $A + b \rightarrow a + B$ nuclear reaction. Here $A = a + \nu$ and $B = b + \nu$.

The differential cross section for the rearrangement nuclear reaction within the DWBA, in particular for prior form, may be written as

$$\frac{d\sigma}{d\Omega_{DWBA}} = \frac{\mu_\beta \mu_\alpha}{2\pi\hbar^2} \left(\frac{k_\beta}{k_\alpha} \right) |T_{prior}(\mathbf{k}_\beta, \mathbf{k}_\alpha)|^2 \quad (3.42)$$

For the post form the differential cross section is written analogously replacing the corresponding transition amplitude.

In accordance with the DWBA calculations the main ingredients are the internal wave functions for the initial (ϕ_A, ϕ_b) and final (ϕ_a, ϕ_B) nuclei. In this scheme, the valence particle ν is bound to the b core to give the composite B . In the simplest picture, the valence particle can be considered a pure single-particle state. This means that within this model there is only one possible configuration of the core and valence particle to give the nucleus B . Thus, the wave function for the nucleus B can be written as

$$\phi_B^{JM}(\xi, \mathbf{r}) = [\phi_b^I(\xi) \times \phi_{lsj}(\mathbf{r})]_{JM} \quad (3.43)$$

In a more sophisticated model, however, the state of the composite contains components of many single particle states coupled to all possible core states. Therefore, the wave function $\phi_B^{JM}(\xi, \mathbf{r})$ may be built as a superposition of the form

$$\phi_B^{JM}(\xi, \mathbf{r}) = \frac{1}{\sqrt{n_B}} \sum_{Ilj} \mathcal{A}_{lsj}^{IJ} [\phi_b^I(\xi) \times \phi_{lsj}(\mathbf{r})]_{JM} \quad (3.44)$$

where the coefficients \mathcal{A}_{lsj}^{IJ} are the so called coefficients of fractional parentage of

spectroscopic amplitudes. More information about the spectroscopic amplitudes will be introduced in the next section. The squared value of the amplitude is

$$S_{lsj}^{IJ} = |\mathcal{A}_{lsj}^{IJ}|^2, \quad (3.45)$$

which is called spectroscopic factors. The spectroscopic factors can be regarded as the probability of finding the valence particle ν in a single particle state l, s, j coupled to the core with spin I . The quantity n_B is the number of nucleons or clusters in the composite system.

Note that the kernel function $I_{\beta\alpha}(\mathbf{r}_\beta, \mathbf{r}_\alpha)$ involve the overlap between the composite and core wave functions. Using the Eq.3.44 the internal variable ξ in the integral can be explicitly performed giving just unity by normalization

$$(\phi_B^{JM}(\xi, \mathbf{r}), \phi_b^I(\xi)) \equiv \int d\xi \phi_B^{JM}(\xi, \mathbf{r}) \phi_b^I(\xi) = \frac{1}{\sqrt{n_B}} \sum_{Ilj} \mathcal{A}_{lsj}^{IJ} \phi_{lsj}(\mathbf{r}) \quad (3.46)$$

The bound wave function $\phi_{lsj}(\mathbf{r})$ obey the Shrödinger equation

$$(T + V_{\nu b}(\mathbf{r}) + \epsilon - E) \phi_{lsj}(\mathbf{r}) = 0 \quad (3.47)$$

where ϵ is the binding energy of the valence particle.

3.5 Spectroscopic Amplitudes within the Shell Model

Spectroscopic amplitudes for nuclear reactions can be obtained by calculations within the shell model. However, this approach is limited in calculating the amount of transferred particles. The calculation of matrix elements for the creation operator is suitable only in the case of the one-particle transfers, or at best two particle transfer. In nuclear reactions with the cluster transfer, one should use other theoretical approaches, or achieve certain values by fitting for the best reproduction of the observables.

Observables for the removal or addition of a nucleon from a specific initial state to a specific final state are related to the matrix elements of the creation and destruction operators. The creation operator a_{km}^+ is a tensor of rank j since it creates the single-particle state $|km\rangle$. Here, k stands for the set of single-particle quantum numbers nlj . The destruction operator a_{km} is not a tensor of

rank j , however,

$$\tilde{a}_{km} = (-1)^{j+m} [a_{k,-m}^+]^+ = (-1)^{j+m} a_{k,-m} \quad (3.48)$$

is a tensor of rank j . And its inverse is $a_{km} = (-1)^{j-m} \tilde{a}_{k,-m}$. Corresponding matrix elements of these operators can be written as

$$\langle k^{n-1} \omega' J' | | \tilde{a}_k | | k^n \omega J \rangle = (-1)^{j+J'-J} \langle k^n \omega J | | a_k^+ | | k^{n-1} \omega' J' \rangle \quad (3.49)$$

where ω – the harmonic oscillator parameter chosen to reproduce the observed mean-square charge radius, J – total spin of the n -particle state. All many-body matrix elements of a^+ can be reduced to these involving a single k -state.

Wave-function expansion relations and sum-rules for the states in the $n - 1$ particle system can be obtained by operating with the number operator

$$\hat{N}_k = \sum_m a_{km}^+ a_{km} \quad (3.50)$$

on the k^n configuration and then inserting a complete set of states with $n - 1$ particles

$$\begin{aligned} \hat{N}_k |k^n \omega JM\rangle &= \sum_m a_{km}^+ a_{km} |k^n \omega JM\rangle = n |k^n \omega J\rangle = \\ &= \sum_{m \omega' J' M'} a_{km}^+ |k^{n-1} \omega' J' M'\rangle \langle k^{n-1} \omega' J' M'| a_{km} |k^n \omega JM\rangle. \end{aligned} \quad (3.51)$$

The matrix element of $a_{k,m}$ can be reduced with the Wigner-Eckhart theorem

$$\begin{aligned} \langle k^{n-1} \omega' J' M' | a_{km} | k^n \omega JM \rangle &= (-1)^{j-m+J'-M'} \begin{pmatrix} J' & j & J \\ -M' & -m & M \end{pmatrix} \times \\ &\times \langle k^{n-1} \omega' J' | | a_{km} | | k^n \omega J \rangle. \end{aligned} \quad (3.52)$$

By Eq. 3.49 the reduced matrix element of \tilde{a} can be converted in a reduced matrix element of a^+ to obtain the final result:

$$\begin{aligned} \hat{N}_k |k^n \omega JM\rangle &= n |k^n \omega JM\rangle = \sum_{m \omega JM} (-1)^{-m-M'+J} \begin{pmatrix} J' & j & J \\ -M' & -m & M \end{pmatrix} \times \\ &\times a_{km}^+ |k^{n-1} \omega' J' M'\rangle \langle k^n \omega' J | | a_k^+ | | k^{n-1} \omega' J' M \rangle. \end{aligned} \quad (3.53)$$

Denoting the n -particle wave function by

$$|k^n \omega JM\rangle \equiv Z^+(k^n \omega JM) |\rangle \quad (3.54)$$

where Z^+ is the linear combination of a^+ operator which create the antisymmetric n -particle state, we can thus expand the k^n wave function in terms of those in the k^{n-1} basis:

$$|k^n\omega JM\rangle = (-1)^n \sum_{\omega' J'} \frac{\langle k^n\omega' J || a_k^+ || k^{n-1}\omega' J' M \rangle}{n\sqrt{2J+1}} [Z^+(k^{n-1}\omega' J') \times a_k^+]_{JM} | \rangle \quad (3.55)$$

A phase factor $(-1)^{n-1}$ arises from commuting a^+ with the $n - 1$ particles in the state k^{n-1}

A sum rule for the matrix elements of a^+ can be obtained by multiplying both sides of Eq. 3.53 by $\langle k^n\omega'' J'' M'' |$ to obtain

$$\begin{aligned} n\delta_{JJ''}\delta_{MM''}\delta_{\omega\omega''} &= \sum_{m\omega' J' M'} (-1)^{-m-M'+J} \begin{pmatrix} J' & j & J \\ -M' & -m & M \end{pmatrix} \times \\ &\quad \times \langle k^n\omega'' J'' M'' | a_{km}^+ | k^{n-1}\omega' J' M' \rangle \langle k^n\omega J | a_k^+ | k^{n-1}\omega' J' \rangle = \\ &= \sum_{m\omega' J' M'} \begin{pmatrix} J' & j & J \\ -M' & -m & M \end{pmatrix} \begin{pmatrix} J'' & j & J' \\ -M'' & -m & M' \end{pmatrix} \times \\ &\quad \times \langle k^n\omega'' J'' | a_k^+ | k^{n-1}\omega' J' \rangle \langle k^n\omega J | a_k^+ | k^{n-1}\omega' J' \rangle = \\ &= \frac{\delta_{JJ''}\delta_{MM''}}{2J+1} \sum_{\omega' J'} \langle k^n\omega'' J'' | a_k^+ | k^{n-1}\omega' J' \rangle \langle k^n\omega J | a_k^+ | k^{n-1}\omega' J' \rangle. \end{aligned} \quad (3.56)$$

Then, one finds the sum-rule:

$$\sum_{\omega' J'} \langle k^n\omega'' J'' | a_k^+ | k^{n-1}\omega' J' \rangle \langle k^n\omega J | a_k^+ | k^{n-1}\omega' J' \rangle = n(2J+1)\delta_{\omega\omega''}\delta_{JJ''} \quad (3.57)$$

Thus:

$$\sum_{\omega' J'} |\langle k^n\omega J | a_k^+ | k^{n-1}\omega' J' \rangle|^2 = \sum_{\omega' J'} |\langle k^{n-1}\omega' J' | \tilde{a}_k | k^n\omega J \rangle|^2 = n(2J+1) \quad (3.58)$$

The matrix elements in which the sum over final states is normalized to unity are historically called coefficients of fractional parentage [3] (CFP) defined by:

$$\mathcal{A}_{lsj}^{JJ'} = \frac{\langle k^n\omega J | a_k^+ | k^{n-1}\omega' J' \rangle}{\sqrt{n(2J+1)}} \quad (3.59)$$

The reduced matrix elements of the creation and destruction operators are used to define the spectroscopic amplitudes associated with nuclear reactions. They are also the building blocks for the more complicated operators associated with one-body (such as electromagnetic and β -decay) and two-body transition

amplitudes.

3.6 Interaction potentials. The double folding model

The numerical calculations of the elastic scattering can be performed in the framework of the OM with the OM potential given by:

$$U(R) = -V^V(R) - iW^V(R) + V^{SO}(R)(\mathbf{l} \cdot \boldsymbol{\sigma}) + V^C(R), \quad (3.60)$$

where V^V , W^V , V^{SO} , and V^C are real volume, imaginary volume, spin-orbit and Coulomb potentials, respectively. The volume potentials of colliding two spherical nuclei may be represented as parametrized function. For example, in practice the Woods-Saxon potential is often used, and it has the form

$$\begin{aligned} V^V(r) &= V_0^V f_{r_V, a_V}(r), \\ W^V(r) &= V_0^W f_{r_W, a_W}(r), \\ f_{r_0, a_0}(r) &= \frac{1}{1 + \exp\left(\frac{r-r_0}{a_0}\right)}, \end{aligned} \quad (3.61)$$

where V_0 is depth of the potential, r_0 is average distance and a_0 is diffusion parameter.

The spin-orbit term of the OM potential has standard form

$$V^{SO}(r) = V_0^{SO} \left(\frac{\hbar}{m\pi c}\right)^2 \frac{1}{r} \frac{d}{dR} f_{R_{SO}, a_{SO}}(r). \quad (3.62)$$

The Coulomb term has been taken as the interaction of a point-charge with a uniformly charged sphere

$$V^C(r) = \begin{cases} \frac{Z_1 Z_2 e^2}{2r_C} \left(3 - \frac{r^2}{r_C^2}\right), & \text{for } r \leq r_C, \\ \frac{Z_1 Z_2 e^2}{r}, & \text{for } r > r_C. \end{cases}$$

The situation changes, when the colliding nuclei have sophisticated shape. Assume, that there are nuclear matter distribution of the projectile $\rho_a(r_a)$ and target $\rho_A(r_a)$, depending on internal their own radii. Folding the nucleon-nucleon interaction potential V_{nn} over the density distributions of nuclear matter, in the framework of the *Double folding model* an interaction potential may




FIGURE 3.3: Radius vectors for the double folding model.

be constructed as follow

$$\begin{aligned} V^V(r) &= N_f V^{DF}(r) \\ V^{DF}(\mathbf{r}) &= \int \int d\mathbf{r}_a d\mathbf{r}_A \rho_a(\mathbf{r}_a) V_{nn}(r_{aA}) \rho_A(\mathbf{r}_A). \end{aligned} \quad (3.63)$$

where $r_{aA} = |\mathbf{r} + \mathbf{r}_A - \mathbf{r}_a|$ (see Fig. 3.3). The N_f normalization parameter is usually fitted in accordance with the dynamics of nuclear reaction of elastic scattering.

The DF potential may be calculated using the effective M3Y-Paris [4] nucleon-nucleon potential and the nuclear-matter-densities of projectile and target nuclei.

The V_{nn} effective nucleon-nucleon interactions usually are taken as a sum of the three Yukawa potentials, i.e. M3Y-potentials:

$$V_{nn}(r) = \sum_{i=1}^3 N_i \frac{\exp(-\mu_i r)}{\mu_i r}. \quad (3.64)$$

The parameters N_i , μ_i of the M3Y-Paris effective NN-potential [4], often used in the double folding calculations, are given in Tab.3.1.

The dependence $g(E)$ of the folding potential on the collision energy and an additional dependence on the nuclear matter density in the region of their overlap $F(\rho)$ have been phenomenologically established. The folding potential is usually taken as the real part of the optical potential, adjusting its depth to better reproduce the experimental data. The imaginary part of the optical potential is selected in the form of the Woods-Saxon potential, or the same folding potential, while varying its depth.

TABLE 3.1: Parameters of the M3Y-Paris [4] NN-potential used in double-folding calculations.

i	$N_i, \text{ MeV}$	Direct T=0	Exchange T=0	Direct T=1	Exchange T=1	μ_i, fm^{-1}
1	11061.60		-1524.25	313.62	-4118.9	4.0000
2	-2537.5		-518.75	223.5	1054.75	2.5000
3	0.0		-7.84	0.0	2.62	0.7072

The potential depending on the collision energy of the projectile, undergoes a change by multiplying a factor

$$g(E) = 1 - 0.003 \frac{E}{A}. \quad (3.65)$$

An additional dependence of the V_{nn} potential on the total density of colliding nuclei in the region of their overlap may given as

$$F(\rho) = C (1 + \alpha \exp(-\beta \rho) - \gamma \rho(\mathbf{r})). \quad (3.66)$$

with the parameters

$$C = 0.2658 \quad \alpha = 3.8033 \quad \beta = 1.4099 \quad \gamma = 4.0 \quad (3.67)$$

For the particles t , ${}^3\text{He}$, α , which have simple structure, the density distribution of nuclear matter can be represented as parametrized Gaussian function as follow

$$\rho(r) = \rho_0 \exp(-\alpha_0 r^2) \quad (3.68)$$

where the parameters ρ_0 and α_0 are defined in condition to reproduce the rms matter radii

$$\alpha_0 = \frac{3}{2\langle r_m^2 \rangle}, \quad \rho_0 = a \left(\frac{\alpha_0}{\pi} \right)^{3/2} \quad (3.69)$$

here a – atomic number of projectile.

The density distributions in (3.63) are normalized so that

$$\int \rho_A(\mathbf{r}) d\mathbf{r} = A \quad \int \rho_a(\mathbf{r}) d\mathbf{r} = a \quad (3.70)$$

where A, a are the number of nucleons in the respective nuclei.

As regards target nuclei, having cluster structure, the density distribution of nuclear matter are calculated by means of the three body wave function, and its details are set out in the Section 2.5 .

Chapter 4

Results and discussions

4.1 The three-body wave function

The chosen three-body wave function are obtained by variational method, and its parameters for ground state of ${}^6\text{He}$, ${}^6\text{Li}$ and ${}^9\text{Be}$ are listed in Application B. The process of minimization of eigen-energy of the three-body systems was done with the RSC potential [5] for nucleon-nucleon interaction, the SBB potential [6] for α - N interaction and the BFW potential [7] for α - α interaction. In Tab. 4.1 the used and obtained data of variational calculations are listed¹.

The obtained results of eigen-energies of the three-body systems in the ground state are equal to -0.228 MeV for ${}^6\text{He}$, -3.258 MeV for ${}^6\text{Li}$ and -1.417 MeV for ${}^9\text{Be}$. These theoretical data are differ from experimental data slightly. For example slightly difference has ${}^9\text{Be}$, its experimental established energy in the three-body system is $E_{exp} = -1.417$ MeV. The significant differences between variational eigen-energy were in ${}^6\text{He}$ and ${}^6\text{Li}$. Corresponding value of the energy are $E_{exp} = -0.975$ MeV for ${}^6\text{He}$, $E_{exp} = -3.700$ MeV for ${}^6\text{Li}$. Such kind of discrepancy of variational calculations with experimental data are probably in the non-completeness of the basis function. The basis function taken in this work has no symmetrization, and doesn't enfold an alternative coordinate space.

¹The obtained data of the variational calculations are not the subject of our research, since the results, perhaps better ones, have already been published and discussed. The purpose of the section result of the three-body wave function is to familiarize with what wave function we are dealing with, and its features will be needed in further discussions.

TABLE 4.1: Results of the search for the minimum energy of three-body systems found by means of the variational approach: ^6He , ^6Li and ^9Be . The obtained minimum energy E_{min} , used set of quantum numbers $\gamma \equiv \lambda lLS$, the dimension D and the probability P of each γ are listed.

^6He			^6Li			^9Be		
$E_{min} = -0.228$			$E_{min} = -3.258$			$E_{min} = -1.417$		
γ	D	P	γ	D	P	γ	D	P
0 0 0 0	9×9	0.884	0 0 0 1	8×8	0.898	0 1 1 1	7×7	0.403
1 1 1 1	9×9	0.102	2 0 2 1	7×7	0.073	2 1 1 1	7×7	0.328
2 2 0 0	5×5	0.010	1 1 1 0	7×7	0.021	2 1 2 1	7×7	0.235
3 3 1 1	5×5	0.004	2 2 0 1	5×5	0.005	2 3 1 1	6×6	0.017
			2 2 1 1	5×5	0.002	2 3 2 1	5×5	0.004
						4 3 1 1	4×4	0.009
						4 3 2 1	4×4	0.003




FIGURE 4.1: The correlation function $W(x, y)$ of the three-body wave function of the ground state of ${}^6\text{He}$ in the *a*) 3D-Plot and *b*) contour-plot.

In Figures 4.1, 4.2 and 4.3 the correlation density of the three-body wave function are illustrated in the 3D-Plot for ${}^6\text{He}$, ${}^6\text{Li}$ and ${}^9\text{Be}$. For the purpose of good visibility the contour-plot are given along.

In the Figures 4.1 and 4.2, two main geometric configurations can be distinguished, cigar-shaped within coordinates $(4.0, 1.0)$, and helicopter-shaped within coordinates $(1.8, 3.0)$, both for ${}^6\text{He}$ and ${}^6\text{Li}$ nuclei. The striking highlight of such geometrical configurations is clear evidence of the cluster structure. In particular, we see that in ${}^6\text{He}$, the effect of pairing two nucleons stands out against the background of comparison with ${}^6\text{Li}$. Similar results were described in works [1], and were obtained in the method using hyperspherical harmonics [8].

The Fig. 4.3 shows the locality of the wave function, which indicates the bound state of the ground state of ${}^9\text{Be}$. Moreover, the maximum of the correlation density function is shown at the coordinates $(3.0, 2.0)$, that is $x \simeq 3$, $y \simeq 2$. This allows us to make the assumption that basically the pair of α -particles are relatively farther than the distance of the valence neutron from the center of




FIGURE 4.2: The same caption as in the Fig.4.1, but for ${}^6\text{Li}$.




FIGURE 4.3: The same caption as in the Fig.4.1, but for ${}^9\text{Be}$.

mass of the pair.

4.2 The density distribution function of nuclear matter

On the basis of the three-body wave function in Eq.2.4, the distribution function of the density of nuclear matter was calculated using Equations 2.36, 2.37 and 2.39 for ${}^6\text{He}$, ${}^6\text{Li}$ and ${}^9\text{Be}$ in the ground states. The calculation results are shown in the Figures 4.4, 4.5 and 4.6. The plotted the density functions of ${}^6\text{He}$, ${}^6\text{Li}$ and ${}^9\text{Be}$ are normalized to their own atomic masses.

A distinctive feature of the obtained results is in the extended tail of the density function for ${}^6\text{He}$ in Fig.4.4. This is caused by the nucleons in the three-body system. In particular, the density function of the alpha particle $\rho_{\alpha}^{(k)}(R)$ inside ${}^6\text{He}$ tends to zero rapidly as the radius R increases. The density function of two neutrons $\rho_N^{(p)}(R) + \rho_N^{(q)}(R)$ starts near zero, reaching a maximum near




FIGURE 4.4: The density distribution function of nuclear matter $\rho(R) = \rho_{\alpha}^{(k)} + \rho_N^{(p)} + \rho_N^{(q)}$ illustrated with the densities of the ${}^6\text{He}$ constituents.




FIGURE 4.5: The same caption as in Fig. 4.4, but for ${}^6\text{Li}$.

$R \simeq 2.5$, and decreases more slowly than the density function of the alpha particle. What is obvious is that the start of the value of the density function $\rho_N^{(p)}(R) + \rho_N^{(q)}(R)$ near zero means that these two neutrons lie in the p -shell. Thus, we see that the main contribution to the density function starting from $R \simeq 3.0$ is due to two neutrons.

A similar structure of the nucleus ${}^6\text{Li}$ is seen in Fig. 4.5. The density function of the alpha particle $\rho_\alpha^{(k)}(R)$ inside ${}^6\text{Li}$ is visible, rapidly tending to zero as the radius R increases. And the density function of the two nucleons $\rho_N^{(p)}(R) + \rho_N^{(q)}(R)$, starting around zero, reaches a maximum near $R \simeq 2.5$, and decreases more slowly than the density function of the α -particle $\rho_\alpha^{(k)}(R)$. The beginning of the value of the density function $\rho_N^{(p)}(R) + \rho_N^{(q)}(R)$ near zero means that these two nucleons lie in the p -shell. In fact, we see, again as in the case with ${}^6\text{He}$, that the main contribution to the density function is due to two nucleons starting from $R \simeq 3.0$.

The density function $\rho(R)$ of ${}^9\text{Be}$ has a different shape than the other two




FIGURE 4.6: The density distribution function of nuclear matter $\rho(R) = \rho_N^{(k)} + \rho_\alpha^{(p)} + \rho_\alpha^{(q)}$ illustrated with the densities of the ${}^9\text{Be}$ constituents.

previous nuclei (see Fig.4.6). The alpha particle density function has become much more voluminous. The nuclear matter density distribution function $\rho_\alpha^{(p)} + \rho_\alpha^{(q)}$ tends to zero faster than the valence neutron density function $\rho_N^{(k)}$ with increasing distance R . Starting from the $R \simeq 5.0$ distance, the valence neutron density function $\rho_N^{(k)}$ prevails in the nuclear matter contribution. The spectator in the (k, pq) scheme has such distribution function starting near zero, which is characteristic of p -shell nucleons.

The formulas obtained by us for calculating the distribution function of the density of nuclear matter in the framework of the three-body model were tested using the parameters of the wave function of the ground state of ${}^9\text{Be}$ and ${}^6\text{Li}$ nuclei presented from the sources [9, 10]. In particular, for verification, calculations were made for the rms charge radius. The results obtained $\langle R_{ch}^2 \rangle^{\frac{1}{2}} = 2.54 \text{ fm}$ and $\langle R_{ch}^2 \rangle^{\frac{1}{2}} = 2.34 \text{ fm}$ for ${}^6\text{Li}$ and ${}^9\text{Be}$ nuclei, respectively, are in excellent agreement with the data indicated in those sources, which means that the chosen method is correct. And also, in addition to the fact that all formulas have been carefully checked at every stage, this excludes the presence of errors in the calculation formulas.

The density function of nuclear matter is an excellent subject for studying rms matter radii. Given the function $\rho(R)$, we can represent the formula for calculating the rms matter radii $\langle R_m^2 \rangle$ as

$$\langle R_m^2 \rangle = \frac{\int_0^\infty dR R^4 \rho(R)}{\int_0^\infty dR R^2 \rho(R)}. \quad (4.1)$$

The results of calculated rms matter radii are presented in Tab.4.2 with the values of the binding energy BE_{min} for the three-body system obtained by the variational method for nuclei. The general picture of the obtained radii shows an overestimation in comparison with other sources [11–13]. For example, the radius of the ${}^6\text{He}$ nucleus obtained in our calculations shows 2.97 fm , while the another source gives 0.35 fm less. The difference with another radius in the

TABLE 4.2: Comparisons of the binding energies obtained by means of variational approach BE_{min} with the experimental data BE_{exp} and the rms matter radii $\langle R_m^2 \rangle^{\frac{1}{2}}$ with other sources.

	${}^6\text{He } (0^+)$	${}^6\text{Li } (1^+)$	${}^9\text{Be } (\frac{3}{2}^-)$
BE_{min}	0.228	3.258	1.417
BE_{exp}	0.97	3.70	1.57
$\langle R_m^2 \rangle^{\frac{1}{2}}, \text{ fm}$	2.97	2.69	2.78
	2.62 [11]	2.45 [12]	2.62 [13]

case of ${}^6\text{Li}$ is 0.24 fm , and for ${}^9\text{Be}$ 0.16 fm . That is, if you list the nuclei from the table from left to right, the difference in radii decreases. If we look at the obtained theoretical calculations of the binding energy BE_{min} and compare with the experimental data, there is an opposite tendency in the energy differences – the difference decreases – 0.742 MeV , 0.442 MeV , 0.153 MeV . Thus, it can be assumed that the greater the underestimation in the binding energy, the more overestimation in the rms matter radii.

In accordance with this proposal, the dependence of the binding energy with the $\langle R_m^2 \rangle^{1/2}$ was studied using the example of ${}^6\text{He}$. In order to obtain the binding energy close to the experimental data, the depth of the $\alpha - N$ interaction potential, which has a Gaussian shape, was increased from 47 MeV to 49 MeV . Thus, it was possible to reproduce the experimental value of the binding energy. However, the obtained rms radius $\langle R_m^2 \rangle^{1/2} = 2.93 \text{ fm}$ decreased by only 0.04 fm .

The reason for the discrepancy between our results obtained from calculations of the rms matter radii with other sources could not be explained. Perhaps, the problem lies in the basis function, which has not been symmetrized or antisymmetrized. Moreover, the basis function is not complete, that it should have enclosed the entire basis space. Nevertheless, knowing the very weak dependence of the density distribution function on the binding energy of the three-body systems, we can use this wave function in calculations to construct interaction potentials within the framework of the folding model.

In the process of calculating the rms matter radii, it was found that elements from the set γ , that have a contribution to the wave function below 0.1 (see Tab.4.1), do not affect the obtained values. Therefore, such elements of the γ were excluded from the density function calculations.




FIGURE 4.7: The folding potential $V^{DF}(r)$ in the framework of the three-body model. The potential is for the interaction of the α particle with the constituents of ${}^6\text{He}$.

4.3 The $\alpha + {}^6\text{He}$ nuclear reaction

4.3.1 Elastic scattering

Theoretical calculations of the differential cross section of the elastic colliding of the ${}^6\text{He} + \alpha$ system has been carried out within the framework of the OM. The calculations were done by means of *FRESCO* code [14]. As the interaction potential of the ${}^6\text{He} + \alpha$ system were taken the double folding potential. The potential was built using the three-body wave function of the ground state of ${}^6\text{He}$. The interactions of its α -cluster and two nucleon with the α -projectile are presented in Fig. 4.7. The calculated double folding potential has a specific structure. It is worth noting that the potential of the α cluster with the projectile decreases rapidly as the distance r increases, while the potential of interaction of two nucleons with the projectile decreases slowly.

The potential obtained within the framework of the double folding model based on the three-body wave function was used to calculate the differential cross section for elastic scattering ${}^6\text{He}$ by alpha particles at energy $E_{lab} = 151 \text{ MeV}$. The results obtained using different potentials, including the double folding potentials, are shown in Fig. 4.8.

In addition to the folding potential, two other potentials (WS1 and WS2) are presented as the Woods-Saxon potential taken from [15]. The parameters of the potentials used in the calculations of the theoretical curves within the optical model and the corresponding χ^2 are presented in Tab. 4.3. Comparison of theoretical curves with the experimental data is in good agreement, if we




FIGURE 4.8: *a)* The differential cross section of the elastic scattering of ${}^6\text{He}$ by α -particles at $E_{lab} = 151\text{MeV}$ with the different potentials: the double folding potential (blue solid), the WS2 (gray dashed) [15] and WS1 (gray dotted) [15]. *b)* The cross sections of the elastic transfer reaction in terms of the transfer mechanisms: elastic scattering (solid gray), simultanious transfer of $2N$ (gray dashed), sequantial transfer of $2N$ (dotted) and teit coherent sum (solid blue). Experimanetal data taken from [15].

TABLE 4.3: The optical parameters of the potentials (DF, WS1, WS2) used in calculations of differential cross section of ${}^6\text{He}$ by α -particle at $E_{lab} = 151$ MeV within the optical model.

	$-V_0$, MeV	R_v , fm	a_v , fm	$-W_0$, MeV	R_w , fm	a_w , fm	χ^2/N
DF		$N_r=1.4$			$N_i=0.5$		7.32
WS1	102.5	1.78	0.920	13.0	3.85	0.5	5.75
WS2	102.5	1.54	0.904	7.0	4.28	0.569	6.31

do not take into account the backward angles caused by another interaction mechanism, which will be discussed in the next section. The difference in cross sections with different potentials is insignificant, except for the folding potential, which does not describe the first experimental point.

The advantage of folding potential in comparison with others consists of a less number of adjustable parameters to the experimental data. If these parameters in the both phenomenological potential WS1 and potential WS2 are six, then folding potential depends on only two parameters. The overhead in this case is a small concession in the χ^2 test.

4.3.2 Elastic transfer

In Fig. 4.8.a) theoretical curves were presented to describe the experimental data on elastic scattering. The calculations made within the framework of the optical model showed a good result. However, this is only at the front scattering angles. The backscattering angles are far from the description of optical calculations. Therefore, in order to explain this difference, we propose mechanisms for the transfer of two nucleons: sequential and simultaneous transfer. A schematic representation of the mechanisms can be seen in Fig. 4.9. Thus, the differential cross section of these mechanisms can be written as follows

$$\frac{d\sigma}{d\Omega}(\theta) = |f(\theta)_{OM} + f_{sim}(\pi - \theta) + f_{seq}(\pi - \theta)|^2 \quad (4.2)$$

Calculations for obtaining the differential cross section of the elastic transfer were carried out within the framework of the CRC method. Potential for the input and output channels was taken the double folding potential, and the potential for the intermediate channel was taken the optical potential with global optical parametrizations for α particles [20]. Trial calculations have shown that the results depend very little on the selected potential for the intermediate channel. The results of calculations for elastic transfer made in the framework of CRC are shown in Fig. 4.8.b). The sequential transfer of two nucleons is shown by gray dotted curves, their simultaneous transfer by dashed gray curves, the




FIGURE 4.9: Schematic representation of the elastic transfer nuclear reaction, on the left - the simultaneous transfer of two nucleons, on the right - the sequential transfer of two nucleons.

elastic scattering cross section by straight gray curves, and, finally, their coherent sum is shown as a blue straight curve. The wave function of the bound states was chosen by fitting the potential depth to the binding energy of the composite systems. In particular, the binding energy of one neutron with ${}^6\text{He}$ was 1.8 MeV, neutron with ${}^5\text{He}$ – 0.1 MeV, and two neutrons with α – 0.9 MeV.

As mentioned in the previous section, the elastic collision mechanism prevails at the forward scattering angles. Beginning at an angle of 90 degrees, the contribution to the cross section is mainly due to the simultaneous transfer of two nucleons. It is worth noting here that two neutrons have the nlj $2S_0$ configuration, which, possibly, leads to an oscillatory cross section. Ten times less is the contribution of the sequential transfer of neutrons, having the $1P_{\frac{3}{2}}$ configuration. Note that the spectroscopic amplitudes for two neutrons in ${}^6\text{He}$, one neutron in ${}^5\text{He}$ and in ${}^6\text{He}$ were taken to be equal to unity for the beginning. The results showed a difference in cross-sections for sequential and simultaneous transmissions of about ten. For the best reproduction of the experimental data, the spectroscopic amplitude of two neutrons was taken $\mathcal{A}_{2S_0}^{00} = 1.3$, while the others remained unchanged.

Thus, it was possible to achieve good agreement between the calculated differential cross section, using the double-folding potential and the proposed transfer mechanisms, with the experimental data.




FIGURE 4.10: Folding potential of α interaction with ${}^9\text{Be}$ nucleus with allowance for the Coulomb interaction.

4.4 The $\alpha + {}^9\text{Be}$ nuclear reactions

4.4.1 Elastic scattering

We chose the DDM3YY Paris potential [4] as our effective nucleon–nucleon interaction for calculating the folding potential. The results from calculating the interaction potential for α and ${}^9\text{Be}$ nuclei are shown in Fig. 4.10. The depth of the obtained potential on the whole agrees with the known global parameterizations of the optical potential for α nuclei. It should be noted that the potential diffusivity, which is to a large extent determined by the extended spatial distribution of ${}^9\text{Be}$ (see Fig. ??), is considerable.

The obtained folding potential was used to calculate the differential cross section of elastic scattering for ${}^9\text{Be}(\alpha, \alpha){}^9\text{Be}$ within the optical model using the data base in [16]. Figure 4.11 shows the angular distributions obtained in the range of collision energies from 4.5 to 25.1 MeV/nucleon, compared to the available experimental data [17–19].

In our calculations, folding potential was used as the real part of the optical potential U_{opt} without additional coefficients. The optical potential was added using an imaginary term in the form of a Woods–Saxon function with parameters chosen by matching the theoretical cross section to the experimental data (see Table 1). The curves in Fig. 4.11 are in good agreement with the data in the region of small and intermediate scattering angles. For large scattering angles at these energies, the elastic scattering cross section is influenced by the




FIGURE 4.11: Differential cross sections of elastic scattering ${}^9\text{Be}(\alpha, \alpha){}^9\text{Be}$ for different collision energies, compared to the experimental data in [17–19] : (1) 4.5, (2) 7.25, (3) 10.00, (4) 11.25, (5) 12.62, and (6) 26.00 MeV/nucleon.

mechanism of transfer of heavy ${}^5\text{He}$ clusters, which is not explicitly considered in the optical model.

The parameters found for the imaginary part of the optical potential (see Table 4.4) allow us to study the dependence of the optical potential on the collision energy. In particular, the depth parameter of the imaginary part of the optical potential can be approximately described as

$$W_0(E) = -(23.2 + 0.15E). \quad (4.3)$$

The form of the dependence and the found slope coefficient in expression 4.3 are in good agreement with the corresponding parameters of the known global parameterizations for optical potentials [20].

4.4.2 Inelastic scattering

The obtained optical potentials were used to calculate the differential cross sections of inelastic channels within strong channel coupling using the FRESCO code [14]. The corresponding angular distributions shown in Fig. 4.12.a are in good agreement with the experimental data. Calculations yield an estimate for the parameter of quadrupole deformation of the target nucleus: which is in good agreement with other sources [21–23].

4.4.3 One nucleon transfer reactions

We also calculated the nucleon transfer cross sections within the distorted waves approach using the DWUCK5 code [24]. The parameters of the potentials for the input and output reaction channels are given in Table 4.5. Figure 4.12.b shows the calculated differential cross sections for the transfer reactions ${}^9\text{Be}(\alpha, {}^3\text{He}){}^{10}\text{Be}$ and ${}^9\text{Be}(\alpha, t){}^{10}\text{B}$; these describe experimental data quite well. A comparison with the measured cross section yields information on the value of the spectroscopic factor for the ground state of a final nucleus, ${}^{10}\text{Be} = ({}^9\text{Be}$

TABLE 4.4: Parameters of imaginary part of the optical potential for calculating elastic scattering cross section $\alpha + {}^9\text{Be}$. The a_w parameter for all energies is 0.97 fm

E_{lab} , MeV	$-W_0$, MeV	r_w , fm
18	26.5	1.35
29	27.5	1.3
40	28.5	1.25
45	30.5	1.25
50.5	31.5	1.1
104	39.5	1.05




FIGURE 4.12: Differential cross sections for $E_{\text{lab}} = 16.25$ MeV/nucleon: (a) inelastic scattering ${}^9\text{Be}(\alpha, \alpha){}^9\text{Be}^*$ with excitation of: (1) $5/2^-$ states, (2) $7/2^-$ states; (b) single nucleon transfer reactions: (1) ${}^9\text{Be}(\alpha, t){}^{10}\text{B}$ and (2) ${}^9\text{Be}(\alpha, {}^3\text{He}){}^{10}\text{Be}$. Experimental data are taken from [21–23]

$+ n)$ and ${}^{10}\text{B} = ({}^9\text{Be} + p)$. The obtained values of $({}^{10}\text{Be}) = 1.1$ and $({}^{10}\text{B}) = 0.6$ in particular differ only slightly from the published values [21, 23, 25]. The values of the neutron structure factor in ${}^9\text{Be}$ obtained in [25] by analyzing reactions ${}^9\text{Be}(d, p){}^{10}\text{Be}$ and ${}^{10}\text{B}(d, p){}^{11}\text{B}$ are in excellent agreement with the results obtained in this work.

4.5 The $d+{}^9\text{Be}$ nuclear reactions

4.5.1 The elastic channel

The DF potential was calculated using the effective M3Y-Paris nucleon-nucleon potential and the nuclear-matter-densities of projectile and target nuclei. In order to calculate the ${}^9\text{Be}$ matter distribution we applied the $\alpha+\alpha+n$ three-body model (for more details, see Ref. [26]), while the matter density distribution of the deuteron projectile was chosen to be of the form

$$\rho\left(\frac{1}{2}r\right) = \int |\Psi(\mathbf{r})|^2 d\Omega_r. \quad (4.4)$$

For convenience, in the OM and CC (CRC) calculations the potentials have been fitted by means of the sum of three Woods-Saxon potentials:

$$V^V(R) = \sum_{i=1}^3 V^i f^{R_i, a_i}(R), \quad (4.5)$$

$$f^{R_V, a_V}(R) = \frac{1}{1 + \exp \frac{R - R_V}{a_V}}. \quad (4.6)$$

The parameters of the imaginary part of the optical potential were obtained by fitting the theoretical cross sections to the experimental data at 19.5 MeV and 35 MeV incident energies. As a starting point, the same parameterizations of the real part were used. The obtained potential parameters after fitting are listed in Table 4.6 for both 19.5 MeV and 35.0 MeV incident energies.

TABLE 4.5: Parameters of the optical potentials for calculations using the distorted waves and channel coupling methods

E_{lab}	$-V_0$, MeV	r_v , fm	a_v , fm	$-W_0$, MeV	r_w , fm	a_w , fm	r_c , fm
$\alpha+{}^9\text{Be}$	The folding potential			32.65	1.14	0.97	1.30
${}^3\text{He}+{}^{10}\text{B}$	132.9	1.54	0.57	19.5	1.82	0.22	0.81
$t+{}^{10}\text{B}$	95.0	0.95	0.82	8.0	1.60	0.73	1.07
$t+{}^{10}\text{Be}$							




FIGURE 4.13: The angular distribution of elastic scattering data of d from ${}^9\text{Be}$ at laboratory energy 19.5 MeV in comparison with theoretical calculations within the OM (solid curve).

The comparison of the results of the theoretical calculations with the measured data for elastic scattering at 19.5 MeV and 35.0 MeV energies are plotted in Fig. 4.13. The cross sections obtained in the framework of the OM with the DF potential are shown as solid curves. Theoretical results obtained by means of the OM give an excellent agreement, $\chi^2 \approx 2.5$, with the experimental data. The parameters of parameterized double-folding potential are listed in Table 4.6.

4.5.2 Inelastic scattering

The CC and DWBA approaches have been applied to analyse the measured inelastic scattering data corresponding to the ${}^9\text{Be}(5/2^-, 2.43 \text{ MeV})$ excitation.

TABLE 4.6: Parameterized double-folding potentials of the $d+{}^9\text{Be}$ system used in the OM, CC and DWBA calculations.

E_d , MeV	$i V_0$, MeV	r_v^a , fm	a_v , fm	W_0 , MeV	r_w^a , fm	a_w , fm	V_0^{SO} , MeV	N_R ,	r_C^a , fm	χ^2/N
19.5	16.18	0.328	0.308	3.99	0.328	0.127	3.275	1.22	0.809	2.490
	270.97	0.746	0.831	25.50 (17.5 ^b)						
	30.605	1.491	1.724	0.924						
35.0	15.941	0.328	0.308	7.07	0.612	0.108	3.275	1.17	0.809	2.503
	268.68	0.746	0.831	22.50 (17.5 ^b)						
	30.58	1.491	1.724	0.999						

^a Radii are defined as $R_i = r_i \left(A_P^{1/3} + A_T^{1/3} \right)$.

^b The values are used in CRC calculations.




FIGURE 4.14: The cross sections of inelastic scattering ${}^9\text{Be}(d, d') {}^9\text{Be}^*$ ($E_{exc}=2.43$ MeV) at laboratory energies 19.5 MeV (full circle) and 35 MeV (full triangle). Theoretical curves are described in the text.

Calculations were performed employing the FRESCO code [14] and the DWUCK5 code [24] which are available in the NRV knowledge-base [16].

In order to describe the measured experimental data one has to consider the ${}^9\text{Be}$ target having a quadrupole deformation. Thus, the ${}^9\text{Be}$ spectrum consists of the rotational band including the $3/2^-$ ground state, $5/2^-$ state at 2.43 MeV and $7/2^-$ state at 6.38 MeV. Couplings to these states were taken into account within the coupled-channel approach. The spin reorientations were also taken into account. The coupling interaction has the usual form:

$$V_\lambda(R) = -\beta_\lambda R_V \left| \frac{dV^V}{dR} \right| - i\beta_\lambda R_W \left| \frac{dW^D}{dR} \right|, \quad (4.7)$$

where β_λ is the deformation parameter of λ multipole describing the target-nucleus form. Here, we neglect as usual the contribution of the Coulomb interaction.

The calculated cross sections for inelastic scattering to the $5/2^-$ state at 2.43 MeV are shown in Fig. 4.14. The solid curves correspond to the results obtained within the CC approach, while the dashed and dotted curves were obtained within the DWBA approach using different values of the deformation parameter β_2 . The used potential parameters are listed in Table ??.

All the results in Fig. 4.14 are in good agreement with the experimental data, except for the cross sections around 60° at 19.5 MeV incident energy. The quadrupole deformation parameter $\beta_2 = 0.64$ extracted within the coupled-channel model is consistent with the previous studies [21, 23].




FIGURE 4.15: The target coupling schemes in the ${}^9\text{Be}(d, p){}^{10}\text{Be}$ (upper) and the ${}^9\text{Be}(d, t){}^8\text{Be}$ (lower) nuclear reactions. The bold two-headed arrows indicate $E\lambda$ transitions. The spin re-orientation effects are indicated as back-pointing arrows.

In the case of DWBA calculations, one uses the DF potential (see Table 4.6) for both the entrance and the exit channels. The DWBA angular distributions very well reproduce the structure of experimental data but clearly underestimate them when the deformation parameter $\beta_2 = 0.64$ is used (see the dotted curves in Fig. 4.14). In order to get the best fit the deformation parameter must be increased up to $\beta_2 = 0.93$, which is quite close to the values reported in previous studies (see, for example, [27, 28]).

Thus, one may confirm that channel coupling and the effects of spin reorientation enhance the cross section that results in the reduction of the deformation parameter. However, the DWBA approach takes into account only first-order contributions to the transition amplitude. In particular, it also describes only general features of the angular distributions and overestimates the deformation parameter in order to compensate the difference between the experimental data and the DWBA cross sections.

4.5.3 One-nucleon transfer reactions

The one-neutron pick-up ${}^9\text{Be}(d, t){}^8\text{Be}$ and stripping ${}^9\text{Be}(d, p){}^{10}\text{Be}$ reactions were analyzed here within the framework of the Coupled Reaction Channels (CRC).

The double-folding potential given in Table 4.6 was used in the CRC calculations for the entrance channel and the global optical parameterizations from Ref. [29, 30] were used for the exit channels. The coupling schemes of target and daughter nuclei for the ${}^9\text{Be}(d, p){}^{10}\text{Be}$ and ${}^9\text{Be}(d, t){}^8\text{Be}$ reactions are illustrated in Fig. 4.15. The states of ${}^{10}\text{Be}$, 2^+ and 2^+_1 , as well as the low-lying




FIGURE 4.16: Differential cross sections for the ${}^9\text{Be}(d, p){}^{10}\text{Be}^*$ reactions at 19.5 MeV leading to different final states (labelled in the figure) in ${}^{10}\text{Be}$. The experimental data are shown in comparison with theoretical results obtained within the CRC method.

excited states of ${}^8\text{Be}$, 2^+ and 4^+ , were included in the coupling scheme. Also, the schemes take into account the spin reorientations of states on the condition $J \neq 0$.

In order to construct the bound-state wave functions of the transferred particle in the entrance and exit channels, the common method, i.e. fitting the depth of the corresponding Woods-Saxon potential to the known binding energy, was employed. The reduced radius and diffuseness in this case are set to be $r = 1.25$ fm and $a = 0.65$ fm, respectively. If the transfer takes place to a final unbound state, the depth of the potential for this state was adjusted to yield a binding energy equal to -0.1 MeV in accordance with the procedure used in Ref. [21].

If the core and the composite nuclei have internal excitation energies, a renewed binding energy BE^* of the transferred particle is expressed by the formula:

$$BE^* = BE - E_{com}^* + E_{core}^* \quad (4.8)$$

where BE – the binding energy of the transferred particle, E_{com}^* , E_{core}^* – excitation energies of the composite and core nuclei, respectively.




FIGURE 4.17: Differential cross sections for the ${}^9\text{Be}(d, t){}^8\text{Be}^*$ reactions at 19.5 and 35 MeV leading to different final states (labelled in the figure) in ${}^8\text{Be}$. The experimental data are shown in comparison with theoretical results obtained within the CRC method.

The spectroscopic amplitude S for the addition of a particle to a core with angular momentum J_{core} to form a composite with J_{com} is related to the matrix element of the creation operator \hat{a}^\dagger :

$$S_{Nlj} = \frac{\langle J_{com} \| \hat{a}_{Nlj}^\dagger \| J_{core} \rangle}{\sqrt{2J_{com} + 1}} \quad (4.9)$$

where Nlj is the set of particle quantum numbers. The spectroscopic amplitudes for one particle states were calculated by means of the *ANTOINE* code [31] using the effective Cohen-Kurath interaction for p -shell nuclei [32]. The calculated spectroscopic amplitudes for the one-nucleon transfer reactions are listed in Table 4.7.

Angular distributions of the ${}^9\text{Be}(d, p){}^{10}\text{Be}$ nuclear reaction at $E_d=19.5$ MeV are shown in comparison with the theoretical curves calculated in the framework of the CRC method in Fig. 4.16.

In order to study the couplings of the input channels, the outputs were fixed using the deformation parameter of ${}^8\text{Be}$ from Ref. [35], and for ${}^{10}\text{Be}$ from Ref. [21]. The direct transition from the ground state is indicated by the dotted line (DWBA). The contributions of the transitions from excited states (CC), and from spin reorientations (SR) are indicated by dashed and solid

TABLE 4.7: Spectroscopic amplitudes used in CRC calculations for the Composite = Core + Nucleon system. The one-nucleon spectroscopic amplitudes have been calculated by means of the *ANTOINE* code [31]. The alpha spectroscopic amplitudes were taken from [33, 34].

Com	2J ₁	Core	2J ₂	N	2J	SA	Com	2J ₁	Core	2J ₂	N	2J	SA
⁹ Be	3	⁸ Be	0	n	3	-0.761	⁹ Be	3	⁸ Li	2 ₁	p	1	-0.444
⁹ Be	3	⁸ Be	4	n	3	0.816	⁹ Be	3	⁸ Li	6	p	3	-0.592
⁹ Be	3	⁸ Be	4	n	1	-0.242	⁹ Be	3	⁸ Li	2 ₂	p	3	-0.236
⁹ Be	5	⁸ Be	4	n	3	0.986	⁹ Be	3	⁸ Li	2 ₂	p	1	0.036
⁹ Be	5	⁸ Be	4	n	1	-0.417	⁹ Be	5	⁸ Li	4	p	3	0.593
⁹ Be	5	⁸ Be	8	n	3	-0.374	⁹ Be	5	⁸ Li	4	p	1	0.515
⁹ Be	7	⁸ Be	4	n	3	-0.457	⁹ Be	5	⁸ Li	2 ₁	p	3	-0.672
⁹ Be	7	⁸ Be	8	n	3	0.919	⁹ Be	5	⁸ Li	6	p	3	-0.571
⁹ Be	7	⁸ Be	8	n	1	-0.429	⁹ Be	5	⁸ Li	6	p	1	-0.171
⁸ Be	0	⁷ Li	3	p	3	-1.204	⁹ Be	5	⁸ Li	2 ₂	p	3	0.200
⁸ Be	0	⁷ Li	1	p	1	0.736	⁹ Be	7	⁸ Li	4	p	3	-0.323
⁸ Be	4	⁷ Li	3	p	3	-0.748	⁹ Be	7	⁸ Li	6	p	3	-0.899
⁸ Be	4	⁷ Li	3	p	1	-0.612	⁹ Be	7	⁸ Li	6	p	1	-0.564
⁸ Be	4	⁷ Li	1	p	3	0.667	⁷ Li	3	⁶ Li	2	n	3	0.657
⁸ Be	4	⁷ Li	7	p	3	0.624	⁷ Li	3	⁶ Li	2	n	1	-0.538
⁸ Be	4	⁷ Li	5 ₂	p	3	0.079	⁷ Li	3	⁶ Li	6	n	3	0.744
⁸ Be	4	⁷ Li	5 ₂	p	3	-0.146	⁷ Li	3	⁶ Li	4	n	3	-0.032
⁸ Be	8	⁷ Li	7	p	3	0.864	⁷ Li	3	⁶ Li	4	n	1	0.399
⁸ Be	8	⁷ Li	7	p	1	0.687	⁷ Li	1	⁶ Li	2	n	3	-0.925
⁸ Be	8	⁷ Li	5 ₂	p	3	0.374	⁷ Li	1	⁶ Li	2	n	1	0.197
⁸ Li	4	⁷ Li	3	n	3	-0.988	⁷ Li	1	⁶ Li	4	n	3	-0.555
⁸ Li	4	⁷ Li	3	n	1	0.237	⁷ Li	7	⁶ Li	6	n	3	-0.936
⁸ Li	4	⁷ Li	1	n	3	0.430	⁷ Li	7	⁶ Li	6	n	1	0.645
⁸ Li	4	⁷ Li	7	n	3	-0.496	⁷ Li	7	⁶ Li	4	n	3	-0.456
⁸ Li	4	⁷ Li	5	n	3	-0.665	⁷ Li	5 ₂	⁶ Li	2	n	3	-0.650
⁸ Li	4	⁷ Li	5 ₂	n	1	-0.275	⁷ Li	5 ₂	⁶ Li	6	n	3	0.732
⁸ Li	2 ₁	⁷ Li	3	n	3	0.567	⁷ Li	5 ₂	⁶ Li	6	n	1	0.549
⁸ Li	2 ₁	⁷ Li	3	n	1	0.351	⁷ Li	5 ₂	⁶ Li	4	n	3	0.200
⁸ Li	2 ₁	⁷ Li	1	n	3	0.905	⁷ Li	5 ₂	⁶ Li	4	n	1	-0.114
⁸ Li	2 ₁	⁷ Li	1	n	1	0.331	⁶ Li	2	d	2	α	0	0.907
⁸ Li	2 ₁	⁷ Li	5 ₂	n	3	0.767	⁶ Li	2	d	2	α	4	0.077
⁸ Li	6	⁷ Li	3	n	3	0.581	⁶ Li	6	d	2	α	4	0.943
⁸ Li	6	⁷ Li	5 ₂	n	3	-0.660	⁶ Li	6	d	2	α	8	0.028
⁸ Li	6	⁷ Li	5 ₂	n	1	-0.541	⁶ Li	4	d	2	α	4	0.929
⁸ Li	6	⁷ Li	7	n	3	0.973	⁹ Be	3	⁵ He	3	α	0	-0.925
⁸ Li	6	⁷ Li	7	n	1	-0.404	⁹ Be	3	⁵ He	3	α	4	0.784
⁸ Li	2 ₂	⁷ Li	3	n	3	-0.617	⁹ Be	5	⁵ He	3	α	4	0.974
⁸ Li	2 ₂	⁷ Li	3	n	1	-0.841	⁹ Be	5	⁵ He	3	α	8	-0.260
⁸ Li	2 ₂	⁷ Li	1	n	3	0.178	⁹ Be	7	⁵ He	3	α	4	0.882
⁸ Li	2 ₂	⁷ Li	1	n	1	0.331	⁹ Be	7	⁵ He	3	α	8	-0.737
⁸ Li	2 ₂	⁷ Li	5	n	3	0.231	⁷ Li	3	t	1	α	1	0.970
⁹ Be	3	⁸ Li	4	p	3	-0.947	⁷ Li	1	t	1	α	1	0.961
⁹ Be	3	⁸ Li	4	p	1	-0.319	⁷ Li	7	t	1	α	3	0.952
⁹ Be	3	⁸ Li	2 ₁	p	3	0.454	⁷ Li	5 ₂	t	1	α	3	0.223

lines, respectively. During the analysis, it was found that spin reorientation has a significant contribution in the $p + {}^{10}\text{Be}_{gs}$ channel, especially in the range of 40-60 degrees.

It is interesting to note that we managed to describe within the CRC method the differential cross section of the ${}^9\text{Be}(d, p){}^{10}\text{Be}_{gs}$ reaction at all scattering angles, including the range 40° - 60° , where they were not covered in Refs. [27, 36].

An appreciable contribution of the $3/2^- \rightarrow 2_1^+$, $5/2^- \rightarrow 2_1^+$, $7/2^- \rightarrow 2_1^+$ transitions was observed in the $p + {}^{10}\text{Be}_{3.37}$ channel in the entire range of scattering angles. In the cross section of the $p + {}^{10}\text{Be}_{5.96}$ channel, the theoretical calculation underestimates the experimental data starting from 70° . Possibly, other higher excited states of ${}^9\text{Be}$ should be taken into account.

Figure 4.17 displays the cross sections of the ${}^9\text{Be}(d, t){}^8\text{Be}$ nuclear reaction at both 19.5 MeV and 35 MeV incident energies. As in the case of the (d, p) reactions, the (d, t) reactions also show the strong channel-coupling effects. We see a manifestation of spin-reorientation effects in the $t + {}^8\text{Be}_{gs}$ channels and a significant contribution of the $3/2^- \rightarrow 2^+$, $5/2^- \rightarrow 2^+$, $7/2^- \rightarrow 2^+$ transitions in the $t + {}^8\text{Be}_{3.03}$ channel. Disagreements around 30° in the $t + {}^8\text{Be}_{gs}$ channel for both 19.5 MeV and 35 MeV incident energies and around 60° in the $t + {}^8\text{Be}_{3.03}$ channel for 19.5 MeV incident energy are possibly caused by the uncertainty in the $t + {}^8\text{Be}$ interaction potential.

Theoretical calculations made within the CRC method show, in general, good agreement with the experimental data for both (d, p) and (d, t) reactions. The analysis showed strong coupling effects in both entrance and exit channels. The effects of such couplings were also emphasized in Refs. [21, 37].

4.5.4 Cluster-transfer reaction

Differential cross sections for the nuclear reaction ${}^9\text{Be}(d, \alpha){}^7\text{Li}$ are of particular interest. This is due to the specific behaviour of the cross section at large scattering angles, which indicates a ${}^5\text{He}$ cluster transfer. In addition, the cross section calculated within the DWBA approach underestimates the data even at forward scattering angles. Therefore, in order to understand the difference between theory and experiment, the following transfer mechanisms are suggested (see Fig. 4.18):

- direct transfer of heavy clusters d and ${}^5\text{He}$;
- sequential two-step transfer of $n-p$, $p-n$, $n-\alpha$ and $\alpha-n$;




FIGURE 4.18: The scheme illustrates the reaction mechanisms taken into account in CRC calculations of the cross sections for ${}^9\text{Be}(d, \alpha){}^7\text{Li}$ reaction.

The resulting differential cross section for the ${}^9\text{Be}(d, \alpha){}^7\text{Li}$ reaction has the form of a coherent sum of two amplitudes

$$\frac{d\sigma}{d\Omega}(\theta) = |f_I(\theta) + f_{II}(\theta)|^2, \quad (4.10)$$

where the amplitude

$$f_I(\theta) = f_{^5\text{He}}(\pi - \theta) + f_{n-\alpha}(\pi - \theta) + f_{\alpha-n}(\pi - \theta) \quad (4.11)$$

describes the transfer of the heavy ${}^5\text{He}$ -cluster and sequential two-step transfer of $n-\alpha$ and $\alpha-n$, and the amplitude

$$f_{II}(\theta) = f_d(\theta) + f_{n-p}(\theta) + f_{p-n}(\theta) \quad (4.12)$$

corresponds to the deuteron pick-up and sequential two-step transfer of $n-p$ and $p-n$.

The DF potential (see Table 4.6) for the entrance channel and global optical potential parameterizations from Refs. [30, 38, 39] for intermediate and exit channels were used in the analysis. The prior form for the first coupling and the post form for the second coupling were chosen for two-step transfer reactions in order to avoid the non-orthogonal terms in the calculations of transition amplitudes.

The spectroscopic amplitudes of the d and ${}^5\text{He}$ clusters were taken from Ref. [40], while the alpha-cluster spectroscopic amplitudes given in Table 4.7 were provided by Dr. A. Volya within the method reported in Ref. [34].

The calculated cross sections are shown in Fig. 4.19 with the α -particle angular distributions formed in the ${}^9\text{Be}(d, \alpha){}^7\text{Li}^*$ reaction at incident energies of 19.5 and 35 MeV and corresponding to the low-lying excitation of the ${}^7\text{Li}$




FIGURE 4.19: Differential cross sections for the ${}^9\text{Be}(d,\alpha){}^7\text{Li}$ reactions measured at 19.5 MeV and 35 MeV energy with the ${}^7\text{Li}$ observed in the ground or low-lying excited states in the exit channels.

nucleus in the exit channels. The transfer of the deuteron (dash-dotted curve) provides the dominant contribution in all the channels. Despite the fact that the spectroscopic amplitude of the deuteron $S_{1D_3} = 0.558$ in the ${}^9\text{Be}$ nucleus is not of great importance, a noticeable cross section is due to the large value of the deuteron spectroscopic amplitude $S_{1S_1} = 1.732$ of ${}^4\text{He}$.

The angular distribution of deuteron transfer has a significant cross section also at the backward scattering angles, which is mainly caused by the contribution of the D wave. This symmetrical behaviour of the cross section of D waves is very similar to the cross section of evaporation residues. Tanaka *et al* [41] analyzed the role of the compound process in ${}^9\text{Be}(d,\alpha){}^7\text{Li}$ reaction and claimed the domination of the compound nucleus channels at the energies of 12.17 MeV and 14.43 MeV. However, in Ref. [27] the negligible contribution of the compound-nucleus mechanism was shown at 7 MeV using the DWBA analysis. In this regard, our theoretical results based on the CRC method show that there is no need to take into account the mechanism through the compound-nucleus formation at energies of 19.5 and 35.0 MeV.

Starting from scattering angle $\theta_{c.m.} = 120^\circ$, the transfer of the ${}^5\text{He}$ cluster, labeled as ${}^5\text{He}$ in Fig. 4.19, has a predominant contribution in all channels. It




FIGURE 4.20: The scheme illustrates the energy balance of the different intermediate stages for the two-step mechanisms of ${}^9\text{Be}(d, \alpha){}^7\text{Li}$ transfer reaction. The Q -values for the different intermediate channels are shown near the corresponding lines. The numbers near the arrows correspond to the spectroscopic amplitudes of the heaviest reaction participants. For example, spectroscopic amplitude for the ${}^9\text{Be} = {}^8\text{Li} + p$ configuration is equal $S = -0.947$.

should be noted that a similar result was reported earlier in Ref. [27]. One-step transfer of the ${}^5\text{He}$ cluster was also indicated as a dominant process by Jarczyk *et al* [42] in studying the ${}^{12}\text{C}({}^{11}\text{B}, {}^6\text{Li}){}^{17}\text{O}$ and ${}^{12}\text{C}(d, {}^7\text{Li}){}^7\text{Be}$ reactions.

Using the CRC method, we are able to estimate the contribution of the sequential transfer of ${}^5\text{He}$, which was not studied before. Corresponding cross sections are shown in Fig. 4.19 as curves labeled $n\alpha$ and αn . It turned out that the $n\alpha$ and αn transfer processes provide indeed a contribution more than one order of magnitude smaller in comparison with the one-step ${}^5\text{He}$ transfer. Nevertheless, it should be noted that the contribution of the $n\alpha$ and the αn transfer channels increases with the increase in the ${}^7\text{Li}$ excitation energy, where they should not be ignored.

The two-step $n-p$ transfer is another mechanism providing a noticeable contribution to the cross section. It is due to the prominent cluster structure of the ${}^9\text{Be}$ nucleus having the weakly bound neutron. This structural feature explains also the weakness of the $p-n$ sequential transfer contribution to the cross section corresponding to the ${}^7\text{Li}(\text{g.s.})$ in the exit channel. However, with increasing the ${}^7\text{Li}$ excitation energy these two mechanisms are interchanged in the significance of their contributions, as depicted by the curves in Fig. 4.19, and the $p-n$ transfer begins to play a leading role, providing, in particular, almost 10 times larger contribution in the case of reaction at $E_{\text{lab}} = 35$ MeV with ${}^7\text{Li}^*(7.46 \text{ MeV})$ in exit channel.




FIGURE 4.21: Contributions of the different mechanisms to the cross section of the ${}^9\text{Be}(d, \alpha){}^7\text{Li}_{g.s.}$ reaction. See Fig. 4.18 for explanation of the curve notations.

In Fig. 4.20, the possible scenarios for the n - p and p - n sequential transfer for the reaction under consideration are shown in respect to the Q -values. One may see that all the steps of the n - p sequential transfer have positive Q -values, while the p - n transfer goes through the intermediate channel ${}^8\text{Li} + {}^3\text{He}$ that has a considerably negative Q -value. Together with the large values of the spectroscopic amplitudes (shown near to the arrows in Fig. 4.20), this explains the leading role of the $(d, t; t, \alpha)$ mechanism in populating the ground state of ${}^7\text{Li}$ in the exit channel.

The situation becomes quite different in the case of the ${}^7\text{Li}^*(5/2^-)$ in the exit channel. First, the population of this state through the n - p transfer involves the ${}^9\text{Be} = {}^8\text{Be}^*(2^+) + n$ intermediate configuration where the ${}^8\text{Be}$ cluster has to be in the 2^+ excited state. Note that the ${}^8\text{Be}(0^+)$ ground state is inappropriate because of angular-momentum-coupling mismatch in the entrance and exit configurations. Second, the extremely small spectroscopic amplitude of the ${}^8\text{Be}^*(2^+) = {}^7\text{Li}^*(5/2^-) + p$ configuration, which is $\mathcal{S} = 0.079$, influences the transfer amplitude. These two factors lead to the suppression of the contribution of $(d, t; t, \alpha)$ mechanism in population of the ${}^7\text{Li}^*(5/2^-)$ state in the exit channel. Therefore, the p - n sequential transfer prevails over the n - p one.

Figure 4.21 shows the contributions of all the mechanisms mentioned above to the total cross section of the ${}^9\text{Be}(d, \alpha){}^7\text{Li}_{g.s.}$ reaction (see Fig. 4.18) as a function of the deuteron energy. One may conclude that mainly four mechanisms contribute to the cross section of this reaction. The transfer of the deuteron-cluster is the predominant channel at all collision energies. The sequential n - p

and p - n transfers play a significant role at the high energies. The ${}^5\text{He}$ -cluster transfer gives almost 20% of the cross section at low energies and outdoes the sequential p - n transfer in this energy domain. This allows us to claim that the configurations $n+{}^8\text{Be}$ and $\alpha + {}^5\text{He}$ provide noticeable contributions to the ground-state wave function of the ${}^9\text{Be}$ nucleus. These conclusions agree well with the previous experimental studies [43, 44].

Chapter 5

Conclusion

Appendix A

Some aspects from the quantum theory of angular momenta

A total angular momentum \mathbf{j} are decomposed into two angular momenta \mathbf{j}_1 and \mathbf{j}_2 by means of the Clebsch-Gordan coefficient. For example, to quote a basis $|jm\rangle$ with the angular momentum j with its z -component m , the Clebsch-Gordan coefficient can be represented as follow

$$|jm\rangle = \sum_{m_1 m_2} \langle j_1 m_1 j_2 m_2 | jm \rangle |j_1 m_1\rangle |j_2 m_2\rangle, \quad (\text{A.1})$$

For non-zero values of the coefficient (A.1) vectors \mathbf{j}_1 , \mathbf{j}_2 and \mathbf{j} must satisfy the rule of triangle:

$$\begin{aligned} |j_1 - j_2| &\leq j \leq j_1 + j_2 \\ |j - j_2| &\leq j_1 \leq j + j_2 \\ |j_1 - j| &\leq j_2 \leq j_1 + j \end{aligned}$$

and the condition

$$m = m_1 + m_2.$$

If there are three vectors \mathbf{j}_1 , \mathbf{j}_2 and \mathbf{j}_3 , one can get a total angular momentum \mathbf{j} in two ways

$$\mathbf{j} = (\mathbf{j}_1 + \mathbf{j}_2) + \mathbf{j}_3 = \mathbf{j}_{12} + \mathbf{j}_3 \quad (\text{A.2})$$

$$= \mathbf{j}_1 + (\mathbf{j}_2 + \mathbf{j}_3) = \mathbf{j}_1 + \mathbf{j}_{23} \quad (\text{A.3})$$

The Basis $|(j_1 j_2) j_{12}, j_3; jm\rangle$ and the basis $|j_1, (j_2 j_3); jm\rangle$ corresponding to Eq. (A.2) and Eq. (A.3) are related through a factor $U(j_1 j_2 j_3; j_{12} j_{23})$, which

is the Racah coefficient:

$$|(j_1 j_2) j_{12}, j_3; jm\rangle = \sum_{j_{23}} U(j_1 j_2 j_3; j_{12} j_{23}) |j_1, (j_2 j_3); jm\rangle. \quad (\text{A.4})$$

Four angular momenta, \mathbf{j}_1 , \mathbf{j}_2 , \mathbf{j}_3 and \mathbf{j}_4 , are added into the total momentum \mathbf{j} by

$$\mathbf{j} = (\mathbf{j}_1 + \mathbf{j}_2) + (\mathbf{j}_3 + \mathbf{j}_4) = \mathbf{j}_{12} + \mathbf{j}_{34} \quad (\text{A.5})$$

$$= (\mathbf{j}_1 + \mathbf{j}_3) + (\mathbf{j}_2 + \mathbf{j}_4) = \mathbf{j}_{13} + \mathbf{j}_{24} \quad (\text{A.6})$$

Two basis $|j_1 j_2(j_{12}), j_3 j_4(j_{34}); jm\rangle$ and $|j_1 j_3(j_{13}), j_2 j_4(j_{24}); jm\rangle$, constructed respectively on the scheme Eq. A.5 and Eq. A.6, are related as follow

$$|j_1 j_2(j_{12}), j_3 j_4(j_{34}); jm\rangle = \sum_{j_{13}, j_{24}} \begin{bmatrix} j_1 & j_2 & j_{12} \\ j_3 & j_4 & j_{34} \\ j_{13} & j_{24} & j \end{bmatrix} |j_1 j_3(j_{13}), j_2 j_4(j_{24}); jm\rangle \quad (\text{A.7})$$

where transformation coefficient with square brackets is called a unitary 9j-symbol.

A spacial spherical harmonics is expressed like

$$\mathcal{Y}_{lm}(\mathbf{r}) = r^l Y_{lm}(\hat{r}) \quad (\text{A.8})$$

where $Y_{lm}(\hat{r})$ – spherical function, which is a eigenfunction of angular part the $\Delta_{\hat{r}}$ Laplace operator. For $\mathbf{r} = a\mathbf{r}_1 + b\mathbf{r}_2$ a decomposition of the spacial spherical harmonics $\mathcal{Y}_{lm}(\mathbf{r})$ leads to the following equality

$$\begin{aligned} \mathcal{Y}_{lm}(\mathbf{r} = a\mathbf{r}_1 + b\mathbf{r}_2) &= \sum_{l_1, l_2, m_1, m_2} a^{l_1} b^{l_2} \langle l_1 m_1 | l_2 m_2 | lm \rangle \mathcal{D}(l, l_1, l_2) \times \\ &\quad \times \mathcal{Y}_{l_1 m_1}(\mathbf{r}_1) \mathcal{Y}_{l_2 m_2}(\mathbf{r}_2) \\ &= \sum_{l_1, l_2} a^{l_1} b^{l_2} \mathcal{D}(l, l_1, l_2) [\mathcal{Y}_{l_1}(\mathbf{r}_1) \times \mathcal{Y}_{l_2}(\mathbf{r}_2)]_{lm} \end{aligned} \quad (\text{A.9})$$

with the condition $l = l_1 + l_2$, and $\mathcal{D}(l, l_1, l_2)$ is given by

$$\mathcal{D}(l, l_1, l_2) = \sqrt{\frac{4\pi(2l+1)!}{(2l_1+1)!(2l_2+1)!}} \quad (\text{A.10})$$

Spherical harmonics with the argument are coupled as follow

$$[Y_{l_1}(\hat{r}) \times Y_{l_2}(\hat{r})]_{lm} = \mathcal{C}(l_1, l_2, l) Y_{lm}(\hat{r}) \quad (\text{A.11})$$

where the $\mathcal{C}(l_1, l_2, l)$ coefficient reads as

$$\mathcal{C}(l_1, l_2, l) = \sqrt{\frac{(2l_1 + 1)(2l_2 + 1)}{4\pi(2l + 1)}} \langle l_1 0 \ l_2 0 \ | \ l 0 \rangle \quad (\text{A.12})$$

It would be useful also note a coupling between two spherical hyper harmonics kind of

$$\left[Y_{l_{12}}^{(l_1 l_2)}(\hat{\mathbf{r}}_1, \hat{\mathbf{r}}_2) \times Y_{l_{34}}^{(l_3 l_4)}(\hat{\mathbf{r}}_1, \hat{\mathbf{r}}_2) \right]_{lm} = \sum_{l_{13} l_{24}} E_{l_{13} l_{24}}^{l_1 l_2 l_{12} l_2 l_4 l_{34} l} Y_{lm}^{(l_{13} l_{24})}(\hat{\mathbf{r}}_1, \hat{\mathbf{r}}_2) \quad (\text{A.13})$$

where the coupling coefficient $E_{l_{13} l_{24}}^{l_1 l_2 l_{12} l_2 l_4 l_{34} l}$ is given as

$$E_{l_{13} l_{24}}^{l_1 l_2 l_{12} l_2 l_4 l_{34} l} = \begin{bmatrix} l_1 & l_2 & l_{12} \\ l_3 & l_4 & l_{34} \\ l_{13} & l_{24} & l \end{bmatrix} \mathcal{C}(l_1, l_3, l_{13}) \mathcal{C}(l_2, l_4, l_{24}). \quad (\text{A.14})$$

Appendix B

Parameters of the three body wave function

B.1 Helium-6

TABLE B.1: The three-body wave function parameters found by means of variational approach for the ground state of ${}^6\text{He}$

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
$\gamma \equiv 0 \ 0 \ 0$			
1	0.03872262	0.09444994	-0.1737337132E-04
2	0.03872262	0.28926931	-0.7746568943E-01
3	0.03872262	0.50341072	0.1537953138E+00
4	0.03872262	0.75592149	-0.2068222047E+00
5	0.03872262	1.07956780	0.1915099409E+00
6	0.03872262	1.54178260	-0.1186045848E+00
7	0.03872262	2.31514050	0.4377908070E-01
8	0.03872262	4.02900180	-0.8511508780E-02
9	0.03872262	12.33951600	0.8124669663E-03
10	0.11859473	0.09444994	0.1914598067E+00
11	0.11859473	0.28926931	-0.4142283997E-01
12	0.11859473	0.50341072	-0.9504287755E+00
13	0.11859473	0.75592149	0.5100204701E+00
14	0.11859473	1.07956780	0.4351961872E+00
15	0.11859473	1.54178260	-0.5512501566E+00
16	0.11859473	2.31514050	0.2353573699E+00
17	0.11859473	4.02900180	-0.3888384752E-01
18	0.11859473	12.33951600	0.2142226702E-02

Continued on next page

Table B.1 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
19	0.20638850	0.09444994	-0.7243857878E+00
20	0.20638850	0.28926931	0.1811320023E+01
21	0.20638850	0.50341072	-0.2829679245E+00
22	0.20638850	0.75592149	0.6392623958E-01
23	0.20638850	1.07956780	-0.1629089089E+01
24	0.20638850	1.54178260	0.1304993071E+01
25	0.20638850	2.31514050	-0.4928065738E+00
26	0.20638850	4.02900180	0.1996819496E-01
27	0.20638850	12.33951600	0.1060241013E-01
28	0.30991295	0.09444994	0.1599423834E+01
29	0.30991295	0.28926931	-0.5264844147E+01
30	0.30991295	0.50341072	0.7012917232E+01
31	0.30991295	0.75592149	-0.4714686948E+01
32	0.30991295	1.07956780	-0.4796509839E+00
33	0.30991295	1.54178260	0.2065489512E+01
34	0.30991295	2.31514050	-0.8987555045E+00
35	0.30991295	4.02900180	0.4500233344E+00
36	0.30991295	12.33951600	-0.8218053258E-01
37	0.44260156	0.09444994	-0.1956103683E+01
38	0.44260156	0.28926931	0.7193813517E+01
39	0.44260156	0.50341072	-0.1163049706E+02
40	0.44260156	0.75592149	0.6193322640E+01
41	0.44260156	1.07956780	0.7415949805E+01
42	0.44260156	1.54178260	-0.8938587045E+01
43	0.44260156	2.31514050	0.3391887206E+01
44	0.44260156	4.02900180	-0.1168066181E+01
45	0.44260156	12.33951600	0.1744136293E+00
46	0.63210054	0.09444994	0.1402383982E+01
47	0.63210054	0.28926931	-0.5136700789E+01
48	0.63210054	0.50341072	0.8665026533E+01
49	0.63210054	0.75592149	-0.2933774272E+01
50	0.63210054	1.07956780	-0.8943836753E+01
51	0.63210054	1.54178260	0.8717580886E+01
52	0.63210054	2.31514050	-0.3215399907E+01

Continued on next page

Table B.1 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
53	0.63210054	4.02900180	0.1165125187E+01
54	0.63210054	12.33951600	-0.1665816218E+00
55	0.94916212	0.09444994	-0.5455065421E+00
56	0.94916212	0.28926931	0.1973142226E+01
57	0.94916212	0.50341072	-0.3305486410E+01
58	0.94916212	0.75592149	0.5936725650E+00
59	0.94916212	1.07956780	0.4339113002E+01
60	0.94916212	1.54178260	-0.4005833236E+01
61	0.94916212	2.31514050	0.1671843031E+01
62	0.94916212	4.02900180	-0.6442299484E+00
63	0.94916212	12.33951600	0.8553541645E-01
64	1.65181150	0.09444994	0.1291465276E+00
65	1.65181150	0.28926931	-0.3358554818E+00
66	1.65181150	0.50341072	0.6888942468E+00
67	1.65181150	0.75592149	-0.1350120966E+00
68	1.65181150	1.07956780	-0.1153775517E+01
69	1.65181150	1.54178260	0.1271270643E+01
70	1.65181150	2.31514050	-0.7908870481E+00
71	1.65181150	4.02900180	0.2382269089E+00
72	1.65181150	12.33951600	-0.2472769828E-01
73	5.05895900	0.09444994	-0.1003954377E+00
74	5.05895900	0.28926931	-0.1270223362E+00
75	5.05895900	0.50341072	-0.3653418537E+00
76	5.05895900	0.75592149	0.6548391778E+00
77	5.05895900	1.07956780	-0.1837812114E+00
78	5.05895900	1.54178260	0.2724244480E+00
79	5.05895900	2.31514050	0.5022837020E-01
80	5.05895900	4.02900180	-0.1318577464E-01
81	5.05895900	12.33951600	-0.5334531276E-03
$\gamma \equiv 1\ 1\ 1\ 1$			
1	0.03872262	0.09444994	-0.1209912997E-03
2	0.03872262	0.28926931	-0.2752286691E-03
3	0.03872262	0.50341072	0.8313112741E-03

Continued on next page

Table B.1 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
4	0.03872262	0.75592149	-0.1569396728E-02
5	0.03872262	1.07956780	0.1841931183E-02
6	0.03872262	1.54178260	-0.1329878249E-02
7	0.03872262	2.31514050	0.5732118509E-03
8	0.03872262	4.02900180	-0.1407523317E-03
9	0.03872262	12.33951600	0.2464625078E-04
10	0.11859473	0.09444994	0.3780785946E-03
11	0.11859473	0.28926931	-0.1000393773E-01
12	0.11859473	0.50341072	0.1426933985E-01
13	0.11859473	0.75592149	-0.2639755490E-01
14	0.11859473	1.07956780	0.3639070623E-01
15	0.11859473	1.54178260	-0.3096578932E-01
16	0.11859473	2.31514050	0.1507393647E-01
17	0.11859473	4.02900180	-0.4078109925E-02
18	0.11859473	12.33951600	0.7893561124E-03
19	0.20638850	0.09444994	-0.1675476628E-02
20	0.20638850	0.28926931	0.3933745812E-01
21	0.20638850	0.50341072	-0.1092045673E+00
22	0.20638850	0.75592149	0.1542197135E+00
23	0.20638850	1.07956780	-0.1840525972E+00
24	0.20638850	1.54178260	0.1475223686E+00
25	0.20638850	2.31514050	-0.6976541581E-01
26	0.20638850	4.02900180	0.1855170403E-01
27	0.20638850	12.33951600	-0.3521003985E-02
28	0.30991295	0.09444994	0.4106806042E-02
29	0.30991295	0.28926931	-0.9266191014E-01
30	0.30991295	0.50341072	0.2815735552E+00
31	0.30991295	0.75592149	-0.4603450597E+00
32	0.30991295	1.07956780	0.5000169839E+00
33	0.30991295	1.54178260	-0.3928705377E+00
34	0.30991295	2.31514050	0.1830077652E+00
35	0.30991295	4.02900180	-0.4774102277E-01
36	0.30991295	12.33951600	0.7847255538E-02
37	0.44260156	0.09444994	-0.5723743876E-02

Continued on next page

Table B.1 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
38	0.44260156	0.28926931	0.1267185089E+00
39	0.44260156	0.50341072	-0.4013118748E+00
40	0.44260156	0.75592149	0.6809541150E+00
41	0.44260156	1.07956780	-0.7658315705E+00
42	0.44260156	1.54178260	0.5861381351E+00
43	0.44260156	2.31514050	-0.2712619146E+00
44	0.44260156	4.02900180	0.7108163202E-01
45	0.44260156	12.33951600	-0.1022953848E-01
46	0.63210054	0.09444994	0.4677586570E-02
47	0.63210054	0.28926931	-0.1014186442E+00
48	0.63210054	0.50341072	0.3298241305E+00
49	0.63210054	0.75592149	-0.5616842239E+00
50	0.63210054	1.07956780	0.6411713372E+00
51	0.63210054	1.54178260	-0.4946856323E+00
52	0.63210054	2.31514050	0.2242477697E+00
53	0.63210054	4.02900180	-0.6092825450E-01
54	0.63210054	12.33951600	0.8592838651E-02
55	0.94916212	0.09444994	-0.2194397884E-02
56	0.94916212	0.28926931	0.4740184928E-01
57	0.94916212	0.50341072	-0.1572767288E+00
58	0.94916212	0.75592149	0.2605021995E+00
59	0.94916212	1.07956780	-0.2950306986E+00
60	0.94916212	1.54178260	0.2325275547E+00
61	0.94916212	2.31514050	-0.1002393441E+00
62	0.94916212	4.02900180	0.2887819798E-01
63	0.94916212	12.33951600	-0.4542642070E-02
64	1.65181150	0.09444994	0.5992856734E-03
65	1.65181150	0.28926931	-0.1164675896E-01
66	1.65181150	0.50341072	0.4636939204E-01
67	1.65181150	0.75592149	-0.5909598945E-01
68	1.65181150	1.07956780	0.7349403097E-01
69	1.65181150	1.54178260	-0.6253335023E-01
70	1.65181150	2.31514050	0.2300764414E-01
71	1.65181150	4.02900180	-0.6869029310E-02

Continued on next page

Table B.1 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
72	1.65181150	12.33951600	0.1279464637E-02
73	5.05895900	0.09444994	-0.8739688325E-04
74	5.05895900	0.28926931	0.2659208242E-02
75	5.05895900	0.50341072	-0.4297540353E-02
76	5.05895900	0.75592149	0.1984703845E-01
77	5.05895900	1.07956780	-0.1802355036E-01
78	5.05895900	1.54178260	0.1173642602E-01
79	5.05895900	2.31514050	-0.7529136326E-02
80	5.05895900	4.02900180	0.1375626183E-02
81	5.05895900	12.33951600	-0.2653098521E-03
$\gamma \equiv 2\ 2\ 0\ 0$			
1	0.07010120	0.17098674	0.8287413344E-05
2	0.07010120	0.55006725	-0.2315910360E-03
3	0.07010120	1.07956780	0.2205912189E-03
4	0.07010120	2.11877100	-0.1546351743E-03
5	0.07010120	6.81612260	0.1309976070E-03
6	0.22551676	0.17098674	0.1592019751E-03
7	0.22551676	0.55006725	0.1457399081E-02
8	0.22551676	1.07956780	-0.6213676741E-02
9	0.22551676	2.11877100	0.5178724836E-02
10	0.22551676	6.81612260	-0.4694996433E-02
11	0.44260156	0.17098674	-0.3630365442E-03
12	0.44260156	0.55006725	0.5589096347E-03
13	0.44260156	1.07956780	0.9499448616E-02
14	0.44260156	2.11877100	-0.1099890159E-01
15	0.44260156	6.81612260	0.1241336217E-01
16	0.86865448	0.17098674	0.3996664351E-03
17	0.86865448	0.55006725	-0.4801971131E-03
18	0.86865448	1.07956780	-0.9127801910E-02
19	0.86865448	2.11877100	0.6666082171E-02
20	0.86865448	6.81612260	-0.1471088050E-01
21	2.79447630	0.17098674	-0.3154650054E-03
22	2.79447630	0.55006725	-0.8889072435E-04

Continued on next page

Table B.1 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
23	2.79447630	1.07956780	0.1109393782E-01
24	2.79447630	2.11877100	-0.2003915591E-01
25	2.79447630	6.81612260	0.1595658076E-01
$\gamma \equiv 3\ 3\ 1\ 1$			
1	0.07010120	0.17098674	-0.2579335841E-06
2	0.07010120	0.55006725	-0.2765357271E-05
3	0.07010120	1.07956780	0.6139330639E-05
4	0.07010120	2.11877100	-0.7938375153E-05
5	0.07010120	6.81612260	0.1167298934E-04
6	0.22551676	0.17098674	0.5917355271E-06
7	0.22551676	0.55006725	-0.6483646762E-04
8	0.22551676	1.07956780	0.8695083699E-05
9	0.22551676	2.11877100	-0.4800513075E-05
10	0.22551676	6.81612260	0.6570347080E-06
11	0.44260156	0.17098674	-0.1470428346E-05
12	0.44260156	0.55006725	0.1350176334E-03
13	0.44260156	1.07956780	-0.2583695882E-03
14	0.44260156	2.11877100	0.1700235229E-03
15	0.44260156	6.81612260	-0.2196174841E-03
16	0.86865448	0.17098674	0.2122290583E-05
17	0.86865448	0.55006725	-0.1948661774E-03
18	0.86865448	1.07956780	0.3856831336E-03
19	0.86865448	2.11877100	-0.3277457806E-03
20	0.86865448	6.81612260	0.4749778522E-03
21	2.79447630	0.17098674	-0.3460810077E-05
22	2.79447630	0.55006725	0.3075671947E-03
23	2.79447630	1.07956780	-0.6343122836E-03
24	2.79447630	2.11877100	0.6129185220E-03
25	2.79447630	6.81612260	-0.7981418030E-03

B.2 Lithium-6

TABLE B.2: The three-body wave function parameters found by means of variational approach for the ground state of ${}^6\text{Li}$

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
$\gamma \equiv 0\ 0\ 0\ 1$			
1	0.04011531	0.10896291	-0.1803892906E-03
2	0.04011531	0.33559821	0.3207207170E-01
3	0.04011531	0.59133984	-0.3287679135E-01
4	0.04011531	0.90793256	0.1666301953E-02
5	0.04011531	1.34805360	0.2701001430E-01
6	0.04011531	2.06977730	-0.2223424493E-01
7	0.04011531	3.64704510	0.6998916361E-02
8	0.04011531	11.23264600	-0.9092908423E-03
9	0.12355238	0.10896291	-0.1229642431E+00
10	0.12355238	0.33559821	0.4717347384E+00
11	0.12355238	0.59133984	-0.4651643626E+00
12	0.12355238	0.90793256	0.9962817084E+00
13	0.12355238	1.34805360	-0.1276226929E+01
14	0.12355238	2.06977730	0.7795864921E+00
15	0.12355238	3.64704510	-0.2215678668E+00
16	0.12355238	11.23264600	0.2737675477E-01
17	0.21770511	0.10896291	0.2891126276E+00
18	0.21770511	0.33559821	-0.2411541395E+01
19	0.21770511	0.59133984	0.4451709775E+01
20	0.21770511	0.90793256	-0.5916104705E+01
21	0.21770511	1.34805360	0.6633130630E+01
22	0.21770511	2.06977730	-0.3973171424E+01
23	0.21770511	3.64704510	0.1148495732E+01
24	0.21770511	11.23264600	-0.1428310234E+00
25	0.33426051	0.10896291	-0.5014236271E+00
26	0.33426051	0.33559821	0.4520521450E+01
27	0.33426051	0.59133984	-0.1112526104E+02
28	0.33426051	0.90793256	0.1674630659E+02
29	0.33426051	1.34805360	-0.1709289595E+02
30	0.33426051	2.06977730	0.9959388720E+01

Continued on next page

Table B.2 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
31	0.33426051	3.64704510	-0.2904926089E+01
32	0.33426051	11.23264600	0.3617330676E+00
33	0.49629358	0.10896291	0.3691148992E+00
34	0.49629358	0.33559821	-0.4642975913E+01
35	0.49629358	0.59133984	0.1274670370E+02
36	0.49629358	0.90793256	-0.2118020809E+02
37	0.49629358	1.34805360	0.2169847161E+02
38	0.49629358	2.06977730	-0.1253059589E+02
39	0.49629358	3.64704510	0.3772546645E+01
40	0.49629358	11.23264600	-0.4754319741E+00
41	0.76200024	0.10896291	-0.1809029181E+00
42	0.76200024	0.33559821	0.2500421767E+01
43	0.76200024	0.59133984	-0.7609285593E+01
44	0.76200024	0.90793256	0.1339612438E+02
45	0.76200024	1.34805360	-0.1393810450E+02
46	0.76200024	2.06977730	0.8191177607E+01
47	0.76200024	3.64704510	-0.2652360165E+01
48	0.76200024	11.23264600	0.3460395761E+00
49	1.34268030	0.10896291	-0.3393395929E-01
50	1.34268030	0.33559821	-0.8838357055E+00
51	1.34268030	0.59133984	0.2777870668E+01
52	1.34268030	0.90793256	-0.5070333148E+01
53	1.34268030	1.34805360	0.5647602756E+01
54	1.34268030	2.06977730	-0.3354616735E+01
55	1.34268030	3.64704510	0.1188442606E+01
56	1.34268030	11.23264600	-0.1651377670E+00
57	4.13536210	0.10896291	0.1870036658E+00
58	4.13536210	0.33559821	0.4191765900E+00
59	4.13536210	0.59133984	-0.7254042356E+00
60	4.13536210	0.90793256	0.9549950353E+00
61	4.13536210	1.34805360	-0.1622631305E+01
62	4.13536210	2.06977730	0.8926129595E+00
63	4.13536210	3.64704510	-0.3158854492E+00

Continued on next page

Table B.2 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
64	4.13536210	11.23264600	0.4519872066E-01
$\gamma \equiv 2\ 0\ 2\ 1$			
1	0.04589142	0.12465222	-0.4544993203E-05
2	0.04589142	0.38711777	0.1753871978E-03
3	0.04589142	0.69514632	-0.2908374912E-03
4	0.04589142	1.10631900	0.1937616620E-03
5	0.04589142	1.76069660	-0.5504100412E-04
6	0.04589142	3.16167810	-0.2590345143E-05
7	0.04589142	9.81885300	0.3537294375E-05
8	0.14251960	0.12465222	-0.8283263110E-03
9	0.14251960	0.38711777	0.2882961865E-02
10	0.14251960	0.69514632	-0.7044119057E-02
11	0.14251960	1.10631900	0.1208480243E-01
12	0.14251960	1.76069660	-0.8850892215E-02
13	0.14251960	3.16167810	0.3211420819E-02
14	0.14251960	9.81885300	-0.5615393109E-03
15	0.25592205	0.12465222	0.3778904276E-02
16	0.25592205	0.38711777	-0.1265670820E-01
17	0.25592205	0.69514632	0.5268784537E-01
18	0.25592205	1.10631900	-0.7447374680E-01
19	0.25592205	1.76069660	0.5134143603E-01
20	0.25592205	3.16167810	-0.1939583611E-01
21	0.25592205	9.81885300	0.3609459940E-02
22	0.40729761	0.12465222	-0.1850338246E-01
23	0.40729761	0.38711777	0.1964881039E-01
24	0.40729761	0.69514632	-0.9278266295E-01
25	0.40729761	1.10631900	0.1667969445E+00
26	0.40729761	1.76069660	-0.1109233499E+00
27	0.40729761	3.16167810	0.4768747213E-01
28	0.40729761	9.81885300	-0.9728947184E-02
29	0.64821044	0.12465222	0.2452749104E-01
30	0.64821044	0.38711777	-0.2206628080E-01
31	0.64821044	0.69514632	0.1119515408E+00

Continued on next page

Table B.2 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
32	0.64821044	1.10631900	-0.1861487916E+00
33	0.64821044	1.76069660	0.7751512522E-01
34	0.64821044	3.16167810	-0.4867592996E-01
35	0.64821044	9.81885300	0.1231423886E-01
36	1.16398970	0.12465222	-0.7562685391E-01
37	1.16398970	0.38711777	-0.8661044694E-03
38	1.16398970	0.69514632	-0.1641712051E+00
39	1.16398970	1.10631900	0.4712722505E+00
40	1.16398970	1.76069660	-0.2597092534E+00
41	1.16398970	3.16167810	0.8507230018E-01
42	1.16398970	9.81885300	-0.1807666847E-01
43	3.61486630	0.12465222	-0.1151873725E+00
44	3.61486630	0.38711777	-0.4136720190E-01
45	3.61486630	0.69514632	-0.9558240386E-01
46	3.61486630	1.10631900	0.4532504074E+00
47	3.61486630	1.76069660	-0.1845375904E+00
48	3.61486630	3.16167810	0.1043736131E+00
49	3.61486630	9.81885300	-0.1537030856E-01
$\gamma \equiv 1\ 1\ 1\ 0$			
1	0.04589142	0.12465222	-0.4866938229E-04
2	0.04589142	0.38711777	-0.1458515802E-03
3	0.04589142	0.69514632	0.3281186122E-03
4	0.04589142	1.10631900	-0.4226021938E-03
5	0.04589142	1.76069660	0.2998171719E-03
6	0.04589142	3.16167810	-0.1127048329E-03
7	0.04589142	9.81885300	0.2936410508E-04
8	0.14251960	0.12465222	-0.2504898128E-04
9	0.14251960	0.38711777	-0.5265548964E-02
10	0.14251960	0.69514632	0.4963256286E-02
11	0.14251960	1.10631900	-0.5769305333E-02
12	0.14251960	1.76069660	0.4424222636E-02
13	0.14251960	3.16167810	-0.2011073744E-02
14	0.14251960	9.81885300	0.5788519952E-03

Continued on next page

Table B.2 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
15	0.25592205	0.12465222	0.1851282670E-04
16	0.25592205	0.38711777	0.8034383746E-02
17	0.25592205	0.69514632	-0.3170206581E-01
18	0.25592205	1.10631900	0.2216280794E-01
19	0.25592205	1.76069660	-0.1433958645E-01
20	0.25592205	3.16167810	0.7072212353E-02
21	0.25592205	9.81885300	-0.2351274853E-02
22	0.40729761	0.12465222	0.2014357651E-05
23	0.40729761	0.38711777	-0.1019895539E-01
24	0.40729761	0.69514632	0.4134925630E-01
25	0.40729761	1.10631900	-0.4900724944E-01
26	0.40729761	1.76069660	0.2061304370E-01
27	0.40729761	3.16167810	-0.1128969179E-01
28	0.40729761	9.81885300	0.4373172358E-02
29	0.64821044	0.12465222	-0.7329403869E-05
30	0.64821044	0.38711777	0.6398513513E-02
31	0.64821044	0.69514632	-0.3750326707E-01
32	0.64821044	1.10631900	0.4142645560E-01
33	0.64821044	1.76069660	-0.1634689117E-01
34	0.64821044	3.16167810	0.7968526649E-02
35	0.64821044	9.81885300	-0.3264161578E-02
36	1.16398970	0.12465222	0.4470438404E-04
37	1.16398970	0.38711777	0.8876501864E-03
38	1.16398970	0.69514632	0.2859676625E-01
39	1.16398970	1.10631900	-0.2243748552E-01
40	1.16398970	1.76069660	0.8253615321E-02
41	1.16398970	3.16167810	-0.4010945593E-02
42	1.16398970	9.81885300	0.1415435440E-02
43	3.61486630	0.12465222	0.1027736424E-04
44	3.61486630	0.38711777	0.1488674478E-02
45	3.61486630	0.69514632	0.1023799446E-03
46	3.61486630	1.10631900	0.7482735185E-02
47	3.61486630	1.76069660	-0.4434008119E-02
48	3.61486630	3.16167810	0.1560196173E-02

Continued on next page

Table B.2 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
49	3.61486630	9.81885300	-0.8083446767E-03
$\gamma \equiv 2\ 2\ 0\ 1$			
1	0.06450960	0.17522372	-0.9922676129E-06
2	0.06450960	0.56369770	0.1221705852E-03
3	0.06450960	1.10631900	-0.1523134894E-03
4	0.06450960	2.17127340	0.1088605050E-03
5	0.06450960	6.98502340	-0.9061751882E-04
6	0.20752850	0.17522372	-0.1090065045E-03
7	0.20752850	0.56369770	-0.2692287200E-03
8	0.20752850	1.10631900	0.3087213448E-02
9	0.20752850	2.17127340	-0.2277606332E-02
10	0.20752850	6.98502340	0.2119734920E-02
11	0.40729761	0.17522372	0.2079958375E-03
12	0.40729761	0.56369770	-0.2667376133E-02
13	0.40729761	1.10631900	-0.8877499020E-03
14	0.40729761	2.17127340	-0.1511321584E-03
15	0.40729761	6.98502340	-0.2069870206E-02
16	0.79936658	0.17522372	-0.2263976040E-03
17	0.79936658	0.56369770	0.1939224400E-02
18	0.79936658	1.10631900	-0.3396150888E-03
19	0.79936658	2.17127340	0.1016127706E-01
20	0.79936658	6.98502340	-0.2185699277E-02
21	2.57157590	0.17522372	0.1298152166E-03
22	2.57157590	0.56369770	-0.7626896248E-03
23	2.57157590	1.10631900	-0.8197544068E-02
24	2.57157590	2.17127340	-0.7645842537E-03
25	2.57157590	6.98502340	0.7937033880E-02
$\gamma \equiv 2\ 2\ 1\ 1$			
1	0.06450960	0.17522372	-0.3339597522E-05
2	0.06450960	0.56369770	-0.1132330265E-04
3	0.06450960	1.10631900	0.2267914059E-04
4	0.06450960	2.17127340	-0.2352307209E-04

Continued on next page

Table B.2 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
5	0.06450960	6.98502340	0.1834281885E-04
6	0.20752850	0.17522372	-0.5748859294E-05
7	0.20752850	0.56369770	-0.4563431809E-03
8	0.20752850	1.10631900	0.4295222452E-03
9	0.20752850	2.17127340	-0.3213521201E-03
10	0.20752850	6.98502340	0.2805387071E-03
11	0.40729761	0.17522372	0.3533052459E-05
12	0.40729761	0.56369770	0.2970520213E-03
13	0.40729761	1.10631900	-0.1260175766E-02
14	0.40729761	2.17127340	0.6248126780E-03
15	0.40729761	6.98502340	-0.6892394941E-03
16	0.79936658	0.17522372	-0.2664777462E-04
17	0.79936658	0.56369770	-0.9840418266E-03
18	0.79936658	1.10631900	0.2070404093E-02
19	0.79936658	2.17127340	-0.6012080564E-03
20	0.79936658	6.98502340	0.1244787678E-02
21	2.57157590	0.17522372	-0.4021970008E-04
22	2.57157590	0.56369770	-0.2991833581E-03
23	2.57157590	1.10631900	-0.7585129820E-02
24	2.57157590	2.17127340	-0.1618767394E-02
25	2.57157590	6.98502340	-0.1750811298E-02
$\gamma \equiv 2\ 2\ 2\ 1$			
1	0.08101653	0.22006054	-0.1929412768E-05
2	0.08101653	0.73921874	-0.2006956937E-04
3	0.08101653	1.65572340	0.1507761142E-04
4	0.08101653	5.56184130	-0.3426758073E-04
5	0.27214757	0.22006054	0.3125675258E-04
6	0.27214757	0.73921874	-0.1565966223E-03
7	0.27214757	1.65572340	0.3834815879E-03
8	0.27214757	5.56184130	0.1940567272E-03
9	0.60956396	0.22006054	-0.4781842976E-05
10	0.60956396	0.73921874	-0.4525121801E-03
11	0.60956396	1.65572340	-0.1060836916E-01

Continued on next page

Table B.2 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
12	0.60956396	5.56184130	0.5677338934E-02
13	2.04762340	0.22006054	-0.1457667771E-03
14	2.04762340	0.73921874	-0.9277348020E-03
15	2.04762340	1.65572340	-0.8276319924E-02
16	2.04762340	5.56184130	-0.8204815867E-02
$\gamma \equiv 0\ 2\ 2\ 1$			
1	0.10913507	0.29643729	0.1620767206E-03
2	0.10913507	1.10631900	0.7256729573E-02
3	0.10913507	4.12883880	-0.4323400140E-02
4	0.40729761	0.29643729	-0.1575387211E-02
5	0.40729761	1.10631900	-0.1957717882E-01
6	0.40729761	4.12883880	0.9438138297E-02
7	1.52005540	0.29643729	0.1390639775E-02
8	1.52005540	1.10631900	0.6800024355E-02
9	1.52005540	4.12883880	-0.1569399655E-01

B.3 Berillium-9

TABLE B.3: The three-body wave function parameters found by means of variational approach for the ground state of ${}^9\text{Be}$

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
$\gamma \equiv 0 \ 1 \ 1 \ 1$			
1	0.09222844	0.03467231	-0.5749151068E-03
2	0.09222844	0.10767772	-0.3838851470E-02
3	0.09222844	0.19335659	-0.3343703114E-02
4	0.09222844	0.30772526	0.6152031912E-03
5	0.09222844	0.48974194	0.3127639817E-02
6	0.09222844	0.87942826	-0.2202594466E-02
7	0.09222844	2.73113720	0.7301101473E-04
8	0.28642305	0.03467231	-0.1402006121E-03
9	0.28642305	0.10767772	-0.7091214738E-02
10	0.28642305	0.19335659	0.2305680350E-02
11	0.28642305	0.30772526	-0.1742778302E+00
12	0.28642305	0.48974194	-0.3187167848E-01
13	0.28642305	0.87942826	0.3039506091E-01
14	0.28642305	2.73113720	0.9514809808E-02
15	0.51432908	0.03467231	-0.8527376132E-03
16	0.51432908	0.10767772	0.7747996702E-02
17	0.51432908	0.19335659	-0.4446218448E-01
18	0.51432908	0.30772526	0.4402942877E+00
19	0.51432908	0.48974194	-0.2046128930E+00
20	0.51432908	0.87942826	-0.4056323274E+00
21	0.51432908	2.73113720	-0.7813977210E-01
22	0.81855004	0.03467231	0.5375067316E-02
23	0.81855004	0.10767772	0.2796107394E-01
24	0.81855004	0.19335659	0.1235796374E+00
25	0.81855004	0.30772526	-0.4224621555E+00
26	0.81855004	0.48974194	0.6577748985E+00
27	0.81855004	0.87942826	0.9996163672E+00
28	0.81855004	2.73113720	0.1445247127E+00
29	1.30271490	0.03467231	-0.5298458930E-02
30	1.30271490	0.10767772	-0.3527622920E-01

Continued on next page

Table B.3 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
31	1.30271490	0.19335659	-0.1045218882E+00
32	1.30271490	0.30772526	0.2291556567E+00
33	1.30271490	0.48974194	-0.5780951776E+00
34	1.30271490	0.87942826	-0.8020981495E+00
35	1.30271490	2.73113720	-0.6308805304E-01
36	2.33928160	0.03467231	0.1647190294E-02
37	2.33928160	0.10767772	0.1114473473E-01
38	2.33928160	0.19335659	0.3518997183E-01
39	2.33928160	0.30772526	-0.9067262584E-01
40	2.33928160	0.48974194	0.1819309090E+00
41	2.33928160	0.87942826	0.1997706489E+00
42	2.33928160	2.73113720	-0.1672610367E-01
43	7.26483260	0.03467231	-0.1754592338E-03
44	7.26483260	0.10767772	-0.8593955119E-03
45	7.26483260	0.19335659	-0.6094809212E-02
46	7.26483260	0.30772526	0.2033436508E-01
47	7.26483260	0.48974194	-0.2275729417E-01
48	7.26483260	0.87942826	-0.8355648909E-02
49	7.26483260	2.73113720	0.7660503690E-02
$\gamma \equiv 2\ 1\ 1\ 1$			
1	0.09222844	0.03467231	-0.1924771157E-05
2	0.09222844	0.10767772	-0.2356249922E-04
3	0.09222844	0.19335659	-0.2411396428E-04
4	0.09222844	0.30772526	-0.3277817525E-03
5	0.09222844	0.48974194	0.3510077596E-03
6	0.09222844	0.87942826	-0.1042721976E-03
7	0.09222844	2.73113720	0.7300483783E-05
8	0.28642305	0.03467231	-0.1811185889E-04
9	0.28642305	0.10767772	-0.9263076412E-03
10	0.28642305	0.19335659	-0.1355643769E-02
11	0.28642305	0.30772526	-0.1275242738E-02
12	0.28642305	0.48974194	-0.6633639490E-02
13	0.28642305	0.87942826	0.4570012890E-02

Continued on next page

Table B.3 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
14	0.28642305	2.73113720	-0.6773126034E-03
15	0.51432908	0.03467231	-0.2451069527E-04
16	0.51432908	0.10767772	0.2390967055E-02
17	0.51432908	0.19335659	-0.5125262756E-02
18	0.51432908	0.30772526	-0.5418800266E-02
19	0.51432908	0.48974194	-0.2109364349E-01
20	0.51432908	0.87942826	-0.2515442562E-01
21	0.51432908	2.73113720	0.6601216814E-02
22	0.81855004	0.03467231	0.2286563138E-04
23	0.81855004	0.10767772	-0.5363732756E-02
24	0.81855004	0.19335659	0.2315350945E-01
25	0.81855004	0.30772526	-0.1755538546E-01
26	0.81855004	0.48974194	0.5532754524E-01
27	0.81855004	0.87942826	-0.3382279171E-01
28	0.81855004	2.73113720	-0.1186306105E-01
29	1.30271490	0.03467231	0.1042128190E-03
30	1.30271490	0.10767772	0.6603541558E-02
31	1.30271490	0.19335659	-0.1579342851E-01
32	1.30271490	0.30772526	0.3817737886E-01
33	1.30271490	0.48974194	-0.1880665950E-01
34	1.30271490	0.87942826	0.6354958107E-01
35	1.30271490	2.73113720	-0.1023900583E-01
36	2.33928160	0.03467231	-0.1138017603E-03
37	2.33928160	0.10767772	-0.3549492629E-02
38	2.33928160	0.19335659	-0.2172338237E-02
39	2.33928160	0.30772526	-0.1716855378E-01
40	2.33928160	0.48974194	-0.2158352491E-01
41	2.33928160	0.87942826	-0.2664998049E-02
42	2.33928160	2.73113720	0.1795868885E-01
43	7.26483260	0.03467231	0.5481566912E-04
44	7.26483260	0.10767772	0.1616977160E-02
45	7.26483260	0.19335659	0.2999057062E-02
46	7.26483260	0.30772526	0.5941831686E-02
47	7.26483260	0.48974194	0.2509228284E-01

Continued on next page

Table B.3 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
48	7.26483260	0.87942826	-0.2888442185E-01
49	7.26483260	2.73113720	-0.1306911414E-01
$\gamma \equiv 2\ 1\ 2\ 1$			
1	0.09222844	0.03467231	-0.3010378678E-05
2	0.09222844	0.10767772	-0.3755683361E-04
3	0.09222844	0.19335659	-0.1320445807E-04
4	0.09222844	0.30772526	-0.4112501943E-04
5	0.09222844	0.48974194	0.6953669839E-04
6	0.09222844	0.87942826	-0.2860739308E-04
7	0.09222844	2.73113720	0.5423290871E-05
8	0.28642305	0.03467231	-0.1103139569E-04
9	0.28642305	0.10767772	-0.5090367741E-03
10	0.28642305	0.19335659	-0.7148466875E-03
11	0.28642305	0.30772526	-0.3929303844E-02
12	0.28642305	0.48974194	-0.3977209961E-04
13	0.28642305	0.87942826	0.9213700358E-03
14	0.28642305	2.73113720	-0.1948617319E-03
15	0.51432908	0.03467231	-0.4733213692E-04
16	0.51432908	0.10767772	-0.2857738750E-03
17	0.51432908	0.19335659	-0.2777454876E-02
18	0.51432908	0.30772526	0.3561667784E-02
19	0.51432908	0.48974194	-0.2946736188E-01
20	0.51432908	0.87942826	-0.8666890705E-02
21	0.51432908	2.73113720	0.2748423502E-02
22	0.81855004	0.03467231	0.5631914170E-04
23	0.81855004	0.10767772	-0.7971505555E-03
24	0.81855004	0.19335659	0.5371812194E-02
25	0.81855004	0.30772526	-0.2551441441E-01
26	0.81855004	0.48974194	0.4680352343E-01
27	0.81855004	0.87942826	-0.3854700723E-01
28	0.81855004	2.73113720	-0.1222388447E-01
29	1.30271490	0.03467231	0.6206550924E-04
30	1.30271490	0.10767772	0.3834269660E-02

Continued on next page

Table B.3 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
31	1.30271490	0.19335659	0.4162933824E-03
32	1.30271490	0.30772526	0.4735754022E-01
33	1.30271490	0.48974194	-0.1389393548E-01
34	1.30271490	0.87942826	0.8961263346E-01
35	1.30271490	2.73113720	0.8808594255E-02
36	2.33928160	0.03467231	-0.7647542798E-04
37	2.33928160	0.10767772	-0.2890372198E-02
38	2.33928160	0.19335659	-0.3276057155E-02
39	2.33928160	0.30772526	-0.2749917292E-01
40	2.33928160	0.48974194	-0.4710540922E-02
41	2.33928160	0.87942826	-0.5466921396E-01
42	2.33928160	2.73113720	0.2818082044E-02
43	7.26483260	0.03467231	0.3333979438E-04
44	7.26483260	0.10767772	0.1155051008E-02
45	7.26483260	0.19335659	0.1905609943E-02
46	7.26483260	0.30772526	0.9732835265E-02
47	7.26483260	0.48974194	0.2074495567E-02
48	7.26483260	0.87942826	0.1851242373E-01
49	7.26483260	2.73113720	-0.4869370430E-02
$\gamma \equiv 2\ 3\ 1\ 1$			
1	0.10776416	0.04051280	-0.1718969469E-07
2	0.10776416	0.12746398	-0.1350667625E-05
3	0.10776416	0.23612589	-0.1105295982E-04
4	0.10776416	0.40103536	-0.9776430048E-06
5	0.10776416	0.74291449	0.7818043812E-05
6	0.10776416	2.33740540	-0.1657183025E-05
7	0.33905453	0.04051280	0.5619336286E-07
8	0.33905453	0.12746398	0.5297325790E-06
9	0.33905453	0.23612589	0.2429380314E-04
10	0.33905453	0.40103536	-0.7202852868E-03
11	0.33905453	0.74291449	-0.5017077937E-03
12	0.33905453	2.33740540	0.7685878901E-04
13	0.62809554	0.04051280	-0.2882973360E-06

Continued on next page

Table B.3 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
14	0.62809554	0.12746398	-0.1419326948E-04
15	0.62809554	0.23612589	-0.1971653098E-03
16	0.62809554	0.40103536	0.1367614996E-02
17	0.62809554	0.74291449	-0.2329445855E-03
18	0.62809554	2.33740540	-0.1038875830E-02
19	1.06675520	0.04051280	0.5215988170E-06
20	1.06675520	0.12746398	0.2838726456E-04
21	1.06675520	0.23612589	0.5215310433E-03
22	1.06675520	0.40103536	-0.1687394348E-02
23	1.06675520	0.74291449	0.2978139773E-02
24	1.06675520	2.33740540	0.5095314999E-02
25	1.97615460	0.04051280	-0.4079628548E-06
26	1.97615460	0.12746398	-0.1393485092E-04
27	1.97615460	0.23612589	-0.5466031490E-03
28	1.97615460	0.40103536	0.1777350931E-02
29	1.97615460	0.74291449	-0.3110115061E-02
30	1.97615460	2.33740540	-0.6388951126E-02
31	6.21750490	0.04051280	0.2858299794E-06
32	6.21750490	0.12746398	-0.2515569774E-05
33	6.21750490	0.23612589	0.4286470482E-03
34	6.21750490	0.40103536	-0.1506471110E-02
35	6.21750490	0.74291449	0.1647806248E-02
36	6.21750490	2.33740540	0.3849236254E-02
$\gamma \equiv 2\ 3\ 2\ 1$			
1	0.12964559	0.04873889	0.1693294553E-07
2	0.12964559	0.15679385	0.1049895108E-05
3	0.12964559	0.30772526	0.1228675253E-04
4	0.12964559	0.60394482	0.2771362931E-05
5	0.12964559	1.94290080	-0.2211980330E-04
6	0.41707208	0.04873889	0.3551996642E-07
7	0.41707208	0.15679385	0.1075704259E-04
8	0.41707208	0.30772526	0.8745751123E-04
9	0.41707208	0.60394482	0.9594057967E-03

Continued on next page

Table B.3 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
10	0.41707208	1.94290080	0.8746328794E-03
11	0.81855004	0.04873889	0.6352400316E-07
12	0.81855004	0.15679385	-0.1942454226E-04
13	0.81855004	0.30772526	0.1609826431E-04
14	0.81855004	0.60394482	-0.1726785289E-02
15	0.81855004	1.94290080	-0.4042936946E-04
16	1.60649490	0.04873889	-0.1710826173E-06
17	1.60649490	0.15679385	0.2301228629E-05
18	1.60649490	0.30772526	-0.1979072562E-03
19	1.60649490	0.60394482	0.9738317810E-03
20	1.60649490	1.94290080	-0.3203561255E-02
21	5.16812160	0.04873889	0.1116340388E-06
22	5.16812160	0.15679385	0.1167778442E-04
23	5.16812160	0.30772526	0.1738791058E-03
24	5.16812160	0.60394482	-0.3609843531E-03
25	5.16812160	1.94290080	0.5494840873E-02
$\gamma \equiv 4\ 3\ 1\ 1$			
1	0.16281973	0.06121036	0.1321914819E-08
2	0.16281973	0.20561544	-0.4736845147E-06
3	0.16281973	0.46054339	-0.3319521179E-05
4	0.16281973	1.54703930	0.3553888116E-05
5	0.54693765	0.06121036	-0.6503262677E-07
6	0.54693765	0.20561544	0.1654995181E-05
7	0.54693765	0.46054339	-0.1640576015E-03
8	0.54693765	1.54703930	-0.4829795892E-03
9	1.22504670	0.06121036	0.1390571831E-06
10	1.22504670	0.20561544	-0.2891665504E-05
11	1.22504670	0.46054339	0.3281309143E-03
12	1.22504670	1.54703930	0.8592636208E-03
13	4.11512900	0.06121036	-0.1154820611E-06
14	4.11512900	0.20561544	0.2842324272E-05
15	4.11512900	0.46054339	-0.2326745567E-03
16	4.11512900	1.54703930	-0.5225362461E-03

Continued on next page

Table B.3 – continued from previous page

i	$\alpha_i^{(k)}$	$\beta_i^{(k)}$	C_i
$\gamma \equiv 4\ 3\ 2\ 1$			
1	0.16281973	0.06121036	-0.1950076982E-09
2	0.16281973	0.20561544	-0.2964169729E-06
3	0.16281973	0.46054339	-0.1306289078E-05
4	0.16281973	1.54703930	0.1006880888E-05
5	0.54693765	0.06121036	-0.2314866478E-07
6	0.54693765	0.20561544	-0.1222001577E-06
7	0.54693765	0.46054339	-0.1063250475E-03
8	0.54693765	1.54703930	-0.1370765485E-03
9	1.22504670	0.06121036	0.4339252791E-07
10	1.22504670	0.20561544	0.1146087127E-05
11	1.22504670	0.46054339	0.2039624307E-03
12	1.22504670	1.54703930	0.3150677285E-03
13	4.11512900	0.06121036	-0.2831722723E-07
14	4.11512900	0.20561544	-0.1086792529E-05
15	4.11512900	0.46054339	-0.1370131651E-03
16	4.11512900	1.54703930	-0.2779272754E-03

Bibliography

1. Kukulin, V. & Krasnopol'sky, V. A stochastic variational method for few-body systems. *Journal of Physics G: Nuclear Physics* **3**, 795 (1977).
2. Satchler, G. R. & Love, W. G. Folding model potentials from realistic interactions for heavy-ion scattering. *Physics Reports* **55**, 183–254 (1979).
3. De Shalit, A & Talmi, I. *Nuclear shell theory* 1963.
4. Anantaraman, N, Toki, H & Bertsch, G. An effective interaction for inelastic scattering derived from the Paris potential. *Nuclear Physics A* **398**, 269–278 (1983).
5. Day, B. Three-body correlations in nuclear matter. *Physical Review C* **24**, 1203 (1981).
6. Sack, S, Biedenharn, L. & Breit, G. The Elastic Scattering of Protons by Alpha Particles. *Physical Review* **93**, 321 (1954).
7. Buck, B, Friedrich, H & Wheatley, C. Local potential models for the scattering of complex nuclei. *Nuclear Physics A* **275**, 246–268 (1977).
8. Zhukov, M. *et al.* Bound state properties of Borromean halo nuclei: ${}^6\text{He}$ and ${}^{11}\text{Li}$. *Physics reports* **231**, 151–199 (1993).
9. Voronchev, V., Kukulin, V., Pomerantsev, V., Razikov, K. D. & Ryzhikh, G. Study of the Structure and Properties of Nuclei with $A=9$ (${}^9\text{Be}$ - ${}^9\text{B}$) in Terms of Multicenter Dynamic Model $2\alpha + N$. *Yad. Fiz* **57**, 1964–1980 (1994).
10. Pomerantsev, V. Private communication.
11. Antonov, A. *et al.* Charge and matter distributions and form factors of light, medium, and heavy neutron-rich nuclei. *Physical Review C* **72**, 044307 (2005).
12. Egelhof, P *et al.* in *Exotic Nuclei and Atomic Masses* 61–67 (Springer, 2003).
13. Hirai, M, Kumano, S, Saito, K. & Watanabe, T. Clustering aspects in nuclear structure functions. *Physical Review C* **83**, 035202 (2011).

14. Thompson, I. J. Coupled reaction channels calculations in nuclear physics. *Comp. Phys. Rep.* **7**, 167 –212 (1988).
15. Oganessian, Y. T., Zagrebaev, V. & Vaagen, J. Dynamics of two-neutron transfer reactions with the Borromean nucleus ${}^6\text{He}$. *Physical Review C* **60**, 044605 (1999).
16. Karpov, A. *et al.* NRV web knowledge base on low-energy nuclear physics. *Nucl. Instr. and Meth. Phys. Res.t* **A859**, 112 –124 (2017).
17. Lucas, B., Cosper, S. & Johnson, O. Scattering of 18.4-MeV Alpha Particles by Beryllium. *Physical Review* **133**, B963 (1964).
18. Burtebaev, N & et al. *Vopr. At. Nauki Tekh., Ser.: Fiz. Yad. Reakt.* **2002**, 137 (2002).
19. Hauser, G *et al.* Elastic scattering of 104 MeV alpha particles. *Nuclear Physics A* **128**, 81–109 (1969).
20. Avrigeanu, V, Hodgson, P. & Avrigeanu, M. Global optical potentials for emitted alpha particles. *Physical Review C* **49**, 2136 (1994).
21. Harakeh, M., Van Popta, J., Saha, A & Siemssen, R. Strong coupled-channels effects in the ${}^9\text{Be} (\alpha, t) {}^{10}\text{B}$ reaction. *Nuclear Physics A* **344**, 15–40 (1980).
22. Roy, S. *et al.* Coupled channel folding model description of α scattering from Be 9. *Physical Review C* **52**, 1524 (1995).
23. Lukyanov, S. *et al.* Study of internal structures of 9 , ${}^{10}\text{Be}$ and ${}^{10}\text{B}$ in scattering of ${}^4\text{He}$ from ${}^9\text{Be}$. *Journal of Physics G: Nuclear and Particle Physics* **41**, 035102 (2014).
24. Kunz, P. D. *Computer code DWUCK5* unpublished. <https://www.oecd-nea.org/tools/abstract/detail/nesc9872>.
25. Galanina, L. & Zelenskaya, N. Mechanisms of sequential particle transfer and characteristics of light neutron-excess and oriented nuclei. *Physics of Particles and Nuclei* **43**, 147–186 (2012).
26. Urazbekov, B. A., Denikin, A. S., Sakhiev, S. K. & Burtebaev, N. T. Manifestation of the cluster structure of the ${}^9\text{Be}$ nucleus in direct nuclear reactions. *Bull. Rus. Acad. Sci. Phys.* **80**, 247 (2016).
27. Szczerba, A. *et al.* Mechanism of Reactions Induced by 7 MeV Deuterons on ${}^9\text{Be}[(d, p), (d, d), (d, t), (d, {}^4\text{He})]$. *Z. Phys.* **A333**, 271 (1989).

28. Votava, H. J., Clegg, T. B., Ludwig, E. J. & Thompson, W. J. Proton Scattering from ${}^9\text{Be}$ between 6 and 30 MeV and the Structure of ${}^9\text{Be}$. *Nucl. Phys.* **A204**, 529 (1973).
29. Koning, A. J. & Delaroche, J. P. Local and global nucleon optical models from 1 keV to 200 MeV. *Nucl. Phys.* **A713**, 231 (2003).
30. Li, X., Liang, C. & Cai, C. Global triton optical model potential. *Nucl. Phys.* **A789**, 103 (2007).
31. Caurier, E., Martínez-Pinedo, G., Nowacki, F., Poves, A. & Zuker, A. P. The shell model as a unified view of nuclear structure. *Rev. Mod. Phys.* **77**, 427–488 (2 2005).
32. Cohen, S. & Kurath, D. Effective Interactions for the 1p Shell. *Nucl. Phys.* **73**, 1 (1965).
33. Volya, A. *Private communication*.
34. Kravvaris, K. & Volya, A. Study of Nuclear Clustering from an Ab Initio Perspective. *Phys. Rev. Lett.* **119**, 062501 (2017).
35. Della Rocca, V. & Iachello, F. Cluster shell model: I. Structure of ${}^9\text{Be}$, ${}^9\text{B}$. *Nucl. Phys.* **A973**, 1 (2018).
36. Galanina, L. I. & Zelenskaya, N. S. Mechanisms of sequential particle transfer and characteristics of light neutron-excess and oriented nuclei. *Phys. Part. Nucl.* **43**, 147 (2012).
37. Rudchik, A. T. *et al.* Elastic and inelastic scattering of ${}^{15}\text{N}$ ions by ${}^9\text{Be}$ at 84 MeV. *Nucl. Phys.* **A947**, 161 (2016).
38. Avrigeanu, V., Hodgson, P. E. & Avrigeanu, M. Global Optical Potentials for Emitted Alpha Particles. *Phys. Rev.* **C49**, 2136 (1994).
39. Cook, J. Global Optical-Model Potentials for the Elastic Scattering of ${}^{6,7}\text{Li}$ Projectiles. *Nucl. Phys.* **A388**, 153 (1982).
40. Kwasniewicz, E. & Jarczyk, L. Five-Nucleon Spectroscopic Amplitudes for 1p-Shell Nuclei. *Nucl. Phys.* **A441**, 77 (1985).
41. Tanaka, S. The ${}^9\text{Be}(\text{d}, \text{t}){}^8\text{Be}$ and ${}^9\text{Be}(\text{d}, \alpha){}^7\text{Li}$ Reaction in the Energy Range from 12.17 MeV to 14.43 MeV. *J. Phys. Soc. Jpn.* **44**, 1405 (1978).
42. Jarczyk, L. *et al.* Five-Nucleon Simultaneous and Sequential Transfer in the ${}^{12}\text{C}({}^{11}\text{B}, {}^6\text{Li}){}^{17}\text{O}$ and ${}^{12}\text{C}(\text{d}, {}^7\text{Li}){}^7\text{Be}$ Reactions. *Phys. Rev.* **C54**, 1302 (1996).

43. Brown, T. A. D. *et al.* Decay studies for states in ${}^9\text{Be}$ up to 11 MeV: Insights into the $n + {}^8\text{Be}$ and $\alpha + {}^5\text{He}$ cluster structure. *Phys. Rev.* **C76**, 054605 (2007).
44. Papka, P. *et al.* Decay path measurements for the 2.429 MeV state in ${}^9\text{Be}$: Implications for the astrophysical $\alpha + \alpha + n$ reaction. *Phys. Rev.* **C75**, 045803 (2007).