



All chemical reactions are affected by thermal fluctuations and are stochastic



1. Reactants diffuse to find each other in solution
2. They must overcome the energy barrier of the reaction

Both events are randomly affected by thermal fluctuations: collisions with other molecules.



How should we quantify stochasticity?



Noise is often defined as the coefficient of variation, which is the typical size of a fluctuation relative to the mean:

$$\text{noise} = \frac{\text{standard deviation}}{\text{mean}}$$


Stochasticity is more substantial at low numbers. Why?



Why is stochasticity only substantial when typical numbers of molecules are low?

As a reaction only increases or decreases the number of molecules by one or two, it is only when numbers of molecules are small that stochasticity – the random timing of individual reactions – matters.

Stochasticity can be exploited: persister cells enable a population to be both invasive and tolerant to drugs




probability of being a persister is less than  $10^{-5}$  for *E. coli*

Lewis, *Nat Rev Microbiol* 1977

Stochasticity affects the reliability of biochemical networks by affecting timing and is therefore regulated away

e.g. biological rhythms




Elowitz & Leibler, *Nature* 2000  
Barkai & Leibler, *Nature* 2000

## Probing Gene Expression in Live Cells, One Protein Molecule at a Time


Ji Yu,<sup>1\*</sup> Jie Xiao,<sup>1\*</sup> Xiaojia Ren,<sup>1</sup> Kaiqin Lao,<sup>2</sup> X. Sunney Xie<sup>1†</sup>

Following expression of a fluorescent membrane protein in bacteria over time.



Translation can occur in bursts

Occasionally, one mRNA is transcribed.



Bursts of translated protein.




Yu *et al.*, Science (2006)

Transcription can occur in bursts

# Real-Time Kinetics of Gene Activity in Individual Bacteria

Ido Golding,<sup>1,\*</sup> Johan Paulsson,<sup>2,3</sup> Scott M. Zawilski,<sup>1</sup> and Edward C. Cox<sup>1,\*</sup>


Cell 123, 1025–1036, December 16, 2005




Red: protein  
Green spots: mRNA

scale bar: 1  $\mu\text{m}$

Time course of mRNA numbers: mRNA is produced in bursts



The most common model of gene expression can have both bursts in transcription and translation




To perform stochastic simulations, we typically use the Gillespie algorithm

**Step 1:** choose which reaction will occur

**Step 2:** choose when that reaction will occur

Example: a birth-death process with two reactions



probability of a reaction in time  $\delta t$

$$a_1 \delta t = k \delta t \quad \text{propensity of reaction 1}$$

$$a_2 \delta t = dA \delta t \quad \text{propensity of reaction 2}$$

probability of no reaction

$$P_0(t + \delta t) = P_0(t) \left[ 1 - (a_1 + a_2) \delta t \right] \quad \text{hence} \quad P_0 \sim e^{-(a_1 + a_2)t}$$

probability of a reaction  $i$  at time  $t + \delta t$

$$P_i(t) \delta t = P_0(t) a_i \delta t$$