A

## Virion protein DE
## Virocells (n=21) vs all latent GEVE cells (≥10 UMIs, n=1714)

Dot size ∝ -log₁₀(FDR)

%V vs %L

| Gene (GEVEv2_) | log₂ fold-change | %V vs %L |
|---|---|---|
| 80 | *** | 52% vs 0% |
| 68 | *** | 24% vs 0% |
| 63 | *** | 38% vs 0% |
| 447 | *** | 38% vs 0% |
| 329 | *** | 71% vs 0% |
| 392 | *** | 33% vs 0% |
| 233 | *** | 33% vs 0% |
| 415 | *** | 62% vs 0% |
| 279 | *** | 33% vs 0% |
| 62 | *** | 33% vs 0% |
| 434 | *** | 10% vs 0% |
| 431 | *** | 10% vs 0% |
| 122 | *** | 10% vs 0% |
| 135 | *** | 10% vs 0% |
| 101 | *** | 19% vs 0% |
| 304 | *** | 57% vs 0% |
| 64 | *** | 57% vs 0% |
| 288 | *** | 14% vs 0% |
| 201 | *** | 29% vs 0% |
| 118 | *** | 33% vs 0% |
| 127 | * | 5% vs 0% |
| 152 | *** | 10% vs 0% |
| 39 | *** | 14% vs 0% |
| 132 | ** | 10% vs 0% |
| 88 | *** | 24% vs 0% |
| 121 | * | 5% vs 0% |
| 87 | ** | 10% vs 0% |
| 103 | | 5% vs 0% |
| 348 | | 10% vs 4% |
| 371 | | 0% vs 0% |
| 128 | | 0% vs 0% |
| 83 | | 0% vs 0% |
| 326 | | 0% vs 0% |
| 289 | | 0% vs 0% |
| 476 | | 0% vs 0% |
| 61 | | 0% vs 0% |
| 456 | | 38% vs 41% |
| 37 | ** | 29% vs 62% |
| 284 | | 0% vs 0% |
| 265 | | 0% vs 0% |
| 426 | | 0% vs 0% |
| 448 | | 0% vs 0% |

Higher in Virocell (FDR<0.05, n=27)
Higher in all latent GEVE cells (FDR<0.05, n=1)
Not significant

−10 −5 0 5 10

log₂ fold-change in detection rate
(Virocell / all latent GEVE cells)

B

**Functional category**

Unknown
Transcription, replication
Structural
Mobile elements
Metabolism, miscellaneous

0 2 4 6 8 10 12 14 16 18

Number of genes