Beyond the average phenotype: individual variation alters mechanistic forecasts of climate responses

Urtzi Enriquez-Urzelai[1*], Gioele Pascucci[1,2], and Lumír Gvoždík[1]

[1]Czech Academy of Sciences, Institute of Vertebrate Biology, Brno, Czech Republic

[2]Department of Botany and Zoology, Faculty of Science, Masaryk University, Brno, Czech Republic

E-mails: UEU, uenriquez@proton.me; GP, gioelepascucci@mail.muni.cz; LG, gvozdik@ivb.cz

[*]Corresponding author

Running title: Individual variation alters forecasts

Keywords: biophysical modelling; climate change; Dynamic Energy Budget theory; ecological forecasting; individual variation; intraspecific variation; Jensen's inequality; nonlinear averaging; performance surface

Type of article: Letter

Word counts: abstract, 147; main text, 3945; text boxes, none

Number of references: 73

Number of figures, tables and text boxes: 4 figures; no tables; no text boxes

Correspondence: Urtzi Enriquez-Urzelai, Institute of Vertebrate Biology, Czech Academy of Sciences, Brno, Czech Republic. E-mail: uenriquez@proton.me

Statement of authorship: UEU and LG designed the study. LG provided the individual growth and metabolic data. UEU estimated the DEB parameters, performed the simulations and analyses, and produced the figures. UEU wrote the first draft of the manuscript, and all authors contributed substantially to revisions.

Data accessibility statement: All data and code supporting the results are archived in Zenodo (https://doi.org/10.5281/zenodo.23738540; Enriquez-Urzelai et al. 2026). The individual growth and metabolic data were originally published by Baškiera & Gvoždík (2021).

Abstract

Individuals differ in physiology, and organismal performance depends nonlinearly on both phenotype and environment. Jensen's inequality therefore predicts that the performance of an average individual differs from the average performance of a population, yet how this bias varies across traits and environments has rarely been tested with measured individual variation. We estimated Dynamic Energy Budget parameters for 37 individually monitored alpine newts (*Ichthyosaura alpestris*) and for their population average, and coupled them with biophysical models to predict twelve performance traits across Europe under current climate and +2 °C and +4 °C warming. Average-phenotype predictions diverged systematically from mean individual predictions, with the sign and magnitude of this Jensen bias depending on trait, sex and location; reproduction was the most sensitive. Warming amplified, attenuated or reversed the bias, sometimes non-monotonically, although both approaches agreed on the direction of change. The consequences of individual variation are therefore strongly context dependent.

1 Introduction

Ecologists routinely predict population-level outcomes from mean trait values. Trait-based and mechanistic niche models, for instance, commonly summarise a population or species by an average phenotype (Briscoe et al. 2023; Kearney & Porter 2009; Laughlin & Messier 2015). Yet individuals within populations consistently differ in morphology, physiology, behaviour and life history, and these differences influence resource use, species interactions and demographic rates (Bolnick et al. 2011; Bolnick et al. 2003; Des Roches et al. 2018). Whether an average phenotype adequately represents a variable population depends on how trait variation is translated into performance. When this translation is nonlinear, the performance of the average individual differs from the average performance of individuals, a consequence of Jensen's inequality (Jensen 1906; Ruel & Ayres 1999). Although this principle is well established in theory, how strongly it biases ecological predictions, and how that bias changes across traits and environments, has rarely been quantified with empirically measured individual variation.

Performance emerges from interactions between phenotype and environment. These interactions can be represented as performance surfaces describing how a given component of performance varies across combinations of phenotypes and abiotic conditions (Laughlin & Messier 2015; Li et al. 2021; Fig. 1a). Such surfaces are rarely linear: thermal optima, thresholds and asymmetries are pervasive in ecological systems (Bolnick et al. 2011; Ruel & Ayres 1999). We define Jensen bias as $J = f(\overline{\theta}) - \overline{f(\theta)}$, the difference between performance predicted from the mean phenotype $\overline{\theta}$ and mean performance across phenotypes. To second order, $J \approx -\frac{1}{2} f''(\overline{\theta}) \operatorname{Var}[\theta]$, so bias arises not from trait variation alone but from its interaction with the local curvature of the performance surface (Denny 2017; Denny 2019). Where the surface is concave, the average phenotype overestimates mean performance ($J > 0$); where it is convex, it underestimates it ($J < 0$; Bolnick et al. 2011; Ruel & Ayres 1999). This simple result has three consequences that are rarely considered together (Fig. 1). First, because curvature changes along environmental gradients, the same distribution of phenotypes can produce negligible bias in some environments and substantial bias in others (Ross et al. 2023). Second, because growth, reproduction and

survival need not share the same optima or thresholds (Kontopoulos et al. 2024; Sinclair et al. 2016), different components of performance can be biased by different amounts and even in opposite directions. Third, any environmental change that moves populations to a different region of the surface can amplify, attenuate or reverse the bias (Fig. 1b,c).

Physiology provides a natural setting in which to test these predictions, because it is the mechanistic link between environmental conditions and ecological performance (Seebacher & Franklin 2012), and individuals commonly differ in metabolic rates, thermal sensitivity and energy allocation (Gvoždík 2024; Pettersen & Metcalfe 2024). Mechanistic models make this link explicit by simulating how organisms interact with their microclimates to determine growth, survival and reproduction (Briscoe et al. 2023; Kearney & Porter 2009; Kearney & Porter 2020). These models are increasingly used to forecast responses to climate change and now represent environmental heterogeneity in great detail, yet they almost always represent species by a single average phenotype (but see Enriquez-Urzelai et al. 2020; Kolbe et al. 2010; Riddell et al. 2022). Because their performance surfaces are generated from first principles rather than assumed, mechanistic models allow measured individual variation to be propagated through realistic nonlinearities, while climate warming provides an environmental shift of known magnitude with which to test whether the bias changes as environments change.

Here, we combined longitudinal measurements of growth and metabolism from individually monitored alpine newts (*Ichthyosaura alpestris*; data from Baškiera & Gvoždík 2021) with Dynamic Energy Budget (DEB) and biophysical models to compare predictions from a population-average parameterisation with the mean of predictions from 37 individual parameterisations, under identical environmental simulations across Europe and under warming. We tested three hypotheses: (H1) predictions from the average phenotype diverge systematically from the mean of individual predictions; (H2) the sign and magnitude of this Jensen bias differ among performance components and across environmental gradients, reflecting differences in local performance-surface geometry; and (H3) warming changes Jensen bias in trait- and context-specific ways, including amplification, attenuation and reversal. We found support for all three hypotheses, although the two

parameterisations largely agreed on the direction of the projected climate response.

2 Materials and Methods

2.1 Study system and data

The alpine newt is a temperate amphibian widely distributed across Europe. Juveniles are strictly terrestrial until sexual maturity, and adults spend most of the year on land, outside the breeding season (April–June), where they are predominantly nocturnal or crepuscular (Denoël 2004; Roček et al. 2003). We compiled species-level life-history and growth information from the literature, including lifespan (20 years), snout–vent length (SVL) at metamorphosis (1.83 cm), female and male SVL at sexual maturity (4.9 and 4.4 cm), asymptotic SVL (5.84 cm females, 5.05 cm males), asymptotic body mass (3.16 g females, 2.04 g males), maximum reproductive rate (600 eggs year^{-1}) and published growth trajectories (Gvoždík & Kristín 2017; Miaud et al. 2000). Individual-level data consisted of annual measurements of body mass (g) and standard metabolic rate (mL O_2 h^{-1}) of 37 alpine newts (18 females and 19 males) from a population near Jihlava, Czech Republic (49.2247° N, 16.0579° E; 600 m a.s.l.), monitored from metamorphosis in 2014 until 2018 (Baškiera & Gvoždík 2021). These data were collected in a previous study under the permits and ethical approvals reported therein (Baškiera & Gvoždík 2021); no new animals were sampled for this study.

2.2 DEB parameterisation

DEB theory describes the acquisition, storage and allocation of energy throughout the life cycle (Kooijman 2010). The standard DEB model represents an organism by its structure, reserve, maturity and, in adults, a reproduction buffer, and is governed by 12 primary parameters, each quantifying the rate or efficiency of a specific physiological process (Kooijman 2010; Marques et al. 2018). We estimated parameters by inverse modelling using the covariation method (Lika et al. 2011; Marques et al. 2018), with the AmPtool and DEBtool_M packages in MATLAB (The MathWorks Inc., Natick, MA,

USA), at two levels (Appendix S1). First, a single population-average parameterisation (hereafter, average phenotype) was fitted simultaneously to the species-level information and to all longitudinal observations of the 37 individuals, retained as separate data points. This mirrors standard practice in mechanistic modelling and avoids averaging nonlinear trajectories into a composite that no individual follows. We estimated seven primary parameters, namely the zoom factor z (which scales maximum size and surface-specific assimilation; Sousa et al. 2008), energy conductance $\dot{v}$, allocation fraction κ, volume-specific somatic maintenance $[\dot{p}_M]$, specific cost of structure $[E_G]$ and the maturity thresholds at birth E_H^b and puberty E_H^p, together with a threshold at metamorphosis and a male-specific zoom factor (z_m) and puberty threshold ($E_{H,m}^p$); the remaining parameters were fixed at standard values (Kearney et al. 2013; Kooijman 2010).

Second, we re-estimated three parameters independently for each individual from its own growth and metabolic trajectories: the sex-specific zoom factor (z or z_m), $\dot{v}$ and $[\dot{p}_M]$. These parameters govern the processes captured by the individual data, i.e., energy uptake and body size, reserve mobilisation and maintenance costs (Kooijman et al. 2021). All other parameters were held at their population-average values, and all individuals were assigned the same temperature (12 °C) and food availability during estimation, so that differences among parameter sets reflect differences in growth and metabolism rather than in the conditions experienced. Parameter values are given in Table S2.1.

2.3 Mechanistic simulations

We coupled each DEB parameterisation with biophysical models using the NicheMapR R package (Kearney & Porter 2017; Kearney & Porter 2020). The biophysical model computes hourly body temperature, activity and foraging time from microclimatic conditions and the organism's morphology, physiology and behaviour (Kearney et al. 2018; Porter et al. 1973). Body temperature then scales all DEB rates through the Arrhenius relationship and, with food availability held constant, foraging time determines energy intake (Kearney 2012; Kearney & Porter 2020). Hourly microclimates were computed with `micro_global()` from the 10 arc-minute climatology of New et al. (2002), for 0%

and 90% shade, with moisture and snow routines enabled, for 22 years, of which the first two were discarded as spin-up (Kearney 2020; Kearney & Maino 2018; Kearney et al. 2014).

The `ectotherm()` model was parameterised for a terrestrial newt with 90% of the skin acting as a free water exchanger (Enriquez-Urzelai & Gvoždík 2024). Animals were active at night and at dusk (juveniles also by day; Enriquez-Urzelai et al. 2020) when body temperature was between 4 °C and 20 °C and relative humidity was at least 65% (Bonnefond et al. 2020; Giacometti & Tattersall 2025), selecting shade (0–90%) to approach the preferred body temperature of 17.3 °C (Gvoždík & Kristín 2017); otherwise they retreated underground to depths (2.5 cm–2 m) that kept them within critical thermal limits (−2 °C and 36 °C; Enriquez-Urzelai & Gvoždík 2024). Females reproduced in batches of 200–600 eggs (Pecquerie et al. 2009), whereas males depleted their reproduction buffer continuously during the breeding season. Simulations started with juveniles on 15 August and ran until death or for up to 20 years, the maximum known lifespan (Appendix S1).

Simulations were run for every terrestrial cell of a 10 arc-minute grid covering Europe (10° W–40° E, 35°–70° N), for the female and male average phenotypes and for each of the 37 individual parameterisations. The whole framework was repeated under uniform increases of +2 °C and +4 °C in hourly air temperature, which span projected warming across Europe under intermediate to high emission pathways (Calvin et al. 2023). All other climatic variables were held constant, isolating the thermal effects of warming; because relative humidity was unchanged, warming also increased evaporative demand. Thus, within each grid cell and scenario, predictions differed only in the physiological parameterisation.

2.4 Quantifying Jensen bias

For every grid cell, parameterisation and scenario, we extracted 12 performance metrics: lifespan (year), lifetime reproductive output (eggs in females, hereafter lifetime fecundity; reproductive energy, J, in males), net reproductive rate (R_0), maximum and minimum annual body temperature ($T_{b,max}$, $T_{b,min}$), maximum body mass (g), maximum SVL (cm),

minimum reserve energy (J), and lifetime basking time (h; time above ground without foraging), foraging time (h), food intake (g dry mass) and mean daily water need (g day^{-1}). For each metric and sex, Jensen bias was computed per grid cell as

$$J = P_{\text{pop}} - \overline{P}_{\text{ind}},$$

where P_{pop} is the prediction of the average phenotype and $\overline{P}_{\text{ind}}$ the mean of the individual predictions. Positive values indicate overprediction by the average phenotype and negative values underprediction. To compare metrics with different units, we used the normalised absolute bias: the mean $|J|$ across grid cells divided by the mean $|\overline{P}_{\text{ind}}|$ ($\times 100$). Independently of bias, we quantified among-individual divergence in predictions as the coefficient of variation (CV) of individual predictions in each cell, and compared its median between southern ($<40^\circ$ N), central and northern ($>60^\circ$ N) latitudes.

To test how warming modified bias, we computed $\Delta|J| = |J_{\text{future}}| - |J_{\text{current}}|$, where positive values indicate amplification and negative values attenuation, and summarised it per metric as the relative change, i.e., the mean $\Delta|J|$ divided by the mean $|J_{\text{current}}|$ ($\times 100$). We also report the proportions of cells in which $|J|$ increased or decreased and in which the sign of J reversed. We described how $\Delta|J|$ under $+4\,^\circ$C varied along mean annual air temperature ($\overline{T}_a$) using generalised additive models fitted with `mgcv` (Wood 2025); because model outputs are deterministic and spatially autocorrelated, these served as descriptive smoothers rather than for inference (Appendix S1).

Finally, for three traits summarising survival, growth and reproduction (lifespan, maximum body mass and lifetime fecundity or reproductive energy), we compared predictions of both parameterisations under each scenario and their projected changes ($P_{\text{future}} - P_{\text{current}}$). We report the percentage of cells in which the average phenotype under- or overpredicted the individual mean, the percentage in which both projected the same direction of change, and the median and 95th percentile of absolute relative bias, $(P_{\text{pop}} - \overline{P}_{\text{ind}})/|\overline{P}_{\text{ind}}| \times 100$. All data and code are available in Zenodo (Enriquez-Urzelai et al. 2026).

3 Results

3.1 Individual physiological variation

Individual DEB parameters varied among the 37 individuals, with coefficients of variation ranging from 1.7% (z) to 8% ($[\dot{p}_M]$; Fig. S2.1). Somatic maintenance $[\dot{p}_M]$ was the most variable (CV = 8%), followed by energy conductance $\dot{v}$ (CV = 5.6%) and the zoom factors (z: 1.7%; z_m: 1.9%), and each parameter was approximately normally distributed around the population-average estimate (Fig. S2.2). Both parameterisations reproduced the observed growth and metabolic trajectories accurately (Figs. S2.3, S2.4, S2.5–S2.14).

3.2 Physiological averaging generates Jensen bias (H1, H2)

Under the current climate, predictions of the average phenotype differed systematically from the mean of individual predictions for most metrics. For females, the average phenotype underestimated most metrics (Fig. 2), including basking time ($|J| = 419\,\mathrm{h}$), foraging time ($|J| = 143\,\mathrm{h}$), lifetime fecundity ($|J| = 63\,\mathrm{eggs}$), food intake ($|J| = 3.7\,\mathrm{g}$) and maximum body mass ($|J| = 0.2\,\mathrm{g}$; absolute value of the mean J across cells). Relative to the individual mean, bias was largest for R_0 (normalised $|J| = 13.3\%$), lifetime fecundity (10.1%) and maximum body mass (5.3%), and negligible for body temperatures ($\leq 0.2\%$; mean $|J| < 0.004\,^{\circ}\mathrm{C}$). For males, the pattern was largely opposite: the average phenotype overestimated most metrics, with the largest relative bias for reproductive energy (normalised $|J| = 9.1\%$) and R_0 (7.4%; Fig. S3.1, Appendix S3).

The direction of bias differed among metrics. In females, underestimation dominated across most of the range for body size, reserve, food intake, lifetime fecundity and basking ($J < 0$ in 63–98% of cells), whereas body temperatures were predominantly overestimated ($J > 0$ in 65–83% of cells), and foraging time was split between overestimation (52% of cells), underestimation (46%) and no bias (2%). In males, these directions were reversed ($J > 0$ in 57–96% of cells for the same metrics).

Among-individual divergence in predictions was also geographically structured (Figs. S3.2, S3.3). Compared with central latitudes, the median CV of individual predictions in females

was 3.3-fold higher at the southern margin for lifetime fecundity, 3.1-fold for minimum reserve and 3.4-fold for maximum body length. In males, the largest increases occurred at the northern margin for R_0 (4.3-fold) and reproductive energy (2.2-fold). In contrast, lifespan, basking time, foraging time and body temperatures differed by at most 1.5-fold between marginal and central latitudes in either sex.

3.3 Climate warming reshapes Jensen bias (H3)

Under +4 °C warming, $|J|$ changed for all 12 female metrics (Fig. 3). The strongest relative amplification occurred for minimum body temperature (rel. $\Delta|J| = +568\%$; increase in 72% of cells), maximum body temperature (rel. $\Delta|J| = +182\%$; 73% of cells) and R_0 (rel. $\Delta|J| = +138\%$), whereas lifespan (-15%) and minimum reserve (-6%) were attenuated. The large relative changes in body temperature bias, however, arose from very small absolute biases (mean $|J|$ from 0.0021 to 0.014 °C for $T_{b,min}$ and from 0.0035 to 0.0099 °C for $T_{b,max}$). In males, the strongest amplification occurred for minimum body temperature (rel. $\Delta|J| = +510\%$) and maximum body temperature (rel. $\Delta|J| = +215\%$), whereas lifespan (-1%) was attenuated (Fig. S3.4).

Warming also reversed the direction of bias in some regions. The sign of J changed between current and +4 °C scenarios in 2–49% of cells in females, most often for minimum body temperature (49%), foraging time (47%) and R_0 (44%), and in 5–46% of cells in males. Changes in bias were geographically structured and varied systematically, but in metric-specific ways, along the mean annual temperature gradient (Fig. 3). Responses were not always monotonic: under +2 °C, 11 female metrics showed increased $|J|$ (from +4% for daily water need to +152% for minimum body temperature), and lifespan switched from amplification at +2 °C to attenuation at +4 °C (see Appendix S3 for males).

3.4 Consequences for forecasts

The average phenotype underestimated female and overestimated male performance for several traits, with differences in the magnitude and geographical extent of bias (Figs. 4, S3.5). In females, underprediction of lifetime fecundity became more widespread with

warming, from 63% of cells under the current climate (median relative bias −7.1%) to 71% and 76% under +2 °C and +4 °C. Underpredicted lifespan increased from 76% of cells (median −1.9%) to 86% under +4 °C, while maximum body mass was underpredicted in 83% of cells (median −3.8%). Across 95% of cells, the average phenotype underestimated female lifespan by up to 19% and lifetime fecundity by up to 100%, compared with ±19% for body mass (Fig. S3.6). In males, reproductive energy was overpredicted in 85% of cells under the current climate (median +8.1%) and 95% under +4 °C, by up to 66% of the individual mean (Fig. S3.7). Despite these differences, both parameterisations projected the same direction of change under +4 °C in at least 90% of cells for all three traits in both sexes.

4 Discussion

Jensen's inequality implies that the ecological consequences of individual variation depend on the shape of the relationship between phenotype, environment and performance (Bolnick et al. 2011; Ruel & Ayres 1999). By propagating measured physiological variation through mechanistic performance surfaces, we show that this principle has large, structured consequences for predictions: an average phenotype misrepresented the mean performance of a heterogeneous population for most traits (H1), the bias differed in magnitude and sign among traits, sexes and locations (H2), and warming reshaped it in trait-specific and sometimes non-monotonic ways (H3). Notably, variation in only three energetic parameters was sufficient to generate these patterns at continental scale, extending previous work on nonlinear responses to environmental variability (Denny 2017; Paaijmans et al. 2013) to variability among individuals.

Which traits are sensitive to averaging

Reproductive traits showed the largest relative bias in both sexes, whereas body mass was less sensitive. This difference is consistent with how traits integrate energy over the life cycle. Reproduction reflects acquisition, allocation and maturation history throughout

adulthood (Jusup et al. 2017; Kooijman 2010), so small physiological differences accumulate through time and are amplified by thresholds such as the energy required for a clutch. Body mass, in contrast, is constrained by tight feedbacks between assimilation and maintenance that converge on an asymptotic size (Lika et al. 2019; Nisbet et al. 2012), buffering the effect of individual variation. Traits that integrate cumulative allocation are therefore expected to be more sensitive to averaging than structural traits (Nisbet et al. 2012; Palmer et al. 2024), a prediction that should apply beyond our study system. The opposite direction of bias in females and males further shows that the sign of J is not a property of the trait alone: batch reproduction in females converts energetic differences into discrete clutches, whereas continuous depletion of the reproduction buffer in males smooths them (Muller et al. 2019; Pecquerie et al. 2009), so that the same parameter variation meets performance surfaces of different curvature. Likewise, foraging time switched between over- and underestimation across the range, indicating that local environmental conditions can determine the direction of bias (Martin & Huey 2008).

Where averaging matters

The divergence among individual predictions increased towards the northern and southern margins of the modelled area, particularly for reproduction- and reserve-related traits. This is consistent with the expectation that the consequences of trait variation depend on where populations operate on the performance surface, which changes along environmental gradients (Laughlin & Messier 2015; Li et al. 2021; Polazzo et al. 2024; Ross et al. 2023). Under benign conditions, individuals with different physiologies may achieve similar outcomes, whereas under constraining conditions small differences can determine whether energy suffices for growth, reserves or reproduction. Although the strongest effects occurred partly beyond the current range of *I. alpestris* (IUCN 2021), they emerged precisely where ecological responses are expected to become most sensitive, as populations approach environmental limits (Holt 2003; Sexton et al. 2009; Valladares et al. 2014). Predictions based on average phenotypes are therefore least reliable where predicting persistence, range expansion and local extinction matters most (DeMarche et al. 2019).

Environmental change reshapes the bias

Warming changed Jensen bias for every metric, but not uniformly. It amplified the bias in net reproductive rate in both sexes, the most demographically relevant change, attenuated it for lifespan, and reversed its sign across large parts of the range for some metrics. For lifespan, bias increased under +2 °C but decreased under +4 °C, showing that the effect of individual variation does not scale monotonically with environmental change. These responses are expected when warming moves populations across regions of a nonlinear surface with different curvature (Angilletta Jr. 2009; Kingsolver & Woods 2016; Li et al. 2021): further warming can move a population past an inflection point, weakening rather than strengthening the effect of averaging. Hence, whether an average phenotype over- or underestimates performance is a property of the phenotype distribution, the trait and the environment jointly, and validation of average-phenotype predictions under current conditions does not guarantee their accuracy under future conditions (Buckley et al. 2018; Buckley et al. 2023; Cannistra & Buckley 2024).

Robust trends, biased magnitudes

Despite these biases, both parameterisations agreed on the direction of the projected response in at least 90% of cells: warming reduced lifespan across most of the range (Burraco et al. 2020; Munch & Salinas 2009), and both projected reduced body mass and reproduction towards the north-eastern range limit, where warming improves thermal suitability. Physiological averaging therefore affected quantitative predictions, i.e., the magnitude and spatial pattern of change, without overturning the qualitative interpretation. This distinction between robust trends and biased magnitudes clarifies when average phenotypes are adequate: for the direction of change, our results suggest they generally are, whereas predictions of absolute performance, and particularly of reproduction and population growth, should incorporate individual variation. Mechanistic models are well suited to this task because they represent physiology explicitly and have already been used to explore trait evolution and thermal adaptation using observed or hypothetical variation (De Jong et al. 1996; Enriquez-Urzelai et al. 2020; Kingsolver & Woods 2016;

Riddell et al. 2023; Riddell et al. 2022). Propagating empirical trait distributions requires that individual-level estimates, not only population means, are reported and archived (Wolf & McKechnie 2023).

Several simplifications point to future work. We allowed only three parameters to vary, used individuals from a single population and held trait distributions fixed under warming, whereas variation in other traits, among-population differentiation and plasticity or adaptation are all expected to shape trait distributions across ranges and through time (Chevin et al. 2010; Kingsolver & Buckley 2017; Moran et al. 2016; Sexton et al. 2009). Each of these could amplify, attenuate or reverse the patterns of Jensen bias reported here, and integrating them is an important next step towards biologically realistic predictions.

In conclusion, representing a population by a single average phenotype introduced systematic, spatially structured and climate-dependent biases in predicted performance, because individual variation interacts with nonlinear phenotype–environment–performance relationships. The bias was largest for traits that integrate energy allocation over the life cycle and under environmentally constraining conditions, and it could be amplified, attenuated or reversed by environmental change. More generally, our results show that the ecological consequences of individual variation cannot be inferred from the amount of variation alone, but depend on where populations lie on the underlying performance surface, which is a principle that applies to any prediction made from mean traits in a changing environment.

Acknowledgements

We thank Bas Kooijman and Gonçalo Marques for help with DEB parameter estimation, and Michael R. Kearney for help setting up the sex-specific NicheMapR simulations. Computational resources were provided by the e-INFRA CZ project (ID:90254), supported by the Ministry of Education, Youth and Sports of the Czech Republic. Funding was provided by the Czech Science Foundation (25-15551S to UEU, GP and LG), the Department of Botany and Zoology, Faculty of Science, Masaryk University in Brno,

Czechia (MUNI/A/1615/2025 to GP), and the Institute of Vertebrate Biology of the Czech Academy of Sciences (RVO: 68081766 to LG). The authors declare no conflicts of interest.

5 References

Angilletta Jr., M. J. (2009). *Thermal Adaptation*. Oxford University Press.

Baškiera, S. & Gvoždík, L. (2021). Repeatability and Heritability of Resting Metabolic Rate in a Long-Lived Amphibian. Comp. Biochem. Physiol. A. Mol. Integr. Physiol., 253, 110858.

Bolnick, D. I., Amarasekare, P., Araújo, M. S., Bürger, R., Levine, J. M., Novak, M., et al. (2011). Why Intraspecific Trait Variation Matters in Community Ecology. Trends Ecol. Evol., 26, 183–192.

Bolnick, D. I., Svanbäck, R., Fordyce, J. A., Yang, L. H., Davis, J. M., Hulsey, C. D., et al. (2003). The Ecology of Individuals: Incidence and Implications of Individual Specialization. Am. Nat., 161, 1–28.

Bonnefond, A., Courtois, E. A., Sueur, J., Sugai, L. S. M. & Llusia, D. (2020). Climatic Breadth of Calling Behaviour in Two Widespread Neotropical Frogs: Insights from Humidity Extremes. Glob. Change Biol., 26, 5431–5446.

Briscoe, N. J., Morris, S. D., Mathewson, P. D., Buckley, L. B., Jusup, M., Levy, O., et al. (2023). Mechanistic Forecasts of Species Responses to Climate Change: The Promise of Biophysical Ecology. Glob. Change Biol., 29, 1451–1470.

Buckley, L. B., Cannistra, A. F. & John, A. (2018). Leveraging Organismal Biology to Forecast the Effects of Climate Change. Integr. Comp. Biol., 58, 38–51.

Buckley, L. B., Carrington, E., Dillon, M. E., García-Robledo, C., Roberts, S. B., Wegrzyn, J. L., et al. (2023). Characterizing Biological Responses to Climate Variability and Extremes to Improve Biodiversity Projections. PLOS Clim., 2, e0000226.

Burraco, P., Orizaola, G., Monaghan, P. & Metcalfe, N. B. (2020). Climate Change and Ageing in Ectotherms. Glob. Change Biol., 26, 5371–5381.

Calvin, K., Dasgupta, D., Krinner, G., Mukherji, A., Thorne, P. W., Trisos, C., et al. (2023). *IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (Eds.)]. IPCC, Geneva, Switzerland*. Intergovernmental Panel on Climate Change (IPCC).

Cannistra, A. F. & Buckley, L. B. (2024). Accounting for Nonlinear Responses to Traits Improves Range Shift Predictions. Ecosphere, 15, e4860.

Chevin, L.-M., Lande, R. & Mace, G. M. (2010). Adaptation, Plasticity, and Extinction in a Changing Environment: Towards a Predictive Theory. PLOS Biol., 8, e1000357.

De Jong, P. W., Gussekloo, S. W. S. & Brakefield, P. M. (1996). Differences in Thermal Balance, Body Temperature and Activity Between Non-Melanic and Melanic Two-Spot Ladybird Beetles (*Adalia Bipunctata*) Under Controlled Conditions. J. Exp. Biol., 199, 2655–2666.

DeMarche, M. L., Doak, D. F. & Morris, W. F. (2019). Incorporating Local Adaptation into Forecasts of Species' Distribution and Abundance under Climate Change. Glob. Change Biol., 25, 775–793.

Denny, M. (2017). The Fallacy of the Average: On the Ubiquity, Utility and Continuing Novelty of Jensen's Inequality. J. Exp. Biol., 220, 139–146.

Denny, M. (2019). Performance in a Variable World: Using Jensen's Inequality to Scale up from Individuals to Populations. Conserv. Physiol., 7, coz053.

Denoël, M. (2004). Terrestrial*versus*aquatic Foraging in Juvenile Alpine Newts (*Triturus Alpestris*). Écoscience, 11, 404–409.

Des Roches, S., Post, D. M., Turley, N. E., Bailey, J. K., Hendry, A. P., Kinnison, M. T., et al. (2018). The Ecological Importance of Intraspecific Variation. Nat. Ecol. Evol., 2, 57–64.

Enriquez-Urzelai, U. & Gvoždík, L. (2024). Impacts of Behaviour and Acclimation of Metabolic Rate on Energetics in Sheltered Ectotherms: A Climate Change Perspective. Proc. R. Soc. B Biol. Sci., 291, 20232152.

Enriquez-Urzelai, U., Pascucci, G. & Gvoždík, L. (2026). *Data from: Beyond the Average Phenotype: Individual Variation Alters Mechanistic Forecasts of Climate Responses*. Zenodo. Available at: https://doi.org/10.5281/zenodo.23738540. Zenodo.

Enriquez-Urzelai, U., Tingley, R., Kearney, M. R., Sacco, M., Palacio, A. S., Tejedo, M., et al. (2020). The Roles of Acclimation and Behaviour in Buffering Climate Change Impacts along Elevational Gradients. J. Anim. Ecol., 89, 1722–1734.

Giacometti, D. & Tattersall, G. J. (2025). Behavioural Evidence of a Humidistat: A Temperature-Compensating Mechanism of Hydroregulation in Spotted Salamanders. J. Exp. Biol., 228, jeb250297.

Gvoždík, L. (2024). Individual Variation in Thermally Induced Plasticity of Metabolic Rates: Ecological and Evolutionary Implications for a Warming World. Philos. Trans. R. Soc. B Biol. Sci., 379, 20220494.

Gvoždík, L. & Kristín, P. (2017). Economic Thermoregulatory Response Explains Mismatch between Thermal Physiology and Behavior in Newts. J. Exp. Biol., 220, jeb.145573.

Holt, R. D. (2003). On the Evolutionary Ecology of Species' Ranges. Evol. Ecol. Res., 5, 159–178.

IUCN (2021). *Ichthyosaura Alpestris: IUCN SSC Amphibian Specialist Group: The IUCN Red List of Threatened Species 2022: E.T59472A89702213*. International Union for Conservation of Nature.

Jensen, J. L. W. V. (1906). Sur Les Fonctions Convexes et Les Inégalités Entre Les Valeurs Moyennes. Acta Math., 30, 175–193.

Jusup, M., Sousa, T., Domingos, T., Labinac, V., Marn, N., Wang, Z., et al. (2017). Physics of Metabolic Organization. Phys. Life Rev., 20, 1–39.

Kearney, M. (2012). Metabolic Theory, Life History and the Distribution of a Terrestrial Ectotherm. Funct. Ecol., 26, 167–179.

Kearney, M. & Porter, W. (2009). Mechanistic Niche Modelling: Combining Physiological and Spatial Data to Predict Species' Ranges. Ecol. Lett., 12, 334–350.

Kearney, M. R. (2020). How Will Snow Alter Exposure of Organisms to Cold Stress under Climate Warming? Glob. Ecol. Biogeogr., 29, 1246–1256.

Kearney, M. R. & Maino, J. L. (2018). Can Next-Generation Soil Data Products Improve Soil Moisture Modelling at the Continental Scale? An Assessment Using a New Microclimate Package for the R Programming Environment. J. Hydrol., 561, 662–673.

Kearney, M. R., Munns, S. L., Moore, D., Malishev, M. & Bull, C. M. (2018). Field Tests of a General Ectotherm Niche Model Show How Water Can Limit Lizard Activity and Distribution. Ecol. Monogr., 88, 672–693.

Kearney, M. R. & Porter, W. P. (2017). NicheMapR – an R Package for Biophysical Modelling: The Microclimate Model. Ecography, 40, 664–674.

Kearney, M. R. & Porter, W. P. (2020). NicheMapR – an R Package for Biophysical Modelling: The Ectotherm and Dynamic Energy Budget Models. Ecography, 43, 85–96.

Kearney, M. R., Shamakhy, A., Tingley, R., Karoly, D. J., Hoffmann, A. A., Briggs, P. R., et al. (2014). Microclimate Modelling at Macro Scales: A Test of a General Microclimate Model Integrated with Gridded Continental-scale Soil and Weather Data. Methods Ecol. Evol., 5, 273–286.

Kearney, M. R., Simpson, S. J., Raubenheimer, D. & Kooijman, S. A. L. M. (2013). Balancing Heat, Water and Nutrients under Environmental Change: A Thermodynamic Niche Framework. Funct. Ecol., 27, 950–966.

Kingsolver, J. G. & Buckley, L. B. (2017). Evolution of Plasticity and Adaptive Responses to Climate Change along Climate Gradients. Proc. R. Soc. B Biol. Sci., 284, 20170386.

Kingsolver, J. G. & Woods, H. A. (2016). Beyond Thermal Performance Curves: Modeling Time-Dependent Effects of Thermal Stress on Ectotherm Growth Rates. Am. Nat., 187, 283–294.

Kolbe, J. J., Kearney, M. & Shine, R. (2010). Modeling the Consequences of Thermal Trait Variation for the Cane Toad Invasion of Australia. Ecol. Appl., 20, 2273–2285.

Kontopoulos, D. .-. G., Sentis, A., Daufresne, M., Glazman, N., Dell, A. I. & Pawar, S. (2024). No Universal Mathematical Model for Thermal Performance Curves across Traits and Taxonomic Groups. Nat. Commun., 15, 8855.

Kooijman, S. A. L. M., Lika, K., Augustine, S. & Marn, N. (2021). Multidimensional Scaling for Animal Traits in the Context of Dynamic Energy Budget Theory. Conserv. Physiol., 9, coab086.

Kooijman, S. A. L. M. (2010). *Dynamic Energy Budget Theory for Metabolic Organisation.* 3rd ed. Cambridge: Cambridge university press.

Laughlin, D. C. & Messier, J. (2015). Fitness of Multidimensional Phenotypes in Dynamic Adaptive Landscapes. Trends Ecol. Evol., 30, 487–496.

Li, Y., Jiang, Y., Shipley, B., Li, B., Luo, W., Chen, Y., et al. (2021). The Complexity of Trait–Environment Performance Landscapes in a Local Subtropical Forest. New Phytol., 229, 1388–1397.

Lika, K., Augustine, S. & Kooijman, S. A. (2019). Body Size as Emergent Property of Metabolism. J. Sea Res., 143, 8–17.

Lika, K., Kearney, M. R. & Kooijman, S. A. (2011). The "Covariation Method" for Estimating the Parameters of the Standard Dynamic Energy Budget Model II: Properties and Preliminary Patterns. J. Sea Res., 66, 278–288.

Marques, G. M., Augustine, S., Lika, K., Pecquerie, L., Domingos, T. & Kooijman, S. A. L. M. (2018). The AmP Project: Comparing Species on the Basis of Dynamic Energy Budget Parameters. PLOS Comput. Biol., 14, e1006100.

Martin, T. L. & Huey, R. B. (2008). Why "Suboptimal" Is Optimal: Jensen's Inequality and Ectotherm Thermal Preferences. Am. Nat., 171, E102–E118.

Miaud, C., Guyetant, R. & Faber, H. (2000). Age, Size, and Growth of the Alpine Newt, *Triturus Alpestris* (Urodela: Salamandridae), at High Altitude and a Review of Life-History Trait Variation throughout Its Range. Herpetologica, 56, 135–144.

Moran, E. V., Hartig, F. & Bell, D. M. (2016). Intraspecific Trait Variation across Scales: Implications for Understanding Global Change Responses. Glob. Change Biol., 22, 137–150.

Muller, E. B., Lika, K., Nisbet, R. M., Schultz, I. R., Casas, J., Gergs, A., et al. (2019). Regulation of Reproductive Processes with Dynamic Energy Budgets. Funct. Ecol., 33, 819–832.

Munch, S. B. & Salinas, S. (2009). Latitudinal Variation in Lifespan within Species Is Explained by the Metabolic Theory of Ecology. Proc. Natl. Acad. Sci., 106, 13860–13864.

New, M., Lister, D., Hulme, M. & Makin, I. (2002). A High-Resolution Data Set of Surface Climate over Global Land Areas. Clim. Res., 21, 1–25.

Nisbet, R. M., Jusup, M., Klanjscek, T. & Pecquerie, L. (2012). Integrating Dynamic Energy Budget (DEB) Theory with Traditional Bioenergetic Models. J. Exp. Biol., 215, 1246–1246.

Paaijmans, K. P., Heinig, R. L., Seliga, R. A., Blanford, J. I., Blanford, S., Murdock, C. C., et al. (2013). Temperature Variation Makes Ectotherms More Sensitive to Climate Change. Glob. Change Biol., 19, 2373–2380.

Palmer, M., Moro-Martínez, I., Tomàs-Ferrer, J., Grau, A., López-Belluga, M. D., Herlin, M., et al. (2024). Assessing Between-Individual Variability in Bioenergetics Modelling: Opportunities, Challenges, and Potential Applications. Ecol. Model., 498, 110848.

Pecquerie, L., Petitgas, P. & Kooijman, S. A. (2009). Modeling Fish Growth and Reproduction in the Context of the Dynamic Energy Budget Theory to Predict Environmental Impact on Anchovy Spawning Duration. J. Sea Res., 62, 93–105.

Pettersen, A. K. & Metcalfe, N. B. (2024). Consequences of the Cost of Living: Is Variation in Metabolic Rate Evolutionarily Significant? Philos. Trans. R. Soc. B Biol. Sci., 379, 20220498.

Polazzo, F., Limberger, R., Pennekamp, F., Ross, S. R. P.-J., Simpson, G. L. & Petchey, O. L. (2024). Measuring the Response Diversity of Ecological Communities Experiencing Multifarious Environmental Change. Glob. Change Biol., 30, e17594.

Porter, W. P., Mitchell, J. W., Beckman, W. A. & DeWitt, C. B. (1973). Behavioral Implications of Mechanistic Ecology: Thermal and Behavioral Modeling of Desert Ectotherms and Their Microenvironment. Oecologia, 13, 1–54.

Riddell, E. A., Burger, I. J., Tyner-Swanson, T. L., Biggerstaff, J., Muñoz, M. M., Levy, O., et al. (2023). Parameterizing Mechanistic Niche Models in Biophysical Ecology: A Review of Empirical Approaches. J. Exp. Biol., 226, jeb245543.

Riddell, E. A., Patton, J. L. & Beissinger, S. R. (2022). Thermal Adaptation of Pelage in Desert Rodents Balances Cooling and Insulation. Evolution, evo.14643.

Roček, Z., Joly, P. & Grossenbacher, K. (2003). "Triturus Alpestris (Laurenti, 1768) – Bergmolch". In: *Handbuch Der Reptilien Und Amphibien Europas. Band 4/IIA. Schwanzlurche (Urodela) IIA. Salamandridae II: Triturus 1.* Wiesbaden, AULA, 607–656.

Ross, S. R. P.-J., Petchey, O. L., Sasaki, T. & Armitage, D. W. (2023). How to Measure Response Diversity. Methods Ecol. Evol., 14, 1150–1167.

Ruel, J. J. & Ayres, M. P. (1999). Jensen's Inequality Predicts Effects of Environmental Variation. Trends Ecol. Evol., 14, 361–366.

Seebacher, F. & Franklin, C. E. (2012). Determining Environmental Causes of Biological Effects: The Need for a Mechanistic Physiological Dimension in Conservation Biology. Philos. Trans. R. Soc. B Biol. Sci., 367, 1607–1614.

Sexton, J. P., McIntyre, P. J., Angert, A. L. & Rice, K. J. (2009). Evolution and Ecology of Species Range Limits. Annu. Rev. Ecol. Evol. Syst., 40, 415–436.

Sinclair, B. J., Marshall, K. E., Sewell, M. A., Levesque, D. L., Willett, C. S., Slotsbo, S., et al. (2016). Can We Predict Ectotherm Responses to Climate Change Using Thermal Performance Curves and Body Temperatures? Ecol. Lett., 19, 1372–1385.

Sousa, T., Domingos, T. & Kooijman, S. (2008). From Empirical Patterns to Theory: A Formal Metabolic Theory of Life. Philos. Trans. R. Soc. B Biol. Sci., 363, 2453–2464.

Valladares, F., Matesanz, S., Guilhaumon, F., Araújo, M. B., Balaguer, L., Benito-Garzón, M., et al. (2014). The Effects of Phenotypic Plasticity and Local Adaptation on Forecasts of Species Range Shifts under Climate Change. Ecol. Lett., 17, 1351–1364.

Wolf, B. O. & McKechnie, A. E. (2023). Biophysical Approaches to Predicting Species Vulnerability. Glob. Change Biol., 29, 1421–1422.

Wood, S. (2025). *mgcv: Mixed GAM Computation Vehicle with Automatic Smoothness Estimation*. R package version 1.9-4.




Figure 1: Jensen's inequality and its implications for predicting performance in changing environments. For a nonlinear function, the value at the mean $f(\bar{\theta})$ differs from the mean of the values $\overline{f(\theta)}$, producing a bias (Jensen bias, $J = f(\bar{\theta}) - \overline{f(\theta)}$), where $J \approx -\frac{1}{2} f''(\bar{\theta})\ \mathrm{Var}[\theta]$, whose sign and magnitude follow the local curvature of f (Denny 2017; Jensen 1906). **(a) Representative trait–performance surfaces illustrating common forms of nonlinear responses to temperature**. The blue-to-red bar on the far left represents an environmental temperature gradient (cold to warm). Performance (F) is shown as a function of trait value (θ) and temperature (T) for five illustrative types: concave dome, Gaussian, Gaussian with shifting thermal optimum (T_{opt}), sigmoidal with fixed threshold, and sigmoidal with shifting threshold. Coloured vertical slices show the trait–performance relationship at cold (*blue*), intermediate (*black*) and warm (*red*) temperatures, while *black points* denote the mean trait value. **(b) Temperature-dependent Jensen bias predicted for each response surface**. *Red shading* indicates positive bias ($J > 0$), where predictions based on the mean trait overestimate performance, whereas *blue shading* indicates negative bias ($J < 0$), where the mean trait underestimates performance. The *solid curve* represents the bias under the current climate, $J(T)$, and the *dashed curve* the bias after warming, $J(T + \Delta T)$, for a fixed trait distribution ($\bar{\theta}$ and $\mathrm{Var}[\theta]$ held constant). **(c) Impact of warming on the magnitude of Jensen bias**. The *black curve* shows the change in absolute bias, $|J|$, across the temperature gradient. *Yellow shading* denotes amplification ($|J|$ increases), whereas *purple shading* denotes attenuation ($|J|$ decreases). The same warming can either strengthen or weaken the bias depending solely on the curvature and temperature dependence of the underlying trait–performance relationship.




Figure 2: Jensen bias (J) between population-average and mean individual-level predictions for females under the current climate. Negative values indicate underprediction by the population-average parameterisation, whereas positive values indicate overprediction; values close to zero indicate close agreement. The colour scale is symmetric and centred at zero. The black outline delimits the current distribution of *I. alpestris* (IUCN 2021). T_b, body temperature.




Figure 3: Change in Jensen bias under +4 °C warming for females. Each map displays the change in absolute Jensen bias ($\Delta|J|$, left, with colour bar) for each performance metric. The curve (descriptive GAM smoother) at the right of each map shows how $\Delta|J|$ changes along the mean annual temperature ($\overline{T}_a$) gradient. Positive values (*yellow*) indicate that warming increases the disagreement between population-average and mean individual-level predictions (amplification), whereas negative values (*purple*) indicate reduced disagreement (attenuation). The black outline delimits the current distribution of *I. alpestris* (IUCN 2021). T_b, body temperature.




Figure 4: Population-average and mean individual-level predictions and projected changes of lifespan, maximum body mass and lifetime fecundity for females under the current climate and +2 °C and +4 °C warming. For each trait, the top row shows absolute predictions from the population-average parameterisation and the mean of the individual-level parameterisations under each scenario, on a common trait-specific scale (*green*). The bottom row shows the projected change under warming relative to the corresponding current-climate prediction ($P_{\text{future}} - P_{\text{current}}$), normalised within each trait by the maximum absolute change across all grid cells, parameterisations and scenarios (scale from −1 to 1). Negative values (*red*) indicate decreases under warming, positive values (*blue*) indicate increases. The black outline delimits the current distribution of *I. alpestris* (IUCN 2021).