



Exercise 5.5



1. If $L = \{a, b, c\}$, $M = \{3, 4\}$, then find two binary relations of $L \times M$ and $M \times L$.

Solution: $L = \{a, b, c\}$, $M = \{3, 4\}$

$$L \times M = \{a, b, c\} \times \{3, 4\} \\ = \{(a, 3), (a, 4), (b, 3), (b, 4), (c, 3), (c, 4)\}$$

$$\text{Then } R_1 = \{(a, 3), (b, 4), (c, 3)\}$$

$$R_2 = \{(a, 4), (b, 3), (c, 4)\}$$

Now $M \times L = \{3, 4\} \times \{a, b, c\}$

$$= \{(3, a), (3, b), (3, c), (4, a), (4, b), (4, c)\}$$

$$\text{Here, } R_1 = \{(3, a), (4, a), (4, c)\}$$

$$R_2 = \{(3, b), (4, c)\}$$

2. If $Y = \{-2, 1, 2\}$, then make two binary relations for $Y \times Y$. Also find their domain and range.

Solution: $Y = \{-2, 1, 2\}$

$$Y \times Y = \{-2, 1, 2\} \times \{-2, 1, 2\}$$



$$= \{(-2, -2), (-2, 1), (-2, 2), (1, -2), (1, 1), (1, 2), (2, -2), (2, 1), (2, 2)\}$$

$$R_1 = \{(-2, -2), (-2, 1), (1, 2), (2, 2)\}$$

$$\text{Dom } R_1 = \{-2, 1, 2\}$$

$$\text{Range } R_1 = \{-2, 1, 2\}$$

$$\text{and } R_2 = \{(-2, 1), (1, 1), (-2, 2)\}$$

$$\text{Dom } R_2 = \{1, 2\}$$

$$\text{Range } R_2 = \{1, 2\}$$

3. If $L = \{a, b, c\}$, and $M = \{d, e, f, g\}$, then find two binary relations in each:

Solution:

- (i) $L \times L$

$$L = \{a, b, c\}$$

$$\text{Now } L \times L = \{a, b, c\} \times \{a, b, c\}$$

$$= \{(a, a), (a, b), (a, c), (b, a), (b, b), (b, c), (c, a), (c, b), (c, c)\}$$

$$R_1 = \{(a, a)\}$$

$$R_2 = \{(a, b), (b, b), (c, b)\}$$

- (ii) $L \times M$

$$L = \{a, b, c\}$$

$$M = \{d, e, f, g\}$$

$$\text{Now } L \times M = \{a, b, c\} \times \{d, e, f, g\}$$

$$= \{(a, d), (a, e), (a, f), (a, g), (b, d), (b, e), (b, f), (b, g), (c, d), (c, e), (c, f), (c, g)\}$$

$$\text{Here, } R_1 = \{(a, d), (b, f)\}$$

$$R_2 = \{(b, e), (b, g), (c, d), (c, e)\}$$

- (iii) $M \times M$

$$M = \{d, e, f, g\}$$

$$\text{Now } M \times M = \{d, e, f, g\} \times \{d, e, f, g\}$$

$$= \{(d, d), (d, e), (d, f), (d, g), (e, d), (e, e), (e, f), (e, g), (f, d), (f, e), (f, f), (f, g), (g, d), (g, e), (g, f), (g, g)\}$$

$$\text{Here, } R_1 = \{(d, e), (d, f), (f, f)\}$$

$$R_2 = \{(d, f), (e, d), (e, e), (g, g)\}$$

4. If set M has 5 elements, then find the number of binary relations in M .

Solution:

$$\text{Numbers of elements in } M = 5$$

$$\text{Numbers of elements in } M = 5$$

$$\text{Numbers of binary relations in } M \times M$$

$$= 2^{5 \times 5} = 2^{25}$$

20

5. If $L = \{x | x \in N \wedge x \leq 5\}$, $M = \{y | y \in P \wedge y < 10\}$, then make the following relations from L to M

Solution:

(i) $R_1 = \{(x, y) | y < x\}$

$L = \{x | x \in N \wedge x \leq 5\}$

Thus, $L = \{1, 2, 3, 4, 5\}$

and $M = \{y | y \in P \wedge y < 10\}$ *Prime = 2*

Thus, $M = \{2, 3, 5, 7\}$

Now, $L \times M = \{1, 2, 3, 4, 5\} \times \{2, 3, 5, 7\}$

$= \{(1, 2), (1, 3), (1, 5), (1, 7), (2, 2), (2, 3), (2, 5), (2, 7), (3, 2), (3, 3), (3, 5), (3, 7), (4, 2), (4, 3), (4, 5), (4, 7), (5, 2), (5, 3), (5, 5), (5, 7)\}$

$R_1 = \{(x, y) | y < x\}$

$= \{(3, 2), (4, 2), (4, 3), (5, 2), (5, 3)\}$

Dom $R_1 = \{3, 4, 5\}$

Range $R_1 = \{2, 3\}$

(ii) $R_2 = \{(x, y) | y = x\}$

$R_2 = \{(2, 2), (3, 3), (5, 5)\}$

Dom $R_2 = \{2, 3, 5\}$ *y = x*

Range $R_2 = \{2, 3, 5\}$

(iii) $R_3 = \{(x, y) | y + x = 6\}$

$R_3 = \{(1, 5), (3, 3), (4, 2)\}$

Dom $R_3 = \{1, 3, 4\}$

Range $R_3 = \{5, 3, 2\}$

(iv) $R_4 = \{(x, y) | y - x = 2\}$

$R_4 = \{(1, 3), (3, 5), (5, 7)\}$

Dom $R_4 = \{1, 3, 5\}$

Range $R_4 = \{3, 5, 7\}$

Also write the domain and range of each relation.

6. Indicate relations, into functions, one-one function, on to function, and bijective function from the following. Also find their domain and the range.

Solution:

(i) $R_1 = \{(1, 1), (2, 2), (3, 3), (4, 4)\}$

Bijective

Dom $R_1 = \{1, 2, 3, 4\}$

Range $R_1 = \{1, 2, 3, 4\}$

(ii) $R_2 = \{(1, 2), (2, 1), (3, 4), (3, 5)\}$

Relation

$$\text{Dom } R_2 = \{1, 2, 3\}$$

$$\text{Range } R_2 = \{1, 2, 4, 5\}$$

(iii) $R_3 = \{(b, a), (c, a), (d, a)\}$

Function

$$\text{Dom } R_3 = \{a, b, c\}$$

$$\text{Range } R_3 = \{a\}$$

(iv) $R_4 = \{(a, b), (b, a), (c, d), (d, e)\}$

Onto function

$$\text{Dom } R_4 = \{1, 2, 3, 4, 5\}$$

$$\text{Range } R_4 = \{1, 3, 4\}$$

(v) $R_5 = \{(a, b), (b, a), (c, d), (d, e)\}$

One - one function

$$\text{Dom } R_5 = \{a, b, c, d\}$$

$$\text{Range } R_5 = \{a, b, d, e\}$$


(vi) $R_6 = \{(1, 2), (2, 3), (1, 3), (3, 4)\}$

Relation

$$\text{Dom } R_6 = \{1, 2, 3\}$$

$$\text{Range } R_6 = \{2, 3, 4\}$$


(vii) $R_7 =$



$$\text{Dom } R_7 = \{1, 2, 3\}$$

$$\text{Range } R_7 = \{r, p, s\}$$

(viii) $R_8 =$



Relation

$$\text{Dom } R_8 = \{1, 3, 7\}$$

$$\text{Range } R_8 = \{a, b, c\}$$