



## MATHEMATICS HSSC— SECTION – A (Marks 20)

Time allowed: 25 Minutes

Section – A is compulsory. All parts of this section are to be answered on this page and handed over to the Centre Superintendent. Deleting/overwriting is not allowed.

Do not use lead pencil.

حصہ اول لازمی ہے۔ اس کے جوابات اسی صفحہ پر دے کر نام مرکز کے حوالے کریں۔ کٹ کر دیا جائے گا۔  
لکھنے کی اجازت نہیں ہے۔ لے پینسل کا استعمال ممنوع ہے۔

Version No.			
3	1	1	1

ROLL NUMBER					

- ① ① ① ①  
② ② ② ②  
③ ③ ③ ③  
④ ④ ④ ④  
⑤ ⑤ ⑤ ⑤  
⑥ ⑥ ⑥ ⑥  
⑦ ⑦ ⑦ ⑦  
⑧ ⑧ ⑧ ⑧  
⑨ ⑨ ⑨ ⑨

- ① ① ① ① ① ①  
② ② ② ② ② ②  
③ ③ ③ ③ ③ ③  
④ ④ ④ ④ ④ ④  
⑤ ⑤ ⑤ ⑤ ⑤ ⑤  
⑥ ⑥ ⑥ ⑥ ⑥ ⑥  
⑦ ⑦ ⑦ ⑦ ⑦ ⑦  
⑧ ⑧ ⑧ ⑧ ⑧ ⑧  
⑨ ⑨ ⑨ ⑨ ⑨ ⑨

Answer Sheet No. \_\_\_\_\_

ہر سوال کے سامنے دیے گئے، کریکولم کے مطابق درست دائرہ کو پر کریں۔  
Invigilator Sign. \_\_\_\_\_

Fill the relevant bubble against each question according to curriculum: Candidate Sign. \_\_\_\_\_

Question	Candidate Sign.							
	A	B	C	D				
1. The multiplicative inverse of $-i$ is:	$i$	1	$-i$	-1	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
2. What is the modulus of a complex number $(8-15i)$ ?	$8+15i$	17	$\sqrt{161}$	-15	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
3. The contrapositive of a conditional $p \rightarrow q$ is:	$q \rightarrow p$	$\sim q \rightarrow p$	$\sim q \rightarrow \sim p$	$\sim p \rightarrow q$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
4. Which structure in the following is true for the set of natural numbers under multiplication?	Groupoid	Semi group	Monoid	Group	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
5. Which one of the following matrices is singular?	$\begin{bmatrix} 1 & -4 \\ 2 & 8 \end{bmatrix}$	$\begin{bmatrix} 1 & 4 \\ \sqrt{4} & 8 \end{bmatrix}$	$\begin{bmatrix} 1 & -4 \\ \sqrt{2} & -8 \end{bmatrix}$	$\begin{bmatrix} -1 & 4 \\ 2 & 8 \end{bmatrix}$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
6. Rank of matrix $\begin{bmatrix} -2 \\ 0 \\ -1 \end{bmatrix}$ is:	0	1	2	3	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
7. For what value of $k$ , roots of $kx^2 - 12x + 4 = 0$ are equal?	9	-9	9.5	18	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
8. One of the multiplicative factors of $(x^4 - 5x^2 + 4)$ is:	$x+2$	$x-3$	$x+3$	$x+4$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
9. Which one of the following represents $\frac{x^3 + 2x^2 + 3}{(x^2 + 1)(x + 4)}$ ?	Proper fraction	Improper fraction	Cubic polynomial	Polynomial	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
10. For what value of $x$ , the numbers $\frac{1}{2}, \frac{1}{5}, \frac{1}{x}$ are in harmonic progression?	-10	-8	8	10	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
11. If $\binom{n}{8} = \binom{n}{12}$ , then value of $n$ is:	4	8	12	20	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>



	Question	A	B	C	D	A	B	C	D
12.	The probability of getting same upper face on throwing two fair dice simultaneously is:	$\frac{1}{12}$	$\frac{1}{6}$	$\frac{1}{4}$	$\frac{1}{2}$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
13.	The coefficient of third term in the expansion of $\left(x - \frac{1}{x}\right)^8$ is:	$\binom{8}{0}$	$\binom{8}{1}$	$\binom{8}{2}$	$\binom{8}{3}$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
14.	In which quadrant, terminal side of the angle $-510^\circ$ lies?	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
15.	$(\sin 5x + \sin 7x)$ is expressed in product form as:	$2\sin 6x \cos x$	$2\cos 6x \sin x$	$2\sin 6x \sin x$	$2\cos 6x \cos x$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
16.	The value of $\cos(x + 60^\circ) + \cos(x - 60^\circ)$ is:	$\cos x$	$\sqrt{3} \cos x$	$\cos 2x$	0	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
17.	The period of $\frac{8}{7} \sec(x - \pi)$ is:	$-\pi$	$\pi$	$3\pi$	$\frac{8\pi}{7}$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
18.	What is the area of a triangle $\triangle ABC$ , if $a = 10, b = 20$ and $\gamma = 30^\circ$ ?	$25\sqrt{2}$	$50\sqrt{3}$	50	$\frac{100}{\sqrt{3}}$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
19.	The value of $\cos\left[\frac{\pi}{6} + \cos^{-1}\left(-\frac{1}{2}\right)\right]$ is:	$\frac{1}{2}$	$-\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
20.	The solution of a trigonometric equation $(\sin x + \cos x = 0)$ is:	$-30^\circ$	$-120^\circ$	$45^\circ$	$135^\circ$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

—1HA-I 2311-3111—

ROLL NUMBER					





# MATHEMATICS HSSC-I

Time allowed: 2:35 Hours

Total Marks Sections B and C: 80

NOTE: Attempt any twelve parts from Section 'B' and any four questions from Section 'C' on the separately provided answer book. Write your answers neatly and legibly.

### SECTION - B (Marks 48)

Q. 2 Attempt any TWELVE parts. All parts carry equal marks. (12 x 4 = 48)

- (i) If  $Z_1 = 2 + 3i$  and  $Z_2 = 4 + 2i$ , then show that  $(Z_1\bar{Z}_2 + \bar{Z}_1Z_2)$  is a real number.
- (ii) Construct a truth table of a logical statement  $(p \leftrightarrow q) \wedge (p \rightarrow q)$
- (iii) Solve for  $x$ : 
$$\begin{vmatrix} x & -1 \\ 5 & 1-x \end{vmatrix} = \begin{vmatrix} 1 & 0 & -3 \\ 2 & x & -6 \\ 1 & 3 & x-5 \end{vmatrix}$$
- (iv) If  $\alpha, \beta$  are the roots of  $x^2 + px + q = 0$ , find the quadratic equation whose roots are  $\frac{\alpha}{\beta}$  and  $\frac{\beta}{\alpha}$ .
- (v) Using properties of the cube roots of unity, verify that  $(1 + \omega) + (1 + \omega)^2 + (1 + \omega)^3 = 2\omega$
- (vi) Express  $\frac{125 + 4x - 9x^2}{(x-1)(x+3)(x+4)}$  in partial fractions.
- (vii) Second term of a geometric sequence is 9 and its fourth term is 1. Find sum to infinity.
- (viii) Insert six arithmetic means between 15 and -13.
- (ix) Prove that Sine is a periodic function and its period is  $2\pi$
- (x) A die is thrown twice. Find the probability that sum of the upper face numbers is a prime number or an odd number.
- (xi) Find the value of  $k$ , if the constant term in the expansion of  $(2x^2 + \frac{k}{x})^6$  is 960.
- (xii) If  $\cos \theta = \frac{\sqrt{10}}{10}$  with  $2\pi < \theta < \frac{5\pi}{2}$ , then find values of the remaining five trigonometric ratios.
- (xiii) Verify that:  $\cos 4x \cos x - \sin 6x \sin 3x = \cos 7x \cos 2x$
- (xiv) In an oblique triangle  $\triangle ABC$  (with usual notations)  $a = 6, c = 12$  and  $\beta = 124^\circ$ . Apply law of cosines and law of sines to find the values of 'b',  $\alpha$  and  $\gamma$
- (xv) Verify that:  $2 \tan^{-1} \frac{1}{2} + \tan^{-1} \frac{1}{7} = \tan^{-1} \frac{31}{17}$
- (xvi) Solve the trigonometric equation  $\cos 5\theta + \cos \theta = \cos 3\theta$  where  $\theta \in [0, \pi]$

### SECTION - C (Marks 32)

Note: Attempt any FOUR questions. All questions carry equal marks. (4 x 8 = 32)

- Q. 3 Use Cramer's rule to solve the system of linear equations.  $x + y - z = 3; 2x - y - z = 1; 3x + y + 2z = 0$
- Q. 4 If three consecutive numbers in an arithmetic progression are increased by 1, 2 and 3 respectively, the resulting numbers are in geometric progression. Find the original numbers if their sum is 12.
- Q. 5 If  $y = \frac{1}{4} + \frac{1.3}{4.8} + \frac{1.3.5}{4.8.12} + \dots$  then prove that  $y^2 + 2y - 1 = 0$
- Q. 6 Without using calculator, prove that  $\cos 10^\circ \cdot \cos 30^\circ \cdot \cos 50^\circ \cdot \cos 70^\circ = \frac{3}{16}$
- Q. 7 Solve the following system of equations:  $5x^2 - 14xy + 9y^2 = 0; 4x^2 - 3xy - 16 = 0$
- Q. 8 Solve triangles  $\triangle ABC$  (with usual notations) if,
  - (a)  $\alpha = 60^\circ, \beta = 15^\circ$  and  $b = 33$
  - (b)  $b = 23, c = 24$  and  $\alpha = 75^\circ$



# MATHEMATICS HSSC-I

## SECTION – A (Marks 20)

Time allowed: 25 Minutes

Section – A is compulsory. All parts of this section are to be answered on this page and handed over to the Centre Superintendent. Deleting/overwriting is not allowed.

Do not use lead pencil.

حصہ اول لازمی ہے۔ اس کے جوابات اسی صفحہ پر دے کر تمام مرکز کے حوالے کریں۔ کٹ کر دیا جانے کی اجازت نہیں ہے۔ سیاہ پنسل کا استعمال ممنوع ہے۔

Version No.			
7	1	1	4

ROLL NUMBER					

- ○ ○ ○  
 ① ● ● ①  
 ② ② ② ②  
 ③ ③ ③ ③  
 ④ ④ ④ ●  
 ⑤ ⑤ ⑤ ⑤  
 ⑥ ⑥ ⑥ ⑥  
 ● ⑦ ⑦ ⑦  
 ⑧ ⑧ ⑧ ⑧  
 ⑨ ⑨ ⑨ ⑨

- ○ ○ ○ ○ ○  
 ① ① ① ① ① ①  
 ② ② ② ② ② ②  
 ③ ③ ③ ③ ③ ③  
 ④ ④ ④ ④ ④ ④  
 ⑤ ⑤ ⑤ ⑤ ⑤ ⑤  
 ⑥ ⑥ ⑥ ⑥ ⑥ ⑥  
 ⑦ ⑦ ⑦ ⑦ ⑦ ⑦  
 ⑧ ⑧ ⑧ ⑧ ⑧ ⑧  
 ⑨ ⑨ ⑨ ⑨ ⑨ ⑨

Answer Sheet No. \_\_\_\_\_

Invigilator Sign. \_\_\_\_\_

Fill the relevant bubble against each question according to curriculum: Candidate Sign. \_\_\_\_\_

Question	Candidate Sign.			
	A	B	C	D
1. What is the area of a triangle $\Delta ABC$ , if $a = 1, c = 2$ and $\beta = 60^\circ$ ?	$\sqrt{3}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	1
2. The value of $\sin(\pi + \sin^{-1} x)$ is:	$\pi x$	$x$	$-x$	$-\pi x$
3. Which one of the following is a solution of $\sin 2x + \cos 2x = -1$ ?	$30^\circ$	$45^\circ$	$90^\circ$	$0^\circ$
4. If $z = -2 - 3i$ then, what is the value of $z - \bar{z}$ ?	$6i$	$-4$	4	$-6i$
5. The simplified form of $i^{18}$ is:	$-i$	1	$-1$	$i$
6. The converse of the conditional $\sim q \rightarrow p$ is:	$\sim q \rightarrow p$	$q \rightarrow \sim p$	$\sim p \rightarrow q$	$q \rightarrow p$
7. Which structure in the following is TRUE for the set of natural numbers under addition?	Semi group	Monoid	Group	Groupoid
8. The determinant of a matrix $\begin{bmatrix} i & 0 & \sqrt{3} \\ 0 & i & \sqrt{3} \\ 0 & 0 & i \end{bmatrix}$ is:	$i$	$-1$	1	$-i$
9. Rank of matrix $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ is:	1	2	3	0
10. If $(-1)$ is a root of $x^3 + px^2 - x + 2 = 0$ , then value of $p$ is:	$-4$	0	2	$-2$



	Question	A	B	C	D	A	B	C	D
11.	For what value of $m$ , $(x^2 - 2mx - 3)$ has zero remainder, when divided by $(x - 3)$ ?	1	-2	2	-1	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
12.	For what value of $B$ , $\frac{1}{x^2 - 1} = \frac{A}{x - 1} + \frac{B}{x + 1}$ ?	2	-1	$-\frac{1}{2}$	$\frac{1}{2}$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
13.	If $a - 3, 6, b + 3$ are in arithmetic progression, then value of $(a + b)$ is:	6	12	18	0	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
14.	If ${}^n P_2 = 30$ then, value(s) of $n$ is/are:	6, -5	8	12	6, 5	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
15.	The probability of getting two tails when two coins are tossed is:	$\frac{1}{6}$	$\frac{1}{4}$	$\frac{1}{3}$	$\frac{1}{2}$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
16.	The middle term in the expansion of $\left(x - \frac{1}{2x}\right)^{12}$ is:	6 <sup>th</sup>	7 <sup>th</sup>	8 <sup>th</sup>	5 <sup>th</sup>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
17.	If $\sin^2 \theta = \frac{1}{7}$ the value of $\sec^2 \theta$ is:	$\frac{6}{7}$	$\frac{7}{6}$	$\frac{8}{7}$	7	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
18.	$2\sin 7x \cos 3x$ is expressed in sum or difference form as:	$\sin 10x - \sin 4x$	$\sin 5x - \sin 2x$	$\cos 5x + \cos 2x$	$\sin 10x + \sin 4x$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
19.	The value of $\sin(90^\circ + x) + \sin(90^\circ - x)$ is:	0	$-2\sin x$	$-2\cos x$	$2\cos x$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
20.	The period of $3\cos \frac{x}{5}$ is:	$10\pi$	$30\pi$	$\frac{\pi}{5}$	$5\pi$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

—1HA-I 2311-7114 HA —

ROLL NUMBER					





# MATHEMATICS HSSC-I

36

Time allowed: 2:35 Hours

Total Marks Sections B and C: 80

NOTE: Attempt any twelve parts from Section 'B' and any four questions from Section 'C' on the separately provided answer book. Write your answers neatly and legibly.

### SECTION - B (Marks 48)

Q. 2 Attempt any TWELVE parts. All parts carry equal marks. (12 x 4 = 48)

- (i) If  $Z_1 = 3 + 4i$  and  $Z_2 = 4 + 3i$ , then show that  $\left| Z_1 + \frac{1}{Z_2} \right| \cong 5$ .
- (ii) Construct truth table of a logical statement  $(\sim p \rightarrow \sim q) \wedge p$
- (iii) Show that 
$$\begin{vmatrix} y+z & x & y \\ z+x & z & x \\ x+y & y & z \end{vmatrix} = (x+y+z)(z-x)^2$$
- (iv) If  $(-5)$  is a root of the equation  $2x^2 + px - 15 = 0$  and the equation  $p(x^2 + x) + k = 0$  has equal roots, find values of 'p' and 'k'.
- (v) Using properties of the cube roots of unity, Show that  $(x+y)^2 + (x\omega + y\omega^2)^2 + (x\omega^2 + y\omega)^2 = 6xy$
- (vi) Express  $\frac{x^2 + 2x - 1}{(x)(x+2)(2x-1)}$  in partial fractions.
- (vii) Which term of the arithmetic sequence 4, -4, -12, -20, -28, ..... is  $(-172)$ ?
- (viii) Show that  $S_n = \sum_{k=1}^n (6k^2 + 4k - 1) = n(n+2)(2n+1)$
- (ix) Prove that Tangent is a periodic function and its period is  $\pi$
- (x) A fair cubical die is rolled three times.
  - (a) Find the probability of getting a six all three times.
  - (b) Find the probability of getting no sixes.
- (xi) Find the value of  $k$ , if the constant term in the expansion of  $\left(4x^2 + \frac{k}{2x}\right)^9$  is 84.
- (xii) If  $\sin \theta = \frac{-\sqrt{15}}{15}$  and  $\pi < \theta < \frac{3\pi}{2}$ , then find values of the remaining trigonometric ratios.
- (xiii) If  $\operatorname{Cosec} \alpha = \frac{5}{3}$ , where  $0 < \alpha < \frac{\pi}{2}$  and  $\operatorname{Sec} \beta = \frac{13}{12}$ , where  $\frac{3\pi}{2} < \beta < 2\pi$ , then find the value of  $\cos(\alpha - \beta)$
- (xiv) Two escribed circles of radii  $r_2$  and  $r_3$  are connected with triangle  $\triangle ABC$ . Using usual notations, prove that  $(r_2 + r_3) \tan \frac{\alpha}{2} = a$
- (xv) Prove that:  $2 \tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{7} = \frac{\pi}{4}$
- (xvi) Solve the trigonometric equation  $\sin 2x = \sqrt{3} \cos x$ , where  $0 < x < 2\pi$

### SECTION - C (Marks 32)

Note: Attempt any FOUR questions. All questions carry equal marks. (4 x 8 = 32)

- Q. 3 Use Cramer's rule to solve the system of linear equations  $x + 5y - 3z = -36$  ;  $x + 4y + 2z = -11$  ;  $2x - y = 7$
- Q. 4 If 1, 4 and 3 are added to three consecutive terms of a geometric progression, the resulting numbers are in arithmetic progression. What are the numbers if their sum is 13?
- Q. 5 If  $y = \frac{3}{4} + \frac{3.5}{4.8} + \frac{3.5.7}{4.8.12} + \dots$  then prove that  $y^2 + 2y - 7 = 0$
- Q. 6 Without using calculator, prove that  $\cos 40^\circ \cdot \cos 80^\circ \cdot \cos 120^\circ \cdot \cos 160^\circ = \frac{1}{16}$
- Q. 7 Solve the following system of equations  $6x^2 + 9xy + 3y^2 = 0$  ;  $x^2 - 5xy + 6y^2 = 26$
- Q. 8 Solve triangle  $\triangle ABC$  (with usual notations) if,
  - (a)  $a = 7$ ,  $b = 10$  and  $c = 13$
  - (b)  $a = 25$ ,  $b = 24$  and  $\gamma = 120^\circ$

Version No.			
3	1	1	1

ROLL NUMBER					



- 0 0 0 0
- 1 ● ● ●
- 2 2 2 2
- 3 3 3
- 4 4 4 4
- 5 5 5 5
- 6 6 6 6
- 7 7 7 7
- 8 8 8 8
- 9 9 9 9

- 0 0 0 0 0 0
- 1 1 1 1 1 1
- 2 2 2 2 2 2
- 3 3 3 3 3 3
- 4 4 4 4 4 4
- 5 5 5 5 5 5
- 6 6 6 6 6 6
- 7 7 7 7 7 7
- 8 8 8 8 8 8
- 9 9 9 9 9 9

Answer Sheet No. \_\_\_\_\_

Sign. of Candidate \_\_\_\_\_

Sign. of Invigilator \_\_\_\_\_

Section – A is compulsory. All parts of this section are to be answered on this page and handed over to the Centre Superintendent. Deleting/overwriting is not allowed. Do not use lead pencil.

**MATHEMATICS HSSC–I**  
**SECTION – A (Marks 20)**  
**Time allowed: 25 Minutes**

حصہ اول لازمی ہے۔ اس کے جوابات اسی صفحہ پر دے کر ناظم مرکز کے حوالے کریں۔ کٹ کر دوبارہ لکھنے کی اجازت نہیں ہے۔ لیڈ پینسل کا استعمال ممنوع ہے۔

Fill the relevant bubble against each question:

ہر سوال کے سامنے دیے گئے درست دائرہ کو پر کریں۔

1. If  $a, b, c$  are real numbers such that  $a < b$ ,  $c < 0$ ,  $a \neq 0$ ,  $b \neq 0$ , then which of the following inequalities holds:
 

$ac > bc$         $ac^2 > bc^2$         $\frac{c}{a} > \frac{c}{b}$         $ac < bc$

---

2. What is the converse of  $p \rightarrow q$  ?
 

$\sim p \rightarrow \sim q$         $q \rightarrow p$         $\sim q \rightarrow \sim p$         $p \leftrightarrow q$

---

3. The set of non-zero rational numbers is a group under the operation of:
 

Addition       Subtraction       Multiplication       Division

---

4. For what value of  $\lambda$  is the matrix  $\begin{bmatrix} 1 & 0 & 0 \\ 2 & \lambda & 0 \\ 1 & 2 & 3 \end{bmatrix}$  singular?
 

1       0       3       -4

---

5. If  $A$  is a skew-symmetric matrix then:
 

$A = A'$         $A = -A'$         $A = (\bar{A})'$         $A = -(\bar{A})'$

---

6. If the polynomial  $f(x)$  is divided by  $x+2$ , the quotient is  $x-2$  and the remainder is 2, then  $f(x)$  will be:
 

$x^2 - 4$         $x^2 + 4$         $x^2 - 2$         $x^2 + 2$

---

7. If  $w$  is a cube root of unity, then which of the following equations is true?
 

$1+w=0$         $1+w^2=0$         $w+w^2=0$         $1+w+w^2=0$

---

8. What is the partial fractions of  $\frac{x^2+2x-1}{x^2-1}$  ?
 

$1 + \frac{1}{x+1} - \frac{1}{x-1}$         $1 + \frac{1}{x-1} - \frac{1}{x+1}$         $1 - \frac{1}{x+1} - \frac{1}{x-1}$         $1 + \frac{1}{x-1} + \frac{1}{x+1}$

---

9. Find the second term of the sequence whose general term is  $a_n = 2n^2 - 3$ 

-1       13       5       11

10. If  $s_{\infty} = \frac{2}{3}$  and  $a = \frac{2}{7}$  in an infinite geometric progression, then the common ratio is:   $-\frac{4}{7}$    $\frac{4}{7}$    $\frac{2}{7}$    $-\frac{2}{7}$

11. For what values of  $x$ , the binomial expansion of  $\left(1 - \frac{x}{2}\right)^{-1}$  is convergent (valid)?   $x > 2$    $|x| > 2$    $|x| < 2$    $x < 1$

12. What is radius of the circle whose part of arc-length of measure 4 is with central angle  $\frac{\pi}{2}$ ?   $\frac{8}{\pi}$    $\frac{4}{\pi}$    $\frac{2}{\pi}$    $\frac{\pi}{2}$

13. If  $D(-5, 5\sqrt{2})$  lies on the terminal side of  $\theta$ , then find the value of  $\tan \theta$    $-\frac{1}{\sqrt{2}}$    $\frac{1}{\sqrt{2}}$    $\sqrt{2}$    $-\sqrt{2}$

14. If  ${}^n C_4 = {}^n C_{10}$ , then  $n = \dots$   4  10  14  6

15. How many distinct three-digit numbers can be formed from the integers 1,2,3,4,5,6 if each digit is used at most once?  360  120  20  10

16. What is the middle term in the expansion of  $(x + x^{-1})^{14}$   6th term  7th term  8th term  9th term

17.  $\sin\left(\frac{3\pi}{2} - \alpha\right) =$    $\sin \alpha$    $\cos \alpha$    $-\sin \alpha$    $-\cos \alpha$

18. What is the primary period of  $\frac{\sin 2x}{1 + \cos 2x}$    $2\pi$    $\pi$    $\frac{\pi}{2}$    $4\pi$

19. A ladder makes angle  $30^\circ$  with the wall of height  $8m$ . What is the length of the ladder?   $16m$    $8m$    $4m$    $12m$

20. What is the value of  $\sin^{-1}\left(-\frac{1}{2}\right)$ ?   $-\frac{\pi}{6}$    $\frac{\pi}{6}$    $-\frac{\pi}{3}$    $\frac{\pi}{3}$

—1HA-I 2211-3111 (L)—

ROLL NUMBER

--	--	--	--	--	--



# MATHEMATICS HSSC-I

36

Time allowed: 2:35 Hours

Total Marks Sections B and C: 80

NOTE: Attempt any twelve parts from Section 'B' and any four questions from Section 'C' on the separately provided answer book. Use supplementary answer sheet i.e. Sheet-B if required. Write your answers neatly and legibly. Graph paper will be provided on request.

## SECTION - B (Marks 48)

Q. 2 Attempt any TWELVE parts. All parts carry equal marks.

(12 x 4 = 48)

- (i) Separate  $\frac{(2-3i)^2}{1-i}$  into real and imaginary parts.
- (ii) Determine whether  $p \rightarrow (q \rightarrow p)$  is a tautology, a contingency or an absurdity.
- (iii) If  $A = \{1, 2, 3, 4\}$ , state the domain and range of the relation  $R = \{(x, y) | x + y = 5\}$
- (iv) Under the operation "\*", complete the following table to obtain a semigroup
- |   |  |     |     |   |
|---|--|-----|-----|---|
| * |  | a   | b   | c |
| a |  | c   | a   | b |
| b |  | ... | ... | c |
| c |  | b   | c   | a |
- (v) Find the matrix  $A$  if  $\begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} A = \begin{bmatrix} 0 & -3 & 8 \\ 3 & 3 & -7 \end{bmatrix}$
- (vi) Find the inverse of matrix  $A = \begin{bmatrix} 2i & i \\ i & -i \end{bmatrix}$ , hence show that  $AA^{-1} = I_2$
- (vii) If  $\alpha, \beta$  are roots of  $3x^2 - 2x + 4 = 0$ , then find the value of  $\frac{\alpha}{\beta} + \frac{\beta}{\alpha}$
- (viii) Resolve  $\frac{3x-11}{(x+3)(x^2+1)}$  into partial fractions.
- (ix) If  $y = 1 - \frac{x}{2} + \frac{x^2}{4} - \dots$ , then show that  $x = 2\left(\frac{1-y}{y}\right)$
- (x) Find values of  $n$  and  $r$ , when  ${}^nC_r = 10$  and  ${}^nP_r = 60$ .
- (xi) There are 9 green and 6 red balls in a box. A ball is drawn (taken out). What is the probability that  
(i) the ball is green (ii) the ball is red.
- (xii) Expand and simplify  $(2+i)^4 - (2-i)^4$
- (xiii) Find the remaining trigonometric functions if  $\cos \theta = -\frac{1}{2}$  and the terminal arm of angle  $\theta$  is in quad-III.
- (xiv) Show that  $\frac{\sin(\alpha - \beta)}{\sin(\alpha + \beta)} = \frac{\tan \alpha - \tan \beta}{\tan \alpha + \tan \beta}$
- (xv) Find the measure of smallest angle of the triangle whose sides are 16, 20 and 33
- (xvi) Show that  $2 \cos^{-1} \frac{4}{5} = \sin^{-1} \frac{24}{25}$

## SECTION - C (Marks 32)

Note: Attempt any FOUR questions. All questions carry equal marks.

(4 x 8 = 32)

- Q. 3 Find the real and imaginary parts of the complex number  $\frac{(\sqrt{3}-i)^5}{(\sqrt{3}+i)^5}$
- Q. 4 Find the value of  $\lambda$  for which the system  $2x + y - \lambda z = 0$  has a non-trivial solution. Also solve the system for  $x + 2y - 2z = 0$   
that value of  $\lambda$
- Q. 5 (a) Resolve  $\frac{x^2}{(x^2+4)(x+2)}$  into partial fractions (b) Prove that  ${}^nC_k + {}^nC_{k-1} = {}^{n+1}C_k$
- Q. 6 Expand  $(1-2x)^{\frac{1}{3}}$  to four terms and apply it to evaluate  $(0.8)^{\frac{1}{3}}$  correct to three places of decimal.
- Q. 7 If  $\sin \alpha = \frac{4}{5}$  and  $\sin \beta = \frac{12}{13}$ , where  $\frac{\pi}{2} < \alpha < \pi$  and  $\frac{\pi}{2} < \beta < \pi$ . Find (i)  $\cos(\alpha + \beta)$  (ii)  $\sin(\alpha - \beta)$
- Q. 8 (a) Show that  $R = \frac{abc}{4\Delta}$   
(b) Solve the equation  $\sqrt{3} \tan x - \sec x - 1 = 0$  for its general solution

37

Version No.			
7	1	1	1

ROLL NUMBER					



- 0
- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9

- 0
- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9

Answer Sheet No. \_\_\_\_\_

Sign. of Candidate \_\_\_\_\_

Sign. of Invigilator \_\_\_\_\_

Section – A is compulsory. All parts of this section are to be answered on this page and handed over to the Centre Superintendent. Deleting/overwriting is not allowed. Do not use lead pencil.

**MATHEMATICS HSSC–I**  
**SECTION – A (Marks 20)**  
**Time allowed: 25 Minutes**

حصہ اول لازمی ہے۔ اس کے جوابات اسی صفحہ پر دے کر ناظم مرکز کے حوالے کریں۔ کٹ کر دوبارہ لکھنے کی اجازت نہیں ہے۔ ایڈیشنل کا استعمال ممنوع ہے۔

ہر سوال کے سامنے دیے گئے درست دائرہ کو پر کریں۔

Fill the relevant bubble against each question:

1. For a complex number  $z$ , all the following formulas are true EXCEPT:   $\frac{z}{z} = z$    $z\bar{z} = |z|^2$    $z^2 = |z|^2$    $|z| = |\bar{z}|$

2. Which of the following sets forms an abelian group under the operation of multiplication?  Set of rational numbers  Set of integers  Set of natural numbers  Set of non-zero real numbers

3. Suppose the number of players that play cricket and hockey are 15 and 13 respectively. If the total number of players is 21, what is the number of players that play both the games?  6  8  7  28

4. If  $A$  is a matrix of order  $3 \times 4$ , then which of the following equalities is TRUE?   $AI_3 = A$    $I_4A = A$    $AA^t = I_4$    $AI_4 = A$

5.  $\begin{vmatrix} 1 & 0 & 0 \\ 2 & -i & 0 \\ 3 & -2 & i \end{vmatrix} =$   1  -1   $i$    $-i$

6. If  $f(x)$  is a polynomial with only two roots 1 and 2, then  $f(x) =$    $x^2 + 3x - 2$    $x^2 + 3x + 2$    $x^2 - 3x + 2$    $x^2 - 3x - 2$

7. If one root of the equation  $f(x) = 0$  is  $-1$ , then  $5 - f(-1) =$   6  4  5   $-6$

8. The partial fraction of  $\frac{1}{1-x^3}$  will be in the form of:   $\frac{A}{1-x} + \frac{Bx+C}{1-x^2}$    $\frac{A}{1-x} + \frac{Bx+C}{1+x+x^2}$    $\frac{A}{1-x} + \frac{Bx+C}{(1-x)^2}$    $\frac{A}{1-x} + \frac{B}{1-x+x^2}$

9. If  $a_1 = -1$  in a sequence with general term  $a_n = n + a_{n-1}$  then sum of first two terms  $S_2$  is:  0  1   $-1$   2

10. If  $b$  is  $a$  harmonic mean between  $-2$  and  $4$  then  $b = \dots$   8   $-8$   1   $-1$

11.  $\binom{8}{7} + \binom{8}{6} =$   72  48  63  36

12. If a fair die is rolled, then what is the probability that the top is an even number?   $\frac{1}{2}$    $\frac{1}{3}$    $\frac{1}{6}$   1

13. Which of the following expressions is sum of the series  $1 - x + x^2 - x^3 + \dots$    $\frac{1}{1+x}$    $\frac{1}{1-x}$    $\sqrt{1+x}$    $\frac{1}{\sqrt{1-x}}$

14. What is the length of the arc that subtends an angle of measure  $60^\circ$  at the centre of a circle with radius 6?   $3\pi$    $2\pi$    $6\pi$    $\pi$

15.  $\sin\left(\frac{7\pi}{6}\right) =$    $-\frac{\sqrt{3}}{2}$    $-\frac{1}{2}$    $\frac{1}{2}$    $\frac{\sqrt{3}}{2}$

16. Which of the following trigonometric expressions is identically equal to  $1 - \cos 2\theta$    $2\cos^2 \theta$    $2\sin^2 \theta$    $2\sin^2 2\theta$    $2\cos^2 2\theta$

17. What is the primary period of  $\tan\left(\frac{x}{3}\right)$ ?   $3\pi$    $\frac{\pi}{3}$    $\frac{\pi}{2}$    $\pi$

18. The circumradius  $R$  of a triangle with sides  $a, b, c$  is equal to:   $\frac{abc}{\Delta}$    $\frac{abc}{4\Delta}$    $\frac{4abc}{\Delta}$    $\frac{4\Delta}{abc}$

19. For what value of  $x$ ,  $\tan(x - 30^\circ) = \cot x$    $90^\circ$    $60^\circ$    $120^\circ$    $150^\circ$

20. What is the solution of  $\sec x = 2$  in the interval  $[0, \pi]$ ?   $\left\{\frac{\pi}{6}\right\}$    $\left\{\frac{\pi}{3}\right\}$    $\left\{\frac{\pi}{3}\right\}$    $\left\{\frac{\pi}{6}\right\}$

—1HA-I 2211-7111 (HA)—

ROLL NUMBER					



# MATHEMATICS HSSC-I

38

Time allowed: 2:35 Hours

Total Marks Sections B and C: 80

NOTE: Attempt any twelve parts from Section 'B' and any four questions from Section 'C' on the separately provided answer book. Use supplementary answer sheet i.e. Sheet-B if required. Write your answers neatly and legibly. Graph paper will be provided on request.

### SECTION - B (Marks 48)

Q. 2 Attempt any TWELVE parts. All parts carry equal marks. (12 x 4 = 48)

- (i) Simplify  $\frac{9}{\sqrt{5}+\sqrt{-4}}$  in the form of  $a+bi$
- (ii) If  $U$  = the set of the English alphabets,  $A$  and  $B$  are subsets of  $U$ , where  $A = \{x | x \text{ is a vowel}\}$ ,  $B = \{y | y \text{ is a consonant}\}$ , then verify the de Morgan's Laws (i)  $(A \cup B)' = A' \cap B'$  (ii)  $(A \cap B)' = A' \cup B'$
- (iii) Construct the truth table for the biconditional  $p \leftrightarrow q$
- (iv) If  $A = [1 \ 1+i \ i]$ , then find  $(\overline{A})' A$
- (v) Without expansion, show that  $\begin{vmatrix} 2 & 3 & -1 \\ 1 & 1 & 0 \\ 2 & -3 & 5 \end{vmatrix} = 0$
- (vi) Find the numerical value of  $k$  if polynomial  $x^3 + kx^2 - 7x + 6$  has remainder 4 when divided by  $x - 2$
- (vii) Find the two consecutive numbers whose product is 72
- (viii) If 5, 8 are two arithmetic means between  $a$  and  $b$ , then find  $a$  and  $b$
- (ix) Find 9th term of the harmonic sequence  $-\frac{1}{5}, -\frac{1}{3}, -1, \dots$
- (x) Find values of  $n$  and  $r$ , when  ${}^n C_r = 56$  and  ${}^n P_r = 336$
- (xi) If  $x$  is so small that its square and higher powers can be neglected, then show that  $\frac{\sqrt{4+x}}{(1+x)^3} \cong 2 - \frac{23}{4}x$
- (xii) Show that the area of a sector of a circular region of radius  $r$  is  $\frac{1}{2}r^2\theta$ , where  $\theta$  is the circular measure of the central angle of the sector.
- (xiii) If  $\cot \theta = \frac{4}{3}$  and the terminal arm of the angle is not in the quadrant-I, find the values of  $\cos \theta$  and  $\operatorname{cosec} \theta$
- (xiv) Show that  $\frac{\cos(\pi + \theta) \sec(\pi - \theta)}{\sin^2(\pi + \theta) \cdot \tan(\pi - \theta)} = -\cot \theta \cdot \operatorname{cosec}^2 \theta$
- (xv) Prove that  $\cot 2x = \frac{\sin x - \sin 3x}{\cos 3x - \cos x}$
- (xvi) Show that  $\tan^{-1}\left(\frac{27}{11}\right) - \tan^{-1}\frac{8}{19} = \frac{\pi}{4}$

### SECTION - C (Marks 32)

Note: Attempt any FOUR questions. All questions carry equal marks. (4 x 8 = 32)

- Q. 3 Solve the following system by reducing their augmented matrix to the echelon form
 
$$\begin{matrix} x_1 + 4x_2 + 2x_3 = 2 \\ 2x_1 + x_2 - 2x_3 = 9 \\ 2x_1 + 2x_2 - 2x_3 = 12 \end{matrix}$$
- Q. 4 Solve the system of simultaneous equations:
 
$$\begin{matrix} 3x + 2y = 7 \\ 3x^2 = 25 + 2y^2 \end{matrix}$$
- Q. 5 (a) Resolve  $\frac{2x^4}{(x+3)(x-2)^2}$  into partial fractions  
 (b) Find the sum  $S_n$  of the Arithmetic Series  $a + (a+d) + (a+2d) + \dots + (a+(n-1)d)$
- Q. 6 Find the sum of the following series to n-terms:  $1 + (1+2) + (1+2+3) + \dots$
- Q. 7 If  $2y = \frac{1}{2^2} + \frac{1 \cdot 3}{2! \cdot 2^4} + \frac{1 \cdot 3 \cdot 5}{3! \cdot 2^6} + \dots$  then prove that  $4y^2 + 4y - 1 = 0$
- Q. 8 Without using calculator/table prove that  $\sin 10^\circ \sin 30^\circ \sin 50^\circ \sin 70^\circ = \frac{1}{16}$