

# Investigating cold light: The *R* package Luminescence - signal, statistics and dating of environmental dynamics -

Sebastian Kreutzer<sup>1,2,\*</sup>, Steve Grehl<sup>3</sup>, Michael Dietze<sup>4</sup>, Christoph Burow<sup>5</sup>, Margret C. Fuchs<sup>6</sup>,  
Manfred Fischer<sup>7</sup>, Christoph Schmidt<sup>7</sup>

1. IRAMAT-CRP2A, Université Bordeaux-Montaigne, Maison de l'Archéologie, Esplanade des Antilles, 33607 Pessac Cedex, France
  2. Department of Geography, Justus-Liebig-University Giessen, Senckenbergstrasse 1, 35390 Giessen, Germany
  3. Freiberg Instruments GmbH, Delfter Str. 6, 09599 Freiberg, Germany
  4. Section 5.1 Geomorphology, GFZ German Research Centre for Geosciences, 14473 Potsdam, Germany
  5. Institute for Geography, University of Cologne, 50923 Cologne, Germany
  6. Alfred Wegener Institute for Polar and Marine Research, Department of Periglacial Research, Telegraphenberg A43, 14473 Potsdam, Germany
  7. Geographical Institute, Geomorphology, University of Bayreuth, 95440 Bayreuth, Germany
- \*Contact author: [sebastian.kreutzer@geogr.uni-giessen.de](mailto:sebastian.kreutzer@geogr.uni-giessen.de)

**Keywords:** Luminescence Dating, Geosciences, Dosimetry, Signal Analysis, Dynamic GUI

Earth surface processes decisively shape our planet. They are the primary mediators of environmental change and directly affect human societies. To decipher the timing and rates of Earth surface processes, throughout the last 250,000 years one numerical dating method has reached paramount importance: Luminescence dating. This method provides robust numerical data on environmental changes (a) due to the fact that the luminescence signal is reset by daylight exposure or heating and (b) the advantage of using nearly ubiquitously available mineral grains of quartz or feldspar.

During the last decades more and more ages based on luminescence dating have been requested and the method has been considerably enhanced. However, an increasing data complexity demands for a flexible and scalable software solution for data analysis. For more innovative measurements, existing software solutions (e.g. 'Analyst', Duller, 2007) are limited, especially regarding new experimental measurements.

Therefore, in 2012 the *R* package **Luminescence** has been introduced (Kreutzer et al., 2012, Dietze et al., 2013, Fuchs et al., *subm.*). The package is a toolbox intended to provide customised solutions for a variety of requirements (e.g. data import, statistical analysis, graphical output). The used algorithms and statistical treatments are always transparent and the user remains in control, of combining and adjusting algorithms by taking advantage of the wide range of functions available in *R*.

Our contribution (1) summarizes the concept of the *R* package **Luminescence** and focusses on some conceptual aspects and selected practical examples. (2) We present a sneak preview on a dynamic graphical user interface written in *Java* to make the functions of the package available to users who are not familiar with *R* but wants to access the package functionality.

## References

- Dietze, M., Kreutzer, S., Fuchs, M.C., Burow, C., Fischer, M., Schmidt, C. (2013). A practical guide to the *R* package Luminescence, *Ancient TL* 31, 11–18.
- Duller, G.A.T. (2007). *Analyst*. Unpublished manual.
- Fuchs, M.C, Kreutzer, S., Burow, C., Dietze, M., Fischer, M., Schmidt, C., Fuchs, M. (*subm.*). A practical workflow for data analyses in luminescence dating using the *R* package 'Luminescence': A case study from the Panj River, Pamir, submitted to *Quaternary International*
- Kreutzer, S., Schmidt, C., Fuchs, M.C., Dietze, M., Fischer, M., Fuchs, M. (2012). Introducing an *R* package for luminescence dating analysis. *Ancient TL* 30, 1–8.