

EMMAgeo – end-member modelling analysis of grain-size data

Michael Dietze^{1*}, Elisabeth Dietze²

1. GFZ German Research Centre for Geosciences

Section 5.1 Geomorphology

Telegrafenberg, F427

14473 Potsdam, Germany

2. GFZ German Research Centre for Geosciences

Section 5.2 Climate Dynamics and Landscape Evolution

Telegrafenberg, F451

14473 Potsdam, Germany

*Contact author: mdietze@gfz-potsdam.de

Keywords: EMMA, end-member modelling, grain-size data, Earth surface process, eigenspace

Earth surface processes decisively shape our planet. Identifying and quantifying their rates and impact is one of the major challenges of current geoscientific research. The changing grain-size distributions of deposits (e.g. lake sediments, ocean floor deposits, landforms accumulated by wind-blown sediment) reflect their formation by distinct sediment transport processes and provide a fruitful possibility to unravel the contribution of such processes. However, interpreting sediment transport processes from grain-size data in terrestrial archives runs into problems when source- and process-related grain-size distributions become mixed during deposition. A powerful approach to overcome this ambiguity is to statistically “unmix” the samples. Typical algorithms use eigenspace decomposition and techniques of dimension reduction.

This contribution presents the package **EMMAgeo** (Dietze & Dietze, 2013) for the free statistical software *R*. It bases on an end-member modelling algorithm originally presented as *Matlab*-script (Dietze et al., 2012) and contains several extensions and added functionality. The package comprises 14 functions. It supports simple modelling of grain-size end-member loadings and scores (eigenspace extraction, factor rotation, data scaling, non-negative least squares solving) along with several measures of model quality. It also provides pre-processing tools (grain-size scale conversions, weight factor limit inference, determination of minimum, optimum and maximum number of meaningful end-members) and allows to model data sets with user-defined end-member loadings. **EMMAgeo** also supports uncertainty estimation from a series of plausible model runs and determination of robust end-members.

The contribution depicts important package functions, thereby illustrating how large data sets of artificial and natural grain-size samples from different depositional environments can be analysed to infer and quantify process-related proxies (Dietze et al., 2014) that can be used to better reconstruct environmental conditions in the past (and to learn for future environmental change).

References

- Dietze, M., E. Dietze (2013). EMMAgeo: End-member modelling algorithm and supporting functions for grain-size analysis. R package version 0.9.1. <http://CRAN.R-project.org/package=EMMAgeo>.
- Dietze, E., K. Hartmann, B. Diekmann, J. Ijmker, F. Lehmkuhl, S. Opitz, G. Stauch, B. Wünnemann, A. Borchers (2012). An end-member algorithm for deciphering modern detrital processes from lake sediments of Lake Donggi Cona, NE Tibetan Plateau, China. *Sedimentary Geology*, 243-244, 169-180.
- Dietze, E., F. Maussion, M. Ahlborn, B. Diekmann, K. Hartmann, K. Henkel, T. Kasper, G. Lockot, S. Opitz, T. Haberzettl (2014). Sediment transport processes across the Tibetan Plateau inferred from robust grain-size end members in lake sediments. *Climate of the Past*, 10, 1, 91-106.