

# Integrating R-INLA with R spatial packages and ggplot2

Thomas Jagger and James Elsner

Florida State University

\*Contact author: [Thomas Jagger](#)

**Keywords:** Bayesian, Spatial, Graphics

We develop a model for the space time distribution of tornado count data aggregated yearly over counties in Kansas in the USA. We use a Bayesian model to analyze local counts. A hierarchical model is chosen, in which the local counts are assumed to have a negative binomial distribution when conditioned on the distribution's mean and count parameters. Next in the hierarchy, the mean and count parameters each have a distribution, where the mean is assumed to be a spatial Gaussian process. The spatial Gaussian process is composed of a mean that is linearly regressed onto local and global covariates. Both the intercept and the coefficients are allowed to vary spatially using intrinsic conditional autoregressive (ICAR) priors [1]. Finally, the model is fit by an INLA (Integrated Nested Laplace Approximation) [4] using the **R-INLA** [3].

We demonstrate the use of the R package **rgdal** to read in the tornado shape files, with R packages **maps** and **maptools** to generate a counties shape file. We use functions from the R **sp** and **spdep** packages to overlay tornado tracks onto the counties and generate a spatial neighborhood list [2], followed by `ddply()` from the **plyr** R package to generate yearly county tornado counts. We perform our analysis using **R-INLA** package to estimate posterior densities and means. We generate publication quality plots of these results using the **ggplot2** R package [5]. The example code and presentation will be created using *R-Studio* and published on [Rpubs](#).

## References

- [1] Besag, J., J. York, and A. Mollie (1991). Bayesian image restoration, with two applications in spatial statistics. *Annals of the Institute of Statistical Mathematics* 43(1), 1–59.
- [2] Bivand, R., E. Pebesma, and V. Gomez-Rubio (2008). *Applied Spatial Data Analysis with R*. New York: Springer.
- [3] Rue, H. (2012). *The R-INLA project*. R-INLA <http://www.r-inla.org>.
- [4] Rue, H., S. Martino, and N. Chopin (2009). Approximate Bayesian inference for latent Gaussian models using integrated nested Laplace approximations (with discussion).
- [5] Wickham, H. (2009). *ggplot2: elegant graphics for data analysis*. Springer New York. GGPlot2 <http://had.co.nz/ggplot2/book>.