

SUPPORTIVE CARE

DL-Chol Plus Powder

Water Soluble Powder

For veterinary use only



GENERAL INFORMATION

Product name	DL-Chol Plus Powder
Dosage form	Water Soluble Powder
Target species	Poultry
Route	Oral, via drinking water / feed where stated

COMPOSITION

Each gram contains

Choline Chloride	80 mg	L-Lysine	35 mg
DL Methionine	55 mg	Zinc Bacitracin (10%)	15 mg
Vitamin E	3 mg	Nicotinic Acid	3 mg

(As per innovator's specifications)

PRODUCT DESCRIPTION

Role of Choline Chloride, L-lysine & DL Methionine

Choline chloride, L-Lysine, and DL-methionine are vital in poultry.

Choline chloride supports liver function and nerve signaling, while L-Lysine aids protein synthesis and immune function.

DL-methionine is crucial for feather growth and overall development.

These supplements ensure optimal health, growth in poultry and livestock and egg production in poultry.

INDICATIONS

- DL-CHOL Plus Powder is a balanced vitamin and amino acid used to treat heat stress, decrease chick mortality during rapid growth and prevent the accumulation of fat in livers and kidneys.
- It also helps to prevent decrease egg size and production percentage.
- Vitamin E in DL- CHOL PLUS is a fat-soluble compound and antioxidant needed for a strong immunity and skeletal strength
- Birds suffering from growth and development Birds suffering from Muscle and Tissue Problems. Problems.

DOSAGE & ADMINISTRATION

- Poultry
 - Prevention: 1 gm in 2 liter of drinking water for 7 days.Treatment: 1gm in 1 liter of drinking water for 7 days
 - Calves, goats: 5 g per animal daily, during 1 week.
 - Horses: 10 g per animal daily, during 1 week.
 - Dairy cattle: 8 g per animal daily, during 1 week.
- Prepare fresh medicated water and ensure uniform access for the flock.

PRECAUTIONS

Use DL-Chol Plus Powder under veterinary supervision only. Do not exceed the recommended dosage. Avoid administering to birds with known hypersensitivity to any of the components in the formulation.

WITHDRAWAL PERIOD

Adhere to the specified withdrawal period before consuming bird products from treated birds, as directed by regulatory authorities.

STORAGE

Store in a cool, dry place, protected from direct sunlight and moisture. Keep out of reach of children.

PACKAGING

Refer to the approved commercial pack label for pack size, batch details, manufacturing date and expiry date.

IMPORTANT: This sheet summarizes supplied product literature. Always follow the current approved pack label, veterinary direction and applicable regulatory requirements.