

ANTIBACTERIAL

Floxy N- Mak Water Soluble Powder

Water Soluble Powder

For veterinary use only



GENERAL INFORMATION

Product name	Floxy N- Mak Water Soluble Powder
Dosage form	Water Soluble Powder
Target species	Poultry
Route	Oral, via drinking water / feed where stated

COMPOSITION

Each gram contains

Neomycin Sulphate 150 mg

Florfenicol 100 mg

Oxytetracycline HCL 300 mg

(As per innovator's specifications)

PRODUCT DESCRIPTION

Floxy-N-Mak is a water-soluble veterinary medicine containing a balanced blend of neomycin sulfate, florfenicol, and oxytetracycline HCl, targeting bacterial infections in animals.

Neomycin sulfate provides broad-spectrum antibiotic action, while florfenicol and oxytetracycline HCl enhance efficacy against diverse bacterial strains.

This specialized combination ensures versatility and reliability in treating various infections.

Its water-soluble formulation enables convenient administration and rapid absorption for swift veterinary care.

INDICATIONS

- Floxy-N-Mak is a versatile veterinary medicine combating a broad spectrum of bacterial infections in animals, including respiratory, gastrointestinal, urinary, and reproductive tract infections, as well as skin and soft tissue disorders.
- It effectively treats conditions like CRD, CCRD, air sacculitis, pericarditis, and perihepatitis associated Mycoplasmosis, enteritis, urinary tract infections, and reproductive tract infections, aiding in the recovery and management of these ailments.
- Floxy-N-Mak's efficacy extends to bacterial skin infections and soft tissue disorders, promoting healing and preventing further infection spread.
- It's Important to note that Floxy-N-Mak should be administered under the guidance of a veterinarian, who will assess the specific condition of the animal and prescribe the appropriate dosage and duration of treatment based on the nature and severity of the infection.
- Birds suffering from

DOSAGE & ADMINISTRATION

- Poultry
- Prevention: 1 gm in 6 liters of drinking water for 4-5 days.
- Treatment: 1 gm in 4 liters of drinking water for 4-5 days.

Prepare fresh medicated water and ensure uniform access for the flock.

PRECAUTIONS

Use under veterinary supervision. Do not exceed the recommended dosage.

WITHDRAWAL PERIOD

Meat & Eggs 5 days.

INSTRUCTION

STORAGE

Store in a cool, dry place, protected from direct sunlight and moisture. Keep out of reach of children.

PACKAGING

Refer to the approved commercial pack label for pack size, batch details, manufacturing date and expiry date.

IMPORTANT: This sheet summarizes supplied product literature. Always follow the current approved pack label, veterinary direction and applicable regulatory requirements.