

ANTIBACTERIAL

MAKFLOR-23 oral Liquid

Oral Liquid

For veterinary use only



GENERAL INFORMATION

Product name	MAKFLOR-23 oral Liquid
Dosage form	Oral Liquid
Target species	Poultry
Route	Oral, via drinking water

COMPOSITION

Each ml contains
Florfenicol.....230 mg
(As per innovator's specifications)

PRODUCT DESCRIPTION

ROLE OF FLORFENICOL

Florfenicol is a broad-spectrum antibiotic used to treat bacterial infections in poultry, particularly respiratory and enteric infections.

Makflor-23 Oral Liquid, containing florfenicol, that inhibits bacterial protein synthesis by binding to the 50S subunit of the bacterial ribosome, thus preventing peptide bond formation and ultimately leading to bacterial cell death

INDICATIONS

- Makflor-23 oral liquid is indicated to treat broad spectrum antibiotic covering gram +ve and gram-ve bacteria in respiratory and gastrointestinal infection in poultry including pneumonia, Mycoplasma, CRD, CCRD, fowl cholera, fowl typhoid, coryza, E-coli infection, necrotic enteritis, perihepatitis, pericarditis and endocarditis.
- Hepatitis hydropericardium in poultry Birds affected by the respiratory infections

DOSAGE & ADMINISTRATION

- Prevention: 1 ml per 4 liters of drinking water for 3-5 days
- Treatment: 1 ml per 2 liters of drinking water for 3-5 days. (15-30 mg/kg body weight)

Prepare fresh medicated water and ensure uniform access for the flock.

PRECAUTIONS

Precautions: Use Makflor-23 Oral Liquid under veterinary supervision only. Do not exceed the recommended dosage. Avoid use in birds with known hypersensitivity to florfenicol.

WITHDRAWAL PERIOD

Withdrawal Period: Adhere to the specified withdrawal period before consuming animal products from treated animals, as directed by regulatory authorities.

STORAGE

Store in a cool, dry place, protected from direct sunlight and moisture. Keep out of reach of children.

PACKAGING

Refer to the approved commercial pack label for pack size, batch details, manufacturing date and expiry date.

IMPORTANT: This sheet summarizes supplied product literature. Always follow the current approved pack label, veterinary direction and applicable regulatory requirements.