

ANTIBACTERIAL

Sulpha T- Mak Suspension

Oral Liquid

For veterinary use only



GENERAL INFORMATION

Product name	Sulpha T- Mak Suspension
Dosage form	Oral Liquid
Target species	Poultry
Route	Oral, via drinking water

COMPOSITION

Each ml contains

Sulphadiazine 400 mg

Trimethoprim 80 mg

(As per innovator's specifications)

PRODUCT DESCRIPTION

Role of Sulphadiazine

Sulphadiazine is a sulfonamide antibiotic that is used to treat a broad range of bacterial infections in animals. Its mechanism of action involves inhibiting bacterial synthesis of folic acid, which is essential for bacterial growth and replication. It is also effective against urinary tract infections, rheumatic fever, and some respiratory infections.

Role of Trimethoprim

Trimethoprim is an antibiotic that belongs to the diaminopyrimidine class, primarily used to treat bacterial infections by inhibiting the bacterial enzyme dihydrofolate reductase.

This inhibition blocks the production of tetrahydrofolic acid, a form of folic acid that bacteria require for synthesizing nucleic acids and proteins.

INDICATIONS

- SULPHA T-MAK Suspension is indicated for infections caused by microorganisms susceptible to trimethoprim and/or sulfamethoxazole, such as gastrointestinal, respiratory, or urogenital infections, as well as general infections in poultry.
- Birds suffering from Birds suffering from Respiratory Tract Birds suffering from
- Urinary tract Infections Infections. Grastointestinal Infections.

DOSAGE & ADMINISTRATION

- Amounts to be administered and administration route
Dose
- The recommended daily dose rate should be 15 mg active ingredients per 1kg bodyweight daily. This may be achieved, for example, by administering the suspension to the flock at the rate of 1 ml per 5 litres (or one gallon) of drinking water daily for six-week-old broilers under average conditions. Treatment should continue for five days.
- Administration: By drinking water medication.
Prepare fresh medicated water and ensure uniform access for the flock.

PRECAUTIONS

Use under veterinary supervision. Do not exceed the recommended dosage.

WITHDRAWAL PERIOD

Follow the approved product label and applicable regulatory requirements.

STORAGE

Store in a cool, dry place, protected from direct sunlight and moisture. Keep out of reach of children.

PACKAGING

Refer to the approved commercial pack label for pack size, batch details, manufacturing date and expiry date.

IMPORTANT: This sheet summarizes supplied product literature. Always follow the current approved pack label, veterinary direction and applicable regulatory requirements.