M. A. Kamil Farma (Pvt.) Limited

PRODUCT INFORMATION SHEET

PIS-MAK-032

ANTIBACTERIAL

Xinmak oral liquid

Oral Liquid

For veterinary use only

GENERAL INFORMATION

Product name	Xinmak oral liquid
Dosage form	Oral Liquid
Target species	Poultry
Route	Oral, via drinking water

COMPOSITION

Each gram contains

Tylosin (as tartrate) ……. 100 mg

Doxycycline as HCl ……. 200 mg

Colistin Sulphate …….500,000 MIU

Bromhexine HCl…….. 5 mg

(As per innovator's specifications)

PRODUCT DESCRIPTION

Role of Doxycycline as HCl

Doxycycline hydrochloride, a tetracycline antibiotic, treats bacterial infections including respiratory infections, enteritis, and chlamydiosis. Its broad spectrum of activity against Gram-positive and Gram-negative bacteria is valuable in controlling infectious diseases, especially in poultry.

Role of Tylosin (as Tartrate)

Tylosin tartrate is widely used to treat respiratory infections, particularly those caused by Mycoplasma species. It is effective against a variety of Gram-positive bacteria and has been utilized to control respiratory diseases such as chronic respiratory disease (CRD) in poultry.

Role of Colistin Sulphate

INDICATIONS

- Birds suffering from respiratory Birds suffering from Birds suffering from Mycoplasma
- tract Infections. Enteric Infections. Infections.

DOSAGE & ADMINISTRATION

- Treatment: 1 ml in 2 liters of drinking water for 3- 5 days.
- Prevention: 1 ml in 4 liters of drinking water for 3- 5 days.
- Duration
- Continue treatment with Xinmak oral liquid for the prescribed duration or as recommended by the veterinarian to effectively combat bacterial and mycoplasma infections.
- Mix the prescribed amount of Xinmak oral liquid thoroughly in clean drinking water, ensuring uniform distribution. Provide the medicated water as the sole source of drinking water for the birds during the treatment period.

Prepare fresh medicated water and ensure uniform access for the flock.

PRECAUTIONS

Use Xinmak oral liquid under veterinary supervision only. Do not exceed the recommended dosage. Avoid administering to birds with known hypersensitivity to any of the components in the formulation.

WITHDRAWAL PERIOD

Adhere to the specified withdrawal period before consuming bird products from treated birds, as directed by regulatory authorities.

STORAGE

Store in a cool, dry place, protected from direct sunlight and moisture. Keep out of reach of children.

PACKAGING

Refer to the approved commercial pack label for pack size, batch details, manufacturing date and expiry date.

IMPORTANT: This sheet summarizes supplied product literature. Always follow the current approved pack label, veterinary direction and applicable regulatory requirements.

Manufactured by: M. A. Kamil Farma (Pvt.) Limited
Plot No. H-17, Super Highway SITE-II, Karachi, Pakistan

+92 335 2249111
info@makamilfarma.com
www.makamilfarma.com