

PEFLOXKAM-10

Solution
Pefloxacin 10%



**The Trusted Guard
Against Poultry Threats.**

PEFLOXKAM-10

Solution

پیفلوکس کیم - ۱۰ (سلوٹن)

COMPOSITION:

Each ml contains:

Pefloxacin Methanesulfonate 139.6 mg eq. to

Pefloxacin 100 mg

(As per innovator's specifications)

Role of Pefloxacin Methanesulfonate:

Pefloxacin methanesulfonate, is a potent fluoroquinolone antibiotic used primarily in veterinary medicine. It works by inhibiting bacterial DNA gyrase and topoisomerase IV, essential enzymes for DNA replication and transcription, leading to the rapid death of susceptible bacteria. Pefloxacin is effective against a broad spectrum of Gram-negative and Gram-positive bacteria and is used to treat infections such as those affecting the urinary tract, respiratory system, and gastrointestinal tract in animals.

Indications:

Pefloxkam-10 Solution is indicated for the treatment of CRD, CCRD, Pullorum disease, Fowl typhoid, Colibacillosis, Fowl cholera, Infectious coryza and staphylococcus infection in poultry.



Birds suffering from skin and soft tissues infection.



Birds suffering from Gastrointestinal infections.



Birds suffering from Respiratory tract Infections.

Treatment Protocol with Pefloxkam-10 Solution:

Dosage and Administration:

Administer 1 ml per 1-2 liters of drinking water for 3-4 days (10mg/kg BW).

Chicken-Dilute 5, 10ml (as 500, 1000mg of Pefloxacin) with 10L drinking water and administer orally for 3-4 days.

Note:

Always consult a veterinarian before starting treatment with Pefloxkam-10 Solution to ensure it is appropriate for the specific type of infection and bird species involved. Proper diagnosis and sensitivity testing are recommended to optimize treatment outcomes and prevent the development of antibiotic resistance.

Packaging: Pefloxkam-10 available in 1 Litre.

علامات:

پیفلوکس کیم - ۱۰ سلوٹن سی آر ڈی، سی سی آر ڈی، پلورم جیسی بیماریوں، فائول ٹائیفائیڈ، کولیسیلوسس، فائول کالیرہ، کورائیڈ اور اسٹیفیلوکوکس انفیکشن کے علاج کیلئے تجویز کی جاتی ہے۔

خوراک و طریقہ استعمال:

ایلی لیرنی ۱ سے ۲ لیر پینے کے پانی میں ۳ سے ۴ دن (۱۰ ایلی گرام فی کلو گرام جسمانی وزن کے حساب سے) دیں۔
چکن ڈالوٹ ۵ سے ۱۰ ایلی لیر (۵۰۰ سے ۱۰۰۰ ایلی گرام فی کلوکسیس کے طور پر)
ایلی لیر پینے کے پانی کے ساتھ اور ۳ سے ۴ دن تک دیں۔

Manufactured By:
M. A. Kamil Farma (Pvt.) Limited
Karachi, Pakistan.

mak
M. A. Kamil Farma (Pvt.) Limited