

TYLOKAM-100

Soluble Powder

Tylosin (as tartrate) 100%



TYLOKAM-100

Defence against Poultry Infections
like Mycoplasma, C.R.D., C.C.R.D.

TYLOKAM-100

Soluble Powder

ٹائلوکیم-۱۰۰

(حل پذیر پاؤڈر)



COMPOSITION:

Each gram contains:
Tylosin (as tartrate) 1000 mg
(As per innovator's specification).

Product Description:

The veterinary medicine "Tylokam-100 Soluble Powder," with each gram containing Tylosin as Tartrate at a concentration of 1000mg, is indicated for the treatment of bacterial infections in poultry. Tylosin tartrate, a macrolide antibiotic, inhibits bacterial protein synthesis by binding to the 50S ribosomal subunit. This interference halts peptide chain elongation, preventing bacterial growth.

INDICATIONS

Tylokam -100 Soluble Powder is used for prevention and treatment of infections caused by micro-organisms susceptible to Tylosin, like in Mycoplasma, C.R.D, C.C.R.D, Sinusitis, Airsacculitis and Gastrointestinal infections in poultry.



Birds suffering from
Enteric Infections.



Birds suffering from
respiratory tract



Birds suffering from
Mycoplasma Infections.



Birds suffering from
Chronic Respiratory

Treatment Protocol with Tylokam-100 Soluble Powder for Birds:

DOSAGE:

Poultry: 1 gm per 2 liter of water for 3-5 days.
In feed: 300-500 gm/ 1000 kg feed for 3-5 days.
Calves: 1 gm per 40 kg body weigh

Precautions:

Use **Tylokam-100 Soluble Powder** under veterinary supervision only.
Do not exceed the recommended dosage. Avoid administering to birds with known hypersensitivity to tylosin or other macrolide antibiotics.

Withdrawal Period:

Adhere to the specified withdrawal period before consuming bird products from treated birds, as directed by regulatory authorities.
Always consult with a veterinarian for accurate diagnosis, proper treatment recommendations, and guidance on the use of **Tylokam-100 Soluble Powder** for birds

علامات:

ٹائلوکیم-۱۰۰ حل پذیر پاؤڈر مرغیوں میں مانگو پلازمہ، سی آر ڈی، سی سی آر ڈی، ایئر سیکولائٹس، سائنوسائٹس، اور آئنوں میں موجود ضرر رساں بیکٹیریا اور دیگر جراثیم کے خلاف مؤثر ہے۔

خوراک و طریقہ استعمال:

پولٹری: 1 گرام فی 2 لیٹر پانی میں تین سے پانچ دن تک دیں۔
300 سے 500 گرام **ٹائلوکیم-۱۰۰** حل پذیر پاؤڈر فی ٹن فیڈ میں ملا کر تین سے پانچ دن تک دیں۔



Manufactured By:
M. A. Kamil Farma (Pvt.) Limited
Karachi, Pakistan.



M. A. Kamil Farma (Pvt.) Limited