



Logistic Regression Inference

Julie Deeke

Statistics with Python Course Developer



Cartwheel Data


Random sample of 25 adults attempted cartwheels

Primary Variable of interest: Cartwheel completion



Based on age, can we predict whether a cartwheel is completed?

Logistic Regression Line



Generalized Linear Model Regression Results

Dep. Variable:	CompleteGroup	No. Observations:	25
Model:	GLM	Df Residuals:	23
Model Family:	Binomial	Df Model:	1
Link Function:	logit	Scale:	1.0
Method:	IRLS	Log-Likelihood:	-12.534
Date:	Tue, 27 Nov 2018	Deviance:	25.068
Time:	16:11:20	Pearson chi2:	22.4
No. Iterations: 6			

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

Confidence Interval

Best Estimate \pm Margin of Error

Sample slope \pm “a few” \cdot estimated standard error

Confidence Interval

Best Estimate \pm Margin of Error

Sample slope \pm “a few” \cdot estimated standard error

$$b_1 \pm t^* \text{se}(b_1)$$

Confidence Interval

Best Estimate \pm Margin of Error

Sample slope \pm “a few” \cdot estimated standard error

$$b_i \pm t^* \text{se}(b_i)$$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

Confidence Interval

Best Estimate \pm Margin of Error

Sample slope \pm “a few” \cdot estimated standard error

$b_1 \pm t^* se(b_1)$


	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

Confidence Interval

Best Estimate \pm Margin of Error

Sample slope \pm “a few” \cdot estimated standard error

$b_1 \pm t^* se(b_1)$

 $df = n-2$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

Confidence Interval

Best Estimate \pm Margin of Error

Sample slope \pm “a few” \cdot estimated standard error

$df = n - 2$

$b_1 \pm t^* se(b_1)$

$0.2096 \pm 2.069 (0.171)$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

Confidence Interval

Best Estimate \pm Margin of Error

Sample slope \pm “a few” \cdot estimated standard error

$df = n - 2$

$b_1 \pm t^* se(b_1)$

$0.2096 \pm 2.069 (0.171)$

$(-0.1442, 0.5634)$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

Confidence Interval

Best Estimate \pm Margin of Error

Sample slope \pm “a few” \cdot estimated standard error

$df = n - 2$

$b_1 \pm t^* se(b_1)$

$0.2096 \pm 2.069 (0.171)$

$(-0.1442, 0.5634)$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

IVQ

Does it seem reasonable that there is a significant slope?

We'll continue working through the hypothesis testing framework before coming back to the answer of the IVQ.

Hypothesis Testing

Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

$$\text{Test statistic: } \frac{b_1 - 0}{\text{s.e.}(b_1)}$$

Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

Test statistic: $\frac{b_1 - 0}{\text{s.e.}(b_1)}$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

Test statistic: $\frac{b_1 - 0}{\text{s.e.}(b_1)} = \frac{0.2096}{0.171}$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

Test statistic: $\frac{b_1 - 0}{\text{s.e.}(b_1)} = \frac{0.2096}{0.171} = 1.225$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

Test statistic: $\frac{b_1 - 0}{\text{s.e.}(b_1)} = \frac{0.2096}{0.171} = 1.225$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

Test statistic: $\frac{b_1 - 0}{\text{s.e.}(b_1)} = \frac{0.2096}{0.171} = 1.225$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

With our p-value of about 0.221, we would fail to reject the null hypothesis and cannot conclude that we have a significantly linear relationship between age and the log odds of the probability of successfully completing a cartwheel.

Test statistic: $\frac{b_1 - 0}{\text{s.e.}(b_1)} = \frac{0.2096}{0.171} = 1.225$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

Summary

Logistic regression for a predicted variable with two options
~correct/incorrect or success/failure

Summary

Logistic regression for a predicted variable with two options

~correct/incorrect or success/failure

Confidence Intervals and Hypothesis Tests follow similar format as linear regression

~changing interpretation to the logistic context

Summary

Logistic regression for a predicted variable with two options
~correct/incorrect or success/failure

Confidence Intervals and Hypothesis Tests follow similar format as Linear Regression

~changing interpretation to the logistic context

Coming up, we'll look at an example from NHANES using blood pressure and smoking