



# Interpretations & Assumptions for Two Population Proportion Intervals

---

Julie Deeke

Statistics Course Developer



# Swimming Lessons Confidence Interval

**“range of reasonable values for our parameter”**

With 95% confidence, the population proportion of parents with white children who have taken swimming lessons is 11.23 to 24.77% higher than the population proportion of parents with black children who have taken swimming lessons.

# Intervals for Differences

Is there a difference between two parameters?

If parameters are equal  $\rightarrow$  difference is **0**

If parameters are unequal  $\rightarrow$  difference is not **0**

Look for **0** in the range of reasonable values

# Assumptions

We need to assume that we have **two independent random samples**.

We also need **large enough sample sizes** to assume that the distribution of our estimate is normal. That is, we need  $n_1\hat{p}_1$ ,  $n_1(1-\hat{p}_1)$ ,  $n_2\hat{p}_2$ , and  $n_2(1-\hat{p}_2)$  to all be at least 10.

# Assumptions

We need to assume that we have two independent random samples.

We also need large enough sample sizes to assume that the distribution of our estimate need  $n_1\hat{p}_1$ ,  $n_1(1-\hat{p}_1)$ ,  $n_2\hat{p}_2$ , and  $n_2(1-\hat{p}_2)$  to be at least 10. **In other words, we need at least 10 yes's and at least 10 no's for each sample.**