

Setting up a Test for a Population Proportion

Reed Coots
Statistics Course Developer







Hypothesis Testing

Why do we do Hypothesis Tests?





Hypothesis Testing

Why do we do Hypothesis Tests?

Could the value of the parameter be ____?





Hypothesis Testing

Why do we do Hypothesis Tests?

Could the value of the parameter be ____?

Use data to help support that claim





C.S. Mott Children's Hospital Poll

C.S. Mott Children's Hospital conducted a national poll on an issue in children's health, sleep habits. We will be looking at an example about lack of sleep in teens.





Research Question

In previous years 52% of parents believed that electronics and social media was the cause of their teenager's lack of sleep. Do more parents today believe that their teenager's lack of sleep is caused due to electronics and social media?



Research Question

In previous years 52% of parents believed that electronics and social media was the cause of their teenager's lack of sleep. Do more parents today believe that their teenager's lack of sleep is caused due to electronics and social media?

Population - Parents with a teenager (age 13-18)

Parameter of Interest - p



Research Question

In previous years 52% of parents believed that electronics and social media was the cause of their teenager's lack of sleep. Do more parents today believe that their teenager's lack of sleep is caused due to electronics and social media?

Population - Parents with a teenager (age 13-18)

Parameter of Interest - p

Test for a significant increase in the proportion of parents with a teenager who believe that electronics and social media is the cause for lack of sleep.







$$H_0: p = 0.52$$



 $H_0: p = 0.52$

 $H_a: p? 0.52$



IVQ

What symbol should replace the ?



 $H_0: p = 0.52$ $H_a: p > 0.52$

"Significant Increase"



 $H_0: p = 0.52$

 $H_a: p > 0.52$

Where p is the population proportion of parents with a teenager who believe that electronics and social media is the cause of their teenager's lack of sleep



 $H_0: p = 0.52$

 $H_a: p > 0.52$

Where p is the population proportion of parents with a teenager who believe that electronics and social media is the cause of their teenager's lack of sleep

$$\alpha = 0.05$$





Survey Results

A random sample of **IOI8** parents with a teenager was taken and **56**% said they believe electronics and social media was the cause of their teenager's lack of sleep.





We need a random sample of parents





We need a **random sample** of parents
We also need a **large enough sample size** to ensure our distribution of sample proportions is normal



We need a **random sample** of parents
We also need a **large enough sample size** to ensure our distribution of sample proportions is normal

That is: n·p be at least 10 n·(I-p) be at least 10



We need a **random sample** of parents

We also need a **large enough sample size** to ensure our distribution of sample proportions is normal

That is: $\mathbf{n} \cdot \mathbf{p}$ be at least $10 \longrightarrow \mathbf{n} \cdot \mathbf{p}$

That is: $\mathbf{n} \cdot \mathbf{p}$ be at least $10 \longrightarrow \mathbf{n} \cdot \mathbf{p}_{o}$ $\mathbf{n} \cdot (\mathbf{l} - \mathbf{p})$ be at least $10 \longrightarrow \mathbf{n} \cdot (\mathbf{l} - \mathbf{p}_{o})$






Random Sample





Random Sample V





Random Sample V



$$\mathbf{u} \cdot (\mathbf{I} - \mathbf{b}^{\circ})$$

$$P_{o} = 0.52$$



Random Sample V

$$n \cdot p_o = 1018 \cdot (0.52) = 529$$

 $n \cdot (1-p_o) = 1018 \cdot (1-0.52) = 489$







Testing a One Population Proportion

Reed Coots
Statistics Course Developer





 $H_0: p = 0.52$

 $H_a: p > 0.52$

Best Estimate of p is $\hat{p} = 0.56$

Where p is the population proportion of parents with a teenager who believe that electronics and social media is the cause of their teenager's lack of sleep

$$\alpha = 0.05$$





$$\frac{\hat{p}-p_o}{s.e.}$$



$$\frac{\hat{p}-p_o}{s.e.}$$

$$s.\,e.\,(\hat{p})=\sqrt{rac{p\cdot(1-p)}{n}}$$



$$\frac{\hat{p}-p_o}{s.e.}$$

$$s.\,e.\,(\hat{p})=\sqrt{rac{p\cdot(1-p)}{n}}$$
 \Longrightarrow $s.\,e.\,(\hat{p})=\sqrt{rac{p_0\cdot(1-p_o)}{n}}$



$$\frac{\hat{p}-p_o}{s.e.}$$
 Null $s.e.$ $(\hat{p})=\sqrt{rac{p_o\cdot(1-p_o)}{n}}$



$$\frac{\hat{p}-p_o}{s.e.}$$

Null
$$s.e.(\hat{p}) = \sqrt{\frac{p_0\cdot(1-p_o)}{n}}$$

$$Z = 0.56 - 0.52$$
 0.0157

$$Z = 2.555$$



Test Statistic Interpretation

$$Z = 2.555$$

That means that our observed sample proportion is 2.555 null standard errors above our hypothesized population proportion



• A Z test statistic is another random variable! It has a distribution.



- A Z test statistic is another random variable! It has a distribution.
- The Z test statistic will always follow a N(0, I)



- A Z test statistic is another random variable! It has a distribution.
- The Z test statistic will always follow a N(0, I)
- This is due to us centering and scaling our original data



- A Z test statistic is another random variable! It has a distribution.
- The Z test statistic will always follow a N(0,1)
- This is due to us centering and scaling our original data

$$\hat{p}-p_o \ s.e.(\hat{p})$$




- A Z test statistic is another random variable! It has a distribution.
- The Z test statistic will always follow a N(0,1)
- This is due to us centering and scaling our original data

$$\hat{p}-p_o$$
 Centers Data $s.e.(\hat{p})$



- A Z test statistic is another random variable! It has a distribution.
- The Z test statistic will always follow a N(0,1)
- This is due to us centering and scaling our original data





The P-Value





Conclusions

p-value = $0.0053 < \alpha = 0.05$



Conclusions

p-value = $0.0053 < \alpha = 0.05$ Reject the null hypothesis (H_0 : p = 0.52)



Conclusions

p-value = $0.0053 < \alpha = 0.05$

Reject the null hypothesis (H_0 : p = 0.52)

There is sufficient evidence to conclude that the population proportion of parents with a teenager who believe that electronics and social media is the cause for lack of sleep is greater than 52%.



Summary

- 4 main steps to a hypothesis test
 - Stating hypothesis & select significance level (α)
 - Checking assumptions
 - Calculating a test statistic and getting a p-value from the test statistic
 - Drawing a conclusions from the p-value



Summary

- 4 main steps to a hypothesis test
 - Stating hypothesis & select significance level (α)
 - Checking assumptions
 - Calculating a test statistic and getting a p-value from the test statistic
 - Drawing a conclusions from the p-value
- The Z test statistic distribution is N(0,1)