

Correlation and Convolution in neural networks derivation

Nasy

Jan 27, 2023

Outline

- 1 Introduction
 - Correlation and Convolution
- 2 CNN
- 3 Conv and Corr Calculations
- 4 Conclusion

Two operators \star and \ast

- $(f \star g)(\tau) := \int_{-\infty}^{\infty} f(t)g(t + \tau) dt$
- $(f \ast g)(\tau) := \int_{-\infty}^{\infty} f(t)g(\tau - t) dt$

And for discrete functions:


- $(f \star g)(\tau) := \sum_{-\infty}^{\infty} f(t)g(t + \tau)$
- $(f \ast g)(\tau) := \sum_{-\infty}^{\infty} f(t)g(\tau - t)$

Why?

Problem 1

Two signals f and g are given. They are from a same audio source, but there is a difference in the time when the recording was started. We want to align them.


- $f(x) = \alpha \sin(x)$
- $g(x) = \beta \cos(x)$



Problem 1

Two signals f and g are given. They are from the same piece of audio, but there is a difference in the time when the recording was started. We want to align them.

- Same source, same crests and troughs.
- Keep f , shift g to align with f .
- $\arg \max_{\Delta t} \sum_{-\infty}^{\infty} f(t)g(t + \Delta t)$



Problem 2

There is a function that describes the case where Mr. N is hit.

$$f(t) = 0, 0, 0, 1, 0, 0, 0, \dots$$

And a function describing the change in pain after being hit.

$$g(t) = 3, 2, 1, 0$$

We want a function $h(t)$ to describe the pain of Mr. N.

Problem 2

There is a function that describes the case where Mr. N is hit.

$$f(t) = 0, 0, 0, 1, 0, 0, 0, \dots$$

And a function describing the change in pain after being hit.

$$g(t) = 3, 2, 1; \quad t \in \{0, 1, 2\}$$

$$g(t) = 0; \quad t \in \text{others}$$

We want a function $h(\tau)$ to describe the pain of Mr. N.

- $\text{Pain} = \text{hit} * \text{pain}$
- At the first time hit, where $f(t) = f(3) = 1$, the pain is $g(0) = 3$.
- Next, the pain is $g(1) = 2$, and so on.

Obviously

$$h(\tau) = 0, 0, 0, 3, 2, 1, 0, 0, \dots$$

What if $f(t) = 0, 0, 1, 1, 0, 0, \dots$?

$$h(\tau) = 0, 0, 3, 5, 3, 1, 0, 0, \dots$$

- $h(\tau) = \sum f(t)g(\tau - t)$

Forward (1D)

Correlation $\text{Input}(a) \star \text{kernel}(\text{filter})(w) = \text{output}(z)$

$$[0, 1, 2, 3] \star [1, 2] = [0 \times 1 + 1 \times 2, 1 \times 1 + 2 \times 2, 2 \times 1 + 3 \times 2] = [2, 5, 8]$$

$$[a_1, a_2, a_3, a_4] \star [w_1, w_2, w_3] = [a_1 \times w_1 + a_2 \times w_2 + a_3 \times w_3, a_2 \times w_1 + a_3 \times w_2 + a_4 \times w_3] = [z_1, z_2]$$

Backward (1D)

The δ_i is the gradient from the next layer at point i . And J is the loss function.

$$\frac{\partial J}{\partial a_i} = \sum_j \frac{\partial J}{\partial z_j} \frac{\partial z_j}{\partial a_i} = \sum_j \delta_j w_j$$

Thus,

$$\frac{\partial J}{\partial a_1} = \delta_1 w_1$$

$$\frac{\partial J}{\partial a_2} = \delta_1 w_2 + \delta_2 w_1$$

$$\frac{\partial J}{\partial a_3} = \delta_1 w_3 + \delta_2 w_2$$

$$\frac{\partial J}{\partial a_4} = \delta_2 w_3$$

$$[\delta_1 w_1, \delta_1 w_2 + \delta_2 w_1, \delta_1 w_3 + \delta_2 w_2, \delta_2 w_3]$$

Backward (1D)

$$\frac{\partial J}{\partial a_1} = 0w_3 + 0w_2 + \delta_1 w_1$$

$$\frac{\partial J}{\partial a_2} = 0w_3 + \delta_1 w_2 + \delta_2 w_1$$

$$\frac{\partial J}{\partial a_3} = \delta_1 w_3 + \delta_2 w_2 + 0w_1$$

$$\frac{\partial J}{\partial a_4} = \delta_2 w_3 + 0w_2 + 0w_1$$

$$\begin{aligned} & [0, 0, \delta_1, \delta_2, 0, 0] * [w_1, w_2, w_3] \\ &= [0, 0, \delta_1, \delta_2, 0, 0] \star [w_3, w_2, w_1] = [\delta_1 w_1, \delta_1 w_2 + \delta_2 w_1, \delta_1 w_3 + \delta_2 w_2, \delta_2 w_3] \end{aligned}$$

2D correlation and convolution

Correlation $\text{Input} \star \text{kernel}(\text{filter}) = \text{output}$

$$\begin{array}{|c|c|c|} \hline 0 & 1 & 2 \\ \hline 3 & 4 & 5 \\ \hline 6 & 7 & 8 \\ \hline \end{array} \star \begin{array}{|c|c|} \hline 0 & 1 \\ \hline 2 & 3 \\ \hline \end{array} = \begin{array}{|c|c|} \hline 19 & 25 \\ \hline 37 & 43 \\ \hline \end{array}$$

The shadow part:

$$0 \times 0 + 1 \times 1 + 3 \times 2 + 4 \times 3 = 19$$

Convolutions $\text{Input} * \text{kernel}(\text{filter}) = \text{Input} \star \text{flip}(\text{kernel}(\text{filter})) = \text{output}$

$$\begin{array}{|c|c|c|} \hline 0 & 1 & 2 \\ \hline 3 & 4 & 5 \\ \hline 6 & 7 & 8 \\ \hline \end{array} * \begin{array}{|c|c|} \hline 0 & 1 \\ \hline 2 & 3 \\ \hline \end{array} = \begin{array}{|c|c|} \hline 3 & 2 \\ \hline 1 & 0 \\ \hline \end{array}$$

Forward (2D)

Correlation $\text{Input}(a) \star \text{kernel}(\text{filter})(w) = \text{output}(z)$

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \star \begin{bmatrix} w_{11} & w_{12} \\ w_{21} & w_{22} \end{bmatrix} = \begin{bmatrix} z_{11} & z_{12} \\ z_{21} & z_{22} \end{bmatrix} \quad (1)$$

$$z_{11} = a_{11}w_{11} + a_{12}w_{12} + a_{21}w_{21} + a_{22}w_{22}$$

$$z_{12} = a_{12}w_{11} + a_{13}w_{12} + a_{22}w_{21} + a_{23}w_{22}$$

$$z_{21} = a_{21}w_{11} + a_{22}w_{12} + a_{31}w_{21} + a_{32}w_{22}$$

$$z_{22} = a_{22}w_{11} + a_{23}w_{12} + a_{32}w_{21} + a_{33}w_{22}$$

Backward (2D)

$$\nabla a^{l-1} = \frac{\partial e}{\partial a^{l-1}} = \frac{\partial e}{\partial z^l} \frac{\partial z^l}{\partial a^{l-1}} = \delta^l \frac{\partial z^l}{\partial a^{l-1}} = \delta^l W$$

$$\nabla a_{11} = \delta_{11} w_{11}$$

$$\nabla a_{12} = \delta_{11} w_{12} + \delta_{12} w_{11}$$

$$\nabla a_{13} = \delta_{12} w_{12}$$

$$\begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & \delta_{11} & \delta_{12} & 0 \\ 0 & \delta_{21} & \delta_{22} & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \star \begin{bmatrix} w_{22} & w_{21} \\ w_{12} & w_{11} \end{bmatrix} = \begin{bmatrix} \nabla a_{11} & \nabla a_{12} & \nabla a_{13} \\ \nabla a_{21} & \nabla a_{22} & \nabla a_{23} \\ \nabla a_{31} & \nabla a_{32} & \nabla a_{33} \end{bmatrix} \quad (2)$$

Conv and Corr General Calculation

$$\text{Input}(a) \star \text{kernel}(\text{filter})(w) = \text{output}(z)$$

Basic method Direct calculation

$$\sum_{i=1}^n a_i w_i = z$$

It's an inner product! $\langle A, W \rangle$. And the time complexity is $O(n^2) = O(N_A M_W)$.

Faster one FFT

$$y[n] = f[n] * g[n] \leftrightarrow Y[f] = F[f] G[f]$$

The time complexity is $O(n \log_2 n)$

Conclusion

- Correlation
- Convolution
- Forward and Backward
- Conv and Corr Calculations