

Cystatin C and Long-Term Mortality Among Subjects With Normal Creatinine-Based Estimated Glomerular Filtration Rates

NHANES III (Third National Health and Nutrition Examination Survey)

Cho-Kai Wu, MD,*† Jou-Wei Lin, MD, MPH, PhD,*† James L. Caffrey, PhD,‡
Man-Huei Chang, MPH,|| Juey-Jen Hwang, MD, PhD,*† Yu-Sheng Lin, ScD§
Dou-Liou City and Taipei, Taiwan; Fort Worth, Texas; and Atlanta, Georgia

Objectives	The objective was to test the association of cystatin C (Cys-C) with long-term mortality risk in the subjects with normal creatinine-based estimated glomerular filtration rates (eGFR).
Background	Cys-C has been proposed as a sensitive indicator of renal dysfunction that is associated with cardiovascular events. The predictive value of Cys-C for mortality risk (both cardiovascular and noncardiovascular) and its utility among persons with normal kidney function remains unclear.
Methods	The analysis included 2,990 subjects over 40 years of age with normal eGFR who participated in NHANES III (Third National Health and Nutrition Examination Survey). Normal eGFR was defined by Modification of Diet in Renal Disease (MDRD) equation ≥ 60 ml/min/1.73 m ² . Serum Cys-C was categorized as high, medium, or low. In 1 analysis, the high and low groups were the top and bottom 10%, and in the second analysis, they were the upper and lower thirds. All-cause and cause-specific mortality were obtained from the NHANES III-linked follow-up file through December 31, 2006. Multivariate Cox regression models were applied to assess the association of interest.
Results	Within an average of 13.7 years follow-up, 488 cardiovascular and 719 noncardiovascular deaths occurred. When the first and last deciles were compared, the relative risks were all increased statistically as follows: all-cause, 4.36 (95% confidence interval [CI]: 2.52 to 7.82); cardiovascular, 7.44 (95% CI: 3.06 to 18.1); cancer, 2.45 (95% CI: 0.85 to 7.04); and noncardiovascular 3.15 (95% CI: 1.53 to 6.49) mortalities. Relative risks all moderated to lower values when the comparisons were expanded to include the upper and lower thirds. Similar associations were still present when Cys-C was modeled on a continuous scale, suggesting a linear relationship between Cys-C and mortality outcomes.
Conclusions	Serum Cys-C is prognostic of long-term mortality in the subjects with relatively normal renal function, independent of MDRD eGFR and albuminuria. (J Am Coll Cardiol 2010;56:1930–6) © 2010 by the American College of Cardiology Foundation

Cystatin C (Cys-C) is a 13-kDa protein and member of the family of competitive lysosomal cysteine protease inhibitors. The continual synthesis of Cys-C in nucleated cells and its association with pre-clinical renal disease have prompted the use of Cys-C to estimate renal function. For instance, high serum Cys-C is reportedly predictive for all-cause mortality, cardiovascular disease, and the prognosis for congestive heart failure (1,2).

Cys-C is freely filtered by the glomerulus and has potential advantages over creatinine when estimating glomerular filtration rate (GFR) in that its production is not

See page 1937

dependent on muscle mass. In addition, it is unaffected by age, fever, or exogenous agents (3). In comparison, serum

From the *Cardiovascular Center and Health Management Center, National Taiwan University Hospital Yun-Lin Branch, Dou-Liou City, Taiwan; †Department of Medicine, College of Medicine, National Taiwan University, Taipei, Taiwan; ‡Department of Integrative Physiology and Cardiovascular Research Institute, University of North Texas Health Science Center, Fort Worth, Texas; §Department of Environmental and Occupational Health, University of North Texas Health Science Center Fort Worth, Texas; and the ||Office of Public Health Genomics, Centers for Disease Control and Prevention, Atlanta, Georgia. This work was supported in part by the G62024

Interdisciplinary Research Grant from the University of North Texas Health Science Center at Fort Worth and by grants from National Taiwan University Hospital Yun-Lin Branch (97.S005 and 98.X002). The findings and conclusions in this report are those of the author(s) and do not necessarily represent the views of the Centers for Disease Control and Prevention. The authors have reported that they have no relationships to disclose. The first two authors contributed equally to this paper.

Manuscript received December 10, 2009; revised manuscript received March 10, 2010, accepted April 6, 2010.

creatinine, the primary tool for evaluation of kidney function in clinical practice, can be affected by extrarenal factors including age, body weight, nutritional status, ethnicity, and gender (4). In addition, it is insensitive to modest decreases in kidney function (5). As a result, Cys-C has been suggested as a more accurate estimate of GFR than creatinine-based equations and may also predict declining kidney function even when the GFR is actually near the normal range (6).

Although renal dysfunction appears to be the most plausible link between increased Cys-C and impaired cardiovascular outcomes, the predictive value of Cys-C for long-term mortality outcomes among populations with normal kidney function has not been studied extensively (7). In addition, whereas most studies on Cys-C focus on cardiovascular outcomes, it remains unclear whether elevated Cys-C is also associated with increased mortality of cancer and other noncardiovascular disorders. In the current study, cohort follow-up data were analyzed to test the hypothesis that Cys-C is associated with long-term mortality for all-cause, cardiovascular disease, cancer, and noncardiovascular death among subjects with normal creatinine-based GFR. The data were obtained from a large national dataset.

Methods

Study design and population. NHANES III (Third National Health and Nutrition Examination Survey) was conducted by the National Center for Health Statistics (NCHS) from 1988 to 1994 using a stratified, multistage, and cluster sampling design to obtain a representative sample of the noninstitutionalized civilian U.S. population. The NHANES III was reviewed and approved by the NCHS Institutional Review Board. Data collection included in-depth, in-person interviews, physical and physiological examinations, and laboratory tests. This study was restricted to 2,990 non-Hispanic Whites, non-Hispanic Blacks, and Mexican-American participants aged 40 years or older with normal kidney function, defined as Modification of Diet in Renal Disease (MDRD) estimated glomerular filtration rate (eGFR) ≥ 60 ml/min/1.73 m², and valid measurements of serum Cys-C (8). Serum Cys-C was measured in NHANES III in all participants age 60 years and in a 25% random sample of participants age 12 to 59 years. A sample weight variable for Cys-C was thus used to adjust for the unequal probabilities of selection to represent the U.S. population (9).

Measurements of serum Cys-C. Serum from fasting blood samples was stored at -70°C until analyzed. Cys-C was then measured using a particle-enhanced immunonephelometric assay (N Latex Cys-C, Dade Behring, Deerfield, Illinois) at the Cleveland Clinical Research Laboratory. Cys-C samples with this assay had a range of 0.23 to 7.25 mg/l (17.2 to 543.0 nmol/l) and an intra-assay imprecision

of 2.0% to 3.0% coefficient of variation and an interassay imprecision of 3.2% to 4.4% coefficient of variation (10).

Baseline demographic and clinical factors. Data were collected by trained personnel according to standardized procedures (11). Information for age, sex, and smoking was self-reported. Current cigarette smoking was coded as a dichotomous “yes/no” variable. Hypertension was defined as an average systolic blood pressure ≥ 140 mm Hg, a diastolic blood pressure ≥ 90 mm Hg, a physician’s diagnosis or the use of antihypertensive medication (12). Diabetes mellitus was assessed by self-reported diagnosis of diabetes, the use of diabetic medications (insulin or oral agents), a nonfasting plasma glucose ≥ 11.1 mmol/l (200 mg/dl), or a fasting plasma glucose ≥ 7 mmol/l (126 mg/dl). Serum triglycerides, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, fibrinogen, uric acid, hemoglobin, blood urea nitrogen, C-reactive protein (CRP), urinary albumin, and creatinine were analyzed by laboratory methods reported elsewhere (13). Urinary albumin-to-creatinine ratios (UACR, in mg/mmol) were calculated accordingly (14,15). The GFR was calculated from MDRD (Modification of Diet in Renal Disease) study equation: $(\text{ml/min}/1.73 \text{ m}^2) = 175 \times (\text{Scr})^{-1.154} \times (\text{Age})^{-0.203} \times (0.742 \text{ if female}) \times (1.210 \text{ if African American})$ (16). Serum creatinine was measured by means of the modified kinetic Jaffé reaction, followed by calibration using an enzymatic creatinine assay traceable to a gold-standard reference method as described previously (17).

Mortality follow-up. The underlying cause of death was coded using the International Classification of Diseases-10th Revision (ICD-10) (18). The association of Cys-C with long-term mortality was evaluated for deaths from all-cause, cardiovascular disease (ICD codes: I00 to I78), malignant neoplasm (ICD-10 codes: C00 to C97), and noncardiovascular mortality. Noncardiovascular mortality was defined as mortality cases not belonging to the category of cardiovascular causes. Length of follow-up for each participant was calculated as the time from the NHANES III examination date to the end of follow-up (date of death or December 31, 2006). Thus, those found alive were right-censored on the last date known alive or at the end of the study, and likewise those who died from other causes were also right-censored at the time of death for cause-specific analysis. Age-adjusted all-cause and cause-specific mortality rates (per 1,000 person-years) were calculated using direct standardization according to the 2000 U.S. standard population (19).

Statistical analysis. NHANES used a complex sampling design and a complex survey design, including nonresponse

Abbreviations and Acronyms

AUC	= area under the curve
Cys-C	= cystatin C
eGFR	= estimated glomerular filtration rate
GFR	= glomerular filtration rate
ROC	= receiver-operating characteristic
TG	= triglyceride
UACR	= urinary albumin-to-creatinine ratio

and unequal sampling probabilities, and provided sampling weights for constructing national estimates. SUDAAN 9.03 (2004, Research Triangle Institute, Research Triangle Park, North Carolina) was used with the Taylor-series linearization to obtain unbiased standard errors for statistical estimates in all analyses. Demographic and clinical characteristics were compared across 3 categories of Cys-C level with analysis of variance (ANOVA) for continuous variables and the Cochran-Mantel-Haenszel chi-square test for categorical variables. Subjects were classified into the high (>90th percentile) (Cys-C >1.08 mg/l), medium (10th to 90th percentile) (Cys-C 0.7 to 1.08 mg/l), and low (<10th percentile) (Cys-C <0.7 mg/l) and also into tertiles (Cys-C <0.80, 0.80 to 0.93, and >0.93 mg/l). Time to event was first examined using the Kaplan-Meier product limit survival method, followed by univariate and multivariate Cox proportional hazard model with Efron method to calculate relative risk (RR), also known as the hazard ratios, to evaluate mortality risk associated with serum Cys-C for all-cause and cause-specific deaths. Potential confounding factors such as MDRD, UACR, and CRP were accordingly examined and adjusted in the models based on p values from univariate analyses (included when $p < 0.10$) and prior knowledge regarding the importance of risk factors. The level of significance was set at 0.05.

The receiver-operating characteristic (ROC) curves were used to assess the discriminative power of Cys-C as a continuous variable to differentiate the occurrence of all-cause mortality, cardiovascular mortality, cancer mortality, and noncardiovascular mortality. Area under curve (AUC)

was calculated. The optimal cutoff point, defined as the least $(1 - \text{sensitivity})^2 + (1 - \text{specificity})^2$ or the shortest distance from the left upper corner to the ROC curve, was reported.

Results

Baseline demographic and clinical variables are shown in Table 1. The sample of 2,990 participants was equivalent to an estimated U.S. population of 63,192,373 persons 40+ years of age after adjusting for sample weights. The overall mean (\pm SE) serum Cys-C of the study population was 0.89 ± 0.01 mg/l (5th to 95th percentile range 0.66 to 1.16 mg/l), with the average concentrations, respectively, for subjects in the high (highest decile, $n = 577$), medium (2nd to 9th decile, $n = 2,199$), and low (lowest decile, $n = 214$) groups were 1.19 ± 0.008 mg/l, 0.87 ± 0.004 mg/l, and 0.65 ± 0.005 mg/l (Fig. 1). Except fasting glucose, all demographic and clinical variables were statistically different across Cys-C categories ($p < 0.05$, ANOVA and chi-square tests for continuous and categorical outcomes, respectively). Overall, the participants in the high Cys-C group were more often older, male, and hypertensive. Likewise, CRP increased with Cys-C concentrations (Spearman's correlation: 0.092, $p < 0.001$). In addition, there was a significant but nonlinear relationship between Cys-C and UACR (Spearman's correlation: 0.164, $p < 0.001$, data not shown).

Within an average of 13.7 years of follow-up (range 0.5 to 18 years), 1,206, 487, 320, and 719 subjects died, respectively, from all-cause, cardiovascular, cancer, and noncar-

Table 1 Baseline Demographic and Clinical Characteristics of Participants* ($n = 2,990$)

Characteristics	Cystatin C		
	<10th Percentile (<0.70 mg/l) ($n = 214$)	10th–90th Percentile (0.70–1.08 mg/l) ($n = 2,199$)	>90th Percentile (>1.08 mg/l) ($n = 577$)
Males, %*	20.6 (5.02)	53.3 (1.96)	62.1 (3.82)
Microalbuminuria, %	12.2 (4.41)	10.6 (1.15)	22.3 (2.94)
Diabetes, %	14.7 (4.31)	6.77 (1.19)	11.3 (1.97)
Hypertension, %	25.0 (4.39)	39.9 (1.69)	51.9 (3.54)
Current smoking, %	20.4 (4.09)	24.8 (1.84)	35.5 (4.70)
Age, yrs	46.8 (0.73)	53.4 (0.56)	64.3 (1.17)
BMI, kg/m ²	25.0 (0.62)	27.4 (0.19)	28.0 (0.28)
Hemoglobin, g/dl	13.5 (0.19)	14.3 (0.07)	14.3 (0.07)
TG, mg/dl	117.7 (10.5)	170.0 (10.7)	184.8 (9.52)
HDL, mg/dl	58.5 (2.38)	50.0 (0.79)	44.9 (0.99)
Fibrinogen, mg/dl	310.1 (13.5)	295.5 (4.34)	322.8 (7.57)
C-reactive protein, mg/l	0.35 (0.03)	0.42 (0.03)	0.62 (0.06)
Uric acid, mg/dl	4.16 (0.13)	5.35 (0.06)	5.91 (0.09)
Fasting glucose, mg/dl	113.3 (8.09)	101.6 (1.90)	103.9 (2.83)
Blood urea nitrogen, mg/dl	12.3 (0.37)	14.2 (0.22)	16.0 (0.35)
Serum creatinine, mg/dl	0.92 (0.01)	1.03 (0.01)	1.08 (0.01)
eGFR, ml/min/1.73 m ²	79.4 (1.40)	73.5 (0.38)	68.6 (0.61)
UACR, mg/mmol	4.85 (2.59)	1.77 (0.20)	7.62 (3.29)

All demographic and clinical variables, except fasting glucose, are statistically different across cystatin C categories ($p < 0.05$). Data are expressed as either mean or percentage with standard error. *With adjustment for sample weight.

eGFR = glomerular filtration rate calculated from the Modification of Diet in Renal Disease (MDRD) study equation; HDL = high-density lipoprotein; LDL = low-density lipoprotein; TG = triglyceride; UACR = urinary albumin-to creatinine ratio.




Figure 1 Histogram of Serum Cys-C Concentrations

The histogram shows the distribution of serum cystatin C (Cys-C) concentrations. Kolmogorov-Smirnov Ztest indicates a deviation from normal distribution ($p < 0.001$). The overall mean (\pm SE) serum Cys-C of the study population was 0.89 ± 0.01 mg/l (5th to 95th percentile range: 0.66 to 1.16 mg/l). Solid curve = the normality plot.

di cardiovascular causes. This corresponds to age-adjusted mortality rates (year 2000 U.S. standard population) of 19.8, 7.7, 5.7, and 12.1 per 1,000 person-years in that order (data not shown). The results from Kaplan-Meier survival curves stratified by Cys-C categories for all-cause (Fig. 2A) and cause-specific mortality (Figs. 2B to 2D) showed that, when compared with persons with low Cys-C, those with medium or high levels of Cys-C concentrations were generally at higher risk for all-cause and cause-specific mortality. Similar results were obtained from univariate analyses (Table 2). For instance, the relative risks for all-cause mortality for high and medium Cys-C were 17.6 (95% CI: 10.1 to 30.9) and 5.64 (95% CI: 3.11 to 10.2), respectively, as compared with those with low levels of serum Cys-C. Despite the decrease in relative risks following adjustment for age, sex, diabetes mellitus, hypertension, smoking, CRP, UACR, and MDRD eGFR, the associations between Cys-C and mortality risk remained significant for all-cause, cardiovascular, and noncardiovascular disease but not for cancer (Table 2). Although an apparent increased risk of cancer mortality was still observed in individuals with high and medium Cys-C, the results did not attain statistical significance ($p = 0.18$). Similar trends were observed when the population was divided into tertiles based on serum Cys-C concentrations. The associations were statistically significant for all-cause and cause-specific mortalities in the univariate analyses but did not reach statistical significance after adjustment for covariates (Online Table 1).

Of the covariates examined in multivariate-adjusted analyses, age and current smoking were 2 leading risk factors for

mortality. Excluding cancer, MDRD eGFR and UACR were important predictors of all-cause and cause-specific deaths ($p < 0.05$). Hypertension and CRP are likewise related to increased mortality risk, but are only significant in all-cause mortality and cancer death (hypertension only). Sex and diabetes were also significantly associated with all-cause and cardiovascular mortality ($p < 0.05$) (Table 2, Online Table 1).

The multivariate-adjusted relative risks by modeling mortality using log-transformed serum Cys-C concentrations on a continuous scale were significant for all-cause, 3.96 (95% CI: 2.17 to 7.23, $p < 0.001$), cardiovascular disease, 6.51 (95% CI: 1.96 to 21.6, $p = 0.003$), noncardiovascular disease, 2.77 (95% CI: 1.14 to 6.71, $p = 0.03$) but not for cancer, 2.26 (95% CI: 0.57 to 9.04, $p = 0.24$) (data not shown). The comparable findings suggested a dose-responsive relationship between Cys-C and mortality outcomes. ROC analysis showed that the AUC for all-cause mortality, cardiovascular mortality, cancer mortality, and noncardiovascular mortality were 0.72, 0.74, 0.64, and 0.67, respectively. The estimated optimal cutoff values of Cys-C were 0.95 mg/l for all-cause and cardiovascular mortality and 0.97 mg/l for cancer and noncardiovascular mortality (Online Fig. 1).

Discussion

The present study demonstrates that an elevated serum Cys-C concentration is associated with all-cause, cardiovascular, and noncardiovascular mortality adjusted for other risk factors (e.g., MDRD eGFR and albuminuria) in a general population with creatinine-based eGFR ≥ 60 ml/min/1.73 m². In general, the results support previous findings suggesting that Cys-C is linked to long-term all-cause and cardiovascular mortality independent of kidney function (20).

Although the mechanisms through which elevated Cys-C relates to long-term mortality remain undetermined, it is generally assumed that serum Cys-C reflects the balance of its primary physiological determinants: cellular generation, renal filtration, and subsequent renal degradation (9). Also, the generation of Cys-C by nucleated cells is constant, and the concentration is not influenced by age, sex, height, and body composition (21). An increased Cys-C concentration identifies early deviations in GFR and becomes a sensitive indicator of “pre-clinical” renal disease, which thus may be associated with adverse cardiovascular outcomes (6). In addition, inflammatory status, thyroid disease, serum CRP, and current smoking were all associated with Cys-C concentrations (22). Inflammatory cytokines associated with atherosclerosis may alter the relationship between lysosomal cathepsins and their endogenous inhibitors like Cys-C in favor of remodeling and atherogenesis (23). Elastolytic cysteine proteases and their inhibitors, including Cys-C, are suggested to be involved in the pathogenesis of atherosclerosis (24). This may explain in part the




Figure 2 Survival Curves for All-Cause and Cause-Specific Mortality

Kaplan-Meier cumulative survival curves for risk for (A) all-cause, (B) cardiovascular disease, (C) cancer, and (D) noncardiovascular cause stratified by cystatin C (Cys-C) levels: high, >90th percentile; medium, 10th to 90th percentile; low, <10th percentile. **x-axis** indicates time (in years), **y-axis** indicates cumulative survival (%).

mechanism underlying the statistical link between Cys-C and cardiovascular outcomes in subjects with normal creatinine-based eGFR. Thus, increases in serum Cys-C may be sensitive harbingers of cardiovascular disease that are associated with inflammation (e.g., CRP) and atherosclerosis (25). Alternatively, the connection may be related to other pathophysiologic pathways that amplify the mortality risk independent of GFR (26).

Previous studies have demonstrated that Cys-C predicts all-cause mortality and cardiovascular events among elderly cohorts, subjects with chronic kidney disease, or patients with coronary heart disease (1,27,28). A recent study on 4,663 elderly persons (age >65 years) by Shlipak et al. (27) reported that higher Cys-C was associated with higher risk for cardiovascular events and mortality. The current study reinforces the prior findings that Cys-C is

Table 2 RR of All-Cause and Cause-Specific Mortality by 10th to 90th Percentiles of Cys-C Levels and Other Confounding Risk Factors

Predictor Variable	All-Cause		Cardiovascular		Cancer		Noncardiovascular	
	RR (95% CI)	p Value	RR (95% CI)	p Value	RR (95% CI)	p Value	RR (95% CI)	p Value
Univariate analysis								
Serum Cys-C level		<0.001		<0.001		<0.001		<0.001
>90th percentile	17.6 (10.1–30.9)		24.9 (11.2–55.3)		8.66 (3.94–19.1)		13.8 (7.10–26.7)	
10th–90th percentile	5.64 (3.11–10.2)		5.98 (2.68–13.4)		4.01 (1.70–9.44)		5.45 (2.76–10.7)	
<10th percentile	1.00 (reference)		1.00 (reference)		1.00 (reference)		1.00 (reference)	
Multivariate analysis								
Serum Cys-C level		<0.001		<0.001		0.18		0.006
>90th percentile	4.36 (2.52–7.82)		7.44 (3.06–18.1)		2.45 (0.85–7.04)		3.15 (1.53–6.49)	
10th–90th percentile	3.45 (2.04–5.82)		4.59 (1.99–10.6)		2.36 (0.95–5.88)		2.99 (1.57–5.72)	
<10th percentile	1.00 (reference)		1.00 (reference)		1.00 (reference)		1.00 (reference)	
Males	1.41 (1.06–1.89)	0.02	1.67 (1.11–2.56)	0.02	1.25 (0.72–2.13)	0.42	1.27 (0.93–1.72)	0.14
Diabetes, yes	1.35 (1.03–1.76)	0.03	1.72 (1.23–2.42)	0.002	0.95 (0.58–1.55)	0.82	1.12 (0.77–1.63)	0.56
Hypertension, yes	1.36 (1.04–1.78)	0.03	1.38 (0.96–1.98)	0.08	1.75 (1.15–2.65)	0.01	1.35 (0.99–1.83)	0.06
Current smoking, yes	2.00 (1.54–2.58)	<0.001	1.79 (1.19–2.71)	0.01	2.32 (1.34–4.02)	0.004	2.12 (1.47–3.06)	<0.001
CRP, mg/l*	1.13 (1.00–1.28)	0.04	1.12 (0.90–1.39)	0.30	1.09 (0.86–1.38)	0.45	1.14 (0.97–1.33)	0.10
UACR ratio, mg/mmol*	1.17 (1.10–1.25)	<0.001	1.21 (1.08–1.34)	0.001	1.04 (0.90–1.21)	0.59	1.14 (1.05–1.23)	0.002
Age, per yr	1.10 (1.09–1.11)	<0.001	1.11 (1.09–1.13)	<0.001	1.07 (1.04–1.09)	<0.001	1.09 (1.08–1.11)	<0.001
eGFR, ml/min/1.73 m ² *	4.25 (1.76–10.3)	0.002	9.51 (2.26–40.1)	0.003	1.04 (0.25–4.36)	0.96	2.50 (1.00–6.25)	0.05

*Log-transformed.

CI = confidence interval; CRP = C-reactive protein; Cys-C = cystatin C; RR = relative risk; other abbreviations as in Table 1.

associated with increased risk of all-cause and cardiovascular mortality among the elderly, and expands these findings to include noncardiovascular disorders in a younger (age >40 years) population with average age of 53 years and “clinically normal” kidney function. The ROC analysis by Astor et al. showed that Cys-C–based eGFR was predictive of all-cause and cardiovascular mortality, with an AUC of 0.80, in the general U.S. population with creatinine-based eGFR >15 ml/min/1.73 m² (29). In the current study, we used a simple dichotomized approach to assess the mortality risk associated with serum Cys-C concentrations and found that Cys-C could still predict increased mortality in the subjects with normal creatinine-based eGFR who were previously thought to belong to a low-risk group.

Increased risk of noncardiovascular illness may reflect associations between Cys-C and declining physical function mediated by inflammation (30). Such subjects with declining health might reasonably have additional comorbidities, such as aspiration pneumonia and bedsores, and therefore are prone to noncardiovascular mortality. Alternatively, high Cys-C may contribute to noncardiovascular mortality risks indirectly through cognitive impairments mediated by Cys-C–associated vascular disease (31).

Study limitations. There are some limitations in the current study. First, the selection of subjects used eGFRs based on single creatinine determinations. This potential concern is mitigated in part by the observation that repeated creatinine measurements in a subpopulation of 1,919 NHANES III subjects showed a reasonable agreement between the 2 measurements (32). Thus, errors in selection of subjects as “normal” seem unlikely to account

for the better prognostic value of Cys-C. On the other hand, it is not unlikely that the association of interest could be confounded by other risk factors, such as thyroid disease, which may affect both Cys-C concentrations and mortality outcomes (33). Also, several single nucleotide polymorphisms have been associated with altered Cys-C concentrations (34). The current study, however, does support Cys-C concentrations as an indicator of increased risk of long-term mortality independent of residual renal function. Furthermore, a recent study showed that Cys-C was associated with an increased risk in the composite end point of death or rehospitalization in patients hospitalized for heart failure (35). However, the present analysis dealt with long-term survival. Thus, the association between Cys-C and the incidence of nonfatal cardiovascular events may have been missed.

Conclusions

This study has demonstrated that Cys-C, an alternative predictor of renal function, is significantly associated with all-cause, cardiovascular, and even noncardiovascular mortality in the general population with normal kidney function. Thus, Cys-C has a unique prognostic potential regarding long-term mortality. It remains to be elucidated as to whether some novel combination of measurements will provide even greater sensitivity and predictive power including perhaps for cancer.

Acknowledgments

The authors are grateful to Mr. Peter Meyer and Ms. Karen E. Davis from the National Center for Health Statistics,

Centers for Disease Control and Prevention, and Dr. Gordon Brown from RTI International for their support and advice.

Reprint requests and correspondence: Dr. Yu-Sheng Lin, Department of Environmental and Occupational Health, University of North Texas Health Science Center, Fort Worth, Texas 76107. E-mail: Yu-Sheng.Lin@unthsc.edu; or Dr. Jou-Wei Lin, Cardiovascular Center, National Taiwan University Hospital Yun-Lin Branch, 579 Yun-Lin Road, Section 2, Dou-Liou City, Yun-Lin County 640, Taiwan. E-mail: jouweilin@yahoo.com.

REFERENCES

- Fried LF, Katz R, Sarnak MJ, et al. Kidney function as a predictor of noncardiovascular mortality. *J Am Soc Nephrol* 2005;16:3728–35.
- Shlipak MG, Wassel Fy CL, Chertow GM, et al. Cystatin C and mortality risk in the elderly: the health, aging, and body composition study. *J Am Soc Nephrol* 2006;17:254–61.
- Newman DJ, Thakkar H, Edwards RG, et al. Serum cystatin C measured by automated immunoassay: a more sensitive marker of changes in GFR than serum creatinine. *Kidney Int* 1995;47:312–8.
- Ishida K, Ishida H, Narita M, et al. Factors affecting renal function in 119 985 adults over three years. *QJM* 2001;94:541–50.
- Kassirer JP. Clinical evaluation of kidney function—glomerular function. *N Engl J Med* 1971;285:385–9.
- Shlipak MG, Katz R, Sarnak MJ, et al. Cystatin C and prognosis for cardiovascular and kidney outcomes in elderly persons without chronic kidney disease. *Ann Intern Med* 2006;145:237–46.
- Taglieri N, Fernandez-Berges DJ, Koenig W, et al. Plasma cystatin C for prediction of 1-year cardiac events in Mediterranean patients with non-ST elevation acute coronary syndrome. *Atherosclerosis* 2010;209:300–5.
- Poggio ED, Nef PC, Wang X, et al. Performance of the Cockcroft-Gault and modification of diet in renal disease equations in estimating GFR in ill hospitalized patients. *Am J Kidney Dis* 2005;46:242–52.
- Kottgen A, Selvin E, Stevens LA, et al. Serum cystatin C in the United States: the Third National Health and Nutrition Examination Survey (NHANES III). *Am J Kidney Dis* 2008;51:385–94.
- Newman DJ. Cystatin C. *Ann Clin Biochem* 2002;39:89–104.
- Centers for Disease Control and Prevention. Plan and operation of the Third National Health and Nutrition Examination Survey, 1988–94. Series 1: programs and collection procedures. *Vital Health Stat* 1994;1–407.
- Mancia G, Grassi G. Joint National Committee VII and European Society of Hypertension/European Society of Cardiology guidelines for evaluating and treating hypertension: a two-way road? *J Am Soc Nephrol* 2005;16 Suppl 1:S74–7.
- Gunter EW, Lewis BL, Koncikowski SM. Laboratory Methods used for the Third National Health and Nutrition Examination Survey (NHANES III), 1988–1994. In: CD-ROM 6-0178, NHANES III Reference Manuals and Reports. Hyattsville, MD: Centers for Disease Control and Prevention, 1996.
- Coresh J, Astor BC, Greene T, et al. Prevalence of chronic kidney disease and decreased kidney function in the adult US population: Third National Health and Nutrition Examination Survey. *Am J Kidney Dis* 2003;41:1–12.
- Coresh J, Selvin E, Stevens LA, et al. Prevalence of chronic kidney disease in the United States. *JAMA* 2007;298:2038–2047.
- Levey AS, Coresh J, Greene T, et al. Using standardized serum creatinine values in the modification of diet in renal disease study equation for estimating glomerular filtration rate. *Ann Intern Med* 2006;145:247–54.
- Selvin E, Manzi J, Stevens LA, et al. Calibration of serum creatinine in the National Health and Nutrition Examination Surveys (NHANES) 1988–1994, 1999–2004. *Am J Kidney Dis* 2007;50:918–26.
- Centers for Disease Control and Prevention (CDC). Third National Health and Nutrition Examination Survey (NHANES III) linked mortality file restricted-use file layout. Hyattsville, MD: US Department of Health and Human Services, Centers for Disease Control and Prevention, 2009. Available at: http://www.cdc.gov/nchs/data/datalinkage/file_layout_nh3.pdf. (Accessed October 7, 2009).
- Klein RJ, Schoenborn CA. Age Adjustment Using the 2000 Projected US Population: Healthy People Statistical Notes No. 20. Hyattsville, MD: National Center for Health Statistics, 2001.
- Stevens LA, Levey AS. Chronic kidney disease in the elderly—how to assess risk. *N Engl J Med* 2005;352:2122–4.
- Finney H, Newman DJ, Price CP. Adult reference ranges for serum cystatin C, creatinine and predicted creatinine clearance. *Ann Clin Biochem* 2000;37:49–59.
- Fricker M, Wiesli P, Brandle M, et al. Impact of thyroid dysfunction on serum cystatin C. *Kidney Int* 2003;63:1944–7.
- Taglieri N, Koenig W, Kaski JC. Cystatin C and cardiovascular risk. *Clin Chem* 2009;55:1932–43.
- Sukhova GK, Shi GP, Simon DI, et al. Expression of the elastolytic cathepsins S and K in human atheroma and regulation of their production in smooth muscle cells. *J Clin Invest* 1998;102:576–83.
- Leung-Tack J, Tavera C, Gensac MC, et al. Modulation of phagocytosis-associated respiratory burst by human cystatin C: role of the N-terminal tetrapeptide Lys-Pro-Pro-Arg. *Exp Cell Res* 1990;188:16–22.
- Knight EL, Verhave JC, Spiegelman D, et al. Factors influencing serum cystatin C levels other than renal function and the impact on renal function measurement. *Kidney Int* 2004;65:1416–21.
- Shlipak MG, Sarnak MJ, Katz R, et al. Cystatin C and the risk of death and cardiovascular events among elderly persons. *N Engl J Med* 2005;352:2049–60.
- Ix JH, Shlipak MG, Chertow GM, et al. Association of cystatin C with mortality, cardiovascular events, and incident heart failure among persons with coronary heart disease: data from the Heart and Soul Study. *Circulation* 2007;115:173–9.
- Astor BC, Levey AS, Stevens LA, et al. Method of glomerular filtration rate estimation affects prediction of mortality risk. *J Am Soc Nephrol* 2009;20:2214–22.
- Odden MC, Chertow GM, Fried LF, et al. Cystatin C and measures of physical function in elderly adults: the Health, Aging, and Body Composition (HABC) Study. *Am J Epidemiol* 2006;164:1180–9.
- Yaffe K, Lindquist K, Shlipak MG, et al. Cystatin C as a marker of cognitive function in elders: findings from the health ABC study. *Ann Neurol* 2008;63:798–802.
- Coresh J, Astor BC, McQuillan G, et al. Calibration and random variation of the serum creatinine assay as critical elements of using equations to estimate glomerular filtration rate. *Am J Kidney Dis* 2002;39:920–9.
- Manetti L, Pardini E, Genovesi M, et al. Thyroid function differently affects serum cystatin C and creatinine concentrations. *J Endocrinol Invest* 2005;28:346–9.
- Hwang SJ, Yang Q, Meigs JB, et al. A genome-wide association for kidney function and endocrine-related traits in the NHLBI's Framingham Heart Study. *BMC Med Genet* 2007;8 Suppl 1:S10.
- Campbell CY, Clarke W, Park H, Haq N, Barone BB, Brotman DJ. Usefulness of cystatin C and prognosis following admission for acute heart failure. *Am J Cardiol* 2009;104:389–92.

Key Words: cystatin C ■ glomerular filtration rate ■ nutrition survey ■ survival analysis.

APPENDIX

For a supplementary table and figure, please see the online version of this article.