

The Prognostic Significance of Leukocyte Count on All-Cause and Cardiovascular Disease Mortality: A Systematic Review and Meta-Analysis



Cheri Park^a, Kanghee Yoo^a, Soohyun Lee^a, Hyerin Kim, MD^b, Eunjeong Son, MD, PhD^c, Dongjun Lee, PhD^d, Dai Sik Ko, MD, PhD^e, Kihun Kim, MD^{f,g,**}, and Yun Hak Kim, MD, PhD^{f,g,*}

White blood cells (WBCs) act as mediators of inflammatory responses and are commonly measured in hospitals. Although several studies have reported a relation between WBC count and mortality, no systematic review or meta-analysis has been conducted. This study aimed to identify an association between WBC count and mortality. We conducted a systematic search on Embase using keywords such as “white blood cell” and “mortality.” We analyzed the hazard ratios (HRs) for WBC count of 1.0×10^9 cells/L regarding 2 criteria: the cause of mortality and the follow-up period. A total of 13 of 222 articles comprising a total of 62,904 participants were included in this study, meeting the criteria set. A positive association was observed between WBC count and mortality, as indicated by an HR of 1.10 (95% confidence interval [CI] 1.08 to 1.13). In additionally, WBC count emerged as a significant predictor of mortality in both groups, with an HR of 1.10 (95% CI 1.07 to 1.12) for patients with cardiovascular disease and an HR of 1.12 (95% CI 1.07 to 1.17) for the general population or patients with COVID-19. Furthermore, a higher WBC count demonstrated a significant association with long-term all-cause mortality (HR 1.09, 95% CI 1.07 to 1.12) and long-term cardiovascular mortality (HR 1.05, 95% CI 1.02 to 1.07). Similarly, a significant association was found between higher WBC count and short-term all-cause mortality (HR 1.12, 95% CI 1.09 to 1.16) and cardiovascular mortality (HR 1.12, 95% CI 1.07 to 1.17). Further research is necessary to explore the relation between WBC count and disease progression or death and to establish causality between elevated WBC count and disease progression. © 2023 Elsevier Inc. All rights reserved. (Am J Cardiol 2023;203:226–233)

The leukocyte count serves as mediators of inflammatory responses and play a crucial role in the host's defense. Therefore, they are routinely used as a nonspecific biomarker for acute inflammatory responses. Furthermore, several studies suggest that white blood cell (WBC) count could serve as a

predictive marker for certain diseases, such as coronary heart disease^{1,2} and a marker for disease severity in conditions such as chronic obstructive pulmonary disease.^{3,4} The WBC count is a common component of routine blood tests in most hospitals. It is an easily accessible test because of its low cost and straightforward procedure. As such, changes in WBC count, if clinically significant, could be valuable in diagnosing and monitoring a patient's condition. An increase in WBC count is generally indicative of active inflammation, whereas leukopenia may be an epiphenomenon or may indicate an underlying systemic disease.^{5,6} Moreover, WBC count has been shown to be associated with the prognosis of patients with certain diseases and the general population. It was reported that higher total WBC counts are an independent predictor of all-cause mortality.⁷ In addition, an increase in total WBC count has been linked to poor outcomes in patients with stable coronary disease, acute coronary syndromes, and acute myocardial infarction.^{8–10} An acute increase in the WBC count typically reflects systemic inflammation and is associated with increased mortality. Furthermore, a chronic elevation in the WBC count indicates danger. Prolonged elevation of WBCs can indicate the sustained activation of their cytotoxic mechanism, leading to tissue damage.¹¹ WBC count is an essential indicator because of its accessibility. However, no systematic review or meta-analysis has established the relation between the WBC count and short-term or long-term mortality. Hence, the objective of this study was to investigate all-cause and cardiovascular-specific mortality in relation to the differences in the WBC count.

^aSchool of Medicine, Pusan National University, Yangsan, Republic of Korea; ^bDepartment of Laboratory Medicine, Pusan National University Hospital, Busan, Republic of Korea; ^cDivision of Respiratory and Allergy, Department of Internal Medicine, Pusan National University Yangsan Hospital, Yangsan, Republic of Korea; ^dDepartment of Convergence Medicine, School of Medicine, Pusan National University, Yangsan, Republic of Korea; ^eDivision of Vascular Surgery, Department of Surgery, Gachon University Gil Medical Center, Incheon, Republic of Korea; ^fDepartment of Biomedical Informatics, School of Medicine, Pusan National University, Yangsan, Republic of Korea; and ^gDepartment of Anatomy, School of Medicine, Pusan National University, Yangsan, Republic of Korea. Manuscript received March 21, 2023; revised manuscript received and accepted June 29, 2023.

Drs. Cheri Park, Kanghee Yoo, and Soohyun Lee contributed equally as first authors.

This work was supported by the Medical Research Center program [grant number NRF-2018R1A5A2023879], Basic Science Research Program [grant number RS-2023-00207946], the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI) (HI22C1377). This work was supported by KREONET (Daejeon, Republic of Korea).

See page 231 for Declaration of Competing Interest.

*Corresponding author: Tel: +82-51-510-8091; fax: +82-51-510-8049.

**Corresponding author: Tel: +82-51-510-8091; fax: +82-51-510-8049.

E-mail address: yunhak10510@pusan.ac.kr (Y.H. Kim).

Methods

This study was conducted according to the PRISMA (Preferred Reporting Items for Systematic reviews and Meta-Analyses) guideline.¹² The protocol of the article is registered with international prospective register of systematic reviews (registration number: CRD42022329742).

We conducted a search of the Embase and Medline databases on January 31, 2022, to identify studies examining the relation between WBC count and mortality. To establish our search strategy, we used medical subject headings terms and entry terms related to WBC and mortality in PubMed. We limited our search to titles and abstracts for a more precise retrieval. To ensure relevance to our study, we excluded nonarticle types and nonhuman subjects using the appropriate search options. The search strategy used for Embase and Medline is as follows: ('white blood cell*:ti OR 'white cell*:ti OR 'blood corpuscle*:ti OR wbc:ti OR leukocyte*:ti) AND (mortalit*:ti OR 'case fatality rate*:ti OR 'death rate*:ti) AND ([article]/lim OR [article in press]/lim) AND [humans]/lim.

The following criteria were used to exclude studies: non-articles, articles in press, nonhuman subjects, systematic reviews or meta-analyses, irrelevant studies, studies that were not randomized controlled trials or cohort studies, and studies that did not include hazard ratios (HRs) for WBC count as a continuous variable. Cohort studies were chosen owing to the lack of relevant randomized controlled trials. Cross-sectional studies, case-control studies, case series, case reports, and review articles were excluded. To reduce heterogeneity between studies, only those that treated WBC count as a continuous variable were included. In addition, HR was selected as the effect measure, and only studies that adjusted for at least age and gender were included. Therefore, any studies that did not include the adjusted HR for the WBC count as a continuous variable were excluded.

A total of 3 authors (Park, Yoo, and Lee) conducted an independent review of each article. The articles underwent a 3-stage review process, which included title, abstract, and full-text review. After the initial review, the articles were evaluated as a group through discussion. Disagreements were resolved through discussion, and decisions were made unanimously.

A total of 3 authors (Park, Yoo, and Lee) conducted a full review of the included studies and extracted basic information such as author, year of publication, research design, country, number of subjects, target population, research period, and HRs.

The risk of bias in the cohort studies was assessed using the Newcastle–Ottawa scale. A total of 3 authors (Park, Yoo, and Lee) evaluated all articles, and any disagreements were resolved through discussion, with unanimous decisions made.

To conduct a quantitative analysis, it is necessary to standardize the measurement unit of the WBC count to ensure that the HR is comparable across studies. By unifying the measurement unit to 1.0×10^9 cells/L, it allows the extraction of the adjusted HR of mortality based on a consistent change in leukocyte count of 1.0×10^9 cells/L.

The statistical analysis was performed using RevMan version 5.4.1, the Cochrane Collaboration's review

manager (Cochrane Training, United Kingdom). A total of 3 authors (Park, Yoo, and Lee) independently reviewed all procedures using the same tool and resolved discrepancies through discussion. Statistical heterogeneity was assessed using the I^2 statistic, and a random-effects model was used when the I^2 value was $>50\%$, whereas a fixed-effects model was used when the I^2 value was $<50\%$.¹³ A meta-analysis was conducted using the inverse variance method for weighting. The results were presented using a forest plot to visualize the integrated HR estimates.

A subgroup analysis was conducted based on 2 criteria: period and cause of mortality. A total of 4 subgroups were created based on whether the period was short term or long term (short term defined as <1 year) and whether the cause of mortality was all-cause or cardiovascular disease (CVD)—specific. The 4 subgroups were analyzed separately.

Publication bias was evaluated by visually inspecting the funnel plots. In addition, the Egger regression test was conducted to statistically assess publication bias.¹⁴

The grading of recommendations assessment, development, and evaluation approach was used to assess the certainty of the analysis.¹⁵ A total of 8 factors were thoroughly considered, including the risk of bias, inconsistency, indirectness, imprecision, and publication bias as factors reducing the quality of evidence and large magnitude of an effect, plausible residual confounding, and dose-response gradient as factors improving the quality of evidence. The 3 authors (Park, Yoo, and Lee) evaluated all factors, and decisions were made unanimously.

Results

A Medline and Embase search yielded 222 articles based on their titles and abstracts. During the screening process, 143 records were identified, excluding duplicates, nonarticle studies, and nonhuman studies. Of the remaining 121 articles, those that were irrelevant, had inadequate data, or used inappropriate study designs were excluded. Finally, 13 studies were included based on the selection criteria outlined in [Figure 1](#). These studies were conducted between 1992 and 2020, with 8 targeting patients with CVD and 5 targeting either the general population or patients with COVID-19. A study each on disabled older patients and older population were categorized under “general population” for analysis. Except for 2 studies conducted worldwide across 20 different countries, the remaining 5 were conducted in the United States; 2 in China; and 1 each in Taiwan, Korea, the United Kingdom, and Italy. All included studies measured WBC count as a continuous variable. The characteristics of the included studies are listed in [Table 1](#).

[Figure 2](#) presents the results of the overall analysis on the correlation between WBC count and mortality. WBC count was found to be a significant predictor of mortality rate (HR 1.10, 95% confidence interval [CI] 1.08 to 1.12, heterogeneity $I^2 = 72\%$, $p < 0.001$). In addition, WBC count was also found to be a significant predictor of mortality in both groups, with an HR of 1.10 (1.07 to 1.12, $I^2 = 80\%$) for patients with CVD and an HR of 1.12 (1.07 to 1.17, $I^2 = 0\%$) for the general population or patients with COVID-19.





Figure 1. Study selection flow chart of included studies.

Table 1

Characteristics of the studies included in the meta-analysis

Author, year	Study design	Country	Sample size	Target population	Study duration (mean or median)
Bhatt D.L., 2003 ⁴¹	Cohort	U.S.A.	10,480	CVD	6 months
Fan X., 2015 ²⁶	Cohort	China	570	CVD	30 days, 1.89 years
Gurm H.S., 2003 ⁴²	Cohort	20 different countries	2,279	CVD	3 years
Huang, Z.-S., 2003 ²⁹	Cohort	Taiwan	8,447	General population	65.5 months
Kim K-I, 2013 ²⁸	Cohort	Korea	9,996	Older population	44.9 months
Leng S.X., 2005 ¹⁶	Cohort	U.S.A.	624	Disabled older population	5 years
Newall N., 2007 ⁴³	Cohort	U.K.	3,024	CVD	1 year
Palmerini T., 2011 ⁴⁴	Cohort	U.S.A.	3,345	CVD	30 days, 1 year
Palmerini T., 2013-2 ⁴⁵	Cohort	U.S.A.	13,678	CVD	30 days, 1 year
Taglieri N., 2013 ⁴⁶	Cohort	Italy	1,315	CVD	3 years
Weiss S.T., 1995 ³⁰	Cohort	USA	1,956	General population	22.5 years
Yen M.H., 2003 ⁴⁷	Cohort	20 different countries	2,282	CVD	12 months
Zhu B., 2021 ²⁷	Cohort	China	163	COVID-19	NS

CVD = cardiovascular disease; NS = not specified.



Footnotes

(1) Fan X., 2015 (1) shows the data of long-term overall mortality.

(2) Palmerini T., 2011 (2) shows the data of short-term cardiovascular disease mortality.

(3) Palmerini T., 2013-2 (3) shows the data of short-term cardiovascular disease mortality.

Figure 2. Forest plot for an overall meta-analysis on the effect of WBC count on mortality.

A subgroup analysis was performed based on the overall mortality, cardiovascular mortality, and short-term and long-term mortality. Studies with mortality follow-up of more than 1 year were categorized as long term, whereas studies with mortality follow-up of less than 1 year were categorized as short term. As listed in Table 2, the results indicate that WBC count is a significant predictor of mortality in all subgroup analyses. The HR was 1.09 (95% CI 1.07 to 1.12, $I^2 = 39\%$) for the long-term overall mortality, and the HR was 1.05 (95% CI 1.02 to 1.07, $I^2 = 16\%$) for long-term cardiovascular mortality. The HR was 1.12 (95% CI 1.09 to 1.16, $I^2 = 78\%$) for the short-term overall mortality and the HR was 1.12 (95% CI 1.07 to 1.17, $I^2 = 52\%$) for short-term cardiovascular mortality.

Based on the Newcastle–Ottawa scale, the 13 included studies were categorized as being of good, fair, or poor quality. A total of 7 studies were evaluated as fair quality, 4 as good quality, and 2 as poor quality. Detailed quality assessments are listed in Table 3.

A funnel plot for the overall analysis between WBC count and mortality is presented in Figure 3. The plot shows that there is no obvious asymmetry, suggesting the absence of significant publication bias. This result is supported by the Egger regression test, which also indicates no significant publication bias ($p = 0.124$).

According to our assessment using the grading of recommendations assessment, development, and evaluation approach, the certainty of the outcome of interest, mortality, is very low (Table 4). This is because of the presence of several factors that reduce the quality of evidence, including risk of bias and inconsistency. Specifically, most of the included studies had a fair quality, and there was considerable heterogeneity across the studies.

Discussion

This study aimed to investigate the relation between WBC count and mortality. As demonstrated in Figure 2, it suggests that mortality increases proportionally with the number of WBCs. The subgroup analysis showed that the WBC count was a significant predictor of both short-term and long-term mortality, and the risk was particularly prominent in the short term.

An elevated WBC count has been identified as a prognostic factor that affects mortality, alongside risk factors such as tobacco use, total cholesterol levels, and high-density lipoprotein cholesterol levels.¹⁶ The mechanisms underlying the association between elevated WBC counts and increased mortality are not fully understood, but several hypotheses have been proposed. A hypothesis suggests that

Table 2
Subgroup analysis of mortality according to cause and period

	Long-term all-cause mortality	Short-term all-cause mortality	Long-term CVD-cause mortality	Short-term CVD-cause mortality
Number of studies	5	7	2	2
Hazard Ratio (HR)	1.09	1.12	1.05	1.12
95% CI	1.07 – 1.12	1.09 – 1.16	1.02 – 1.07	1.07 – 1.17
heterogeneity (I^2)	39%	78%	16%	52%

Table 3
Newcastle-Ottawa quality assessment of included in the meta-analysis

Author, year	Selection				Comparability		Outcomes			Quality*
	Representativeness of exposed cohort	Selection of non-exposed cohort	Ascertainment of exposure	Outcome not present at the start	Main factor (age)	Additional factor	Assessment of outcomes	Length of follow up	Adequacy of follow up	
Bhatt D.L., 2003 ⁴¹	0	1	1	0	1	1	-	1	1	Fair
Fan X., 2015 ²⁶	0	1	1	0	1	1	1	1	1	Fair
Gurm H.S., 2003 ⁴²	0	1	1	0	1	1	1	1	1	Fair
Huang, Z-S., 2003 ²⁹	1	1	1	1	1	1	1	1	-	Good
Kim K-I, 2013 ²⁸	1	1	1	1	1	1	1	1	1	Good
Leng S.X., 2005 ¹⁶	0	1	1	0	1	1	1	1	1	Fair
Newall N., 2007 ⁴³	0	1	1	0	1	1	1	1	1	Fair
Palmerini T., 2011 ⁴⁴	0	1	1	0	1	1	1	1	1	Fair
Palmerini T., 2013 ⁴⁵	0	1	1	0	1	1	1	1	1	Fair
Taglieri N., 2013 ⁴⁶	0	1	1	1	1	1	1	1	1	Good
Weiss S.T., 1995 ³⁰	0	1	1	1	-	-	1	1	1	Poor
Yen M.H., 2003 ⁴⁷	0	1	1	1	1	1	1	1	1	Good
Bhatt D.L., 2003 ⁴¹	0	1	-	0	1	1	1	-	1	Poor

* Good quality:3 or 4 stars in the selection domain AND 1 or 2 stars in the compatibility domain AND 2 or 3 stars in the outcome/exposure domain; fair quality:2 stars in the selection domain AND 1 or 2 stars in the comparability domain AND 2 or 3 stars in the outcome/exposure domain; poor quality:0 or 1 star in the selection domain OR 0 stars in the comparability domain OR 0 or 1 stars in the outcome/exposure domain

chronic systemic inflammation resulting from increased activity of pro-inflammatory factors, particularly, interleukin-6, activates the immune system and contributes to the increased mortality.¹⁷ Another potential mechanism is the increased mortality associated with clinical or subclinical diseases that could alter WBC levels.¹⁸

A previous studies have established a positive and linear association between elevated WBC count and all-cause mortality.¹⁹ Furthermore, several studies have shown that when WBC count was treated as a categorical variable, the group with the highest WBC count had a higher risk of all-cause mortality than other groups.^{20–22} Similarly, other studies have reported a positive association between WBC count and CVD-related mortality, with WBC count serving as an independent predictor.^{20,23–25}

A total of 7 of the studies included in our analysis were conducted on patients with ischemic heart disease. Fan et al²⁶ (2015) was conducted on patients with acute aortic dissection (AAD), Zhu et al²⁷ (2021) on patients with COVID-19, Leng et al¹⁶ (2005) on disabled older patients, Kim et al²⁸ (2013) on older patients, and Huang et al²⁹ (2003) and Weiss et al³⁰ (1995) on the general population. Each of these diseases was associated with an increased number of WBC after an inflammatory response. These findings support the significance of our results.

Ischemic heart disease is a condition in which blood supply to the myocardium is insufficient. It is most caused by an atherosclerotic plaque in the coronary artery. The pathogenesis of atherosclerosis is closely related to an inflammatory response.³¹ Hypercholesterolemia, a risk factor for coronary




Figure 3. Funnel plot for an overall meta-analysis on the effect of WBC count on mortality.

Table 4

GRADE approach for certainty assessment of overall analysis between WBC count and mortality

Outcomes	Certainty assessment							No. of participants	Effect Relative (95% CI)	Certainty
	No. of studies	Study design	Risk of bias	Inconsistency	Indirections	Imprecision	Other considerations			
WBC-mortality	13	Observational	Serious*	Serious†	Not serious	Not serious	Dose-response gradient	62,904‡	HR 1.10 (1.08-1.12)	Low

CI = confidence interval; HR = hazard ratio.

* Not all prognostic factors were controlled and the adjusted factors across studies does not match.

† Heterogeneity 72%.

‡ WBC count was treated as continuous variable.

artery disease, promotes low-density lipoprotein accumulation in the intima, which initiates an inflammatory response in the arterial wall.³² Endothelial cells activated by inflammation express leukocyte adhesion molecules, which attract leukocytes.³³ Recruited monocytes mature into macrophages, which then turn into foam cells and produce inflammatory cytokines.³⁴ Infarction of the coronary artery induces systemic inflammatory response, and the myocardium is damaged. This condition is called acute myocardial infarction (AMI). After AMI occurs, the number of peripheral blood mononuclear cells increases, and these cells infiltrate into the damaged region. Peripheral blood mononuclear cells are known to be associated with left ventricular remodeling after AMI.³⁵ Percutaneous coronary intervention, a mechanical way of reperfusion in patients with AMI, is also related to the provocation of inflammatory response. Coronary angioplasty elevates neutrophil, monocyte, and platelet adhesion molecule expression.³⁶

Although acute aortic syndromes are predisposed by many different risk factors, atherosclerosis is mainly related to AAD in the older population. The weakening of the intima of the aorta because of predisposing risk factors leads to a tear. Blood moves into the middle tunica through the tear, which results in the dissection or rupture of the aorta.³⁷ In an induced aortic dissection model, Kurihara et al³⁸ (2012) showed that infiltration of neutrophils and their secretion of matrix metalloproteinase-9 may play a role in the initiation of AAD.

COVID-19 is a relatively recent issue, and further studies are needed to determine the underlying pathophysiology. Current studies suggest that a higher WBC count is associated with mortality in patients with COVID-19,³⁹ but there is also a hypothesis that survival depends on lymphocyte restoration.⁴⁰

There are several limitations in our study. First, although more than 13 studies were conducted on WBC count and mortality, studies in which leukocyte counts were divided by categorical variables could not be combined and were excluded in the selection process. WBC is a highly nonspecific biomarker because elevated levels would simply indicate the presence of infection, inflammation, or a sick patient. Moreover, among the selected studies, 8 focused on patients with CVD and other 5 studies included general population or patients with COVID-19. Because of the heterogeneity of the chosen subjects across these studies, there may be limitations in generalizing the findings of this study.

Furthermore, given that the certainty of outcome was assessed as low, it is important to exercise caution when

interpreting the results of the study. Further research may be necessary to increase the certainty of the findings and to gain a better understanding of the relation between the variables in question. Despite the limitations, our research process yielded a significant correlation.

The objective of this meta-analysis was to determine whether a clinically significant relation exists between WBC count and mortality. Although the study is limited by the fact that only age and gender were the commonly adjusted factors, the results indicate that a higher WBC count is associated with increased mortality across various populations, regardless of time. This positive correlation is more pronounced in short-term mortality than in long-term mortality. Consequently, WBC count should be considered when predicting prognosis or determining a treatment plan. Further research is required to establish the specific relation between WBC count and disease progression or death, and the causality between the elevation of WBC count and disease progression should be thoroughly examined. Although the study was limited to all-cause and CVD mortality, the relevance of WBC count to mortality from more diverse causes should be explored. Further research that evaluates WBC count together with other biomarkers may reveal a stronger prognostic factor.

Declaration of Competing Interest

The authors have no conflicts of interest to declare.

Data availability statement

Data sharing is not applicable to this article as no datasets were generated or analyzed in this study.

1. Wu TH, Chien KL, Lin HJ, Hsu HC, Su TC, Chen MF, Lee YT. Total white blood cell count or neutrophil count predict ischemic stroke events among adult Taiwanese: report from a community-based cohort study. *BMC Neurol* 2013;13:7.
2. Margolis KL, Manson JE, Greenland P, Rodabough RJ, Bray PF, Safford M, Grimm RH, Howard BV, Assaf AR, Prentice R, Women's Health Initiative Research Group. Leukocyte count as a predictor of cardiovascular events and mortality in postmenopausal women: the Women's Health Initiative observational study. *Arch Intern Med* 2005;165:500-508.
3. Koo HK, Kang HK, Song P, Park HK, Lee SS, Jung H. Systemic white blood cell count as a biomarker associated with severity of chronic obstructive lung disease. *Tuberc Respir Dis (Seoul)* 2017;80:304-310.
4. Nerpin E, Jacinto T, Fonseca JA, Alving K, Janson C, Malinovschi A. Systemic inflammatory markers in relation to lung function in NHANES. 2007-2010. *Respir Med* 2018;142:94-100.

5. Abramson N, Melton B. Leukocytosis: basics of clinical assessment. *Am Fam Phys* 2000;62:2053–2060.
6. Bargetzi MJ. Leukopenia/neutropenia [Leukopenie/neutropenie] [in German]. *Ther Umsch* 2006;63:78–82.
7. de Labry LO, Campion EW, Glynn RJ, Vokonas PS. White blood cell count as a predictor of mortality: results over 18 years from the Normative Aging Study. *J Clin Epidemiol* 1990;43:153–157.
8. Danesh J, Collins R, Appleby P, Peto R. Association of fibrinogen, C-reactive protein, albumin, or leukocyte count with coronary heart disease: meta-analyses of prospective studies. *JAMA* 1998;279:1477–1482.
9. Danesh J, Whincup P, Walker M, Lennon L, Thomson A, Appleby P, Gallimore JR, Pepys MB. Low grade inflammation and coronary heart disease: prospective study and updated meta-analyses. *BMJ* 2000;321:199–204.
10. Yarnell JW, Baker IA, Sweetnam PM, Bainton D, O'Brien JR, Whitehead PJ, Elwood PC. Fibrinogen, viscosity, and white blood cell count are major risk factors for ischemic heart disease. The Caerphilly and Speedwell collaborative heart disease studies. *Circulation* 1991;83:836–844.
11. Trivedi A, Khan MA, Bade G, Talwar A. Orchestration of neutrophil extracellular traps (nets), a unique innate immune function during chronic obstructive pulmonary disease (COPD) development. *Biomedicines* 2021;9:53.
12. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Int J Surg* 2021;88:105906.
13. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–1558.
14. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–634.
15. Goldet G, Howick J. Understanding GRADE: an introduction. *J Evid Based Med* 2013;6:50–54.
16. Leng SX, Xue QL, Huang Y, Ferrucci L, Fried LP, Walston JD. Baseline total and specific differential white blood cell counts and 5-year all-cause mortality in community-dwelling older women. *Exp Gerontol* 2005;40:982–987.
17. Chmielewski PP, Strzelec B. Elevated leukocyte count as a harbinger of systemic inflammation, disease progression, and poor prognosis: a review. *Folia Morphol* 2018;77:171–178.
18. Nilsson G, Hedberg P, Öhrvik J. White blood cell count in elderly is clinically useful in predicting long-term survival. *J Aging Res* 2014;2014:475093.
19. Ruggiero C, Metter EJ, Cherubini A, Maggio M, Sen R, Najjar SS, Windham GB, Ble A, Senin U, Ferrucci L. White blood cell count and mortality in the Baltimore Longitudinal Study of Aging. *J Am Coll Cardiol* 2007;49:1841–1850.
20. Johnson DW, Wiggins KJ, Armstrong KA, Campbell SB, Isbel NM, Hawley CM. Elevated white cell count at commencement of peritoneal dialysis predicts overall and cardiac mortality. *Kidney Int* 2005;67:738–743.
21. Zhang C, Fu Z, Bai H, Lin G, Shi R, Chen X, Xu Q. Admission white blood cell count predicts post-discharge mortality in patients with acute aortic dissection: data from the MIMIC-III database. *BMC Cardiovasc Disord* 2021;21:462.
22. Ahmadi-Abhari S, Luben RN, Wareham NJ, Khaw KT. Seventeen year risk of all-cause and cause-specific mortality associated with C-reactive protein, fibrinogen and leukocyte count in men and women: the EPIC-Norfolk study. *Eur J Epidemiol* 2013;28:541–550.
23. Hsu CW, Lin JL, Lin-Tan DT, Yen TH, Chen KH. White blood cell count predicts all-cause, cardiovascular disease-cause and infection-cause one-year mortality of maintenance hemodialysis patients. *Ther Apher Dial* 2010;14:552–559.
24. Abete I, Lu Y, Lassale C, Verschuren M, van der Schouw Y, Bueno-de-Mesquita B. White cell counts in relation to mortality in a general population of cohort study in the Netherlands: a mediating effect or not? *BMJ (Open)* 2019;9:e030949.
25. Huang WY, Peng TI, Weng WC, Chien YY, Wu CL, Lee M, Chen KH. Higher leukocyte count is associated with higher risk of 3-year mortality in non-diabetic patients with first-ever ischemic stroke. *J Neurol Sci* 2012;316:93–98.
26. Fan X, Huang B, Lu H, Zhao Z, Lu Z, Yang Y, Zhang S, Hui R. Impact of admission white blood cell count on short- and long-term mortality in patients with type A acute aortic dissection: an observational study. *Medicine* 2015;94:e1761.
27. Zhu B, Feng X, Jiang C, Mi S, Yang L, Zhao Z, Zhang Y, Zhang L. Correlation between white blood cell count at admission and mortality in COVID-19 patients: a retrospective study. *BMC Infect Dis* 2021;21:574.
28. Kim KI, Lee J, Heo NJ, Kim S, Chin HJ, Na KY, Chae DW, Kim CH, Kim S. Differential white blood cell count and all-cause mortality in the Korean elderly. *Exp Gerontol* 2013;48:103–108.
29. Huang ZS, Chien KL, Yang CY, Wang CH, Chang TC, Chen CJ. Peripheral differential leukocyte counts and subsequent mortality from all diseases, cancers, and cardiovascular diseases in Taiwanese. *J Formos Med Assoc* 2003;102:775–781.
30. Weiss ST, Segal MR, Sparrow D, Wager C. Relation of FEV1 and peripheral blood leukocyte count to total mortality. the Normative Aging Study. *Am J Epidemiol* 1995;142:493–498. discussion 499.
31. Libby P. The Pathogenesis, Prevention, and Treatment of Atherosclerosis. In: Jameson JL, Fauci AS, Kasper DL, Hauser SL, Longo DL, Loscalzo J, eds. *Harrison's Principle of Internal Medicine*. 20th ed, New York, NY: McGraw-Hill; 2018:2252–2258.
32. Tabas I, Williams KJ, Borén J. Subendothelial lipoprotein retention as the initiating process in atherosclerosis: update and therapeutic implications. *Circulation* 2007;116:1832–1844.
33. Ley K, Laudanna C, Cybulsky MI, Nourshargh S. Getting to the site of inflammation: the leukocyte adhesion cascade updated. *Nat Rev Immunol* 2007;7:678–689.
34. Christodoulidis G, Vittorio TJ, Fudim M, Lerakis S, Kosmas CE. Inflammation in coronary artery disease. *Cardiol Rev* 2014;22:279–288.
35. Aoki S, Nakagomi A, Asai K, Takano H, Yasutake M, Seino Y, Mizuno K. Elevated peripheral blood mononuclear cell count is an independent predictor of left ventricular remodeling in patients with acute myocardial infarction. *J Cardiol* 2011;57:202–207.
36. Serrano CV Jr, Ramires JA, Venturini M, Arie S, D'Amico E, Zweier JL, Pileggi F, da Luz PL. Coronary angioplasty results in leukocyte and platelet activation with adhesion molecule expression. Evidence of inflammatory responses in coronary angioplasty. *J Am Coll Cardiol* 1997;29:1276–1283.
37. Wu D, Shen YH, Russell L, Coselli JS, LeMaire SA. Molecular mechanisms of thoracic aortic dissection. *J Surg Res* 2013;184:907–924.
38. Kurihara T, Shimizu-Hirota R, Shimoda M, Adachi T, Shimizu H, Weiss SJ, Itoh H, Hori S, Aikawa N, Okada Y. Neutrophil-derived matrix metalloproteinase 9 triggers acute aortic dissection. *Circulation* 2012;126:3070–3080.
39. Henry BM, de Oliveira MHS, Benoit S, Plebani M, Lippi G. Hematology, biochemical and immune biomarker abnormalities associated with severe illness and mortality in coronavirus disease 2019 (COVID-19): a meta-analysis. *Clin Chem Lab Med* 2020;58:1021–1028.
40. Henry BM. COVID-19, EMO, and lymphopenia: a word of caution. *Lancet Respir Med* 2020;8:e24.
41. Bhatt DL, Chew DP, Lincoff AM, Simoons ML, Harrington RA, Ommen SR, Jia G, Topol EJ, PURSUIT Investigators. Effect of revascularization on mortality associated with an elevated white blood cell count in acute coronary syndromes. *Am J Cardiol* 2003;92:136–140.
42. Gurm HS, Bhatt DL, Lincoff AM, Tchong JE, Kereiakes DJ, Kleiman NS, Jia G, Topol EJ. Impact of preprocedural white blood cell count on long term mortality after percutaneous coronary intervention: insights from the EPIC, EPILOG, and EPISTENT trials. *Heart* 2003;89:1200–1204.
43. Newall N, Grayson AD, Oo AY, Palmer ND, Dihmis WC, Rashid A, Stables RH. Preoperative white blood cell count is independently associated with higher perioperative cardiac enzyme release and increased 1-year mortality after coronary artery bypass grafting. *Ann Thorac Surg* 2006;81:583–589.
44. Palmerini T, Mehran R, Dangas G, Nikolsky E, Witzensbichler B, Guagliumi G, Dudek D, Genereux P, Caixeta A, Rabbani L, Weisz G, Parise H, Fahy M, Xu K, Brodie B, Lansky A, Stone GW. Impact of leukocyte count on mortality and bleeding in patients with myocardial infarction undergoing primary percutaneous coronary interventions: analysis from the Harmonizing Outcome with revascularization

- and Stent in acute myocardial infarction trial. *Circulation* 2011;123:2829–2837.
45. Palmerini T, G  n  reux P, Mehran R, Dangas G, Caixeta A, Riva DD, Mariani A, Xu K, Stone GW. Association among leukocyte Count, Mortality, and Bleeding in Patients with non–ST-Segment Elevation Acute Coronary Syndromes (from the Acute Catheterization and Urgent Intervention Triage StrategY [ACUITY] Trial). *Am J Cardiol* 2013;111:1237–1245.
46. Taglieri N, Bacchi Reggiani ML, Palmerini T, Cinti L, Saia F, Guastaroba P, Marrozzini C, Moretti C, Montefiori M, Rosmini S, Alessi L, Vagnarelli F, Branzi A, Rapezzi C, Marzocchi A. Baseline white blood cell count is an independent predictor of long-term cardiovascular mortality in patients with non-ST-segment elevation acute coronary syndrome, but it does not improve the risk classification of the GRACE score. *Cardiology* 2013;124:97–104.
47. Yen MH, Bhatt DL, Chew DP, Harrington RA, Newby LK, Ardisino D, Van de Werf F, White JA, Moliterno DJ, Topol EJ. Association between admission white blood cell count and one-year mortality in patients with acute coronary syndromes. *Am J Med* 2003;115:318–321.