

Find solutions for your homework

Search

home / study / math / advanced math / advanced math questions and answers / 1. let  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  be a convex function that is  $f(tx_1 + (1-t)x_2)$

## Question: 1. Let $f: \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function that is $f(tx_1 + (1-t)x_2)$

1. Let  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  be a convex function that is  $f(tx_1 + (1-t)x_2) \leq tf(x_1) + (1-t)f(x_2)$ . Let  $\mathbf{x}$  be a random vector with joint PDF  $p(\mathbf{x})$ . If  $f$  is a convex function, then show that

$$\mathbb{E}_{\mathbf{x} \sim p}[f(\mathbf{x})] \geq f(\mathbb{E}_{\mathbf{x} \sim p}[\mathbf{x}]).$$

Show transcribed image text

## Expert Answer



Amitakshar Biswas answered this  
476 answers

Was this answer helpful?



We know that  $f(x)$  is a convex function.

Now let,  $L(x) = a + bx$  is a linear function tangential to  $f(x)$  at point  $E[x]$ .

Hence, since  $f$  is convex and  $L$  is tangential to  $f$ , we know by definition that:

$$f(x) \geq L(x) \quad \forall x \in X$$

Thus,

$$\mathbb{E}[f(X)] \geq \mathbb{E}[L(X)] \geq \mathbb{E}[a + bX] \geq a + b\mathbb{E}[X] \geq L(\mathbb{E}[X]) \geq f(\mathbb{E}[X])$$

The proof is quite straightforward. If one function is always greater than or equal to another function, then the unconditional expectation of the first should be at least as large as that of the second. The interior of the proof follow from the definition of  $L$ , the linearity of expectations, and another application of the definition of  $L$ .

The finally, by the definition of the straight line  $L$ , we know that  $L(\mathbb{E}[X])$  is tangential with  $f$  at  $\mathbb{E}[X]$ .  $\mathbb{E}[L(X)] = L(\mathbb{E}[X]) = f(\mathbb{E}[X])$ .

Proved!

Comment >

## Questions viewed by other students

Q: 2. Consider that  $p = N(\mu, \Sigma)$  and  $q = N(\mu_0, \Sigma_0)$ . Here  $\Sigma = \text{diag}(\sigma_1^2, \dots, \sigma_k^2)$ . Then, shown that the KL divergence between  $p$  and  $q$  is defined as below.  $D_{\text{KL}}(p||q) = \frac{1}{2} \sum_{i=1}^k (\frac{\sigma_i^2}{\sigma_{i0}^2} + \ln \frac{\sigma_i^2}{\sigma_{i0}^2})$

A: [See answer](#)

100% (1 rating)



Snap a photo from your phone to post a question

We'll send you a one-time download link

888-888-8888

Text n

By providing your phone number, you agree to receive a automated text message with a link to get the app. Standard messaging rates may apply.

---

[COMPANY](#) ▾

---

[LEGAL & POLICIES](#) ▾

---

[CHEGG PRODUCTS AND SERVICES](#) ▾

---

[CHEGG NETWORK](#) ▾

---

[CUSTOMER SERVICE](#) ▾

---

