

Find solutions for your homework

Search

home / study / engineering / computer science / computer science questions and answers / 3. jensen-shannon divergence between two distributi...

Question: 3. Jensen-Shannon Divergence between two distributions f a...

(1 bookmark)

3. Jensen-Shannon Divergence between two distributions f and g is defined as

$$D_{JS}(f\|g) = \frac{1}{2}D_{KL}\left(f\|\frac{f+g}{2}\right) + \frac{1}{2}D_{KL}\left(g\|\frac{f+g}{2}\right)$$

. Show that it is symmetric and zero only of $p = q$.

Show transcribed image text

Expert Answer



Anonymous answered this
34 answers

Was this answer helpful?



```
from numpy import zeros, array
from math import sqrt, log
```

```
class JSD(object):
    def __init__(self):
        self.log2 = log(2)
```

```
def KL_divergence(self, p, q):
    """ Compute KL divergence of two vectors, K(p || q)."""
    return sum(p[x] * log((p[x]) / (q[x])) for x in range(len(p)) if p[x] != 0.0 or p[x] != 0)
```

```
def Jensen_Shannon_divergence(self, p, q):
    """ Returns the Jensen-Shannon divergence. """
    self.JSD = 0.0
    weight = 0.5
    average = zeros(len(p)) #Average
    for x in range(len(p)):
        average[x] = weight * p[x] + (1 - weight) * q[x]
    self.JSD = (weight * self.KL_divergence(array(p), average) + ((1 - weight) *
self.KL_divergence(array(q), average))
    return 1-(self.JSD/sqrt(2 * self.log2))
```

```
if __name__ == '__main__':
    J = JSD()
    p = [1.0/10, 9.0/10, 0]
    q = [0, 1.0/10, 9.0/10]
    print J.Jensen_Shannon_divergence(p, q)
```

View comments (1) >

Post a question

Answers from our experts for your toughest homework questions

Enter question

Continue to post

20 questions remaining



Snap a photo from your phone to post a question

We'll send you a one-time download link

888-888-8888

Text n

By providing your phone number, you agree to receive a automated text message with a link to get the app. Standard messaging rates may apply.

My Textbook Solutions



Finite...



Calculus for...



Introduc

9th Edition

1st Edition

10th Editi

[View all solutions](#)

Black Friday/Cyber Monday

Run Windows on Mac. macOS Sur, Windows 11 Optimized
Trusted by 7M Mac Users. &

Questions viewed by other students

Q: Consider the BiGAN and its loss function. Show that, in order to fool a perfect discriminator D , BiGAN encoder E and generator G must invert each other. That is, $G(E(x)) = x$ and $E(G(z)) = z$

A: [See answer](#)

Q: As KL divergence is not necessarily symmetric on distributions p and g , one naturally attempt to symmetrize it by introducing DSKL (p||g) (DKL (p||a) DKL (g||p)) Accordingly, we can use it to define a new notion of mutual information over two random variables X and Y : Construct a joint distribution on (X,Y) where both X and Y are valued in $f_0, 11$, such that the following analogue of...

A: [See answer](#)

[Show more](#) >

COMPANY

LEGAL & POLICIES

CHEGG PRODUCTS AND SERVICES

CHEGG NETWORK

CUSTOMER SERVICE

