

Challenge 1: The Hadwiger Conjecture

Definition 1 (Hadwiger Number). Given a graph G , the *Hadwiger number* of G is the largest integer n such that K_n is a minor of G ; here K_n denotes the complete graph on n vertices. We denote the Hadwiger number by $h(G)$.

Conjecture 1.1 (Hadwiger's Conjecture [1, Conjecture 1]). Let $r \in \mathbb{N}$. For every finite simple graph G , we have:

$$r \leq \chi(G) \implies r \leq h(G)$$

While Conjecture 1.1 is not difficult for every small values of r , already for $r = 4$ it is equivalent to the 4-colour theorem. Robertson, Seymour and Thomas proved the above for $r = 5$ [2].

References

- [1] R. Steiner. “Hadwiger’s conjecture and topological bounds”. In: *European Journal of Combinatorics* 122 (2024), p. 104033. ISSN: 0195-6698. DOI: <https://doi.org/10.1016/j.ejc.2024.104033>. URL: <https://www.sciencedirect.com/science/article/pii/S0195669824001185>.
- [2] N. Robertson, P. Seymour, and R. Thomas. “Hadwiger’s conjecture for K_6 -free graphs”. In: *Combinatorica* 13 (1993), pp. 279–361. DOI: 10.1007/BF01202354.