CLASSIFICATION OF BIPARTITE OBSERVABLES II: NIP OBSERVABLES AND TODORČEVIĆ'S TRICHOTOMY

EDUARDO DUEÑEZ AND JOSÉ IOVINO

Abstract. We classify bipartite observables through the regularity of their idealized responses, the expectation values of Alice's limiting preparations regarded as functions of Bob's probes, with Bob's system idealized first. Continuing the study of stability in Part I, we consider bounded self-adjoint observables on tensor products of separable Hilbert spaces. The absence of an infinite sequence of Alice's preparations on which Bob's probes realize every finite pattern at a fixed positive gap, called NIP, is equivalent to all idealized responses being of the first Baire class, that is, pointwise limits of sequences of continuous functions. These responses then form a separable Rosenthal compactum. Probabilistic mixtures make this compactum convex, so hereditary separability forces metrizability and the full response compacta fall into three of the four general classes associated with Todorčević's trichotomy. Physical order comparisons realize all three classes, while selected trees of preparations realize the seven canonical families of the Argyros–Dodos–Kanellopoulos heptachotomy. For the ferromagnetic transverse-field Ising model at positive temperature, the finite-volume magnetization family with offset longitudinal fields is stable when spontaneous magnetization vanishes; otherwise its closure is first countable but not hereditarily separable.

Contents

1. Introduction 2
2. Main results 7
3. A family of continuous functions is NIP exactly when its closure consists of Baire-1 functions 11
4. Separable Rosenthal compacta fall into finitely many canonical classes 14
5. The idealized responses of an observable form a compact set of functions 18
6. Stability, NIP and the independence property are continuity, Baire class 1 and $\beta\mathbb{N}$ 21
7. Three physical orders realize the three classes of Todorčević's trichotomy that occur for observables 27
8. The seven minimal families in physics 37
9. Spontaneous symmetry breaking is a split pair 43
10. Chiral symmetry breaking and metallic conduction are instabilities 50
11. Questions 54
Acknowledgments 55
Declarations 55

Date: October 27, 2026.

2020 *Mathematics Subject Classification.* Primary 03C45, 54D30, 81P15; Secondary 03C66, 03E15, 26A21, 46L30, 54C35, 82B10, 82B20, 82B26.

Key words and phrases. Bipartite observables, stability, NIP, dividing lines, Rosenthal compacta, Todorčević's trichotomy, Argyros–Dodos–Kanellopoulos heptachotomy, Baire class one, orders of limits, Lee–Yang theorem, transverse-field Ising model.

The most recent version of this paper is available at https://utsa-quantum-math.github.io/resources/bipartite-observables-2-nip/; it may supersede the version deposited on arXiv.

References 55

1. Introduction

We extend the study of commuting idealizations in Part I to a classification of idealized responses, the expectation values of limiting preparations regarded as functions of probes. We determine the regularity these functions retain when the two orders of idealization give different answers. As in Part I [DI26], two quantum systems, Alice's and Bob's, are prepared independently, and a bipartite observable, a physical quantity of the joint system, is measured. We work with separable Hilbert spaces and bounded self-adjoint observables; an actual preparation is described by a density operator, a positive operator of trace 1. Every density operator defines a state of the algebra of bounded operators, that is, a positive linear functional with value 1 at the identity, and we use the remaining states as a mathematical model of idealized preparations. Such states arise as limits of actual preparations in the topology of convergence of expectation values, and provide an algebraic framework for the analysis of idealization. An observable is *stable* when its expectation value does not depend on which system is idealized first. In Part I we identify this condition with properties from analysis, topology, probability and model theory. Stable examples include the swap test, which estimates the overlap of two prepared states, Hong–Ou–Mandel coincidences, in which two photons meeting at a beam splitter leave by different ports, and discrete equality tests, such as "equal photon numbers". These observables are even uniformly stable, with quantitative bounds independent of the dimensions of the Hilbert spaces.

Other observables, no less natural, are of a different kind: for a quantity of Alice's system, given by a self-adjoint operator A, and a quantity of Bob's, given by a self-adjoint operator B, the order comparison $[A < B]$ is the orthogonal projection that represents the event "Bob's value of B exceeds Alice's value of A". The comparison $[A < B]$ is unstable whenever A and B have pure point spectra, that is, orthonormal bases of eigenvectors, and their eigenvalues are unbounded above, as for the observables "Bob's system has more photons than Alice's" and "Bob's oscillator has the higher energy": if Alice and Bob prepare eigenstates of ever higher energy, the probability that Bob's energy is the higher is 1 when Bob's system is idealized first and 0 when Alice's system is. The order of the idealizations matters because an order relation is built into the observable, and observables of this kind supply the principal examples of this paper.

The comparison $[A < B]$ illustrates the distinction between order and independence. Call a sequence of Alice's preparations *independent* for an observable if there are real numbers $a < b$ such that, for any two disjoint finite sets of terms of the sequence, a single preparation of Bob's system, a *probe*, gives expectation values at most a on the first set and at least b on the second. The existence of such a sequence is the *independence property for Alice's responses*; its absence is *NIP for Alice's responses*, where NIP abbreviates "no independence property". Unqualified NIP will refer to this direction, since exchanging Alice and Bob can change the property (see Corollary 8.7). In Section 7 we show that $[A < B]$ is uniformly NIP for all self-adjoint operators A and B: for each gap $b - a$, the size of a finite set of preparations on which probes realize every pattern is bounded by a number depending only on the gap. An independent sequence occurs instead for the fringe observable, whose idealized local detector response follows an interference pattern (see Proposition 7.13). If Alice's preparations produce fringes of wavenumbers $8, 8^2, 8^3, \dots$ on Bob's screen, in units

in which the screen has length 2π, then every finite choice of bright and dark responses is realized by a position of Bob's detector.

The uniform condition also has a statistical interpretation. After normalizing responses to $[0, 1]$, consider independently sampled probe–outcome pairs from an unknown joint distribution. In agnostic learning with absolute loss, one seeks a predictor whose mean absolute error is within a prescribed tolerance of the infimum over the response class, with a prescribed probability of success. For the response classes considered here, distribution-free sample bounds of this kind exist exactly when the fat-shattering dimension is finite at every positive gap [ABDCBH97]. This dimension bounds the size of a finite set of probes on which responses realize every high/low pattern around thresholds that may vary from probe to probe. Its finiteness at every gap is equivalent to uniform NIP by Assouad's duality argument (see Remark 3.8). Part I instead studies prediction along an adaptive sequence of probes, with a threshold fixed within each protocol and mistake bounds uniform in that threshold. Uniform NIP is symmetric in the two systems and implies NIP, whereas NIP for Alice's responses need not be uniform (see Remark 3.8 and Corollary 8.7).

The division of first-order theories, sets of axioms in a formal language, by properties such as stability and NIP, which model theorists call dividing lines, is the subject of Shelah's classification theory [She90]. Part I and the present paper study these properties for the expectation predicate of a fixed observable in a specified preparation structure: stability in Part I and the weaker condition NIP here. Uniform stability and uniform NIP require this predicate to have the corresponding property in every model of the theory of the preparation structure. Stability or NIP of the entire theory requires the corresponding condition for every formula.

The classical theory-level examples illustrate the distinction between stability and NIP. In model theory, an infinite linear order is the prototype of an unstable NIP structure, that is, of a set equipped with relations and operations whose theory is unstable and NIP. The rational order $(\mathbb{Q}, <)$ is unstable, yet it is NIP: over a finite set of parameters it has only polynomially many types in the size of the set, a type being a complete description of a single element in terms of the parameters. The ordered field of real numbers is likewise unstable and NIP, and so, more generally, is every o-minimal structure, that is, every ordered structure in which every set of elements defined by a formula with parameters (a definable set) is a finite union of points and intervals; the o-minimal structures are the best-known unstable NIP structures.[1]

In these model-theoretic examples the formulas are two-valued, even when the underlying set is a continuum. For observables the formulas are real-valued because they are expectation values, and a fixed positive gap separates the high and low values in the independence property. Real-valued forms of NIP already occur in quantum theory through the fat-shattering dimension governing the learnability of quantum states and measurements in finite dimensions [Aar07, CHY15]. Here the object of classification is the entire compactum of idealized responses in the topology of pointwise convergence. The relevant structure theory is that of Rosenthal compacta, compact families of functions of the first Baire class on a Polish space, a separable, completely metrizable space. The passage from finite patterns to the regularity of this compactum is the dictionary on which the classification rests.

The dictionary. Fix an observable, and regard its expectation value for an idealized preparation of Alice's system as a function of Bob's probe, idealized probes included, with Bob's

[1] Other standard examples are the ordered group of integers $(\mathbb{Z}, +, <)$, whose theory is Presburger arithmetic, and, more generally, ordered abelian groups [GS84]; the fields of p-adic numbers, for every prime p; and algebraically closed valued fields (see [Sim15a, Chapter 2]). For o-minimal structures see [PS86, KPS86]; the real field with the exponential function is o-minimal by a theorem of Wilkie [Wil96].

system idealized first; we call this function an *idealized response*. To obtain the domain of these responses, average the observable against each of Alice's actual preparations, obtaining effective observables on Bob's Hilbert space, and take the smallest unital norm-closed algebra containing these operators and their adjoints. A probe is a state of this algebra, a positive linear functional taking the value 1 at the identity. Thus Bob's states are retained through their restrictions to the algebra generated by the fixed observable's effective observables. This algebra is norm-separable, and its state space is compact and metrizable in the topology of convergence of values on the algebra (see Definition 5.1 and Lemma 5.2). In Section 5 we show that the idealized responses form a compact set of functions on this probe space, with the topology of pointwise convergence; in Theorem 6.1 we identify the observable-level conditions with degrees of regularity of these functions:

- the observable is stable exactly when every idealized response is continuous;
- it is NIP exactly when every idealized response is of the first Baire class, that is, a pointwise limit of a sequence of continuous functions; the idealized responses then form a *Rosenthal compactum*, a compact set of functions of the first Baire class, and every one of them is the limit of a sequence of responses of actual preparations (see [BFT78, Theorem 3F]);
- otherwise, some sequence of Alice's actual preparations is independent, and the pointwise closure of the responses of a subsequence is a copy of $\beta\mathbb{N}$, the space of ultrafilters on the natural numbers (see Theorem 6.1(c)).

The fine classification. The compactum of idealized responses is separable (see Lemma 6.3), and separable Rosenthal compacta form a vast class; a priori there is no reason why the failure of such a compactum to be metrizable should be witnessed by anything simple. Todorčević's trichotomy [Tod99] shows that a simple witness always exists: a non-metrizable separable Rosenthal compactum is either hereditarily separable (every subspace has a countable dense subset), or first countable (every point has a countable base of neighborhoods) but not hereditarily separable, or not first countable, and in these three cases it contains, respectively, a copy of the split interval, a copy of the Alexandroff duplicate of the Cantor set or of its analogue for the split Cantor set, and a copy of the one-point compactification of a discrete set of size continuum. The split interval is the unit interval with each point doubled into two adjacent points, in the order topology, and the split Cantor set is the Cantor set doubled in the same way; the Alexandroff duplicate of the Cantor set adds to the Cantor set an isolated twin of each of its points, in such a way that the twins of a sequence of points converging to another point converge to that point as well. In this the trichotomy resembles Rosenthal's ℓ_1 theorem [Ros74]: a bounded sequence in a real Banach space has either a weakly Cauchy subsequence, one whose values under each continuous linear functional form a Cauchy sequence, or a subsequence equivalent to the unit vector basis of ℓ_1, the space of absolutely summable real sequences. Argyros, Dodos and Kanellopoulos extracted from the methods of Todorčević's paper, with the help of Ramsey theory for trees, a refinement of the trichotomy to the level of families, the Argyros–Dodos–Kanellopoulos heptachotomy [ADK08]. Index a pointwise bounded family of pairwise distinct functions on a Polish space by the dyadic tree, the set of finite sequences of zeros and ones, and suppose that its pointwise closure consists of functions of the first Baire class. Then there is a dyadic subtree, a set of nodes that, ordered by extension, forms a copy of the dyadic tree, such that the subfamily indexed by it, reindexed by the dyadic tree through the isomorphism that also preserves the lexicographic order, is equivalent to one of exactly seven minimal families; two families indexed by the same set are *equivalent* when the correspondence between them extends to a homeomorphism of their pointwise closures, and a family is *minimal* when it is equivalent to its

restriction to every dyadic subtree, reindexed in the same way. For a NIP observable, a tree of Alice's preparations with pairwise distinct responses gives such a family, for instance a tree of successively sharper localizations of a particle, and the heptachotomy shows that, up to passing to a subtree, there are seven ways, and only seven, in which the responses of such a tree can approach their idealized limits. Through the connection among NIP, the Vapnik–Chervonenkis dimension (the two-valued form of the fat-shattering dimension) and Rosenthal compacta, the trichotomy also classifies NIP formulas whose parameters range over a compact metrizable space [DIMW+26, Section 4.2].

For observables, the classification is coarser, for a reason that comes from probability. Alice can mix two preparations, that is, make one or the other at random with given probabilities, and the response of the mixture is the corresponding convex combination of the two responses; hence the idealized responses form a convex set (see the proof of Corollary 6.8). In Theorem 6.6 we give a direct proof of a special case of a known result on locally connected Rosenthal compacta: a separable convex Rosenthal compactum that is hereditarily separable is metrizable. Consequently, the compactum of idealized responses of a NIP observable is metrizable, or first countable but not hereditarily separable, or not first countable (see Corollary 6.8 and Theorem 6.9), and NIP observables fall into three classes of Todorčević's trichotomy, the class of metrizable compacta among them. As we show in Section 7, three physical order relations lie in these three classes, respectively: the comparison of photon numbers, the comparison of positions or of quadratures of light (the two field amplitudes of a mode, which behave like a position and a momentum), and the causal order of a spacetime with one dimension of space. The compactum of idealized responses of the causal order fails to be first countable because an event that occurs at the time of Bob's detection can influence a detection only at its own position. The comparison of two phases on a circle, we prove in Theorem 7.12, lies with the causal order in the last class: on a circle, for each pair of antipodal points there is an idealized response that differs from the constant response of the uniform distribution only on probes that give weight to one of the two points. Trees of preparations, nevertheless, realize all seven minimal families, as we show in Theorem 8.5 (see Figure 2): the closure of the responses of a tree is a closed subset of the compactum of idealized responses, and it need not be convex.

The canonical witnesses in this classification also exhibit a role for set-theoretic methods in the analysis of response spaces. Todorčević's proof of the trichotomy uses forcing [Tod99]: the required configuration is constructed in a generic extension, a larger universe of sets, and an absoluteness argument transfers the conclusion back to the original universe. The resulting theorem is a theorem of ZFC, the Zermelo–Fraenkel axioms with the axiom of choice. To our knowledge, no proof of the trichotomy that avoids forcing is known [DIMW+26]. In this paper the trichotomy supplies Theorems 4.7, 6.6 and 6.9, Corollary 6.8 and the canonical subfamilies in Theorems 7.8, 7.10 and 7.12. The location of each physical example in its topological class is proved directly; forcing enters through the structure theorem that supplies canonical witnesses and through its consequences for convex response compacta.

Orders of limits and the quantum Ising model. The same classification applies to response families arising from thermodynamic limits, even when no single bipartite observable is specified. For a ferromagnet, the responses are the magnetizations of finite samples as functions of the applied field, and the probes are the field values. The familiar order of limits has a precise mathematical reading: a spontaneous magnetization can remain when the volume tends to infinity before the external field is switched off, while reversing these limits gives zero (see the proof of Theorem 9.2). A NIP family with separable closure is NIP_1, NIP_2 or NIP_3 when its closure is, respectively, first countable, hereditarily separable or metrizable

(see Section 4); in particular, NIP_3 means that one countable set of probes distinguishes all idealized responses (see Propositions 4.5 and 4.6). The family of finite-volume magnetizations is stable exactly when spontaneous magnetization vanishes. To distinguish the finer levels, we also allow an offset field, which translates the field argument and moves the transition point. When spontaneous magnetization is positive, this enlarged family is NIP_1 but not NIP_2, while the family without offsets remains NIP_3 (see Theorems 9.2 and 9.6). At each translated transition point, the responses of the two oppositely magnetized pure phases form a split pair, and the response of their equal mixture is its midpoint. These midpoints give the same obstruction to hereditary separability as in Theorem 6.6. For the ferromagnetic Ising model in a transverse field, a lattice of spins coupled along one axis in a field perpendicular to it, we establish this classification at positive temperature in Theorem 9.6. The model-specific input is differentiability of the pressure at every nonzero longitudinal field, the field along the coupling axis. The pressure is the limit per site of the logarithm of the partition function, the normalizing constant of the thermal equilibrium state. We derive differentiability, together with real analyticity there, from a Lee–Yang statement asserting that the finite-volume partition functions have no zeros off the imaginary axis of the complex longitudinal field (see Propositions 9.4 and 9.5). We include a proof of this statement and discuss its relation to the quantum Lee–Yang theorems of Asano [Asa70], Suzuki and Fisher [SF71], and their formulation by Fröhlich and Rodriguez [FR12]. Two further orders of limits receive a stability interpretation in Section 10 (see Theorems 10.3 and 10.4): the Banks–Casher relation [BC80], tying the chiral condensate of quarks to the density of near-zero eigenvalues of the Dirac operator governing quarks in a gluon field, and the Scalapino–White–Zhang criterion [SWZ93], distinguishing superconductors, metals and insulators by their current responses. For the spectral families considered here, stability is equivalent to the vanishing of the valence condensate in the specified ensemble, or to agreement of the static and dynamic current-response limits, respectively.

Contributions of the paper. The results concern the relation between finite patterns of expectation values, the regularity of their idealized limits and the topology of the resulting response spaces. The physical orders in Section 7 realize the three classes available to full convex NIP response compacta, while the trees in Section 8 realize all seven canonical families inside such spaces. The examples involve photon numbers, positions and quadratures, planar rotors (particles on a ring, whose positions are angles), causal order and chains of spins; their response functions display the features that distinguish the classes. For the offset magnetization family of the transverse-field Ising model, the same distinction is expressed by the split pairs and their midpoints at translated transition points (see Theorem 9.6). This is a classification of response functions and their closures, with interpretations in terms of preparations, probes and orders of limits. Its logical formulation must therefore retain both the real values of expectation functions and their affine dependence on mixtures.

The logical setting. The classification concerns real-valued expectation predicates, with high and low responses separated by a fixed positive gap, and preserves their affine dependence on probabilistic mixtures. We formulate it in Keisler's model theory of real-valued structures [Kei23], in which no metric is prescribed and each formula has its own bound. For the bounded observables considered here, the continuous logic of Ben Yaacov and Usvyatsov [BYU10] would also serve. The relation between these frameworks, and the treatment of unbounded quantities through restrictions with bounded expectations, are explained in Section 5.2.

Related work. Model theory has met operator algebras and quantum theory along other routes. Farah, Hart and Sherman developed the continuous model theory of C*-algebras,

the algebras of bounded operators on a Hilbert space that are closed under adjoints and in the norm, and of tracial von Neumann algebras, those that are closed under adjoints and in the weak operator topology (the topology of convergence of all matrix elements) and carry a trace, a faithful normal state that vanishes on commutators; in that setting they studied stability [FHS13]; there the structures are the algebras themselves, and stability is a property of their theories, whereas here the object is a single observable together with its idealized responses. Zilber gave a model-theoretic semantics of the canonical commutation relation [Zil16]. On the side of real-valued NIP, Ben Yaacov showed that NIP is preserved under Keisler randomization, the passage from a theory to a theory of random variables with values in its models [Ben09], whence averages of NIP relations over random parameters are again NIP; Proposition 7.3 is an elementary, quantitative instance of this phenomenon for relations between quantities measured by Alice and by Bob. Hrushovski and Pillay studied invariant measures in NIP theories [HP11], Simon connected NIP with Rosenthal compacta [Sim15b], and Glasner and Megrelishvili used Todorčević's trichotomy to build a hierarchy of tame dynamical systems [GM22]. To our knowledge, none of these works considers the orders of limits of a bipartite observable, the compactum of its idealized responses, or the place of physical observables in Todorčević's trichotomy and in the Argyros–Dodos–Kanellopoulos heptachotomy; Part I concerns the first of these topics, and the present paper concerns all three.

The paper is organized as follows. We state the main results in Section 2. The facts about Rosenthal compacta that we need, and a classification of countable NIP families of continuous functions that we derive from them (see Theorem 4.7), occupy Sections 3 and 4. In Sections 5 and 6 we attach to each observable its compactum of idealized responses, prove Theorem 6.1, and show that, for the compactum of idealized responses of a NIP observable, hereditary separability implies metrizability (see Theorem 6.6 and Corollary 6.8). The examples follow in Sections 7 and 8; spontaneous magnetization, classical and quantum, is the subject of Section 9, and two further orders of limits are treated in Section 10. Open questions are collected in Section 11. Standard arguments are cited, and sketched where the adaptation is not routine.

2. Main results

The classification of observables and of families of responses is drawn in Figure 1, with the physical systems that inhabit each class and the minimal families that each class contains, and in Figure 2 each of the seven minimal families of the heptachotomy is followed to its idealized limits. The classes of the classification, among them those of Todorčević's trichotomy, are listed in Table 1 and the minimal families of the heptachotomy in Table 2, each with a physically natural example and with the feature of the example that the class or family isolates.

Throughout, "Alice's responses" are the expectation values of Alice's actual preparations as functions of Bob's probes; NIP for an observable refers to this direction. Her idealized responses, with Bob's system idealized first, form the pointwise closure of these functions (see Lemma 5.4). A NIP family with separable closure is NIP_1, NIP_2 or NIP_3 when its closure is, respectively, first countable, hereditarily separable or metrizable. The two tables distinguish the classification of a whole response space from that of a selected tree of responses: mixtures make the full compactum of an observable convex, forcing NIP_2 to coincide with NIP_3, while selected trees can realize all seven minimal families because their closures need not be convex (see Corollary 6.8 and Theorem 8.5).




FIGURE 1. The classes of observables and of families of responses, with their physical examples and the minimal families of the heptachotomy that they contain. The labels $D_1, \ldots, D_7$ name the seven minimal families of the heptachotomy, and the label $\hat{A}(2^{\mathbb{N}})$ names the compactum that extends the one-point compactification of the Cantor set, regarded as a discrete space (see Tables 1 and 2). The regions are nested: uniformly stable $\subseteq$ stable $\subseteq$ NIP$_3$ $\subseteq$ NIP$_2$ $\subseteq$ NIP$_1$ $\subseteq$ NIP. The innermost examples are those of Part I [DI26].

The results behind the tables are the following:

(A) *The dictionary.* An observable is stable exactly when its idealized responses are continuous, and NIP exactly when they are of the first Baire class; when it is not NIP, some sequence of responses of Alice's actual preparations has a closure homeomorphic to $\beta\mathbb{N}$ (see Theorem 6.1).

(B) *Mixtures force metrizability.* A separable convex Rosenthal compactum (a compact set of functions of the first Baire class on a Polish space) that is hereditarily separable is metrizable (see Theorem 6.6); hence NIP$_2$ observables are NIP$_3$ (see Corollary 6.8), and the fine classification of NIP observables has three classes of Todorčević's trichotomy (see Theorem 6.9).

(C) *Natural examples.* The comparison of photon numbers is NIP$_3$, the comparison of positions or of quadratures of light (the two field amplitudes of a mode) is NIP$_1$ but not NIP$_2$, and the causal order of a spacetime with one dimension of space and the comparison of two phases are NIP but not NIP$_1$ (see Theorems 7.8, 7.10 and 7.12 and Proposition 7.9); each minimal family is realized by a tree of preparations of one of these observables or

Class	Topological criterion or witness	Meaning for responses	Physical setting	Result
stable	continuous	idealizations commute	swap test; gapped transverse-current response	10.4
NIP_3	metrizable	one countable set of probes separates all responses	photon-number comparison; spin-chain scanning	7.8
$\mathrm{NIP}_2 \setminus \mathrm{NIP}_3$	split interval	hereditarily separable but nonmetrizable	absent for full observable compacta	6.6
$\mathrm{NIP}_1 \setminus \mathrm{NIP}_2$	duplicate compactum	each response has its own countable determining probes	positions; quadratures; Ising with offsets	9.6
$\mathrm{NIP} \setminus \mathrm{NIP}_1$	$\hat{A}(2^{\mathbb{N}})$	some response has no countable determining set of probes	causal order; phases on a circle	7.10, 7.12
not NIP	$\beta\mathbb{N}$	one infinite preparation sequence admits every finite high/low pattern at a fixed positive gap	interference fringes	7.13

TABLE 1. From the regularity of idealized responses to their physical interpretation. The NIP levels are nested: $\mathrm{NIP}_3 \subseteq \mathrm{NIP}_2 \subseteq \mathrm{NIP}_1 \subseteq \mathrm{NIP}$; the stable row lies within NIP_3. The four rows from NIP_3 to $\mathrm{NIP} \setminus \mathrm{NIP}_1$ distinguish the four general classes associated with Todorčević's trichotomy. For the full response compactum of an observable, mixtures exclude the row $\mathrm{NIP}_2 \setminus \mathrm{NIP}_3$ (see Corollary 6.8). A determining set of probes identifies one response among all responses in the closure; a separating set distinguishes every pair (see Proposition 4.6). The second column gives either a property of the responses or their closure, or a canonical compactum embedded in that closure (see Section 4): the split interval, the Alexandroff duplicate of the Cantor set or the extended duplicate of the split Cantor set, the compactum $\hat{A}(2^{\mathbb{N}})$, which extends the one-point compactification of the Cantor set $2^{\mathbb{N}}$ regarded as discrete, or the space $\beta\mathbb{N}$ of ultrafilters on $\mathbb{N}$. The Ising entry refers to the finite-volume magnetization family with offset longitudinal fields and positive spontaneous magnetization (see Theorem 9.6). The transverse-current entry concerns the family of Theorem 10.4 under (S1) and (S2), with its spectral measures supported above a common positive energy for all sufficiently small wave vectors; this persistent gap gives $D = D_s$ and hence stability. Physical settings in the fourth column are examples of the indicated class; they are not equivalences defining it. The final column points to the corresponding results.

of an observable that probes a chain of spins, by reading it site by site, by matching it with a template, or both (see Theorem 8.5). The scanning observable, which reads the chain site by site, is NIP for Alice's responses but not for Bob's, so NIP is not symmetric for observables (see Corollary 8.7), although uniform NIP is (see Remark 3.8).

Family	Closure	Pattern of idealization	Selected preparations
D_1	$A(2^{<\mathbb{N}})$	escape to infinity	photon number states
D_2	$2^{\leq\mathbb{N}}$	infinite configurations	a spin chain read site by site
D_3	$\hat{S}_+(2^{\mathbb{N}})$	approach from the right	a particle in a box
D_4	$\hat{S}_-(2^{\mathbb{N}})$	approach from the left	a particle in a box
D_6	$\hat{D}(2^{\mathbb{N}})$	twins over a Cantor set	a spin chain, sites and templates
D_7	$\hat{D}(S(2^{\mathbb{N}}))$	approach from both sides	a particle in a box, half on each side
D_5	$\hat{A}(2^{\mathbb{N}})$	a point-supported limit	light cones; templates

TABLE 2. The seven minimal families $D_1, \dots, D_7$ of the heptachotomy and their closures, the canonical compacta of Section 4, grouped by class of Todorčević's trichotomy as in Figure 2 (see Theorem 8.5). Each row describes the closure of the responses of a selected tree of preparations, which need not be convex. The closures for D_1, D_2 are metrizable; those for D_3, D_4 are hereditarily separable but not metrizable; those for D_6, D_7 are first countable but not hereditarily separable; and that for D_5 is not first countable. Thus all seven families can occur inside observable response spaces even though the full convex NIP compacta have only three classes. Here $2^{<\mathbb{N}}$ is the dyadic tree of finite sequences of zeros and ones, $2^{\mathbb{N}}$ is the Cantor set of infinite such sequences, and $2^{\leq\mathbb{N}}$ is the compact space of all such sequences, finite and infinite, in which the finite ones are isolated. The other closures are the one-point compactification $A(2^{<\mathbb{N}})$ of the countable discrete set $2^{<\mathbb{N}}$, the extended split Cantor sets $\hat{S}_+(2^{\mathbb{N}})$ and $\hat{S}_-(2^{\mathbb{N}})$, the separable extension $\hat{D}(2^{\mathbb{N}})$ of the Alexandroff duplicate of the Cantor set, the extended duplicate $\hat{D}(S(2^{\mathbb{N}}))$ of the split Cantor set $S(2^{\mathbb{N}})$, and the compactum $\hat{A}(2^{\mathbb{N}})$ of Table 1.

(D) *The quantum Ising model.* For the Ising model in a transverse field, the family of magnetizations of finite systems, as functions of the longitudinal field (the field along the axis of the couplings) and with an offset field that moves the transition point (the field at which the magnetization changes sign), is stable, with a metrizable closure, when the spontaneous magnetization vanishes, and it is NIP_1 but not NIP_2 when that magnetization is positive (see Theorem 9.6). The proof uses the real analyticity of the pressure (the limit per site of the logarithm of the partition function) away from zero longitudinal field, which we derive from a Lee–Yang theorem for this model: the partition function of each finite system has no zeros off the imaginary axis of the complex longitudinal field (see Propositions 9.4 and 9.5).

(E) *Orders of limits.* For the classical and quantum Ising models considered here at positive temperature, the family of finite-volume magnetizations, with or without offset fields, is stable exactly when spontaneous magnetization vanishes. For the spectral families considered here, the Banks–Casher relation identifies the stability obstruction with the limiting density of Dirac eigenvalues at zero, while the current-response criterion of Scalapino, White and Zhang identifies it with the dynamic current-response limit minus the static one (see Theorems 9.2, 9.6, 10.3 and 10.4).




FIGURE 2. The seven minimal families of the heptachotomy, arranged by the class of Todorčević's trichotomy to which their closures belong, each with the physical picture of its realization (see Theorem 8.5): how a tree of preparations approaches its idealized limits. In the panels of D_3, D_4 and D_7, the bumps, the distributions of successive preparations of the particle, approach a point a of the box; in that of D_7, each preparation is split into halves, one on each side of a, and in that of D_6, the duplicate is the Alexandroff duplicate of the Cantor set.

The theorem on convex compacta in (B), Theorem 6.6, is a special case of a known result (see Remark 6.7). To our knowledge, its consequences for observables in (B), result (D), the realizations in (C) and the asymmetry of nonuniform NIP for observables are new.

3. A family of continuous functions is NIP exactly when its closure consists of Baire-1 functions

We begin with the topological dichotomy on which the paper rests: a pointwise bounded family of continuous functions on a compact metrizable space either has the independence property or has its pointwise closure inside the functions of the first Baire class. In most of the paper the functions will be responses of Alice's preparations and the points will be Bob's probes.

3.1. **Functions of the first Baire class.** For a topological space Y, let $C_p(Y)$ be the set of continuous functions $Y \to \mathbb{R}$ with the topology of pointwise convergence. A function $f\colon Y \to \mathbb{R}$ is of the *first Baire class*, or *Baire-1*, if it is the pointwise limit of a sequence of continuous functions, and we write $B_1(Y)$ for the set of these functions, again with the topology of pointwise convergence. For $\mathcal{R} \subseteq \mathbb{R}^Y$, we write $\overline{\mathcal{R}}$ for the closure of $\mathcal{R}$ in $\mathbb{R}^Y$, and we call $\mathcal{R}$ *pointwise bounded* if $\sup_{r\in\mathcal{R}} |r(y)| < \infty$ for every $y \in Y$.

We use the classical characterizations of Baire-1 functions on Polish spaces, the separable, completely metrizable spaces: condition (3) of Fact 3.1 below is Baire's characterization, and condition (2) concerns F_σ sets, the countable unions of closed sets.

Fact 3.1 (see [Tod97, Section 10]**).** *Let Y be a Polish space and $f\colon Y \to \mathbb{R}$ a function. The following are equivalent:*

(1) *the function f is Baire-1;*
(2) *for every open $U \subseteq \mathbb{R}$, the set $f^{-1}[U]$ is F_σ;*
(3) *for every nonempty closed $F \subseteq Y$, the restriction $f|_F$ has a point of continuity.*

Moreover, if f is not Baire-1, then there are nonempty countable sets $Z_0, Z_1 \subseteq Y$ with $\overline{Z_0} = \overline{Z_1}$ and reals $a < b$ such that $f \leq a$ on Z_0 and $f \geq b$ on Z_1.

The last clause of Fact 3.1 lets us test membership in $B_1(Y)$ on countable sets, which is what makes pointwise closures manageable.

Lemma 3.2. *Let Y be a Polish space and $\mathcal{R} \subseteq B_1(Y)$ a pointwise bounded family. The following are equivalent:*

(i) *the family $\mathcal{R}$ is relatively compact in $B_1(Y)$;*
(ii) *every sequence in $\mathcal{R}$ has a cluster point in $B_1(Y)$;*
(iii) *we have $\overline{\mathcal{R}} \subseteq B_1(Y)$.*

Sketch. This is the Bourgain–Fremlin–Talagrand argument [BFT78]. The implication (i)⇒(ii) is clear, and (iii)⇒(i) is Tychonov's theorem. For (ii)⇒(iii), suppose that some $f \in \overline{\mathcal{R}}$ is not Baire-1, and take Z_0, Z_1, a and b as in Fact 3.1. A sequence in $\mathcal{R}$ converging to f on the countable set $Z_0 \cup Z_1$ has no cluster point in $B_1(Y)$, because such a point would agree with f on $Z_0 \cup Z_1$ and would therefore have no point of continuity on $\overline{Z_0} = \overline{Z_1}$. □

3.2. **The independence property.** In Theorem 3.6 below, we show that the obstruction to Baire-1 closures is combinatorial: a sequence of functions whose sublevel and superlevel sets meet in every finite pattern.

Definition 3.3. Let Y be a set. A family $\mathcal{R} \subseteq \mathbb{R}^Y$ has the *independence property* if there are a sequence $(r_n)_{n\in\mathbb{N}}$ in $\mathcal{R}$ and reals $a < b$ such that, for all disjoint finite sets $E, F \subseteq \mathbb{N}$, there is $y \in Y$ with $r_n(y) \leq a$ for $n \in E$ and $r_n(y) \geq b$ for $n \in F$. Otherwise, the family $\mathcal{R}$ is *NIP*.

On the Cantor space $2^{\mathbb{N}}$ of infinite sequences of zeros and ones, the family of coordinate functions p_n, given by $p_n(y) = y(n)$ for $n \in \mathbb{N}$, has the independence property, and the closure of $\{p_n : n \in \mathbb{N}\}$ in $\mathbb{R}^{2^{\mathbb{N}}}$ is a copy of $\beta\mathbb{N}$, the space of ultrafilters on $\mathbb{N}$. The next two facts say that this example is the only obstruction.

Fact 3.4 (Rosenthal's dichotomy; see [Ros74, Tod97]**).** *Let Y be a Polish space and $(r_n)_{n\in\mathbb{N}}$ a pointwise bounded sequence in $C_p(Y)$. Then the sequence (r_n) has either a pointwise convergent subsequence or a subsequence whose closure in $\mathbb{R}^Y$ is homeomorphic to $\beta\mathbb{N}$.*

Fact 3.5 (Bourgain–Fremlin–Talagrand; see [BFT78, Theorem 4G]**).** *Let Y be a Polish space and $\mathcal{R} \subseteq C_p(Y)$ a pointwise bounded family. Then $\overline{\mathcal{R}} \subseteq B_1(Y)$ if and only if, for every sequence $(r_n)_{n\in\mathbb{N}}$ in $\mathcal{R}$ and all reals $a < b$, there is $I \subseteq \mathbb{N}$ such that no $y \in Y$ satisfies $r_n(y) \leq a$ for all $n \in I$ and $r_n(y) \geq b$ for all $n \notin I$.*

When the space Y is compact, the condition in Fact 3.5 reduces to a statement about finite patterns, and we obtain the dichotomy announced at the beginning of this section.

Theorem 3.6. *Let Y be a compact metrizable space and $\mathcal{R} \subseteq C_p(Y)$ a pointwise bounded family. Then exactly one of the following holds:*

(i) *we have $\overline{\mathcal{R}} \subseteq B_1(Y)$;*
(ii) *the family $\mathcal{R}$ has the independence property.*

The following proof is the classical passage from infinite to finite patterns by compactness (see [DIMW+26, Section 1]).

Proof. For a sequence (r_n) in $\mathcal{R}$, reals $a < b$ and $I \subseteq \mathbb{N}$, the sets $\{y : r_n(y) \leq a\}$ for $n \in I$ and $\{y : r_n(y) \geq b\}$ for $n \notin I$ are closed, because each function r_n is continuous. By compactness of Y, these closed sets have empty intersection if and only if finitely many of them do. Hence the condition in Fact 3.5 fails for a given sequence (r_n) and given reals $a < b$ exactly when every finite pattern is realized, that is, exactly when (r_n), a and b witness the independence property. By Fact 3.5, alternative (i) therefore fails exactly when alternative (ii) holds. □

We will also need to replace a space of probes by a dense subset; in Lemma 3.7 below, we show that the replacement does not affect the independence property.

Lemma 3.7. *Let Y be a topological space, $\mathcal{R} \subseteq C_p(Y)$ a family and $L \subseteq Y$ a dense set. Then the family $\mathcal{R}$ is NIP if and only if the family $\{r|_L : r \in \mathcal{R}\}$ of restrictions is NIP.*

Sketch. If the restrictions have the independence property, then so does $\mathcal{R}$. Conversely, if (r_n) and $a < b$ witness the independence property of $\mathcal{R}$, and $a < a' < b' < b$, then each finite pattern holds strictly, with a' and b', on a nonempty open set, which meets L; hence the restrictions witness the independence property with the reals $a' < b'$. □

Remark 3.8 (Symmetry). For a formula of a first-order theory, NIP is symmetric in the two variables: the formula $\varphi(x, y)$ is NIP if and only if the formula φ^{opp}, given by $\varphi^{\mathrm{opp}}(y, x) = \varphi(x, y)$, is NIP (see [She90, Chapter II, Section 4] and [Sim15a, Lemma 2.5]); if φ has Vapnik–Chervonenkis (VC) dimension d, that is, if d is the largest size of a finite set of elements every subset of which is cut out by some instance $\varphi(x, c)$ of the formula, for a parameter c that depends on the subset, then the formula φ^{opp} has VC dimension less than 2^{d+1} [Ass83]. Assouad's argument is finite and combinatorial, and it applies, with the reals $a < b$ fixed, to real-valued functions. For uniform NIP, which bounds, for each gap $b - a$, the size of the finite sets of functions on which every pattern of values at most a and at least b is realized at some point (see Remark 6.2 below), the roles of Alice and Bob are therefore interchangeable. The notion of Definition 3.3, which asks for an infinite sequence inside a single family, is not symmetric: the relation $n \in u$ between numbers n and finite sets $u \subseteq \mathbb{N}$ has the independence property in the variable n but not in the variable u.

3.3. **Rosenthal compacta.** A *Rosenthal compactum* is a compact space homeomorphic to a compact subset of $B_1(Y)$, with the topology of pointwise convergence, for some Polish space Y. By Theorem 3.6 and Lemma 3.2, if Y is compact metrizable and $\mathcal{R} \subseteq C_p(Y)$ is pointwise bounded and NIP, then the closure $\overline{\mathcal{R}}$ is a Rosenthal compactum. The classification of Rosenthal compacta (see Section 4) will therefore classify the NIP families that arise below.

3.4. **Historical remarks.** Rosenthal's dichotomy grew out of his ℓ_1 theorem [Ros74]; Bourgain, Fremlin and Talagrand proved Fact 3.5 in 1978 [BFT78], and Godefroy named Rosenthal compacta [God80] (see also [Ai92, Tka11]). The independence property was isolated by Shelah [She71, She90]; the polynomial bound behind its $\{0, 1\}$-valued case, the Sauer–Shelah

lemma, was found independently by Sauer [Sau72] and Shelah [She72], and, a little earlier, by Vapnik and Chervonenkis [VC71, VC74], in their work on the uniform convergence of frequencies to probabilities, which became a foundation of learning theory [Val84]. The connection between NIP and Rosenthal compacta is due to Simon [Sim15b], and its use for limits of computations goes back to our joint work with Matos-Wiederhold, Salvetti and Tall [DIMW$^+$26].

4. Separable Rosenthal compacta fall into finitely many canonical classes

The Baire-1 characterization reduces the analysis of NIP families to the topology of their pointwise closures. For a countable, pointwise bounded NIP family of continuous functions on a compact metrizable space, this closure is a separable Rosenthal compactum. Todorčević's trichotomy distinguishes the nonmetrizable cases by hereditary separability and first countability; the canonical form of the trichotomy identifies minimal families whose closures embed into the corresponding compacta. We state these results together with the Argyros–Dodos–Kanellopoulos heptachotomy, following their notation [ADK08], and formulate the resulting levels of NIP for use with response families.

4.1. **Three topological invariants.** Recall that a space is hereditarily separable if each of its subspaces has a countable dense subset, and first countable if each of its points has a countable base of neighborhoods. Every compact metrizable space is hereditarily separable, and by a theorem of Pol every hereditarily separable Rosenthal compactum is first countable (see [Deb13, Section 10]). For a Rosenthal compactum K we therefore have:

$$K \text{ metrizable} \implies K \text{ hereditarily separable} \implies K \text{ first countable}.$$

For separable Rosenthal compacta, the failure of each converse is witnessed by a canonical subspace, and these canonical subspaces are pointwise closures of families indexed by the dyadic tree, the set of finite sequences of zeros and ones (see Fact 4.3 below).

4.2. **The canonical compacta.** Let $2^{<\mathbb{N}}$ be the set of finite binary sequences, with $|t|$ the length of a sequence $t \in 2^{<\mathbb{N}}$, and let $2^{\mathbb{N}}$ be the Cantor space. For $t \in 2^{<\mathbb{N}}$, let v_t be the characteristic function of the clopen set $V_t = \{\sigma \in 2^{\mathbb{N}} : t \sqsubset \sigma\}$ of extensions of t, and let $t^\frown 0^\infty$ and $t^\frown 1^\infty$ be the sequences that extend t by zeros and by ones. Let $\prec$ be the lexicographic order of $2^{\mathbb{N}}$. For $\sigma \in 2^{\mathbb{N}}$, let x^+_σ and x^-_σ be the characteristic functions of $\{\tau \in 2^{\mathbb{N}} : \sigma \preceq \tau\}$ and $\{\tau \in 2^{\mathbb{N}} : \sigma \prec \tau\}$. For functions f and g on spaces X_1 and X_2, let (f, g) be the function on the disjoint sum $X_1 \oplus X_2$ that is f on X_1 and g on X_2; a point of $2^{\mathbb{N}}$ is regarded as a function on $\mathbb{N}$.

The following compacta appear in the classification:

- the *one-point compactification* $A(2^{\mathbb{N}})$ of a discrete set of size continuum, realized as $\{\delta_\sigma : \sigma \in 2^{\mathbb{N}}\} \cup \{0\}$, where δ_σ is the characteristic function of $\{\sigma\}$, and its separable extension $\hat{A}(2^{\mathbb{N}})$, the pointwise closure of $\{v_t : t \in 2^{<\mathbb{N}}\}$, which equals $\{v_t : t \in 2^{<\mathbb{N}}\} \cup A(2^{\mathbb{N}})$;
- the *split Cantor set* $S(2^{\mathbb{N}}) = \{x^+_\sigma, x^-_\sigma : \sigma \in 2^{\mathbb{N}}\}$, the *split interval* $S(I)$, that is, $I \times \{0, 1\}$ for the unit interval $I = [0, 1]$, ordered lexicographically, with the order topology, and the extended split Cantor sets $\hat{S}_+(2^{\mathbb{N}})$ and $\hat{S}_-(2^{\mathbb{N}})$ defined below; the space $\hat{S}_+(2^{\mathbb{N}})$ is homeomorphic to a closed subspace of $S(I)$;
- the *Alexandroff duplicate* $D(2^{\mathbb{N}})$ of the Cantor set and its separable extension $\hat{D}(2^{\mathbb{N}})$, and the extended duplicate $\hat{D}(S(2^{\mathbb{N}}))$ of the split Cantor set, the last two being defined below; the duplicate $D(2^{\mathbb{N}})$ is the set $2^{\mathbb{N}} \times \{0, 1\}$ in which the points $(\sigma, 1)$ are isolated and the basic

neighborhoods of a point $(\sigma,0)$ are the sets $(U\times\{0,1\})\setminus\{(\sigma,1)\}$, for open neighborhoods U of σ in $2^{\mathbb{N}}$.

Among these, the space $S(2^{\mathbb{N}})$ is hereditarily separable but not metrizable, the space $D(2^{\mathbb{N}})$ is first countable but not separable, and the space $A(2^{\mathbb{N}})$ is not first countable (see [ADK08]).

4.3. **The heptachotomy.** A *dyadic subtree* of $2^{<\mathbb{N}}$ is a subset of $2^{<\mathbb{N}}$ that, ordered by extension, is isomorphic to $2^{<\mathbb{N}}$; we write it as $(s_t)_{t\in 2^{<\mathbb{N}}}$, using its canonical enumeration, the unique such isomorphism $t\mapsto s_t$ that also preserves the lexicographic order (see [ADK08, Section 2]). A dyadic subtree is *regular* if each of its levels is contained in a level of $2^{<\mathbb{N}}$. Fix a regular dyadic subtree $\Theta=(s_t)_{t\in 2^{<\mathbb{N}}}$ such that the sequences $s^\frown 0^\infty$, for $s\in\Theta$, are pairwise distinct, and so are the sequences $s^\frown 1^\infty$; a family indexed by the nodes s of Θ is regarded as indexed by $2^{<\mathbb{N}}$ through the canonical enumeration $t\mapsto s_t$.

Definition 4.1. Let X be a Polish space.

(1) Two pointwise bounded families $\{f_j : j\in J\}$ and $\{g_j : j\in J\}$ of real functions, indexed by the same countable set J, are *equivalent* if the map $f_j\mapsto g_j$ extends to a homeomorphism between their pointwise closures.
(2) A family of pairwise distinct functions $\{f_t : t\in 2^{<\mathbb{N}}\}$ that is relatively compact in $B_1(X)$ is *minimal* if it is equivalent to $\{f_{s_t} : t\in 2^{<\mathbb{N}}\}$ for every dyadic subtree (s_t) of $2^{<\mathbb{N}}$.

The seven minimal families are the following, and the pointwise closure of each of them is a separable Rosenthal compactum that contains the family as a discrete set (see [ADK08]):

(1) $D_1=\{\frac{1}{|t|+1}v_t : t\in 2^{<\mathbb{N}}\}$, whose closure $A(2^{<\mathbb{N}})$ is the one-point compactification of a countable discrete set;
(2) $D_2=\{s^\frown 0^\infty : s\in\Theta\}$, a family of points of $2^{\mathbb{N}}\subseteq\mathbb{R}^{\mathbb{N}}$, whose closure is denoted $2^{\leq\mathbb{N}}$;
(3) $D_3=\{x^+_{s^\frown 0^\infty} : s\in\Theta\}$, whose closure is $\hat{S}_+(2^{\mathbb{N}})$;
(4) $D_4=\{x^-_{s^\frown 1^\infty} : s\in\Theta\}$, whose closure is $\hat{S}_-(2^{\mathbb{N}})$;
(5) $D_5=\{v_t : t\in 2^{<\mathbb{N}}\}$, whose closure is $\hat{A}(2^{\mathbb{N}})$;
(6) $D_6=\{(v_t,t^\frown 0^\infty) : t\in 2^{<\mathbb{N}}\}$, whose closure is $\hat{D}(2^{\mathbb{N}})$;
(7) $D_7=\{(v_s,x^+_{s^\frown 0^\infty}) : s\in\Theta\}$, whose closure is $\hat{D}(S(2^{\mathbb{N}}))$.

The closures in (1) and (2) are metrizable, those in (3) and (4) are hereditarily separable but not metrizable, those in (6) and (7) are first countable but not hereditarily separable, and the closure in (5) is not first countable (see [ADK08]).

Fact 4.2 (Heptachotomy; see [ADK08, Theorem 2]**).**

(a) *Up to equivalence, there are exactly seven minimal families, namely $D_1,\dots,D_7$, and they are pairwise non-equivalent.*
(b) *Let X be a Polish space and $\{f_t : t\in 2^{<\mathbb{N}}\}$ a family of pairwise distinct functions that is relatively compact in $B_1(X)$. Then there is a regular dyadic subtree (s_t) of $2^{<\mathbb{N}}$ such that $\{f_{s_t} : t\in 2^{<\mathbb{N}}\}$ is equivalent to one of $D_1,\dots,D_7$.*

4.4. **Todorčević's trichotomy.** Let X be a Polish space, $K\subseteq B_1(X)$ a separable Rosenthal compactum and $\{f_n : n\in\mathbb{N}\}$ a countable dense subset of K. An injection $\iota\colon 2^{<\mathbb{N}}\to\mathbb{N}$ is *increasing* if $\iota(s)<\iota(t)$ whenever s precedes t in the order of $2^{<\mathbb{N}}$ by length and, within each length, by the lexicographic order. A minimal family D_i *canonically embeds* into K (with respect to $\{f_n\}$) if there is an increasing injection $\iota\colon 2^{<\mathbb{N}}\to\mathbb{N}$ such that $\{f_{\iota(t)} : t\in 2^{<\mathbb{N}}\}$ is equivalent to D_i. In that case the closure of D_i is homeomorphic to a closed subspace of K.

Fact 4.3 (Todorčević's trichotomy in canonical form; see [Tod99] and [ADK08, Theorem 3]). *Let X be a Polish space, $K \subseteq B_1(X)$ a separable Rosenthal compactum and $\{f_n : n \in \mathbb{N}\}$ a countable dense subset of K.*

(a) *If K consists of bounded functions, is metrizable and is not separable in the supremum norm, then the family D_2 canonically embeds into K.*
(b) *If K is hereditarily separable but not metrizable, then the family D_3 or the family D_4 canonically embeds into K.*
(c) *If K is first countable but not hereditarily separable, then the family D_6 or the family D_7 canonically embeds into K.*
(d) *If K is not first countable, then the family D_5 canonically embeds into K.*

In particular, a non-metrizable separable Rosenthal compactum contains a copy of $\hat{S}_+(2^{\mathbb{N}})$ or $\hat{S}_-(2^{\mathbb{N}})$, of $\hat{D}(2^{\mathbb{N}})$ or $\hat{D}(S(2^{\mathbb{N}}))$, or of $\hat{A}(2^{\mathbb{N}})$, according as it is hereditarily separable, first countable but not hereditarily separable, or not first countable.




FIGURE 3. Todorčević's trichotomy as a decision tree, with the canonical compacta and the minimal families of the heptachotomy at its leaves (compare [DIMW+26, Section 4.2]). For observables the dashed leaf merges with the metrizable one (see Theorem 6.6 and Corollary 6.8).

4.5. **Levels of NIP.** We now transfer the classification to NIP families, as in our work on deep computations [DIMW+26, Section 4.2].

Definition 4.4. Let Y be a compact metrizable space and $\mathcal{R} \subseteq C_p(Y)$ a pointwise bounded NIP family whose closure $\overline{\mathcal{R}}$ is separable. The family $\mathcal{R}$ is *NIP$_1$*, *NIP$_2$* or *NIP$_3$* if $\overline{\mathcal{R}}$ is, respectively, first countable, hereditarily separable or metrizable.

By the implications among the three invariants, we have NIP$_3$ $\Rightarrow$NIP$_2$ $\Rightarrow$NIP$_1$ $\Rightarrow$NIP.

The top level has a concrete meaning. A set $Z \subseteq Y$ *separates* a set K of functions on Y if any two distinct members of K differ at some point of Z.

Proposition 4.5. *Let Y be a set and $K \subseteq \mathbb{R}^Y$ a set that is compact in the topology of pointwise convergence. Then the space K is metrizable if and only if some countable set $Z \subseteq Y$ separates K.*

Proof. Suppose that a countable set Z separates K. Restriction to Z is a continuous injection of K into $\mathbb{R}^Z$, which is metrizable because Z is countable. A continuous injection of a compact space into a Hausdorff space is a homeomorphism onto its image, so the space K is metrizable.

Conversely, suppose that K is metrizable. Being compact, it is second countable, and so is $K \times K$. Hence the open set $W = \{(f, g) \in K \times K : f \neq g\}$ is Lindelöf. For $y \in Y$, the set $U_y = \{(f, g) \in K \times K : f(y) \neq g(y)\}$ is open, and these sets cover W. If $\{U_y : y \in Z\}$ is a countable subcover, then the set Z separates K. □

Thus a NIP family with separable closure is NIP_3 exactly when countably many points of Y distinguish all functions in its closure. When the functions are responses and the points are probes, this says that countably many probes distinguish all idealized responses. The other two levels have readings of the same kind, for which we need two notions. Let K be a set of real functions on a set Y. A member f of K is *countably determined* if there is a countable set $Z_f \subseteq Y$ such that f is the only member of K that agrees with f on Z_f. A set $E \subseteq K$ is *finitely separated* if, for each $f \in E$, there are a finite set $F_f \subseteq Y$ and a real number $\varepsilon_f > 0$ such that every $g \in E \setminus \{f\}$ satisfies $|g(y) - f(y)| \geq \varepsilon_f$ for some $y \in F_f$.

Proposition 4.6. *Let Y be a compact metrizable space and $\mathcal{R} \subseteq C_p(Y)$ a pointwise bounded NIP family whose closure $K = \overline{\mathcal{R}}$ is separable.*

(a) *The family is NIP_3 if and only if some countable set $Z \subseteq Y$ separates K.*
(b) *The family is NIP_1 if and only if every member of K is countably determined.*
(c) *The family is NIP_2 if and only if no subset of K of cardinality $2^{\aleph_0}$ is finitely separated.*

Proof. Part (a) is Proposition 4.5. (b) If a member f of K is countably determined, then the set $\{f\}$ is the intersection of the countably many open sets $\{g \in K : |g(y) - f(y)| < \frac{1}{m}\}$, for $y \in Z_f$ and $m \geq 1$, and in a compact Hausdorff space a point that is a countable intersection of open sets has a countable base of neighborhoods. Conversely, if $(U_k)_{k \in \mathbb{N}}$ is a countable base at f and each U_k contains a basic neighborhood defined by a finite set $F_k \subseteq Y$, then the set $Z_f = \bigcup_k F_k$ works: a member of K that agrees with f on Z_f lies in every U_k. (c) A set is finitely separated exactly when it is a discrete subspace, and a separable Rosenthal compactum is hereditarily separable if and only if it has no discrete subspace of cardinality $2^{\aleph_0}$, by a theorem of Pol, which also follows from Todorčević's trichotomy (see [ADK08] and [GM22, Theorem 0.4]). □

The three levels thus measure how many points of Y are needed to determine a limit: one countable set for all of them (NIP_3), a countable set for each of them (NIP_1), or no continuum of limits that finitely many points tell apart one by one (NIP_2). When limits are taken as types and points as parameters, the levels say how many instances of a formula determine a type. This partly answers a question, raised in our work on deep computations [DIMW+26], that asks for a non-topological characterization of the levels; a finitary characterization, in the spirit of the fat-shattering dimension (the supremum, at a given gap, of the sizes of the finite sets of points at which the functions realize every pattern of high and low values, around thresholds that may vary from point to point), remains open (see Question (5)).

Theorem 4.7. *Let Y be a compact metrizable space and $\{r_n : n \in \mathbb{N}\} \subseteq C_p(Y)$ a pointwise bounded NIP family. Then exactly one of the following holds:*

(i) *the family is NIP$_3$;*
(ii) *the family is NIP$_2$ but not NIP$_3$, and there is an increasing injection $\iota\colon 2^{<\mathbb{N}} \to \mathbb{N}$ such that $\{r_{\iota(t)} : t \in 2^{<\mathbb{N}}\}$ is equivalent to D_3 or to D_4;*
(iii) *the family is NIP$_1$ but not NIP$_2$, and there is an increasing injection $\iota\colon 2^{<\mathbb{N}} \to \mathbb{N}$ such that $\{r_{\iota(t)} : t \in 2^{<\mathbb{N}}\}$ is equivalent to D_6 or to D_7;*
(iv) *the family is not NIP$_1$, and there is an increasing injection $\iota\colon 2^{<\mathbb{N}} \to \mathbb{N}$ such that $\{r_{\iota(t)} : t \in 2^{<\mathbb{N}}\}$ is equivalent to D_5.*

The following proof merely combines the facts above.

Proof. By Theorem 3.6 and Lemma 3.2, the closure K of the family is a Rosenthal compactum, and the family is a countable dense subset of it. The four cases are mutually exclusive by the definitions and cover all possibilities. In cases (ii)–(iv), the required injection is provided by Fact 4.3(b)–(d), applied to K and the dense subset $\{r_n\}$. □

Thus a non-metrizable NIP closure is always explained by a subfamily of the original functions, arranged along the dyadic tree and equivalent to one of five canonical families. Applied to realized types, the complete descriptions of elements of the structure in terms of the parameters, the classification of Theorem 4.7 yields that of NIP formulas described in the introduction (see [DIMW+26, Section 4.2]). There the realized types are viewed as continuous functions on a compact metrizable space of parameters, and this proviso matters: the space of types of a formula over a countable set of parameters, with no topology on the parameters, is always metrizable, since the countable set of parameters separates the types (see Proposition 4.5).

4.6. **Historical remarks.** Todorčević proved his trichotomy in 1999 [Tod99], using forcing and absoluteness; Argyros, Dodos and Kanellopoulos extracted the canonical form and the heptachotomy from the methods of his paper in 2008 [ADK08], using Ramsey theory for trees [HL66, Mil81, LSV93]. Glasner and Megrelishvili used the trichotomy to build a hierarchy of tame dynamical systems [GM22], and the levels NIP$_1$, NIP$_2$ and NIP$_3$ were introduced for limits of computations in our work on deep computations [DIMW+26].

The hierarchy obtained in this paper differs from these. In the setting of Glasner and Megrelishvili [GM22] and in that of deep computations [DIMW+26], Todorčević's trichotomy is applied to families of functions that carry no linear structure, such as enveloping semigroups (pointwise closures of the transformations of dynamical systems) or sets of computations, and all its classes can occur. By contrast, the idealized responses of an observable (its expectation values for Alice's idealized preparations, as functions of Bob's probes, with Bob's system idealized first) form a convex set (see the proof of Corollary 6.8), and convexity collapses NIP$_2$ onto NIP$_3$ (see Theorem 6.6 and Remark 6.7), whence the hierarchy has three levels. At the level of families indexed by the dyadic tree, all seven minimal families occur (see Theorem 8.5). The difference is a feature of quantum, or more generally of probabilistic, response spaces.

5. The idealized responses of an observable form a compact set of functions

To apply the classification to a bipartite observable, we retain from Bob's operator algebra the effective observables produced by Alice's preparations. The states of the smallest unital

C*-algebra containing them form a compact metrizable probe space, on which each actual preparation has a continuous response. We identify the pointwise closure of these responses with the responses obtained from all states of Alice's operator algebra, with Bob's system idealized first. The compactum to be classified thus records preparations through their responses to the fixed observable.

5.1. **Observables and responses.** We work in the setting of Part I [DI26] and explain its operator theory briefly. Alice's and Bob's systems are described by separable complex Hilbert spaces H_1 and H_2. A *density operator* on a Hilbert space H is a positive trace-class operator of trace 1; density operators are the *actual states* of the system, and we write $\mathfrak{D}(H)$ for the set they form. The trace-class operators, those bounded operators x with $\operatorname{Tr}|x| < \infty$, where $|x| = (x^*x)^{1/2}$, form a Banach space $\mathfrak{T}(H)$ under the trace norm $\|x\|_1 = \operatorname{Tr}|x|$, and we write $B(H)$ for the algebra of bounded operators on H. A bounded self-adjoint operator C on $H_1 \otimes H_2$ is an *observable*, and its expectation when the two systems are prepared independently in actual states $\rho \in \mathfrak{D}(H_1)$ and $\sigma \in \mathfrak{D}(H_2)$ is

$$\varphi_C(\rho, \sigma) = \operatorname{Tr}\big((\rho \otimes \sigma)C\big).$$

Fixing Alice's state ρ produces an observable on Bob's system alone, the *effective observable* $T_C(\rho) = \operatorname{Tr}_1((\rho \otimes I)C)$, obtained by tracing out Alice's system. It is characterized by $\varphi_C(\rho, \sigma) = \operatorname{Tr}(\sigma\, T_C(\rho))$ for all $\sigma \in \mathfrak{D}(H_2)$. The map $\rho \mapsto T_C(\rho)$ extends linearly to a bounded operator $T_C \colon \mathfrak{T}(H_1) \to B(H_2)$ with $\|T_C\| \le \|C\|$, the *slice map*.

Idealized states are described through C*-algebras. A *C*-algebra* is a norm-closed algebra of bounded operators closed under adjoints; we use only unital ones, which contain the identity. A *state* of a unital C*-algebra $\mathfrak{A}$ is a linear functional ω on $\mathfrak{A}$ that is positive, in the sense that $\omega(a^*a) \ge 0$ for all $a \in \mathfrak{A}$, and satisfies $\omega(I) = 1$. The set $\mathcal{S}(\mathfrak{A})$ of states, with the weak* topology (pointwise convergence on $\mathfrak{A}$), is compact; it is metrizable when $\mathfrak{A}$ is norm-separable. Every density operator ρ defines a state $a \mapsto \operatorname{Tr}(\rho a)$; the remaining states of $B(H)$, which no density operator represents, are the *idealized* states, and each of them is the weak* limit of some net of actual states (see [DI26, Lemma 2.2]).

5.2. **Real-valued predicates and bounded confinements.** The operator-algebraic setting determines the real-valued predicates to which we apply the independence property. For expectation predicates, the fixed positive gap $b - a$ in Definition 3.3 separates responses at a prescribed resolution, while affine dependence on the state records probabilistic mixing. We work in Keisler's model theory of real-valued structures [Kei23], in which no metric is given a priori and each formula carries its own bound, rather than in the continuous logic of Ben Yaacov and Usvyatsov [BYU10], in which each sort, that is, each kind of object, carries a metric given in advance and all formulas take values in $[0, 1]$. For the bounded observables of this paper either setting would serve. The distinction becomes relevant when one also considers unbounded quantities: the position Q and the momentum P of a particle satisfy the canonical commutation relation $QP - PQ = i\hbar I$ on a common dense invariant domain, where $\hbar$ is the reduced Planck constant and I is the identity operator, and no pair of bounded operators satisfies this relation on the whole Hilbert space [Win47, Wie49]. For comparisons, unboundedness is harmless, because $[A < B]$ is a projection, unchanged when A and B are replaced by the same strictly increasing function of themselves (see Proposition 7.9). Unboundedness is not harmless for the expectation value of an unbounded quantity, such as the energy $\rho \mapsto \operatorname{Tr}(\rho H)$, where H, the Hamiltonian of the system, is an unbounded self-adjoint operator, Tr is the trace, and ρ ranges over the density operators for which this trace is defined. On its full domain, such an expectation has no bounded range that can be

normalized to $[0,1]$ by an invertible affine change of scale. A nonlinear rescaling changes the affine dependence on mixtures; another approach uses a point at infinity [Ben08]. In Keisler's setting each formula φ takes its values in an interval $[-r_\varphi, r_\varphi]$, whose radius r_φ depends on the formula. An unbounded quantity therefore enters only through its restrictions to confinements, sets of states on each of which its expectation is bounded: for example, states supported in a bounded spectral interval of that quantity, or, for the energy when $H \geq 0$, states satisfying the constraint $\mathrm{Tr}(\rho H) \leq E$ for a fixed finite energy $E \geq 0$, in the units of H. The confinements play the role of the shards of our joint work with Matos-Wiederhold, Salvetti and Tall [DIMW+26], the parts of a state space on which every predicate stays within prescribed bounds. If a confinement has idealized states in its weak* closure, where the weak* topology is convergence of all bounded-observable expectation values, it cannot be relatively compact in the trace norm, the norm that measures how well two states can be distinguished. When Bob's states are restricted to a set that is relatively compact in the trace norm, the expectation of every bounded observable is stable on that restricted domain, because it is Lipschitz in Bob's state, uniformly in Alice's.

5.3. **The space of probes.** The responses of Alice's preparations involve Bob's system only through the effective observables $T_C(\rho)$. The natural probes are therefore the states of the smallest unital C*-algebra containing them.

Definition 5.1. Let C be an observable on $H_1 \otimes H_2$. The *probe algebra* $\mathfrak{A}_C \subseteq B(H_2)$ is the unital C*-algebra generated by $T_C(\mathfrak{T}(H_1))$, and the *probe space* is $Y_C = \mathfrak{S}(\mathfrak{A}_C)$. For $\rho \in \mathfrak{D}(H_1)$, the *response* of ρ is the function $r_\rho \colon Y_C \to \mathbb{R}$ given by $r_\rho(\tau) = \tau(T_C(\rho))$, and the *response family* is $\mathcal{R}_C = \{r_\rho : \rho \in \mathfrak{D}(H_1)\}$.

Lemma 5.2. *Let C be an observable on $H_1 \otimes H_2$.*

(a) *The probe space Y_C is compact and metrizable.*
(b) *Every response is continuous on Y_C and bounded by $\|C\|$, so the response family $\mathcal{R}_C$ is a pointwise bounded subset of $C_p(Y_C)$.*
(c) *The restrictions to $\mathfrak{A}_C$ of the actual states of Bob's system form a dense subset of Y_C, and for $\rho \in \mathfrak{D}(H_1)$ and $\sigma \in \mathfrak{D}(H_2)$ we have $r_\rho(\sigma|_{\mathfrak{A}_C}) = \varphi_C(\rho, \sigma)$.*

Proof. (a) Because H_1 is separable, so is $\mathfrak{T}(H_1)$; its image under the norm-continuous slice map is therefore separable, and so is the C*-algebra $\mathfrak{A}_C$ that this image generates. The compact space Y_C is thus metrizable.

(b) The response r_ρ is evaluation at the element $T_C(\rho)$ of $\mathfrak{A}_C$, and evaluation at a fixed element is weak* continuous. Since $\rho \geq 0$ and C is self-adjoint, the operator $T_C(\rho)$ is self-adjoint, whence r_ρ is real-valued; moreover, $|r_\rho(\tau)| \leq \|T_C(\rho)\| \leq \|C\|$.

(c) By the Hahn–Banach theorem, every state of $\mathfrak{A}_C$ extends to a state of $B(H_2)$ (see [KR83, Theorem 4.3.13]); restriction to $\mathfrak{A}_C$ is thus a weak*-continuous map of the state space of $B(H_2)$ onto Y_C. The states given by density operators are weak* dense in the state space of $B(H_2)$ (see [DI26, Lemma 2.2]), and their restrictions are therefore dense in Y_C. The formula restates the definition of the effective observable. □

By Lemma 5.2(c) and Lemma 3.7, Bob's actual probes alone decide whether $\mathcal{R}_C$ has the independence property. The family $\mathcal{R}_C$ has it exactly when there are actual states $\rho_1, \rho_2, \ldots$ of Alice's system and reals $a < b$ with the following property: for all disjoint finite $E, F \subseteq \mathbb{N}$, some actual state σ of Bob's system satisfies $\varphi_C(\rho_n, \sigma) \leq a$ for $n \in E$ and $\varphi_C(\rho_n, \sigma) \geq b$ for $n \in F$. A single probe then lets Bob read any finite pattern off Alice's sequence.

Definition 5.3. An observable C on $H_1 \otimes H_2$ is *NIP for Alice's responses* if the response family $\mathcal{R}_C$ is NIP.

5.4. **Idealized responses.** To describe the closure of $\mathcal{R}_C$, we use the bitranspose of the slice map, which extends it from actual to idealized states. Every bounded operator $T\colon V \to W$ between Banach spaces has a *transpose* $T^*\colon W^* \to V^*$, given by $T^*(\lambda) = \lambda \circ T$, and a *bitranspose* $T^{**}\colon V^{**} \to W^{**}$, which extends T and is continuous for the weak* topologies. Here we regard the slice map as a map $T_C\colon \mathcal{T}(H_1) \to \mathfrak{A}_C$. The dual of $\mathcal{T}(H_1)$ is $B(H_1)$; the bidual of $\mathcal{T}(H_1)$ is therefore $B(H_1)^*$, which contains all states of $B(H_1)$, idealized ones included.

For an observable C on $H_1 \otimes H_2$ and a state ω of $B(H_1)$, let $r_\omega\colon Y_C \to \mathbb{R}$ be given by $r_\omega(\tau) = \langle T_C^{**}(\omega), \tau\rangle$. We also recall from Part I [DI26] the two products of a state ω of $B(H_1)$ and a state ω' of $B(H_2)$, evaluated at the observable C: if (ρ_i) and (σ_j) are nets of actual states, indexed by directed sets, with weak* limits ω and ω', then $(\omega \otimes_{\mathsf{A}} \omega')(C) = \lim_j \lim_i \varphi_C(\rho_i, \sigma_j)$ and $(\omega \otimes_{\mathsf{B}} \omega')(C) = \lim_i \lim_j \varphi_C(\rho_i, \sigma_j)$, the subscript naming the system that is idealized first.

Lemma 5.4. *Let C be an observable on $H_1 \otimes H_2$. Then*

$$\overline{\mathcal{R}_C} = \{r_\omega : \omega \in \mathcal{S}(B(H_1))\}.$$

Moreover, if ω is a state of $B(H_1)$, $\tau \in Y_C$ and $\tilde\tau$ is a state of $B(H_2)$ extending τ, then $r_\omega(\tau) = (\omega \otimes_{\mathsf{B}} \tilde\tau)(C)$.

Proof. The map $\omega \mapsto T_C^{**}(\omega)$ is weak* continuous, and on bounded subsets of $\mathfrak{A}_C^{**}$ the weak* topology is the topology of pointwise convergence on Y_C, because states span $\mathfrak{A}_C^*$. The map $\omega \mapsto r_\omega$ is therefore a continuous map of the compact space $\mathcal{S}(B(H_1))$ into $\mathbb{R}^{Y_C}$, and its image K is compact. As T_C^{**} extends T_C, the set K contains $\mathcal{R}_C$, and, being closed, it contains $\overline{\mathcal{R}_C}$. Conversely, actual states are dense in $\mathcal{S}(B(H_1))$ (see [DI26, Lemma 2.2]), whence, by continuity, $K \subseteq \overline{\mathcal{R}_C}$. The last formula is [DI26, Lemma 2.4]; the pairing $\langle T_C^{**}(\omega), \tilde\tau\rangle$ depends only on the restriction of $\tilde\tau$ to $\mathfrak{A}_C$. □

The functions r_ω are the *idealized responses*: the responses of Alice's idealized preparations to Bob's probes, idealized probes included, with Bob's system idealized first. On actual probes the order of idealization does not matter (see [DI26, Corollary 2.5]), and, for every state ω of $B(H_1)$ and every $\sigma \in \mathfrak{D}(H_2)$, the idealized response r_ω satisfies $r_\omega(\sigma|_{\mathfrak{A}_C}) = (\omega \otimes_{\mathsf{B}} \sigma)(C) = (\omega \otimes_{\mathsf{A}} \sigma)(C)$.

6. Stability, NIP and the independence property are continuity, Baire class 1 and $\beta\mathbb{N}$

Stability and NIP admit a common description in terms of the regularity of idealized responses. Recall from Part I [DI26] that an observable is stable when its expectation value is independent of the order in which the two systems are idealized. Here an idealized response is the expectation of an idealized preparation of Alice's system as a function of Bob's probe, with Bob's system idealized first. In Theorem 6.1 below, we characterize stability by continuity of every idealized response and NIP by membership of every idealized response in the first Baire class; when NIP fails, a sequence of actual responses has a closure homeomorphic to the space $\beta\mathbb{N}$ of ultrafilters on the natural numbers. Within NIP, the finer classification concerns the topology of the compactum formed by these functions, and the affine dependence on preparations imposes a further restriction through convexity.

To state the result, we recall the notation of Section 5: an observable C is a bounded self-adjoint operator on the tensor product $H_1 \otimes H_2$ of the Hilbert spaces of Alice's and Bob's systems, its probe space Y_C is the compact metrizable state space of its probe algebra (see Definition 5.1 and Lemma 5.2), and its response family $\mathcal{R}_C$ consists of the responses of

Alice's actual preparations; the closure $\overline{\mathcal{R}_C}$ of this family is the set of idealized responses (see Lemma 5.4).

For stability we need one more fact from operator theory, Kadison's function representation: on the state space of a unital C*-algebra $\mathfrak{A}$, the continuous affine real functions are exactly the evaluations $\tau \mapsto \tau(z)$ at self-adjoint elements $z \in \mathfrak{A}$ (see [Kad51]).

Theorem 6.1. *Let C be an observable on $H_1 \otimes H_2$.*

(a) *The observable C is stable if and only if every idealized response is continuous, that is, $\overline{\mathcal{R}_C} \subseteq C_p(Y_C)$.*

(b) *The observable C is NIP for Alice's responses if and only if every idealized response is Baire-1, that is, $\overline{\mathcal{R}_C} \subseteq B_1(Y_C)$. In that case, the set $\overline{\mathcal{R}_C}$ of idealized responses is a Rosenthal compactum.*

(c) *If the observable C is not NIP for Alice's responses, then there is a sequence $(\rho_n)_{n\in\mathbb{N}}$ of actual states of Alice's system such that the closure of $\{r_{\rho_n}\}$ is homeomorphic to $\beta\mathbb{N}$.*

Proof. Part (a) rests on the characterization of stability by weak compactness (see [DI26, Theorem 5.1]), and parts (b) and (c) rest on the dichotomy of Theorem 3.6.

For part (a), recall that a bounded operator between Banach spaces is *weakly compact* if it maps bounded sets to relatively weakly compact sets, and that, by Gantmacher's theorem, this holds exactly when its bitranspose takes values in the original target space. If C is stable, the slice map is weakly compact as a map into $B(H_2)$, and therefore also as a map into the closed subspace $\mathfrak{A}_C$, whose weak topology is the restriction of that of $B(H_2)$. The bitranspose T_C^{**} then maps $B(H_1)^*$ into $\mathfrak{A}_C$, whence every idealized response r_ω is evaluation at an element of $\mathfrak{A}_C$ and is continuous on Y_C.

Conversely, suppose that every idealized response is continuous, and let ω be a state of $B(H_1)$. The element $T_C^{**}(\omega)$ of $\mathfrak{A}_C^{**}$ is self-adjoint, because the slice map sends self-adjoint trace-class operators to self-adjoint operators and ω is real at self-adjoint operators; the idealized response r_ω is therefore a continuous affine real function on Y_C. By Kadison's function representation, there is a self-adjoint $z \in \mathfrak{A}_C$ with $r_\omega(\tau) = \tau(z)$ for all $\tau \in Y_C$. The elements $T_C^{**}(\omega)$ and z of $\mathfrak{A}_C^{**}$ thus agree on all states, and since states span $\mathfrak{A}_C^*$, they are equal; in particular, the element $T_C^{**}(\omega)$ lies in $\mathfrak{A}_C$, and hence in $B(H_2)$. As ω was arbitrary, the observable C is stable (see [DI26, Theorem 5.1]).

By Lemma 5.2, Theorem 3.6 applies to the family $\mathcal{R}_C \subseteq C_p(Y_C)$, and part (b) follows from it together with Lemma 3.2. For part (c), let $(r_n)_{n\in\mathbb{N}}$ be an independent sequence in $\mathcal{R}_C$, with reals $a < b$. By compactness of Y_C (see the proof of Theorem 3.6), for every $I \subseteq \mathbb{N}$ some probe τ satisfies $r_n(\tau) \leq a$ for $n \in I$ and $r_n(\tau) \geq b$ for $n \notin I$. No subsequence of (r_n) can therefore converge pointwise, and Fact 3.4 yields a subsequence whose closure is homeomorphic to $\beta\mathbb{N}$. □

In terms of preparations and probes, Theorem 6.1 describes how the responses of Alice's preparations depend on Bob's probe at each level:

- for a stable observable, the response of each idealized preparation of Alice's system depends continuously on Bob's probe, idealized probes included;
- for an observable that is NIP but unstable, such as the comparison of photon numbers (see Section 7), these responses can be discontinuous, but each is a pointwise limit of a sequence of continuous functions, and together they form a Rosenthal compactum;

- for an observable with the independence property, a sequence of Alice's actual preparations lets Bob read every finite pattern with a single probe, and every such sequence has a subsequence whose closure is a copy of $\beta\mathbb{N}$ (see the proof of Theorem 6.1(c)).

Remark 6.2. In Part I [DI26] we distinguished stability (no infinite ladder) from uniform stability (bounded ladders at every margin, equivalently stability on ultrapowers). A ladder is a pair of sequences of Alice's and of Bob's preparations, of the same length, finite or infinite, whose expectation values are high when Alice's index does not exceed Bob's and low otherwise, or the reverse, and it has margin δ when every high value exceeds every low value by at least δ. The same distinction appears here. For a real number $\delta > 0$, a finite set of functions on a set Y is *δ-shattered* if there are reals $a < b$ with $b - a \geq \delta$ such that, for any two disjoint subsets E and F of it, some point $y \in Y$ satisfies $r(y) \leq a$ for $r \in E$ and $r(y) \geq b$ for $r \in F$. The notion of NIP in Definition 3.3 forbids an infinite independent sequence, whereas *uniform NIP* bounds, for each $\delta > 0$, the size of the δ-shattered subsets of the family; for a uniformly bounded family, such as a family of responses, this amounts, by pigeonholing the thresholds and by the duality argument of Assouad in Remark 3.8, to the finiteness of the fat-shattering dimension for every gap. Uniform NIP implies NIP. Comparisons $[A < B]$, the projections that represent the event "Bob's value of B exceeds Alice's value of A" for quantities A of Alice's system and B of Bob's, are uniformly NIP, with bounds independent of the Hilbert spaces (see Corollary 7.4), and they are unstable when both quantities have pure point spectra unbounded above, by the ladder argument of Section 7. By Theorem 6.1, their idealized responses form a Rosenthal compactum of Baire-1 functions, not all continuous. As in Part I, the uniform notion is the one that survives ultrapowers: a uniformly bounded family is uniformly NIP if and only if its extension to the ultrapower by every nonprincipal ultrafilter on $\mathbb{N}$ is NIP. Recall that, for an ultrafilter $\mathcal{U}$ on $\mathbb{N}$, every bounded sequence $(x_k)_{k\in\mathbb{N}}$ of complex numbers has a unique ultralimit $\lim_{k,\mathcal{U}} x_k$, the complex number x such that $\{k : |x_k - x| < \varepsilon\} \in \mathcal{U}$ for every $\varepsilon > 0$. Here the extension of a family of functions on Y to the ultrapower by $\mathcal{U}$ consists of the functions that send a sequence $(y_k)_{k\in\mathbb{N}}$ of points of Y to the ultralimit $\lim_{k,\mathcal{U}} r_k(y_k)$, one function for each sequence $(r_k)_{k\in\mathbb{N}}$ of members of the family; each value is a single ultralimit along the common index k, not the value of a limit function at a limit point. By Łoś's theorem, which says that the value of a formula at elements of the ultrapower, given by sequences, is the ultralimit of its values at the terms of the sequences, a δ-shattered set of size m in the ultrapower yields, for every $\delta' < \delta$, a δ'-shattered set of size m in the original family. Hence, if the sizes of the δ-shattered subsets of the original family are bounded for every $\delta > 0$, so are those of its extension, which then has no infinite independent sequence; conversely, if they are unbounded for some $\delta > 0$, then in the ultrapower by any nonprincipal ultrafilter on $\mathbb{N}$, which is countably incomplete, the finite shattered sets combine, by $\aleph_1$-saturation (every countable set of conditions, each finite subset of which is realized in the ultrapower, is realized in it), into an infinite independent sequence.

6.1. **The finer classification of NIP observables.** For NIP observables, the classification of Section 4 applies to the idealized responses. We first check that they form a separable compactum.

Lemma 6.3. *Let C be an observable on $H_1 \otimes H_2$ and $(\rho_n)_{n\in\mathbb{N}}$ a sequence of density operators that is dense in $\mathfrak{D}(H_1)$ for the trace norm. Then the responses r_{ρ_n} are dense in $\overline{\mathcal{R}_C}$.*

Proof. For density operators ρ and ρ', we have $\sup_{\tau\in Y_C} |r_\rho(\tau) - r_{\rho'}(\tau)| \leq \|T_C(\rho - \rho')\| \leq \|C\|\,\|\rho - \rho'\|_1$. Every response is therefore a uniform limit, and hence a pointwise limit, of a

sequence of responses r_{ρ_n}; the closure of $\{r_{\rho_n}\}$ thus contains $\mathcal{R}_C$, and with it $\overline{\mathcal{R}_C}$. Sequences such as (ρ_n) exist, because $\mathcal{T}(H_1)$ is separable. □

Definition 6.4. An observable C that is NIP for Alice's responses is *NIP*$_1$, *NIP*$_2$ or *NIP*$_3$ if the response family $\mathcal{R}_C$ is, respectively, NIP$_1$, NIP$_2$ or NIP$_3$ (see Definition 4.4).

Proposition 6.5. *Every stable observable is NIP*$_3$*: its idealized responses form a compact metrizable space.*

Proof. By Theorem 6.1(a), every idealized response is continuous on the compact metrizable space Y_C, hence Baire-1, and the observable C is therefore NIP for Alice's responses by Theorem 6.1(b); the set of idealized responses is separable by Lemma 6.3. A countable dense set of probes separates continuous functions, and Proposition 4.5 shows that the idealized responses form a metrizable space. □

6.2. **Mixtures force metrizability.** The idealized responses of an observable form a convex set, because responses depend affinely on Alice's state (see the proof of Corollary 6.8). In Theorem 6.6 below, we show that for separable convex Rosenthal compacta hereditary separability implies metrizability. This is a special case of a known result on locally connected Rosenthal compacta (see Remark 6.7); we give a direct proof. The mechanism is visible in the split interval $S(I)$ of Section 4, which is homeomorphic to the set of indicator functions $\mathbf{1}_{(a,1]}$ and $\mathbf{1}_{[a,1]}$ on $[0,1]$, for $0 \le a \le 1$. For each such a, the two indicators form a split pair, two functions that differ only at a, and a convex set of functions that contains both of them contains their midpoint, which takes the value $\frac{1}{2}$ at a. These midpoints form an uncountable discrete set: the midpoint for a point $b \neq a$ takes the value 0 or 1 at a.

Theorem 6.6 (see [Tod99, Theorems 5 and 7]**,** [Gru88, Lemma 2.4] **and Remark 6.7).** *Let Y be a set, and let $K \subseteq \mathbb{R}^Y$ be a separable compact convex set which, with the topology of pointwise convergence, is a Rosenthal compactum. If K is hereditarily separable, then K is metrizable.*

Proof. The idea is the one described before the theorem: if K were hereditarily separable but not metrizable, it would contain a copy of a split interval, and the midpoints of the split pairs of that copy would form a subspace of K that is not separable.

Suppose, then, that K is hereditarily separable but not metrizable. By Fact 4.3(b), there is a homeomorphism h from $\hat{S}_+(2^{\mathbb{N}})$ or $\hat{S}_-(2^{\mathbb{N}})$, the closures of the minimal families D_3 and D_4 of Section 4, respectively, onto a subspace of K. We treat the case $\hat{S} = \hat{S}_+(2^{\mathbb{N}})$. The other case is symmetric: the map c that exchanges the digits 0 and 1 reverses the lexicographic order $\prec$ of $2^{\mathbb{N}}$, and the map $f \mapsto 1 - f \circ c$ carries D_4 to a family of the form of D_3 for the tree $c(\Theta)$, the image under c of the regular dyadic subtree Θ fixed in Section 4. Write $Q = \{s^\frown 0^\infty : s \in \Theta\}$ for the countable set of points indexing D_3, and Σ for the closure of Q in $2^{\mathbb{N}}$. Recall that, for $\sigma \in 2^{\mathbb{N}}$, the functions x^+_σ and x^-_σ are the characteristic functions of $\{\tau \in 2^{\mathbb{N}} : \sigma \preceq \tau\}$ and $\{\tau \in 2^{\mathbb{N}} : \sigma \prec \tau\}$; thus D_3 consists of the functions x^+_q for $q \in Q$.

We first record the structure of $\hat{S}$. If $(\sigma_i)_{i\in\mathbb{N}}$ is a strictly increasing sequence in $2^{\mathbb{N}}$ converging to σ, then $x^\pm_{\sigma_i} \to x^+_\sigma$ pointwise; if it is strictly decreasing, then $x^\pm_{\sigma_i} \to x^-_\sigma$. Indeed, for $\tau \in 2^{\mathbb{N}}$, the value of $x^\pm_{\sigma_i}$ at τ is eventually 1 if $\tau \succ \sigma$, eventually 0 if $\tau \prec \sigma$, and at $\tau = \sigma$ it is eventually 1 in the increasing case and 0 in the decreasing case. Since $\hat{S}$ is Fréchet–Urysohn (every point in the closure of a subset is the limit of a sequence from the subset; see [BFT78, Theorem 3F]) and every sequence of distinct points of $2^{\mathbb{N}}$ has a strictly monotone subsequence, every point of $\hat{S} \setminus D_3$ has the form x^+_η or x^-_η with $\eta \in \Sigma$; and if $x^{e_i}_{\sigma_i} \in \hat{S}$ for a

strictly monotone sequence (σ_i) and signs $e_i \in \{+,-\}$, then the limit computed above lies in $\hat{S}$, because $\hat{S}$ is closed. Note also that $x^+_\sigma = x^-_\eta$ exactly when σ is the immediate successor of η in $\prec$, and that two of these functions with the same sign are equal only when their indices are.

We use the following fact repeatedly: if $A \subseteq 2^{\mathbb{N}}$ is uncountable, then all but countably many points of A are limits of strictly increasing and of strictly decreasing sequences from A. Indeed, if $\sigma \in A$ is neither the least element 0^∞ of $2^{\mathbb{N}}$ nor a limit of a strictly increasing sequence from A, there is $\tau_\sigma \prec \sigma$ with no point of A strictly between them, and the intervals $(\tau_\sigma, \sigma]$ for distinct such σ are disjoint; each of them either contains a basic clopen set or makes τ_σ the immediate predecessor of σ. The first alternative occurs for only countably many σ, because the intervals are disjoint and the basic clopen sets are countable, and so does the second, because only countably many points of $2^{\mathbb{N}}$ have an immediate predecessor. The decreasing case is the same, with the greatest element 1^∞ in place of 0^∞. Let P be the set of $\sigma \in \Sigma$ that are limits of strictly increasing and of strictly decreasing sequences from Q. The set Σ is uncountable, because distinct branches of the dyadic tree Θ have distinct limits. By the same argument applied to Q, all but countably many points of Σ belong to P (an open interval that misses Q also misses Σ); hence P is uncountable. For $\sigma \in P$, both x^+_σ and x^-_σ belong to $\hat{S}$, they differ at σ, and neither equals $x^\pm_\eta$ for $\eta \neq \sigma$, because σ has neither an immediate predecessor nor an immediate successor in $\prec$.

The split pairs indexed by P now supply the midpoints. For $\sigma \in P$, put $F^\pm_\sigma = h(x^\pm_\sigma)$ and $M_\sigma = \frac{1}{2}(F^+_\sigma + F^-_\sigma)$, which lies in K by convexity. Averaging is continuous for the pointwise topology, and so is h; consequently, if (σ_i) is a strictly increasing (respectively, decreasing) sequence in P converging to σ, then $M_{\sigma_i} \to h(x^+_\sigma)$ (respectively, $h(x^-_\sigma)$). The proof that the midpoints form a non-separable subspace takes three steps.

Step 1: each value is taken by at most countably many midpoints. Let $m \in K$ and suppose that $A = \{\sigma \in P : M_\sigma = m\}$ is uncountable. Choose $\sigma \in A$ that is a limit of strictly increasing and of strictly decreasing sequences from A. Then $h(x^+_\sigma) = m = h(x^-_\sigma)$, and since h is injective, $x^+_\sigma = x^-_\sigma$, a contradiction.

Step 2: only countably many midpoints lie on the copy of $\hat{S}$. Let $E = \{\sigma \in P : M_\sigma \in h(\hat{S})\}$. By Step 1 and the countability of D_3, all but countably many $\sigma \in E$ satisfy $M_\sigma = h(x^{e(\sigma)}_{\eta(\sigma)})$ for some sign $e(\sigma)$ and some $\eta(\sigma) \in \Sigma$, and $\eta(\sigma) \neq \sigma$, since a midpoint of two distinct points differs from both. Suppose that uncountably many such σ exist. Fix a metric d on $2^{\mathbb{N}}$ that induces its topology; then for some positive integer n the set E_n of these σ with $d(\sigma, \eta(\sigma)) \geq 1/n$ is uncountable. Choose $\sigma \in E_n$ and a strictly monotone sequence (σ_i) in E_n converging to σ, say increasing. Passing to a subsequence, we may assume that the signs $e(\sigma_i)$ are constant and that the points $\eta(\sigma_i)$ are either constant or strictly monotone, converging to some η^*. Then the midpoints M_{σ_i} converge both to $h(x^+_\sigma)$ and to $h(x^\pm_{\eta^*})$. Since h is injective, $x^+_\sigma = x^\pm_{\eta^*}$, and since $\sigma \in P$ has no immediate predecessor, $\eta^* = \sigma$. Thus $d(\sigma_i, \eta(\sigma_i)) \to 0$, contrary to $\sigma_i \in E_n$. Consequently, the set E is countable.

Step 3: the midpoints form a non-separable subspace. Let $P' = P \setminus E$, an uncountable set, and let $L = \{M_\sigma : \sigma \in P'\}$. Let $Z \subseteq L$ be countable. By Step 1, there is $\sigma \in P'$ with $M_\sigma \notin Z$. If M_σ were in the closure of Z, then, since Rosenthal compacta are Fréchet–Urysohn (see [BFT78, Theorem 3F]), there would be a sequence (σ_i) in P' with $M_{\sigma_i} \in Z$ and $M_{\sigma_i} \to M_\sigma$. As $M_\sigma \notin Z$, no point occurs infinitely often in (σ_i), so a subsequence is strictly monotone and converges to some $\eta \in \Sigma$, and then the midpoints M_{σ_i} converge to $h(x^+_\eta)$ or $h(x^-_\eta)$. Thus $M_\sigma \in h(\hat{S})$, contrary to $\sigma \notin E$. The subspace L is therefore not separable, and K is not hereditarily separable, a contradiction. □

Remark 6.7. The proof uses convexity only to know that midpoints of points of K lie in K. The same midpoints settle a question about the split interval $S(I) = [0,1] \times \{0,1\}$, ordered lexicographically, with the order topology. In a remark following his Proposition 10 [God80], Godefroy states that the space $P(S(I))$ of Borel probability measures on $S(I)$, with the weak* topology, is hereditarily separable. This is not the case: for $0 < a < 1$, let $a^- = (a, 0)$ and $a^+ = (a, 1)$, and let m_a be the midpoint $\frac{1}{2}(\delta_{a^-} + \delta_{a^+})$ of the point masses at a^- and a^+. The set $\{s \in S(I) : s \leq a^-\}$ is clopen, so its indicator f_a is continuous, and $m_a(f_a) = \frac{1}{2}$, while $m_b(f_a) \in \{0, 1\}$ for $b \neq a$. Hence the set $\{m_a : 0 < a < 1\}$ is an uncountable discrete subspace of $P(S(I))$, exactly as in the proof of Theorem 6.6. Todorčević has pointed out to us that Theorem 6.6 is known, and that it holds in the larger class of locally connected Rosenthal compacta, which contains the convex ones: it follows from his Theorems 5 and 7 [Tod99] and from an argument of Gruenhage [Gru88, Lemma 2.4]. The proof above is direct and uses only the midpoints, the same midpoints that refute Godefroy's remark. Together with the examples of Section 7, which show that the other three classes occur among compacta of idealized responses, Theorem 6.6 answers, for convex compacta, the question of which classes of Todorčević's trichotomy occur, in the spirit of the Glasner–Megrelishvili hierarchy of tame systems [GM22].

Corollary 6.8. *Every observable that is NIP$_2$ is NIP$_3$.*

Proof. Let C be an observable on $H_1 \otimes H_2$ that is NIP$_2$. We show that the set $\overline{\mathcal{R}_C}$ of idealized responses satisfies the hypotheses of Theorem 6.6. Since the slice map and evaluation at a probe are linear and $\mathfrak{D}(H_1)$ is convex, the identity $r_{t\rho+(1-t)\rho'} = t\, r_\rho + (1-t)\, r_{\rho'}$, for $\rho, \rho' \in \mathfrak{D}(H_1)$ and $0 \leq t \leq 1$, shows that the family $\mathcal{R}_C$ is convex. For fixed t, the map $(f, g) \mapsto tf + (1-t)g$ is continuous on $\mathbb{R}^{Y_C} \times \mathbb{R}^{Y_C}$ for the pointwise topology and sends $\mathcal{R}_C \times \mathcal{R}_C$ into $\mathcal{R}_C$; it therefore sends $\overline{\mathcal{R}_C} \times \overline{\mathcal{R}_C}$, the closure of $\mathcal{R}_C \times \mathcal{R}_C$, into $\overline{\mathcal{R}_C}$, which is thus convex. By Lemma 6.3 and Theorem 6.1(b), the set $\overline{\mathcal{R}_C}$ is a separable Rosenthal compactum, and it is hereditarily separable because C is NIP$_2$. By Theorem 6.6, it is metrizable, that is, the observable C is NIP$_3$. □

For observables, then, at most three of the four classes of Todorčević's trichotomy can occur; in Section 7 we show that all three do. For families of functions all four occur: the minimal family D_3 of Section 4, formed by the characteristic functions of the sets $\{\tau \in 2^{\mathbb{N}} : \tau \succeq s^\frown 0^\infty\}$ for s in the regular dyadic subtree Θ, consists of continuous functions on the Cantor set, since each of these sets is clopen, and its closure $\hat{S}_+(2^{\mathbb{N}})$ is hereditarily separable but not metrizable, so the family D_3 is NIP$_2$ but not NIP$_3$. This example comes from the lexicographic order of the Cantor set, which is isomorphic to the order that the real line induces on the middle-thirds Cantor set; the members of D_3 are realized types of this order, and they take only the values 0 and 1. Expectations bring in mixtures, and the midpoints of the split pairs turn the split interval into a space with an uncountable discrete subset; for the order of positions, the resulting space is the space of nondecreasing functions from $[0,1]$ to $[0,1]$ that take the value 0 at 0 and 1 at 1, a closed subspace of the Helly space of all nondecreasing functions from $[0,1]$ to $[0,1]$ (see Section 7). This is why, for averages over Alice's mixed preparations, hereditary separability forces metrizability (see Theorem 6.6).

Theorem 6.9. *Let C be an observable on $H_1 \otimes H_2$ that is NIP for Alice's responses, and let $(\rho_n)_{n\in\mathbb{N}}$ be a trace-norm dense sequence in $\mathfrak{D}(H_1)$. Then, with increasing injections, equivalent families and the minimal families D_5, D_6 and D_7 as in Section 4, exactly one of the following holds:*

(i) *the observable C is NIP$_3$;*
(ii) *the observable C is NIP$_1$ but not NIP$_2$, and there is an increasing injection $\iota\colon 2^{<\mathbb{N}} \to \mathbb{N}$ such that the responses $\{r_{\rho_{\iota(t)}} : t \in 2^{<\mathbb{N}}\}$ are equivalent to D_6 or to D_7;*
(iii) *the observable C is not NIP$_1$, and there is an increasing injection $\iota\colon 2^{<\mathbb{N}} \to \mathbb{N}$ such that the responses $\{r_{\rho_{\iota(t)}} : t \in 2^{<\mathbb{N}}\}$ are equivalent to D_5.*

Proof. By Lemma 5.2, the family $\{r_{\rho_n}\}$ is a pointwise bounded family of continuous functions on the compact metrizable space Y_C, and it is NIP because it is contained in $\mathcal{R}_C$. By Lemma 6.3, its closure is $\overline{\mathcal{R}_C}$. Hence Theorem 4.7 applies, and its case (ii) is excluded by Corollary 6.8. □

Physically, when the idealized responses of a NIP observable do not form a metrizable space, the reason is always visible in Alice's actual preparations: a family of them, arranged along the dyadic tree, has idealized limits that reproduce one of three canonical compacta, the closures of the minimal families D_5, D_6 and D_7. In the next section we locate the comparisons $[A < B]$ in this classification.

6.3. **Historical remarks.** Kadison proved the function representation used in Theorem 6.1 in 1951 [Kad51]. Grothendieck's theorem [Gro52], on which the characterizations of stability in Part I [DI26] rest, lies behind the characterization of stability by the continuity of the idealized responses in Theorem 6.1(a); the NIP version of this characterization in Theorem 6.1(b), with Baire-1 functions in place of continuous ones, follows our work on deep computations [DIMW+26].

7. Three physical orders realize the three classes of Todorčević's trichotomy that occur for observables

Each of the three classes of Todorčević's trichotomy in Theorem 6.9, we show here, contains a physically natural order, and all three orders are unstable and NIP. The comparison of photon numbers has a metrizable compactum of idealized responses and is thus NIP$_3$ (see Theorem 7.8). The comparison of the positions of two particles, in a box or on a line, or of the quadratures of two modes of light (a quadrature is one of the two field amplitudes of a mode, which behave like a position and a momentum), has as its compactum of idealized responses a copy of the space of nondecreasing functions from $[0,1]$ to $[0,1]$ that take the value 0 at 0 and 1 at 1, a closed subspace of the Helly space; this space is first countable but not hereditarily separable, and the comparison is therefore NIP$_1$ but not NIP$_2$ (see Theorem 7.8 and Proposition 7.9). The causal order of a spacetime with one dimension of space, "Bob's detection lies in the causal future of Alice's event", is NIP but not NIP$_1$: since a light cone can shrink to a point, its compactum of idealized responses is not first countable (see Theorem 7.10). The comparison of two phases, "Bob's angle is ahead of Alice's by less than half a turn", a cyclic relation rather than an order, falls in the same class, because on a circle the jumps at the two ends of a half-turn can cancel (see Theorem 7.12).

7.1. **Relations between classical quantities.** All three orders relate a classical quantity measured by Alice to one measured by Bob, and their responses can be computed from classical probability distributions. Let (X_1, m_1) and (X_2, m_2) be σ-finite measure spaces such that the Hilbert spaces $H_1 = L^2(X_1, m_1)$ and $H_2 = L^2(X_2, m_2)$ are separable, fix an integer $p \geq 1$, let $\alpha\colon X_1 \to \mathbb{R}^p$ and $\beta\colon X_2 \to \mathbb{R}$ be measurable, and let $R \subseteq \mathbb{R}^p \times \mathbb{R}$ be a Borel set. The *relation observable* C_R is multiplication on $L^2(X_1 \times X_2, m_1 \otimes m_2) = H_1 \otimes H_2$ by the indicator of $\{(s,t) : (\alpha(s), \beta(t)) \in R\}$; it represents the event "Alice's value and Bob's value are related by R". In this section, the letters ρ and σ denote actual states of Alice's and of Bob's systems, that is, density operators on H_1 and on H_2. For a state ρ of Alice's system, the *distribution* of Alice's value in ρ is the Borel probability measure on $\mathbb{R}^p$ given by $\mu_\rho(U) = \operatorname{Tr}\big(\rho\, M_{\mathbf{1}_{\alpha^{-1}(U)}}\big)$ for Borel sets $U \subseteq \mathbb{R}^p$, where M_f denotes multiplication by f, and the distribution μ_σ of Bob's value in a state σ of Bob's system is defined in the same way. For $y \in \mathbb{R}$, let $R^y = \{x \in \mathbb{R}^p : (x, y) \in R\}$.

By the spectral theorem, every self-adjoint operator on a separable Hilbert space is unitarily equivalent to multiplication by a real measurable function on a space $L^2(X, m)$ of this kind. Thus, for self-adjoint operators A on H_1 and B on H_2, realized as multiplication by α and by β, the *comparison* $[A < B]$ is the relation observable C_R for $R = \{(x, y) : x < y\}$; it is the projection $(E_A \otimes E_B)(R)$, where E_A and E_B are the spectral measures, so it does not depend on the realization. When A and B are bounded, the operator $I \otimes B - A \otimes I$ is multiplication by $\beta(t) - \alpha(s)$, and $[A < B]$ is its spectral projection for the interval $(0, \infty)$, or equivalently the spectral projection of $A \otimes I - I \otimes B$ for the interval $(-\infty, 0)$.

Lemma 7.1. *Let C_R be a relation observable. For states ρ of Alice's system and σ of Bob's, the expectation value of C_R when the two systems are prepared independently in these states is*

$$\varphi_{C_R}(\rho, \sigma) = (\mu_\rho \otimes \mu_\sigma)(R) = \int g_\rho \, d\mu_\sigma, \qquad g_\rho(y) = \mu_\rho(R^y),$$

and the effective observable $T_{C_R}(\rho)$, the observable on Bob's system obtained by fixing Alice's state ρ, is multiplication by $g_\rho \circ \beta$.

Proof. For unit vectors $\psi \in H_1$ and $\chi \in H_2$, with distributions μ_ψ and μ_χ in the corresponding pure states, Tonelli's theorem and a change of variables give

$$\langle \psi \otimes \chi, C_R(\psi \otimes \chi)\rangle = \iint \mathbf{1}_R\big(\alpha(s), \beta(t)\big)\, |\psi(s)|^2 |\chi(t)|^2 \, dm_1(s)\, dm_2(t) = (\mu_\psi \otimes \mu_\chi)(R).$$

A density operator is a countable convex combination of pure states, $\rho = \sum_j \lambda_j |\psi_j\rangle\langle\psi_j|$, the series converging in trace norm; here j ranges over a countable set, the weights $\lambda_j \geq 0$ have sum 1, and $|\psi_j\rangle\langle\psi_j|$ is the orthogonal projection onto the line spanned by a unit vector $\psi_j \in H_1$. Accordingly, the distribution of Alice's value in ρ is $\mu_\rho = \sum_j \lambda_j \mu_{\psi_j}$. Both sides of the first identity are affine in each variable and preserve such combinations, so the first identity holds for all states; the second is Tonelli's theorem, and g_ρ is measurable by the same theorem. Finally, for every unit vector $\chi \in H_2$,

$$\langle \chi, T_{C_R}(\rho)\chi\rangle = \varphi_{C_R}(\rho, |\chi\rangle\langle\chi|) = \int g_\rho\big(\beta(t)\big) |\chi(t)|^2 \, dm_2(t),$$

and a bounded operator on a complex Hilbert space is determined by these values. □

In Lemma 7.2 and Proposition 7.3 below, we show that when the functions g_ρ are monotone, or differences of monotone functions, the responses cannot shatter large sets. Recall from Remark 6.2 that a finite set of responses is δ-shattered if every pattern on it is realized with reals $a < b$ such that $b - a \geq \delta$.

Lemma 7.2. *Let $\delta > 0$ be given, and put $m = \lceil 9/\delta \rceil$. Let n be a positive integer, let $h_1, \dots, h_n \colon \mathbb{R} \to \mathbb{R}$ be functions of the form $h_i = u_i - v_i$, where $u_i, v_i \colon \mathbb{R} \to [0,1]$ are nondecreasing, and let $a < b$ be reals with $b - a \geq \delta$. Suppose that for every $S \subseteq \{1, \dots, n\}$ there is a Borel probability measure ν_S on $\mathbb{R}$ with $\int h_i \, d\nu_S \geq b$ for $i \in S$ and $\int h_i \, d\nu_S \leq a$ for $i \notin S$. Then $2^n \leq (2nm+1)^m$. In particular, $n \leq N(\delta)$, where $N(\delta)$, the largest integer n satisfying this inequality, is finite and depends only on δ.*

Proof. We first approximate a measure by m equally weighted points. Let ν be a Borel probability measure on $\mathbb{R}$, with distribution function $F(t) = \nu((-\infty, t])$. For $k = 1, \dots, m$, let $y_k = \min\{t : F(t) \geq (k - \frac{1}{2})/m\}$; the minimum exists because F is nondecreasing and right continuous, with limits 0 and 1 at $-\infty$ and $+\infty$, and $0 < (k - \frac{1}{2})/m < 1$. Let ν' be the average of the m point masses at the points $y_1, \dots, y_m$. Since $y_k \leq t$ exactly when $F(t) \geq (k - \frac{1}{2})/m$, we have $\nu'((-\infty, t]) = \lfloor mF(t) + \frac{1}{2} \rfloor / m$, whence $|\nu((-\infty, t]) - \nu'((-\infty, t])| \leq 1/m$ for every t; by letting s increase to t, we obtain the same bound for the half-lines $(-\infty, t)$. Hence $|\nu(J) - \nu'(J)| \leq 1/m$ for every set J of the form $\emptyset$, $\mathbb{R}$, $[t, \infty)$ or (t, ∞). For a nondecreasing $u \colon \mathbb{R} \to [0,1]$, every set $\{u > s\}$ has this form, and $\int u \, d\nu = \int_0^1 \nu(\{u > s\}) \, ds$; therefore

$$\left| \int u \, d\nu - \int u \, d\nu' \right| \leq \frac{1}{m}. \tag{1}$$

Next we discretize the functions. Let $\tilde{u}_i = \lfloor m u_i \rfloor / m$ and $\tilde{v}_i = \lfloor m v_i \rfloor / m$. These functions are nondecreasing, take values in $\{0, \frac{1}{m}, \dots, 1\}$, and differ from u_i and v_i by at most $\frac{1}{m}$. Each of the $2n$ functions $\tilde{u}_i, \tilde{v}_i$ changes value at most m times along $\mathbb{R}$, so the map $Q(y) = (\tilde{u}_1(y), \dots, \tilde{u}_n(y), \tilde{v}_1(y), \dots, \tilde{v}_n(y))$ takes at most $2nm + 1$ values.

For each S, let $y_1^S, \dots, y_m^S$ be the points constructed from ν_S as above, and let $w(S) = (Q(y_1^S), \dots, Q(y_m^S))$. By (1), applied to u_i and to v_i, and by the bound on the discretization,

$$\left| \int h_i \, d\nu_S - \frac{1}{m} \sum_{k=1}^{m} \big(\tilde{u}_i(y_k^S) - \tilde{v}_i(y_k^S) \big) \right| \leq \frac{4}{m} \leq \frac{4\delta}{9},$$

and the average on the left is determined by $w(S)$. If $S \neq S'$, choose i in one of the two sets but not the other, say $i \in S \setminus S'$. Then $\int h_i \, d\nu_S - \int h_i \, d\nu_{S'} \geq \delta > \frac{8\delta}{9}$, so the two averages differ, and $w(S) \neq w(S')$. Thus the map $S \mapsto w(S)$ is injective, and $2^n \leq (2nm+1)^m$. ☐

Proposition 7.3. *Let C_R be a relation observable and $c \geq 1$ a constant such that, for every state ρ of Alice's system, $g_\rho = c\,(u_\rho - v_\rho)$ for two nondecreasing functions $u_\rho, v_\rho \colon \mathbb{R} \to [0,1]$. Then, for every $\delta > 0$, no set of more than $N(\delta/(2c))$ responses of Alice's states is δ-shattered by points of the probe space. Hence the observable C_R is uniformly NIP, with bounds that do not depend on the Hilbert spaces, and in particular it is NIP.*

Proof. Suppose that the responses $r_{\rho_1}, \dots, r_{\rho_n}$ are δ-shattered with reals $a < b$, the pattern S being realized by a probe τ_S in the probe space Y_{C_R} of C_R. Let $a' = a + \delta/4$ and $b' = b - \delta/4$. The set of probes τ with $r_{\rho_i}(\tau) > b'$ for $i \in S$ and $r_{\rho_i}(\tau) < a'$ for $i \notin S$ is open, by Lemma 5.2(b), and contains τ_S, so, by Lemma 5.2(c), it contains the restriction of an actual state σ_S of Bob's system. By Lemma 7.1, the value of r_{ρ_i} there is $\int g_{\rho_i} \, d\mu_{\sigma_S}$, and Lemma 7.2, applied with $h_i = g_{\rho_i}/c = u_{\rho_i} - v_{\rho_i}$, $\nu_S = \mu_{\sigma_S}$ and the reals $a'/c < b'/c$, yields $n \leq N(\delta/(2c))$. An infinite independent sequence with reals $a < b$ would contain $(b-a)$-shattered sets of every finite size, so the observable C_R is NIP. ☐

Corollary 7.4. *For all self-adjoint operators A on H_1 and B on H_2, the comparison $[A < B]$ is uniformly NIP, with bounds that depend only on the margin, a lower bound for the gap $b - a$ in Definition 3.3, and in particular it is NIP.*

Proof. Here $R^y = (-\infty, y)$, so the function $g_\rho(y) = \mu_\rho((-\infty, y))$ is nondecreasing, with values in $[0, 1]$. Proposition 7.3 therefore applies, with $c = 1$, $u_\rho = g_\rho$ and $v_\rho = 0$. □

Remark 7.5 (Commuting observables and the general framework). The general classification and its commuting realizations have complementary roles. The definitions of Section 5 and the results of Theorems 6.1 and 6.9 and Corollary 6.8 apply to every bounded self-adjoint bipartite observable; for the swap test, for example, the probe algebra contains all compact operators. For every such observable the full compactum of idealized responses is convex; when the observable is NIP, hereditary separability therefore implies metrizability. The orders considered here have commutative probe algebras and realize all three resulting classes; the label-controlled observables of Section 8, in which Alice's label selects a test on Bob's system, also have commutative probe algebras. For these realizations, integration against classical probability distributions makes the responses explicit. For the relation observables satisfying its monotonicity hypothesis, Proposition 7.3 supplies bounds independent of the Hilbert spaces. In Section 9 we pass to parameterized magnetization families, including those of the Ising model in a transverse field, whose Hamiltonian has noncommuting terms; their classification follows from the same analysis of pointwise closures (see Theorem 9.6). The question of which further phenomena arise from noncommutative probe algebras is stated in Question (3).

7.2. **Order observables.** We now fix two model comparisons, one of discrete and one of continuous quantities, whose values lie in a compact set $\Xi \subseteq [0, 1]$ containing 0 and 1. The two cases are the following:

- *discrete*: the set Ξ is $\{1 - \frac{1}{l+1} : l \in \mathbb{N}\} \cup \{1\}$, the Hilbert spaces are $H_1 = H_2 = \ell^2(\mathbb{N})$, and the operator $A = B$ is the diagonal operator with eigenvalue $1 - \frac{1}{l+1}$ on the l-th basis vector; this operator is a bounded relabeling of the number operator, which counts photons and has eigenvalue l on the l-th basis vector, and the order of its eigenvalues is that of the photon numbers;
- *continuous*: the set Ξ is $[0, 1]$, the Hilbert spaces are $H_1 = H_2 = L^2[0, 1]$, and the operator $A = B$ is multiplication by x, the position of a particle confined to the interval $[0, 1]$.

In both cases, let $C = [A < B]$ be the spectral projection of $I \otimes B - A \otimes I$ for the interval $(0, \infty)$, which represents the event "Bob's value exceeds Alice's". For a state ρ, let μ_ρ be the distribution of A in ρ, a Borel probability measure on Ξ. In the discrete case every probability measure on $\Xi \setminus \{1\}$ occurs, and in the continuous case the measures that occur are exactly the absolutely continuous ones (for a probability density p, take the pure state given by the unit vector $\psi = \sqrt{p}$). By Lemma 7.1, the expectation value of C for Alice's state ρ and Bob's state σ is

$$\varphi_C(\rho, \sigma) = (\mu_\rho \otimes \mu_\sigma)\big(\{(x, y) : x < y\}\big) = \int g_\rho \, d\mu_\sigma, \qquad g_\rho(y) = \mu_\rho\big([0, y)\big).$$

The function $g_\rho \colon \Xi \to [0, 1]$ is nondecreasing, with $g_\rho(0) = 0$ and $g_\rho(1) = 1$ in both cases, since μ_ρ assigns no mass to the point 1; in the continuous case it is also continuous. Accordingly, the effective observable is $T_C(\rho) = g_\rho(B)$. By Corollary 7.4, the observable C is uniformly NIP, and in particular NIP. It is unstable: in the discrete case by the interleaving argument of Part I [DI26, Proposition 4.6(3)], which uses only the order of the eigenvalues, and in the

continuous case by the same ladder argument, with states of Alice's and of Bob's particles supported in small disjoint intervals around positions a_k and b_l, respectively, for $k, l \geq 1$, interleaved as $a_1 < b_1 < a_2 < b_2 < \cdots$. Writing ρ_k and σ_l for these preparations of Alice and Bob, respectively, we obtain $\varphi_C(\rho_k, \sigma_l) = 1$ for $k \leq l$ and 0 for $k > l$; the iterated limit taken first in k is 0, whereas the one taken first in l is 1.

7.3. **The probe space and the idealized responses.** For both comparisons of the previous subsection, the discrete and the continuous one, and for every state ρ of Alice's system, the function g_ρ, whose value at a point $y \in \Xi$ is the probability that Alice's value lies below y, is nondecreasing. In Proposition 7.7 below, we show that the pointwise limits of these functions describe all idealized responses. In that proposition we write $C(\Xi)$ for the algebra of continuous complex functions on Ξ, with the supremum norm. Let M_Ξ be the set of nondecreasing functions $g\colon \Xi \to [0,1]$ with $g(0) = 0$ and $g(1) = 1$, with the topology of pointwise convergence. It is a closed subset of $[0,1]^\Xi$, hence compact. Although M_Ξ contains discontinuous functions, integration against a fixed measure is continuous on it, by the following lemma.

Lemma 7.6. *Let ν be a Borel probability measure on Ξ, and let (g_i) be a net in M_Ξ, indexed by a directed set, that converges pointwise to a function $g \in M_\Xi$. Then $\int g_i \, d\nu \to \int g \, d\nu$.*

Proof. Let $\varepsilon > 0$ be given. Choose a positive integer q with $1/q < \varepsilon$. For each $j = 1, \ldots, q-1$, take the least point $x \in \Xi$ with $\nu(\Xi \cap [0,x]) \geq j/q$, and adjoin 0 and 1. List the distinct points obtained as $0 = p_0 < p_1 < \cdots < p_m = 1$. Each open cell $c_k = \Xi \cap (p_{k-1}, p_k)$ has measure at most $1/q < \varepsilon$: a cell of greater measure would contain a point at which one of the intervening levels j/q is first reached, contrary to the choice of consecutive partition points. Put $\eta_i = \max_k |g_i(p_k) - g(p_k)|$, which tends to 0. On c_k both g_i and g take values between their values at p_{k-1} and at p_k, because they are nondecreasing. Hence

$$\Big| \int_{c_k} g_i \, d\nu - \int_{c_k} g \, d\nu \Big| \leq \nu(c_k)\big(g(p_k) - g(p_{k-1}) + 2\eta_i\big),$$

and summing over k, together with the contribution of the points p_k, shows that $|\int g_i \, d\nu - \int g \, d\nu| \leq \varepsilon + 3\eta_i$, since the differences $g(p_k) - g(p_{k-1})$ add up to at most 1. □

Proposition 7.7. *For each of the two comparisons $C = [A < B]$, the discrete and the continuous one, the probe algebra $\mathfrak{A}_C$, the unital C*-algebra generated by the effective observables, is $\{f(B) : f \in C(\Xi)\}$, so the probe space Y_C, its state space, is the space $P(\Xi)$ of Borel probability measures on Ξ with the weak* topology. The map $\Phi\colon r \mapsto \big(r(\delta_y)\big)_{y\in\Xi}$, where δ_y is the point mass at y, is a homeomorphism from the compactum $\overline{\mathcal{R}_C}$ of idealized responses onto M_Ξ, and every idealized response is given by $r(\nu) = \int \Phi(r) \, d\nu$.*

Proof. Each function g_ρ is continuous on Ξ: in the continuous case this follows from absolute continuity of the distribution, and in the discrete case continuity at the only accumulation point follows from $g_\rho(1 - 1/(l+1)) \to 1 = g_\rho(1)$ as $l \to \infty$. The functions g_ρ for states ρ whose distribution has full support in Ξ separate the points of Ξ, and the map $f \mapsto f(B)$ is an isometric $*$-isomorphism from $C(\Xi)$ onto its image, because the spectrum of B is Ξ. By the Stone–Weierstrass theorem, the unital C*-algebra generated by the operators $g_\rho(B)$ is $\{f(B) : f \in C(\Xi)\}$, and the slice map takes values in it. By the Riesz representation theorem, its state space is $P(\Xi)$, and the response of ρ is $r_\rho(\nu) = \int g_\rho \, d\nu$.

The map Φ is continuous, because its coordinates are evaluations. Let $r \in \overline{\mathcal{R}_C}$ be the pointwise limit of a net (r_{ρ_i}). Then the functions $g_{\rho_i} = \Phi(r_{\rho_i})$ converge pointwise to $g =$

$\Phi(r) \in M_\Xi$, and, by Lemma 7.6, $r(\nu) = \lim_i \int g_{\rho_i}\, d\nu = \int g\, d\nu$. Thus the response r is determined by $\Phi(r)$, the map Φ is injective, and, as a continuous injection from a compact space, it is a homeomorphism onto its image. The image is compact and contains the functions g_ρ.

It remains to be shown that the functions g_ρ are dense in M_Ξ. Let $g \in M_\Xi$ and a finite set $F \subseteq \Xi$ containing 0 and 1 be given. In the continuous case, the piecewise linear interpolation $\tilde{g}$ of g on F is continuous and nondecreasing, with values 0 at 0 and 1 at 1; hence $\tilde{g} = g_\rho$ for a state ρ whose distribution has $\tilde{g}$ as its distribution function, which is absolutely continuous, and $\tilde{g}$ agrees with g on F. In the discrete case, write $F = \{\xi_0 < \xi_1 < \cdots < \xi_m\}$, with $\xi_0 = 0$ and $\xi_m = 1$. The probability measure with mass $g(\xi_{j+1}) - g(\xi_j)$ at ξ_j for $0 \le j < m$ is the distribution of a state ρ with finitely many photons, and $g_\rho(\xi_j) = g(\xi_j) - g(\xi_0) = g(\xi_j)$ for every j, so the function g_ρ agrees with g on F. Every neighborhood of g therefore contains some g_ρ. □

7.4. **Location in the classification.** We can now place both comparisons in the classification of Theorem 6.9.

Theorem 7.8. *Let $C = [A < B]$ be the comparison of Alice's and Bob's photon numbers, in the discrete case, or of the positions of their particles in a box, in the continuous case.*

(a) *In the discrete case, the observable C is unstable and NIP$_3$.*

(b) *In the continuous case, the observable C is unstable and NIP$_1$, but not NIP$_2$. Hence, for every trace-norm dense sequence $(\rho_n)_{n\in\mathbb{N}}$ of states of Alice's particle, there is an increasing injection $\iota\colon 2^{<\mathbb{N}} \to \mathbb{N}$ such that the responses $\{r_{\rho_{\iota(t)}} : t \in 2^{<\mathbb{N}}\}$ are equivalent to D_6 or to D_7, the two minimal families of the heptachotomy whose closures are first countable but not hereditarily separable.*

Proof. (a) By Proposition 7.7, the compactum of idealized responses is homeomorphic to a subspace of $[0,1]^\Xi$ with Ξ countable, which is metrizable.

(b) By Proposition 7.7, the compactum of idealized responses is homeomorphic to $M_{[0,1]}$, a copy of the closed subspace of the Helly space described at the start of this section. It is first countable. Given $g \in M_{[0,1]}$, let Z_g be the countable set consisting of the rational points, the jump points of g and the endpoints; we claim that the sets $\{h : |h(p) - g(p)| < 1/n \text{ for } p \in F\}$, with $F \subseteq Z_g$ finite and $n \ge 1$, form a neighborhood base at g. It suffices to show that each set $\{h : |h(y) - g(y)| < \varepsilon\}$, with $y \in [0,1]$ and $\varepsilon > 0$, contains one of them. If $y \in Z_g$, this is clear. If $y \notin Z_g$, then g is continuous at y, and there are rationals $p < y < q$ with $g(q) - g(p) < \varepsilon/2$. Every $h \in M_{[0,1]}$ with $|h - g| < \varepsilon/4$ at p and q satisfies $h(p) \le h(y) \le h(q)$, because h is nondecreasing, and hence $|h(y) - g(y)| < \varepsilon$.

The compactum is not hereditarily separable. For $0 < a < 1$, let h_a be 0 on $[0, a)$, $\frac{1}{2}$ at a and 1 on $(a, 1]$. The set $\{h \in M_{[0,1]} : \frac{1}{4} < h(a) < \frac{3}{4}\}$ is a neighborhood of h_a that contains no other h_b, so the set $\{h_a : 0 < a < 1\}$ is an uncountable discrete subspace.

The observable is therefore NIP$_1$ but not NIP$_2$, and the last statement is Theorem 6.9(ii). □

The idealized responses of the continuous comparison have a direct physical reading, which is also a probabilistic one. The idealized responses are described by arbitrary functions g in $M_{[0,1]}$, where $g(y)$ is the limiting probability that Alice's value lies below Bob's fixed value y. Such a function need not be the distribution function of a single countably additive position measure: its value at a jump also records how the particle approaches the jump point, from

the left, from the right, or both. The function h_a describes a half-and-half mixture of a particle concentrating at a from the left and one concentrating at a from the right, and Bob's idealized probe concentrated at a reads the value $\frac{1}{2}$. In the discrete case no such splitting occurs at an eigenvalue, because each eigenvalue is isolated; the additional limiting behavior is escape to infinity, which may be mixed with an ordinary distribution on the eigenvalues.

7.5. **Unconfined quantities.** Confinement to a box is not what produces this subspace of the Helly space: a comparison depends only on the order of the two outcomes, so it is unchanged when both quantities are replaced by the same strictly increasing function of themselves, and a suitable function turns unconfined quantities into positions in a box (see Proposition 7.9 below).

Proposition 7.9. *Let A be a self-adjoint operator on the Hilbert space H_1 of Alice's system and B one on the Hilbert space H_2 of Bob's, each unitarily equivalent to multiplication by x on $L^2(\mathbb{R})$; for instance, the positions of two particles on a line, or quadratures of two modes of light. Then the comparison $[A < B]$ is unstable, and its idealized responses form a copy of the space of nondecreasing functions $[0,1] \to [0,1]$ taking the value 0 at 0 and 1 at 1, a closed subspace of the Helly space. Hence the comparison is NIP$_1$ but not NIP$_2$.*

Proof. We may assume that A and B are multiplication by x on $L^2(\mathbb{R})$. For physical quantities, x denotes the numerical value in the fixed common unit used for the comparison. Let $\theta\colon \mathbb{R} \to (0,1)$ be the increasing diffeomorphism $\theta(x) = \frac{1}{2}(1 + \tanh x)$. The operator U given by $(U\psi)(u) = \psi(\theta^{-1}(u))\,|(\theta^{-1})'(u)|^{1/2}$ is a unitary from $L^2(\mathbb{R})$ onto $L^2[0,1]$ that carries multiplication by $\theta(x)$ to multiplication by u. Since θ is strictly increasing, $x < y$ exactly when $\theta(x) < \theta(y)$; hence the spectral projection $[A < B]$ equals $[\theta(A) < \theta(B)]$, and $U \otimes U$ carries it to the comparison of the continuous case. The unitary $U \otimes U$ preserves expectation values and carries actual and idealized states to actual and idealized states; it therefore induces a homeomorphism between the two compacta of idealized responses. The statement now follows from Proposition 7.7 and Theorem 7.8(b). □

For the comparisons covered by Proposition 7.9, confinement does not affect the compactum of idealized responses. For quantities whose expectation values matter, such as energies, the rescaling by θ is not available, because it is not linear (see Section 5.2).

7.6. **The causal order is not NIP$_1$.** The compactum of idealized responses of a comparison is first countable because its members are monotone, and a monotone function has at most countably many jumps, each with two sides (see the proof of Theorem 7.8(b)). A relation between Bob's value and two values of Alice's, one on each side, removes this restriction (see Theorem 7.10 below), and the natural example is the causal order of spacetime. We work in a $(1+1)$-dimensional spacetime, with one dimension of time and one of space, in units in which the speed of light is 1. An event (t, ξ), at time t and position ξ, can influence a detection by Bob at the point y at time 0 exactly when $|y - \xi| \le -t$, that is, when y lies between the points $a = \xi + t$ and $b = \xi - t$ at which the two light rays through the event cross Bob's time slice. We describe Alice's event by these null coordinates and confine everything to the causal diamond over Bob's interval $[0,1]$, the set of events whose two light rays cross Bob's slice inside that interval: Alice's Hilbert space is $H_1 = L^2([0,1]^2)$, with variables (a,b), Bob's is $H_2 = L^2[0,1]$, with variable y, and the *causal-order observable* C is multiplication by the indicator of

$$R = \{(a,b,y) : a \le y \le b\}.$$

Events with $a > b$ lie in the future of Bob's slice and influence none of his detections.

For a state ρ of Alice's system, let μ_ρ be the distribution of (a,b) in ρ, an absolutely continuous probability measure on $[0,1]^2$, and similarly for Bob. The observable C is the relation observable C_R for this relation, so, by Lemma 7.1, its expectation value in states ρ and σ of the two systems is $\varphi_C(\rho,\sigma) = (\mu_\rho \otimes \mu_\sigma)(R) = \int g_\rho \, d\mu_\sigma$, where $g_\rho(y) = \mu_\rho(\{(a,b) : a \le y \le b\})$ is the probability that Alice's event can influence a detection at y. The function g_ρ is continuous: the sections of R at y and at y' differ by a set contained in two strips, each of area at most $|y-y'|$, and the μ_ρ-measure of such strips tends to 0 with $|y-y'|$, because μ_ρ is absolutely continuous. The effective observable $T_C(\rho)$ is multiplication by g_ρ.

Theorem 7.10. *The causal-order observable is NIP but not NIP$_1$. Hence, for every trace-norm dense sequence $(\rho_n)_{n\in\mathbb{N}}$ of states of Alice's system, there is an increasing injection $\iota\colon 2^{<\mathbb{N}} \to \mathbb{N}$ such that the responses $\{r_{\rho_{\iota(t)}} : t \in 2^{<\mathbb{N}}\}$ are equivalent to D_5, the minimal family of the heptachotomy whose closure, $\hat{A}(2^{\mathbb{N}})$, is not first countable.*

Proof. For every state ρ, the event $\{a \le y \le b\}$ is the event $\{a \le y\}$ with the event $\{a \le y,\ b < y\}$ removed, so

$$g_\rho(y) = \mu_\rho\big(\{(a,b) : a \le y\}\big) - \mu_\rho\big(\{(a,b) : a \le y,\ b < y\}\big),$$

a difference of two nondecreasing functions with values in $[0,1]$. By Proposition 7.3, whose constant is here 1, the observable C is uniformly NIP, and in particular NIP. By Lemma 6.3 the compactum of idealized responses is separable, so its level is defined.

The probe algebra consists of operators M_f of multiplication by continuous functions f; it is therefore a C*-subalgebra of $\{M_f : f \in C[0,1]\}$, which is isometrically isomorphic to $C[0,1]$, because $\|M_f\| = \sup|f|$. Every state of $\mathfrak{A}_C$ extends to a state of this larger algebra (see [KR83, Theorem 4.3.13]), and, by the Riesz representation theorem, every state τ of $\mathfrak{A}_C$ is thus represented by a probability measure ν on $[0,1]$, in the sense that $\tau(M_f) = \int f \, d\nu$ for $M_f \in \mathfrak{A}_C$; in particular, $r_\rho(\tau) = \int g_\rho \, d\nu$. For each point $c \in (0,1)$, the evaluation $\tau_c(M_f) = f(c)$ is a state of $\mathfrak{A}_C$, Bob's probe localized at c.

Let ρ_0 be a state whose distribution is uniform on $[\frac12, 1] \times [0, \frac12]$, an event after Bob's slice. Then $g_{\rho_0} = 0$, and the zero function is a response. For $c \in (0,1)$ and large n, let $\rho_{c,n}$ be a state whose distribution is uniform on $[c - \frac{2}{n}, c - \frac{1}{n}] \times [c + \frac{1}{n}, c + \frac{2}{n}]$, an event at position near c shortly before Bob's slice. Then $g_{\rho_{c,n}}(c) = 1$, and $g_{\rho_{c,n}}(y) = 0$ when $|y - c| > \frac{2}{n}$. Hence the function $g_{\rho_{c,n}}$ converges pointwise to the indicator of $\{c\}$, and, by dominated convergence, $r_{\rho_{c,n}}(\tau) \to \nu(\{c\})$ for every probe τ and every measure ν representing it. The limit r_c is an idealized response, with $r_c(\tau_c) = 1$.

Suppose that the zero function had a countable base of neighborhoods (U_k) in the compactum of idealized responses. Each U_k contains a set $\{r : |r(\tau)| < \varepsilon_k \text{ for } \tau \in F_k\}$ with F_k finite. Choose a representing measure for each probe in $F = \bigcup_k F_k$; together these measures have countably many atoms, so some $c \in (0,1)$ is an atom of none of them. Then $r_c(\tau) = 0$ for $\tau \in F$, so the response r_c lies in every U_k. Since the space is Hausdorff, the intersection of a neighborhood base at a point is that point; hence $r_c = 0$, contrary to $r_c(\tau_c) = 1$. The compactum is therefore not first countable, and the last statement is Theorem 6.9(iii). □

Physically, the proof says the following. The idealized response r_c is the limit of the responses of events that approach the point c of Bob's time slice from the past: the light cones of these events meet the slice in intervals that shrink to the point c, so in the limit only probes that assign positive weight to the point c itself detect anything. There is one such idealized response for each point c, and countably many of Bob's probes cannot tell all of

them apart from the response of an event that influences nothing; this is the failure of first countability. The relation R is the intersection of two comparisons, $a \le y$ and $y \le b$, one for each light ray. A single comparison has monotone idealized responses; the two rays together allow idealized responses supported on a single point. Here Todorčević's trichotomy is used in full: it turns the failure of first countability into a dyadic tree of actual events whose responses are equivalent to the canonical family D_5. The proof suggests a tree of this kind, namely events whose light cones meet Bob's slice in the intervals of a Cantor construction, each emitted later than its predecessor, although we do not claim that the trichotomy produces this one.

7.7. **The comparison of phases is not NIP$_1$.** Orders on a line give first countable compacta of idealized responses (see Theorem 7.8 and Proposition 7.9). In Theorem 7.12 below, we show that on a circle first countability can fail. Identify the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ with $[0,1)$, so that angles are measured in turns and a half-turn is $\frac{1}{2}$. Alice and Bob each hold a particle on a ring, a planar rotor, with Hilbert spaces $H_1 = H_2 = L^2(\mathbb{T})$, the angle acting by multiplication. The *phase comparison* is the relation observable $C = C_R$ for

$$R = \big\{(x,y) \in [0,1)^2 : \ y - x \in (0, \tfrac{1}{2}) \bmod 1\big\},$$

the event "Bob's angle is ahead of Alice's by less than half a turn". For a state ρ of Alice's rotor, the distribution μ_ρ of her angle is an absolutely continuous probability measure on $\mathbb{T}$, and every such measure occurs. By Lemma 7.1, the phase comparison has, in states ρ and σ of the two rotors, the expectation value $\varphi_C(\rho,\sigma) = \int g_\rho \, d\mu_\sigma$, where $g_\rho(y) = \mu_\rho\big((y - \frac{1}{2}, y)\big)$ is the probability that Alice's angle lies in the open half-circle ending at y; the function g_ρ is continuous on $\mathbb{T}$, because μ_ρ has no atoms.

Lemma 7.11. *For every probability measure μ on $\mathbb{T} = [0,1)$, the function $g(y) = \mu((y - \frac{1}{2}, y))$ on $[0,1)$ is $2(u - v)$ for two nondecreasing functions $u, v \colon [0,1) \to [0,1]$.*

Proof. Let $G(y) = \mu([0,y))$. For $y \in [\frac{1}{2}, 1)$ the half-circle is the interval $(y - \frac{1}{2}, y)$, and $g(y) = G(y) - \mu([0, y - \frac{1}{2}])$; for $y \in [0, \frac{1}{2})$ it is $(y + \frac{1}{2}, 1) \cup [0, y)$, and $g(y) = G(y) + 1 - \mu([0, y + \frac{1}{2}])$. Let $V(y) = \mu([0, y + \frac{1}{2}]) - 1$ for $0 \le y < \frac{1}{2}$ and $V(y) = \mu([0, y - \frac{1}{2}])$ for $\frac{1}{2} \le y < 1$. Then $g = G - V$. The functions G and V are nondecreasing, with values in $[0,1]$ and $[-1,1]$, respectively; indeed, V is nondecreasing on each half, and at $\frac{1}{2}$ it passes from $\lim_{y \uparrow 1/2} V(y) = 0$ to $V(\frac{1}{2}) = \mu(\{0\}) \ge 0$. Put $u = \frac{1}{2}(G + 1)$ and $v = \frac{1}{2}(V + 1)$; then $g = 2(u - v)$. □

Theorem 7.12. *The phase comparison is uniformly NIP, with bounds that depend only on the margin, and it is not NIP$_1$: its compactum of idealized responses is not first countable. Hence, for every trace-norm dense sequence $(\rho_n)_{n \in \mathbb{N}}$ of states of Alice's rotor, there is an increasing injection $\iota \colon 2^{<\mathbb{N}} \to \mathbb{N}$ such that the responses $\{r_{\rho_{\iota(t)}} : t \in 2^{<\mathbb{N}}\}$ are equivalent to D_5.*

Proof. By Lemma 7.11, the function g_ρ agrees on $[0,1)$ with $2(u - v)$ for two nondecreasing functions $u, v \colon [0,1) \to [0,1]$, which we extend to nondecreasing functions on $\mathbb{R}$, constant on $(-\infty, 0)$ and on $[1, \infty)$. Bob's distributions are carried by $[0,1)$, so only these values of g_ρ enter the proof of Proposition 7.3; that proposition therefore applies with the constant 2, and the observable is uniformly NIP.

As in the proof of Theorem 7.10, every probe τ is represented by a probability measure ν on $\mathbb{T}$, with $r_\rho(\tau) = \int g_\rho \, d\nu$, and for $z \in \mathbb{T}$ the evaluation $\tau_z(M_f) = f(z)$ is a probe, Bob's

rotor localized at z. Let ρ_0 be a state in which Alice's angle is uniformly distributed. Every open half-circle has measure $\frac{1}{2}$, so $r_{\rho_0}(\tau) = \frac{1}{2}$ for every probe τ.

For $a \in \mathbb{T}$ and large n, let $\rho_{a,n}$ be a state whose distribution is uniform, with weight $\frac{1}{2}$ each, on the arcs $B_1 = (a - \frac{1}{n}, a)$ and $B_2 = (a + \frac{1}{2}, a + \frac{1}{2} + \frac{1}{n})$: half of the angle just behind a, half just ahead of the antipode. Write $A_y = (y - \frac{1}{2}, y)$. If $y = a$, then the half-circle A_y contains B_1 and B_2, so $g_{\rho_{a,n}}(a) = 1$. If $y = a + \frac{1}{2}$, then the half-circle $A_y = (a, a + \frac{1}{2})$ meets neither, and $g_{\rho_{a,n}}(a + \frac{1}{2}) = 0$. Otherwise the endpoints of A_y are at some positive distance d from a and from $a + \frac{1}{2}$; an open half-circle whose endpoints differ from these two antipodal points contains exactly one of them. Once $\frac{1}{n} < d$, the half-circle A_y therefore contains the arc B_1 just behind a if it contains a, and the arc B_2 just ahead of $a + \frac{1}{2}$ if it contains $a + \frac{1}{2}$, but not both, and $g_{\rho_{a,n}}(y) = \frac{1}{2}$. Hence the function $g_{\rho_{a,n}}$ converges pointwise to $g_a = \frac{1}{2} + \frac{1}{2}\mathbf{1}_{\{a\}} - \frac{1}{2}\mathbf{1}_{\{a+1/2\}}$, and, by dominated convergence, $r_{\rho_{a,n}}(\tau) \to \frac{1}{2} + \frac{1}{2}\nu(\{a\}) - \frac{1}{2}\nu(\{a + \frac{1}{2}\})$ for every probe τ and every measure ν representing it. The limit r_a is an idealized response, with $r_a(\tau_a) = 1$.

Suppose that r_{ρ_0} had a countable base of neighborhoods (U_k) in the compactum of idealized responses. Each U_k contains a set $\{r : |r(\tau) - \frac{1}{2}| < \varepsilon_k \text{ for } \tau \in F_k\}$ with F_k finite. Choose a representing measure for each probe in $F = \bigcup_k F_k$; together these measures have countably many atoms, so there is $a \in \mathbb{T}$ such that neither a nor $a + \frac{1}{2}$ is one of them. Then $r_a(\tau) = \frac{1}{2}$ for every $\tau \in F$, so r_a lies in every U_k, and, the space being Hausdorff, the response r_a equals r_{ρ_0}, contrary to $r_a(\tau_a) = 1$. The compactum is therefore not first countable, and the last statement is Theorem 6.9(iii). □

Here, too, the proof has a physical reading. A state in which the angle is uniformly distributed responds $\frac{1}{2}$ to every probe. For an antipodal pair of points a and $a + \frac{1}{2}$, the idealized limit of preparations that put half of the angle's distribution just behind a and half just ahead of $a + \frac{1}{2}$ also responds $\frac{1}{2}$, except that a probe at a reads 1 and a probe at the antipode reads 0: the jumps at the two ends of the half-turn point in opposite directions and cancel everywhere else. There is one such pair for each point of the circle, and countably many probes cannot distinguish all of them from such a state. On a line this cannot happen, because a monotone response cannot hide a jump (see the proof of Theorem 7.8(b)); on a circle there is no first or last point. The same contrast appears in topological dynamics. The enveloping semigroup of a dynamical system is the pointwise closure of its acting maps. Glasner and Megrelishvili showed that compact metrizable linearly ordered spaces, with their order topologies and order-preserving group actions, have first countable enveloping semigroups, while the circular analogue of the Helly space is not first countable [GM22, Theorem 8.8, Proposition 8.10 and Remark 8.11]. The circular space in that comparison consists of circular-order-preserving maps of the circle, whereas the responses here are real-valued; neither result implies the other.

7.8. **Interference fringes are not NIP.** The relations studied so far in this section are all NIP. The fixed-contrast interference patterns in Proposition 7.13 below show how oscillation at ever finer scales can produce the independence property. Positions on Bob's screen are measured in a fixed unit of length, in which the screen is the interval $[0, 2\pi]$, and wavenumbers are measured in the inverse unit. We model the experiment by a register on Alice's side with orthonormal basis $(e_n)_{n\in\mathbb{N}}$, whose label n selects a fringe pattern with wavenumber $k_n = 8^n$. At position $x \in [0, 2\pi]$, Bob's detector accepts with probability $f_n(x) = \frac{1}{2}(1 + \cos k_n x)$, an idealized local response that is largest on a bright fringe. The *fringe observable* is $C = \sum_n |e_n\rangle\langle e_n| \otimes M_{f_n}$ on $\ell^2(\mathbb{N}) \otimes L^2[0, 2\pi]$: the label n of Alice's state selects the test f_n applied

to Bob's detector, and C is thus a label-controlled observable in the sense of Definition 8.1 below.

Proposition 7.13. *The fringe observable is not NIP: Alice's states e_n form an independent sequence, and the closure of their responses contains a copy of $\beta\mathbb{N}$.*

Proof. Let $E, F \subseteq \mathbb{N}$ be finite and disjoint. Choose $y \in [0,1)$ whose base-8 digit in place $n+1$ is 0 for $n \in F$ and 4 for $n \in E$, and put $x = 2\pi y$. Then $\cos k_n x = \cos(2\pi\{8^n y\})$, where $\{8^n y\}$, the fractional part of $8^n y$, lies in $[0, \frac{1}{8})$ for $n \in F$ and in $[\frac{1}{2}, \frac{5}{8})$ for $n \in E$; hence $f_n(x) \geq \frac{3}{4}$ for $n \in F$ and $f_n(x) \leq \frac{1}{4}$ for $n \in E$. By continuity of the finitely many functions f_n involved, a state σ of Bob's detector concentrated near x has expectation values at least $\frac{5}{8}$ for $n \in F$ and at most $\frac{3}{8}$ for $n \in E$. By Lemma 5.2(c), the restrictions of such states are probes; the sequence (e_n) is therefore independent. As in the proof of Theorem 6.1(c), no subsequence of the responses converges pointwise, and Fact 3.4 yields a subsequence whose closure is homeomorphic to $\beta\mathbb{N}$. □

The contrast between these fringes and the orders of this section displays the finite-pattern obstruction measured by NIP. A comparison asks on which side of a threshold a value lies; the pattern of wavenumber 8^n has 8^n periods on Bob's screen, each with a bright and a dark half. The base-8 construction lets one prescribe bright or dark responses independently for any finite collection of labels. This is independence of finite patterns, rather than statistical independence of the fringe intensities under a distribution of screen positions.

8. The seven minimal families in physics

The classification of a full response compactum and the classification of a selected tree of preparations concern different compact sets. For a NIP observable, the full compactum is convex, and hereditary separability implies metrizability by Corollary 6.8; the closure of a selected tree need not be convex. In Theorem 8.5 below, we realize each of the seven minimal families by such a tree of actual preparations. Photon-number comparisons, position comparisons and spin registers provide the realizations, while the passage from functions of position to responses on probability measures preserves the indexed families and their pointwise closures.

8.1. Label-controlled observables.

Several of our observables are of the following kind. Alice's system is a register that holds a label, and the label selects the test applied to Bob's system: a detector that accepts Bob's system at each point with a probability that depends on the label and varies continuously with the point.

Definition 8.1. Let L be a compact metrizable space, λ a Borel probability measure on L with full support, I a countable set and $(f_i)_{i\in I}$ a family of continuous functions $L \to [0,1]$. The *label-controlled observable* is

$$C = \sum_{i\in I} |e_i\rangle\langle e_i| \otimes M_{f_i} \qquad \text{on} \quad \ell^2(I) \otimes L^2(L,\lambda),$$

where (e_i) is the standard basis of $\ell^2(I)$ and M_f is multiplication by f.

The sum converges strongly, that is, on each vector, because its terms act on orthogonal subspaces, and the operator C lies between 0 and the identity operator. When L contains a countable dense set Z of isolated points, we take $\lambda = \sum_{x\in Z} w_x \delta_x$, a combination of the point masses δ_x with positive weights w_x of sum 1; then the space $L^2(L,\lambda)$ is a weighted copy of $\ell^2(Z)$, and the position of Bob's system takes its values in Z. As in Section 7, we write $C(L)$ for the algebra of continuous complex functions on L, with the supremum norm.

Lemma 8.2. *Let C be as in Definition 8.1, and suppose that the functions f_i separate the points of L. Then the probe algebra $\mathfrak{A}_C$, generated by the effective observables, is $\{M_f : f \in C(L)\}$, the probe space Y_C, its state space, is the space $P(L)$ of Borel probability measures on L with the weak* topology, and the response of the label state e_i is $r_{e_i}(\nu) = \int f_i \, d\nu$.*

Proof. For a density operator ρ, tracing out Alice's system yields $T_C(\rho) = \sum_i \langle e_i, \rho e_i \rangle M_{f_i}$, a norm-convergent sum, so the slice map takes values in $\{M_f : f \in C(L)\}$. Since λ has full support and f is continuous, $\|M_f\| = \sup |f|$; hence the map $f \mapsto M_f$ is an isometric $*$-isomorphism from $C(L)$ onto its image. The unital C*-subalgebra of $C(L)$ generated by the functions f_i is $C(L)$, by the Stone–Weierstrass theorem, because they separate points. It follows that $\mathfrak{A}_C = \{M_f : f \in C(L)\}$, and, by the Riesz representation theorem, its states are the measures $\nu \in P(L)$, acting by $M_f \mapsto \int f \, d\nu$. □

8.2. **Transfer from points to measures.** The responses of our observables are integrals $\nu \mapsto \int g \, d\nu$ of functions g on a compact space against Bob's probes. In Lemma 8.3 below, we show that a family of such responses has a closure homeomorphic to that of the family of integrands, by a homeomorphism that matches the two families term by term.

Lemma 8.3. *Let L be a compact metrizable space and $(f_t)_{t \in T}$ a countable family of continuous functions $L \to [0,1]$ that is relatively compact in the space $B_1(L)$ of Baire-1 functions on L. For $t \in T$ and $\nu \in P(L)$, put $F_t(\nu) = \int f_t \, d\nu$. Then the families $\{F_t\}$, of functions on $P(L)$, and $\{f_t\}$, of functions on L, are equivalent in the sense of Definition 4.1, and $\{F_t\}$ is NIP.*

Proof. Every sequence (f_{t_n}) has a subsequence that converges pointwise on L. Indeed, otherwise, by Fact 3.4, some subsequence would have a closure homeomorphic to $\beta\mathbb{N}$, which is not Fréchet–Urysohn (a space is Fréchet–Urysohn if every point in the closure of a subset is the limit of a sequence from that subset), whereas the closure of $\{f_t\}$ is a Rosenthal compactum and therefore Fréchet–Urysohn (see [BFT78, Theorem 3F]). By the dominated convergence theorem, the corresponding subsequence of (F_{t_n}) converges pointwise on $P(L)$.

The family $\{F_t\}$ is thus NIP. Indeed, if a sequence (F_{t_n}) and reals $a < b$ witnessed the independence property, then, since $P(L)$ is compact and the functions F_t are continuous, for each $E \subseteq \mathbb{N}$ there would be ν with $F_{t_n}(\nu) \leq a$ for $n \in E$ and $F_{t_n}(\nu) \geq b$ for $n \notin E$ (see the proof of Theorem 3.6); taking for E every other index of a pointwise convergent subsequence would yield a ν at which that subsequence does not converge, contrary to the previous paragraph. By Theorem 3.6, the closure of $\{F_t\}$ is contained in $B_1(P(L))$; it is a Rosenthal compactum, hence Fréchet–Urysohn.

Let Φ send a function G on $P(L)$ to the function $x \mapsto G(\delta_x)$ on L. The map Φ is continuous for the pointwise topologies, and $\Phi(F_t) = f_t$. The map Φ is injective on the closure of $\{F_t\}$: if G lies in that closure, then, the closure being Fréchet–Urysohn, the function G is the pointwise limit of a sequence (F_{t_n}); the sequence (f_{t_n}) then converges pointwise to $\Phi(G)$, which is therefore a bounded Borel function, and, by dominated convergence, the value $G(\nu)$ equals $\lim_n \int f_{t_n} \, d\nu = \int \Phi(G) \, d\nu$ for every ν. Thus G is determined by $\Phi(G)$. As a continuous injection on a compact space, the map Φ is a homeomorphism of the closure of $\{F_t\}$ onto its image; this image is compact, contains $\{f_t\}$ and lies in the closure of $\{f_t\}$, so it is that closure. □

We also need to discard points that carry no information about limits.

Lemma 8.4. *Let L be a set, $L_0 \subseteq L$, and $(g_t)_{t \in T}$ a countable pointwise bounded family of real functions on L whose pointwise closure K, a compact space, is Fréchet–Urysohn. Suppose that*

(i) *the restrictions $g_t|_{L_0}$ are pairwise distinct, and none of them lies in the closure of the others;*
(ii) *if two limits of sequences $(g_{t_k})_{k\in\mathbb{N}}$ with pairwise distinct indices agree on L_0, then they are equal.*

Then the families (g_t) and $(g_t|_{L_0})$ are equivalent.

Proof. Restriction $\pi\colon K \to \mathbb{R}^{L_0}$ is continuous, and its image is the closure of $(g_t|_{L_0})$. Let $f \in K$ be different from every g_t. Since K is Fréchet–Urysohn, the function f is the limit of a sequence (g_{t_k}); since K is Hausdorff and f is not a g_t, no index occurs infinitely often, and, after passage to a subsequence, the indices are pairwise distinct. By (i), the map π is injective on the family; moreover, $\pi(g_t) \neq \pi(f)$ for every such limit f and every t, because $\pi(f)$ lies in the closure of $\{g_s|_{L_0} : s \neq t\}$, whereas $g_t|_{L_0}$ does not. By (ii), the map π is also injective on the limits. Hence π is a continuous injection of the compact space K, and therefore a homeomorphism onto its image. □

8.3. **Three settings.** The realizations use three physical settings.

Photon numbers. This is the discrete comparison of Section 7, in which Alice's and Bob's values lie in $\Xi = \{x_l : l \in \mathbb{N}\} \cup \{1\}$, where $x_l = 1 - \frac{1}{l+1}$. For Alice's number state e_l, the state with exactly l photons, the function $g_l = g_{e_l}$, whose value at a point y is the probability that Alice's value lies below y, is the indicator of $\{x_j : j > l\} \cup \{1\}$; the probe space is the space $P(\Xi)$ of Borel probability measures on Ξ, and $r_{e_l}(\nu) = \int g_l \, d\nu$ (see Proposition 7.7).

Positions in a box. This is the continuous comparison of Section 7: the probe space is $P([0,1])$, and a state ρ of Alice's particle with distribution μ_ρ has response $\nu \mapsto \int g_\rho \, d\nu$, where $g_\rho(y) = \mu_\rho([0,y))$. We place a Cantor set in the box. Let $\psi(\sigma) = \frac{3}{4} - \sum_n \sigma(n) 3^{-n-1}$ for $\sigma \in 2^{\mathbb{N}}$; this is an order-reversing homeomorphism of the Cantor space $2^{\mathbb{N}}$, with its lexicographic order $\prec$, onto a Cantor set $C_0 \subseteq [\frac{1}{4}, \frac{3}{4}]$. For a finite binary sequence $s \in 2^{<\mathbb{N}}$, let I_s be the smallest interval containing the image $\psi(V_s)$ of the clopen set V_s of infinite extensions of s; it has length $\ell_s = \frac{1}{2} 3^{-|s|}$, left endpoint $\psi(s^\frown 1^\infty)$ and right endpoint $\psi(s^\frown 0^\infty)$, the images of the extensions of s by ones and by zeros. If τ is the immediate successor of σ in $\prec$, then the interval $(\psi(\tau), \psi(\sigma))$ is a component of $[0,1] \setminus C_0$; the other components are $[0, \frac{1}{4})$ and $(\frac{3}{4}, 1]$. The components adjacent to I_s have length at least $\ell_s/2$: each of them is either $[0, \frac{1}{4})$ or $(\frac{3}{4}, 1]$, of length $\frac{1}{4} \geq \ell_s/2$, or the gap between $I_{u^\frown 0}$ and $I_{u^\frown 1}$ for some proper initial segment u of s, of length $\ell_u/3 \geq \ell_s$.

Registers. Alice's system is a chain of spins of variable finite length, with Hilbert space $\ell^2(2^{<\mathbb{N}})$ over the set $2^{<\mathbb{N}}$ of finite binary sequences; for $t \in 2^{<\mathbb{N}}$, the basis vector e_t is the chain of length $|t|$ whose spin at site n is up exactly when $t(n) = 1$. Bob probes it in two ways. In *scanning*, Bob's probe sits at a site $n \in \mathbb{N}$ and asks whether Alice's spin at that site is up: the test is $a_t(n) = t(n)$ for $n < |t|$ and $a_t(n) = 0$ otherwise. In *template matching*, Bob holds a reference configuration u, finite or infinite, and asks whether Alice's chain coincides with the beginning of his reference: the test is $b_t(u) = 1$ if $t \sqsubseteq u$ and $b_t(u) = 0$ otherwise. We extend a_t to the one-point compactification $L_a = \mathbb{N} \cup \{\infty\}$ by $a_t(\infty) = 0$, and we let L_b be the space of finite and infinite binary sequences, with the ultrametric $d(u,u') = 2^{-|u \wedge u'|}$ for $u \neq u'$, where $u \wedge u'$ is the longest common initial segment; it is compact, the finite sequences are isolated and dense, and the functions b_t are continuous, since $\{u : t \sqsubseteq u\}$ is clopen. The *scanning*, *template* and *combined* observables are the label-controlled observables with labels $2^{<\mathbb{N}}$, on L_a, L_b and their disjoint sum $L_a \oplus L_b$, with the functions a_t, b_t and (a_t, b_t), the last equal to a_t on L_a and to b_t on L_b, and with measures concentrated on $\mathbb{N}$, on the finite sequences, and on both, respectively. These functions separate the points of L_a, of L_b and of

$L_a \oplus L_b$; for instance, the functions of the empty label, $a_\emptyset = 0$ and $b_\emptyset = 1$, separate L_a from L_b. Hence Lemma 8.2 applies to all three observables.

8.4. **The realization.** Together, the three settings realize all seven minimal families.

Theorem 8.5. *Each minimal family is realized by a tree of actual preparations of a physically natural observable: for each $i = 1, \dots, 7$, the observable and the pure states $(\rho_t)_{t\in 2^{<\mathbb{N}}}$ of Alice's system listed below, given by pairwise orthogonal vectors, have responses $\{r_{\rho_t}\}$ equivalent to D_i. Here $(s_t)_{t\in 2^{<\mathbb{N}}}$ is the canonical enumeration of the regular dyadic subtree Θ fixed in Section 4, and the chains e_t, the Cantor set C_0, the homeomorphism ψ and the intervals I_s, of lengths ℓ_s, are those of the three settings described before the theorem.*

D_1: the comparison of photon numbers, with Alice's number states in any order;
D_2: scanning, with Alice's chains e_{s_t}, whose scanning functions are $s_t^\frown 0^\infty$, for $s_t \in \Theta$;
D_3: the comparison of positions, with Alice's particle uniformly distributed over the gap of C_0 immediately to the right of $\psi(s_t^\frown 0^\infty)$;
D_4: the comparison of positions, with Alice's particle uniformly distributed over the gap of C_0 immediately to the left of $\psi(s_t^\frown 1^\infty)$;
D_5: template matching, with Alice's chains e_t;
D_6: the combined observable, with Alice's chains e_t;
D_7: the comparison of positions, with Alice's particle distributed uniformly, half over $[\min I_{s_t} - \frac{\ell}{2}, \min I_{s_t} - \frac{\ell}{4}]$ and half over $[\max I_{s_t} + \frac{\ell}{4}, \max I_{s_t} + \frac{\ell}{2}]$, where $\ell = \ell_{s_t}$.

Moreover, the causal order of spacetime realizes D_5 in the sense of Theorem 7.10.

Proof. In each case the probe space is $P(L)$ for a compact metrizable space L, the responses are $r_{\rho_t}(\nu) = \int g_t \, d\nu$ with g_t continuous, and the closure of $\{g_t\}$ consists of Baire-1 functions; by Lemma 8.3, it suffices to show that $\{g_t\}$ is equivalent to D_i. We use the notation of Section 4: the function v_t is the characteristic function of the set V_t of infinite binary sequences that extend t, and the functions x_σ^+ and x_σ^- are the characteristic functions of $\{\tau \in 2^\mathbb{N} : \sigma \preceq \tau\}$ and $\{\tau \in 2^\mathbb{N} : \sigma \prec \tau\}$, for the lexicographic order $\prec$.

D_1. Index the functions g_l, $l \in \mathbb{N}$, by $2^{<\mathbb{N}}$ through any bijection, and write g_t for the function of index t. Every sequence of pairwise distinct members converges to the indicator of $\{1\}$, which is not a member, and each member is isolated. Hence the closure is a countable compact space with a single non-isolated point, and so is the closure $D_1 \cup \{0\}$ of D_1; the bijection $g_t \mapsto \frac{1}{|t|+1} v_t$, extended by sending the indicator of $\{1\}$ to 0, is a homeomorphism. The space Ξ is countable, so every function on it is Baire-1, every subset of a countable space being an F_σ.

D_2. Here $g_t = a_{s_t}$ is $s_t^\frown 0^\infty$ on $\mathbb{N}$ and 0 at ∞. This is the family D_2, extended continuously to $\mathbb{N} \cup \{\infty\}$; every function in its closure vanishes at ∞, so the extension does not change the closure, and every function on the countable space L_a is Baire-1.

D_3 and D_4. Let $\sigma_t = s_t^\frown 0^\infty$. If $\sigma_t \neq 0^\infty$, it has an immediate predecessor σ_t^-, and $G_t = (\psi(\sigma_t), \psi(\sigma_t^-))$ is a gap; if $\sigma_t = 0^\infty$, let $G_t = (\frac{3}{4}, 1]$. Let ρ_t be uniform on G_t. Then $g_t = 0$ on $[0, \psi(\sigma_t)]$, $g_t = 1$ to the right of G_t, and g_t is linear on G_t. On C_0, the function g_t satisfies $g_t(\psi(\tau)) = 0$ exactly when $\tau \succeq \sigma_t$; that is, $g_t \circ \psi = 1 - x_{\sigma_t}^+$. The sequences σ_t are pairwise distinct, and so are the gaps G_t. We apply Lemma 8.4 with $L_0 = C_0$. The closure of $\{g_t\}$ consists of nondecreasing functions; it is therefore a Rosenthal compactum, and hence Fréchet–Urysohn. Condition (i) holds because D_3 is discrete in its closure. For (ii), let J be a component of $[0,1] \setminus C_0$. For pairwise distinct t_k, eventually $G_{t_k} \neq J$, and then the function g_{t_k} is constant on $\overline{J}$; every such limit f is thus constant on $\overline{J}$, and so is determined on J by its

value at an endpoint of J in C_0. Consequently, the family $\{g_t\}$ is equivalent to $\{g_t|_{C_0}\}$, which is carried to D_3 by the homeomorphism ψ followed by the map $f \mapsto 1-f$, a homeomorphism for the pointwise topology. The case of D_4 is symmetric: with $\sigma_t = s_t^\frown 1^\infty$ and its immediate successor σ_t^+, the gap is $(\psi(\sigma_t^+), \psi(\sigma_t))$, or $[0, \frac{1}{4})$ if $\sigma_t = 1^\infty$, and $g_t \circ \psi = 1 - x^-_{\sigma_t}$.

D_5. Here $g_t = b_t$ on L_b, and its restriction to the infinite sequences is v_t. Let K be the pointwise closure of $\{b_t\}$. For a finite sequence u, only finitely many functions b_t are nonzero at u; hence any member of K that is nonzero at u belongs to this finite set of functions. Every member of $K \setminus \{b_t : t \in 2^{<\mathbb{N}}\}$ therefore vanishes on the finite sequences. The restriction to the closed set $2^{\mathbb{N}}$ of every member of K lies in $\hat{A}(2^{\mathbb{N}}) \subseteq B_1(2^{\mathbb{N}})$. The inverse image of an open set under such a member is therefore an F_σ subset of $2^{\mathbb{N}}$, together with at most countably many isolated points of L_b, and is thus F_σ in L_b. By Fact 3.1(2), each member of K is Baire-1 on L_b. Thus K is a Rosenthal compactum and is Fréchet–Urysohn. We may now apply Lemma 8.4 with $L_0 = 2^{\mathbb{N}}$: condition (i) holds because D_5 is discrete in its closure, and condition (ii) holds because limits of sequences with pairwise distinct indices vanish at every finite sequence. The family $\{b_t\}$ is consequently equivalent to D_5.

D_6. Here $g_t = (a_t, b_t)$ on $L_a \oplus L_b$. Every function on the countable compact metrizable space L_a is Baire-1, while every member of the pointwise closure of $\{b_t\}$ is Baire-1 on L_b, as shown for D_5. The pointwise closure of $\{g_t\}$ therefore consists of Baire-1 functions on the disjoint sum and is a Rosenthal compactum, hence Fréchet–Urysohn. On $L_0 = \mathbb{N} \oplus 2^{\mathbb{N}}$, the function g_t is $(t^\frown 0^\infty, v_t)$, the corresponding member of D_6 with its two summands interchanged. These restrictions are discrete in their closure, and every limit of a sequence with pairwise distinct indices vanishes at ∞ and at each finite sequence in L_b. Thus both conditions of Lemma 8.4 hold, and the family $\{g_t\}$ is equivalent to D_6.

D_7. Let $\ell = \ell_{s_t}$, and let ρ_t be the state whose distribution is half uniform on $A_t = [\min I_{s_t} - \frac{\ell}{2}, \min I_{s_t} - \frac{\ell}{4}]$ and half uniform on $B_t = [\max I_{s_t} + \frac{\ell}{4}, \max I_{s_t} + \frac{\ell}{2}]$; these intervals lie in the components adjacent to I_{s_t}. On C_0, the function g_t equals 0 to the left of I_{s_t}, $\frac{1}{2}$ on I_{s_t} and 1 to the right; that is, $g_t \circ \psi = 1 - x^+_{\sigma_t} + \frac{1}{2} v_{s_t}$ with $\sigma_t = s_t^\frown 0^\infty$, since $\psi(\sigma_t) = \max I_{s_t}$. Each member of the closure of D_7 has the form (v, x), for functions v and x on $2^{\mathbb{N}}$ with $v \le x$ that take only the values 0 and 1; hence the map $(v, x) \mapsto (1 - x + \frac{1}{2}v) \circ \psi^{-1}$ is a continuous injection, and it carries D_7 to $\{g_t|_{C_0}\}$; as D_7 is discrete in its closure, condition (i) of Lemma 8.4 holds. For (ii), note that A_t lies in the component whose right endpoint is $\psi(s_t^\frown 1^\infty)$, and B_t in the component whose left endpoint is $\psi(s_t^\frown 0^\infty)$. These sequences are pairwise distinct for $s_t \in \Theta$, so each component of $[0,1] \setminus C_0$ contains at most one interval A_t and at most one interval B_t. For pairwise distinct t_k, eventually neither A_{t_k} nor B_{t_k} lies in a given component J, and then the function g_{t_k} is constant on $\overline{J}$. Every such limit is therefore constant on $\overline{J}$ and is determined by its values on C_0.

Finally, the photon and register preparations are given by distinct basis vectors. In the box, take the unit vectors $\sqrt{d\mu_{\rho_t}/dy}$. Within each of the D_3 and D_4 realizations, the supports lie in distinct Cantor gaps. For D_7, each gap contains at most one interval A_t and at most one interval B_u, by the distinctness of the endpoint sequences fixed in Section 4. If an internal gap contains both, its length is at least each of ℓ_{s_t} and ℓ_{s_u}; the interval A_t lies in its right half and B_u in its left half, so they meet in at most the midpoint. Each exterior gap contains at most one support interval. Within each realization, the supports of distinct preparations consequently meet in at most finitely many points, whence the vectors are pairwise orthogonal in $L^2[0,1]$. □

The realizations display what distinguishes the families.

- D_1, *escape to infinity*: an oscillator excited ever higher has a single idealized limit, and each actual preparation is isolated from it.
- D_2, *infinite configurations*: the idealized limits of the chains are the infinite spin configurations, a Cantor set of them, and countably many site probes tell them all apart, so the closure is metrizable; but any two configurations differ by 1 at some site, so the closure is not separable in the supremum norm.
- D_3 and D_4, *one-sided localization*: a particle that approaches a point from only one side produces one half of a split pair, and the split Cantor set appears; the closure is hereditarily separable but not metrizable.
- D_5, *point-supported limits*: light cones that shrink to a point on Bob's slice, or templates following longer and longer initial segments of one infinite configuration, yield a point-supported limiting profile; the corresponding idealized response evaluates the mass that a probe assigns to that point, and the selected response closure is not first countable.
- D_6, *twins over a metrizable base*: each infinite configuration has two idealized versions, one that every template misses and one that the template of the configuration itself detects; the scanning probes alone cannot tell the twins apart, and, as seen by these probes, the limits form a Cantor set.
- D_7, *twins over a split base*: a particle that approaches a point from both sides at once produces the midpoint of a split pair, as the symmetric state does in the Ising model (see Section 9 below); the twins of the closure sit over the split Cantor set.

In Proposition 8.6 and Corollary 8.7 below, we show that the scanning and template observables are NIP, which places them in the classification of Theorem 6.9; the combined observable is NIP by Proposition 8.6 as well, since its functions separate points and every sequence of them has a subsequence that converges pointwise on each of the two summands (see the proof of Corollary 8.7). The proof of Proposition 8.6 uses a theorem of Stegall. Recall that a subset of a Banach space is *weakly precompact* if it is bounded and every sequence in it has a weakly Cauchy subsequence, one whose values under each continuous linear functional form a Cauchy sequence, and that the *absolutely convex hull* of a set consists of the finite combinations $\sum_k a_k x_k$ of its elements with $\sum_k |a_k| \leq 1$. Stegall's theorem states that the absolutely convex hull of a weakly precompact set is weakly precompact (see Rodríguez [Rod21] for a quantitative version).

Proposition 8.6. *Let C be a label-controlled observable whose functions f_i separate the points of L and have the property that every sequence of them has a subsequence converging pointwise on L. Then the observable C is NIP for Alice's responses.*

Proof. For a density operator ρ, put $p_i = \langle e_i, \rho e_i \rangle$, a probability distribution on I, and $g_\rho = \sum_i p_i f_i$, a norm-convergent sum in $C(L)$; then $r_\rho(\nu) = \int g_\rho \, d\nu$. By dominated convergence, a sequence of functions f_i that converges pointwise on L is weakly Cauchy in $C(L)$; hence the set $\{f_i\}$ is weakly precompact. By Stegall's theorem, so is its absolutely convex hull, and so is the norm closure of that hull: given y_n in the closure, choose x_n in the hull with $\|x_n - y_n\| < 1/n$ and a weakly Cauchy subsequence of (x_n); the corresponding subsequence of (y_n) is weakly Cauchy. The partial sums of g_ρ lie in the hull, and every function g_ρ therefore lies in this closure. Consequently, every sequence (g_{ρ_n}) has a weakly Cauchy subsequence, and, since every $\nu \in P(L)$ is a continuous linear functional on $C(L)$, the corresponding subsequence of (r_{ρ_n}) converges pointwise on $P(L) = Y_C$. As in the proof of Lemma 8.3, an independent sequence has no such subsequence. □

Corollary 8.7.

(a) *The scanning observable is unstable and NIP$_3$, but not uniformly NIP.*
(b) *The template observable is NIP but not NIP$_1$.*
(c) *The scanning observable is not NIP for Bob's responses, that is, the observable obtained by exchanging the two tensor factors is not NIP for Alice's responses. Hence NIP for Alice's responses does not imply NIP for Bob's.*

Proof. (a) Every sequence of functions a_t on the countable set L_a has a pointwise convergent subsequence, by a diagonal argument, so the observable is NIP by Proposition 8.6. For a trace-norm dense sequence $(\rho_n)_{n\in\mathbb{N}}$, the closure of the responses is that of $\{r_{\rho_n}\}$ (see Lemma 6.3). Every function on the countable compact metrizable space L_a is Baire-1, so Lemma 8.3 applies and identifies this closure with the closure of $\{g_{\rho_n}\}$ in $\mathbb{R}^{L_a}$; since L_a is countable, this space is metrizable. For each positive integer m, the chains $t_1, \dots, t_m$ of length 2^m with $t_i(n)$ the i-th binary digit of n are shattered, with values 0 and 1, by the site probes $n < 2^m$; hence the observable is not uniformly NIP. Conversely, on any first m site probes, the chains of length m realize every binary pattern; the fat-shattering dimension of the response family is therefore infinite at every positive gap at most 1, with the gap convention of Remark 6.2. Finally, the observable is unstable: the functions a_{1^k} of the chains 1^k of k up spins, for $k \in \mathbb{N}$, converge pointwise to the indicator of $\mathbb{N}$, so the idealized response $\nu \mapsto \nu(\mathbb{N})$, which is not continuous on $P(L_a)$, is a limit of responses (see Theorem 6.1(a)).

(b) Let (t_k) be a sequence of labels. By compactness of L_b, a subsequence converges in L_b; if the limit is finite, the subsequence is eventually constant. If the limit is an infinite sequence σ, then $b_{t_k}(u) \to 0$ for finite u, and $b_{t_k}(\tau) \to 0$ for infinite $\tau \neq \sigma$, since eventually t_k agrees with σ beyond the first place where σ and τ differ; and, along a further subsequence, the value $b_{t_k}(\sigma)$ is constant. Hence the observable is NIP by Proposition 8.6. By Theorem 8.5, the closure of its responses contains a closed copy of $\hat{A}(2^{\mathbb{N}})$, which is not first countable.

(c) For $n \in \mathbb{N}$, consider the state of Bob's system in which his probe sits at the site n; these states form an independent sequence: for disjoint finite sets $E, F \subseteq \mathbb{N}$, choose a positive integer q larger than every member of $E \cup F$ and let Alice's chain t of length q have $t(n) = 1$ exactly for $n \in F$. Then $a_t(n) = 1$ for $n \in F$ and $a_t(n) = 0$ for $n \in E$. Alice's actual states restrict to points of the probe space of the exchanged observable (see Lemma 5.2(c)), so the family of Bob's responses has the independence property. This is the relation $n \in u$ of Remark 3.8, realized by a physical observable. □

The scanning observable shows the difference between NIP and uniform NIP (see Remark 6.2): its idealized responses form a metrizable compactum, yet the response family has infinite fat-shattering dimension at a fixed positive gap. In distribution-free learning from independent samples, the aim is to predict an unknown response with small mean absolute error under the probe distribution, with a prescribed probability of success. No sample bound depending only on the required accuracy and probability of success can achieve this for the scanning family uniformly over target responses and probe distributions [ABDCBH97]. The obstruction is already visible on the first m sites, where chains of length m realize every binary pattern.

9. Spontaneous symmetry breaking is a split pair

In this section and the next we show that the levels of Section 4 have a physical meaning beyond bipartite observables. Several notions of physics rest on an order of limits of a family of responses: a preparation, such as a finite sample, has a response, such as its magnetization

as a function of an external field, and the notion depends on whether the sample becomes infinite before or after the field is switched off.

The common setting is a pointwise bounded family $\mathcal{R}$ of continuous functions on a compact metrizable probe space Y; its idealized responses are the members of the pointwise closure $\overline{\mathcal{R}}$. We call the family *stable* when, for every sequence $(r_k)_{k\in\mathbb{N}}$ in $\mathcal{R}$ and every sequence $(y_l)_{l\in\mathbb{N}}$ in Y, the iterated limits $\lim_l \lim_k r_k(y_l)$ and $\lim_k \lim_l r_k(y_l)$ agree whenever both exist. For a uniformly bounded family, as for the magnetizations below, Grothendieck's double limit criterion [Gro52] identifies stability with continuity of every idealized response; countably many probes then separate the closure, which is metrizable by Proposition 4.5. The NIP characterization of Theorem 3.6 applies to pointwise bounded families, and Definition 4.4 assigns their finer levels when the closure is separable. For a fixed observable, convexity follows from allowing all mixed preparations; for the parameterized families considered here, we compute the closures directly from the stated properties of the physical models.

The standard example of an order of limits is spontaneous symmetry breaking: for a ferromagnet below its critical temperature, letting the external field vanish after the volume becomes infinite yields a nonzero magnetization, whereas the other order yields zero [FV17]. In Theorem 9.2 below, we show that this is an instability in our sense, absent above the critical temperature, and that, with an offset field that moves the transition point (the field at which the magnetization changes sign), the family of responses below the critical temperature is NIP_1 but not NIP_2: the idealized responses of the two pure phases, the equilibrium states magnetized in opposite directions, form a split pair, two functions that differ only at the transition point, and the idealized response of the symmetric state, the equal mixture of the two phases, is their midpoint, as in the proof of Theorem 6.6.

9.1. **The Ising model.** We recall the facts we need; all of them can be found in the book of Friedli and Velenik [FV17]. Fix a dimension $d \geq 2$ and a dimensionless inverse temperature $\beta > 0$, the ratio of the coupling energy of two neighboring spins to the thermal energy $k_B T$, where T is the temperature and k_B is the Boltzmann constant. For $N \geq 1$, let $\Lambda_N = \{-N, \dots, N\}^d$ with periodic boundary conditions, and for a real field h let

$$Z_N(h) = \sum_{\sigma \in \{-1,1\}^{\Lambda_N}} \exp\Big(\beta \sum_{x\sim y} \sigma_x \sigma_y + h \sum_x \sigma_x\Big), \qquad p_N(h) = \frac{1}{|\Lambda_N|} \log Z_N(h),$$

where $x \sim y$ ranges over the nearest-neighbor pairs of the torus; in this convention the field h already includes the factor β. The magnetization per site in the Gibbs state, the probability measure on configurations σ proportional to the corresponding term of $Z_N(h)$, is $m_N(h) = p_N'(h)$. The normalizing constant $Z_N(h)$ of the Gibbs state is the *partition function*, and the function p_N is the *pressure* of the finite system.

(F1) The function p_N is smooth, convex and even. Hence the function m_N is continuous, nondecreasing and odd, with $m_N(0) = 0$. The bound $|m_N| \leq 1$ holds because $m_N(h)$ is the mean, in the Gibbs state, of the average spin $|\Lambda_N|^{-1} \sum_x \sigma_x$, which lies in $[-1, 1]$.

(F2) The limit $p(h) = \lim_N p_N(h)$, the pressure, exists for every h; it is convex and even, and it is differentiable at every $h \neq 0$, by the Lee–Yang theorem [LY52]. Since p_N and p are convex, $m_N(h) \to p'(h)$ at every $h \neq 0$, by Griffiths' lemma on derivatives of convex functions. Indeed, convexity bounds $p_N'(h)$ between the secant slopes over $[h-\eta, h]$ and $[h, h+\eta]$ for $\eta > 0$; first let $N \to \infty$, then $\eta \to 0$ at a differentiability point of p.

(F3) The spontaneous magnetization $m^*(\beta) = p'(0^+)$ is the right derivative of p at 0, and $p'(0^-) = -m^*(\beta)$. The critical inverse temperature $\beta_c = \inf\{\beta : m^*(\beta) > 0\}$ is positive, and it is finite by Peierls's argument [Pei36, Gri64]; since $m^*(\beta)$ is the magnetization

of the plus state (the infinite-volume limit of the Gibbs states with all boundary spins fixed up), it is nondecreasing in β, by the Griffiths–Kelly–Sherman inequalities, and $m^*(\beta) > 0$ for $\beta > \beta_c$ (see [FV17, Chapter 3]).

As p' is nondecreasing and odd, (F3) implies that $p'(t) \geq m^*$ for $t > 0$ and $p'(t) \leq -m^*$ for $t < 0$.

9.2. **Preparations and probes.** A preparation consists of a finite system Λ_N and an offset field $a \in [-1, 1]$, that is, a uniform field built into the sample; a probe is an applied field $h \in Y = [-2, 2]$. The system then feels the field $h - a$, and the response of the preparation (N, a) is the function

$$r_{N,a}(h) = m_N(h - a) \qquad (h \in Y).$$

The offset is a device that moves the transition point; without it, the family $\{m_N\}$ has a single transition point, at $h = 0$. Let $\mathcal{R} = \{r_{N,a} : N \geq 1,\ a \in [-1, 1]\}$. Each $r_{N,a}$ is continuous and nondecreasing, and $r_{N,a'} \to r_{N,a}$ uniformly as $a' \to a$, because m_N is uniformly continuous on compact sets. The responses with rational a are therefore dense in $\mathcal{R}$, and the closure $\overline{\mathcal{R}}$ is separable. Every function in $\overline{\mathcal{R}}$ is nondecreasing, and so of the first Baire class; thus $\overline{\mathcal{R}}$ is a Rosenthal compactum, and, by Theorem 3.6, the family $\mathcal{R}$ is NIP.

We write $m^* = m^*(\beta)$, and for $a \in [-1, 1]$ and $c \in [-m^*, m^*]$ we let $g_{a,c} \colon Y \to \mathbb{R}$ be given by $g_{a,c}(h) = p'(h - a)$ for $h \neq a$ and $g_{a,c}(a) = c$. These functions describe the idealized responses.

Lemma 9.1. *The following hold:*

(i) *every $g_{a,c}$ with $-1 < a < 1$ belongs to $\overline{\mathcal{R}}$;*
(ii) *every function in $\overline{\mathcal{R}} \setminus \mathcal{R}$ equals $g_{a,c}$ for some $a \in [-1, 1]$ and $c \in [-m^*, m^*]$.*

Proof. (i) Let $-1 < a < 1$. If $m^* > 0$, then, since $\overline{\mathcal{R}}$ is closed and $g_{a,c} \to g_{a,\pm m^*}$ pointwise as $c \to \pm m^*$, it suffices to take $|c| < m^*$, and by symmetry $0 \leq c < m^*$; if $m^* = 0$, then $c = 0$. If $c = 0$, then $r_{N,a} \to g_{a,0}$, by (F1) at $h = a$ and by (F2) elsewhere. If $0 < c < m^*$, fix $\eta > 0$; by (F2), $m_N(\eta) \to p'(\eta) \geq m^* > c$, so for all large N, since $m_N(0) = 0$, the intermediate value theorem yields $\varepsilon_N \in (0, \eta)$ with $m_N(\varepsilon_N) = c$. Choose such roots in $(0, 1)$ for all sufficiently large N. For each fixed $\eta \in (0, 1)$, eventually $m_N(\eta) > c$, so monotonicity forces every chosen root below η; hence $\varepsilon_N \to 0$. Put $a_N = a - \varepsilon_N$. Then $r_{N,a_N}(a) = m_N(\varepsilon_N) = c$. For $h \neq a$, let $t = h - a$ and fix η' with $0 < \eta' < |t|$; for large N, we have $t - \eta' \leq h - a_N \leq t + \eta'$, and hence, by monotonicity, $m_N(t - \eta') \leq r_{N,a_N}(h) \leq m_N(t + \eta')$. By (F2), the outer terms converge to $p'(t - \eta')$ and $p'(t + \eta')$, which tend to $p'(t)$ as $\eta' \to 0$, because p' is continuous at $t \neq 0$. Finally, as $a > -1$, the offsets a_N lie in $[-1, 1]$ for large N, and $r_{N,a_N} \to g_{a,c}$.

(ii) Let $g \in \overline{\mathcal{R}} \setminus \mathcal{R}$. Since $\overline{\mathcal{R}}$ is a Rosenthal compactum, it is Fréchet–Urysohn [BFT78, Theorem 3F], so the function g is the limit of a sequence r_{N_k,a_k}. Passing to a subsequence, we may assume that $a_k \to a$ and that either N_k is constant or $N_k \to \infty$. In the first case, uniform continuity forces $g = r_{N,a} \in \mathcal{R}$, which is excluded. In the second case, the argument of the previous paragraph shows that $g(h) = p'(h - a)$ for $h \neq a$. Finally, given $\eta > 0$, we have $|a - a_k| < \eta$ for all large k, and then $|r_{N_k,a_k}(a)| = |m_{N_k}(a - a_k)| \leq m_{N_k}(\eta)$, because m_{N_k} is odd and nondecreasing; the right side tends to $p'(\eta)$, and $p'(\eta) \to m^*$ as $\eta \to 0$. Hence $|g(a)| \leq m^*$, and $g = g_{a,g(a)}$. □

9.3. **The classification detects the phase transition.** By Lemma 9.1, the classification of the family $\mathcal{R}$ of responses with offsets reduces to the behavior of p' at 0.

Theorem 9.2. *Let $d \geq 2$ and $\beta > 0$ be given, let $m^*(\beta)$ be the spontaneous magnetization and β_c the critical inverse temperature of (F3), and let $\mathcal{R}$ be the family of responses $r_{N,a}$ with offsets.*

(a) *If $m^*(\beta) = 0$, in particular if $\beta < \beta_c$, then every idealized response is continuous: the family $\mathcal{R}$ is stable, and $\overline{\mathcal{R}}$ is metrizable.*

(b) *If $m^*(\beta) > 0$, in particular if $\beta > \beta_c$, then the family $\mathcal{R}$ is unstable, and $\overline{\mathcal{R}}$ is first countable but not hereditarily separable: the family is NIP$_1$ but not NIP$_2$.*

Already the family $\{m_N\}$ without offsets is stable exactly when $m^*(\beta) = 0$; it is NIP$_3$ for every β.

Proof. (a) If $m^* = 0$, then the pressure p is differentiable everywhere, and its derivative p', being the derivative of a convex function, is continuous. By Lemma 9.1(ii), every idealized response outside $\mathcal{R}$ has the form $g_{a,0}(h) = p'(h - a)$ and is therefore continuous; so are the responses in $\mathcal{R}$. By Grothendieck's theorem [Gro52], a family of continuous functions on a compact space whose pointwise closure consists of continuous functions satisfies the double limit condition, so the family $\mathcal{R}$ is stable. The closure is metrizable by the argument of Proposition 6.5: evaluation at the rational points of Y is a continuous injection on continuous functions.

(b) Let $m^* > 0$. For the family without offsets, take $h_l = 1/l$ for $l \geq 1$ and a sequence $(N_k)_{k\in\mathbb{N}}$ of positive integers with $N_k \to \infty$. By (F2) and (F3), $\lim_l \lim_k m_{N_k}(h_l) = \lim_l p'(1/l) = m^*$, whereas, by continuity of each m_N and (F1), $\lim_k \lim_l m_{N_k}(h_l) = \lim_k m_{N_k}(0) = 0$. The family $\{m_N\}$, and with it the family $\mathcal{R}$, is therefore unstable.

Every function in $\overline{\mathcal{R}}$ is nondecreasing, and the argument in the proof of Theorem 7.8(b) shows that the space of nondecreasing functions $Y \to [-1, 1]$ is first countable; hence so is $\overline{\mathcal{R}}$. For $a \in (-1, 1)$, the neighborhood $\{g : |g(a)| < m^*/2\}$ of $g_{a,0}$ contains no $g_{b,0}$ with $b \neq a$, because $|g_{b,0}(a)| = |p'(a - b)| \geq m^*$. The set $\{g_{a,0} : -1 < a < 1\}$ is thus an uncountable discrete subspace, and $\overline{\mathcal{R}}$ is not hereditarily separable.

The first part of the last statement follows from part (a), since $\{m_N\} = \{r_{N,0}\}$ is a subfamily of $\mathcal{R}$, and from the computation at the beginning of (b). For the second part, the sequence (m_N) converges pointwise, by (F1) and (F2), so its closure is countable, and a countable compact Hausdorff space is metrizable. □

Physically, the proof tells a familiar story. At the transition point $h = a$, the idealized responses take every value c between $-m^*$ and m^*. The extreme values $\pm m^*$ are the magnetizations of the two pure phases, obtained by letting the field reach the transition from one side after the volume has become infinite; they are the values at a of the idealized responses g_{a,m^*} and $g_{a,-m^*}$, which form a split pair. The value $c = (2\lambda - 1)m^*$ is the magnetization of the mixture of the two phases with weights λ and $1 - \lambda$, for $0 \leq \lambda \leq 1$, and it is reached through finite-size rounding of the transition, the smoothing of the jump of the magnetization in a finite sample [BK90], by tuning the offset as the volume grows. The uncountable discrete set that destroys hereditary separability consists of the symmetric responses $g_{a,0}$, the midpoints of the split pairs, exactly as in the proof of Theorem 6.6. Above the critical temperature there is no split pair, and the idealized responses are continuous. In this sense the classification detects the phase transition: crossing β_c moves the family with offsets from the stable, metrizable class to the class of NIP$_1$ families that are not NIP$_2$, the class of the compactum of idealized responses of the comparison of positions in a box, a closed subspace

of the Helly space. Without offsets, the family stays NIP_3, and only stability registers the transition.

9.4. **The quantum Ising model.** In Theorem 9.6 below, we show that the analogue of Theorem 9.2 holds for a Hamiltonian whose terms do not commute: the Ising model in a transverse field. The only ingredient of the classical proof that requires an argument for this model is the differentiability of the pressure away from zero field. It can be derived from the Griffiths–Hurst–Sherman inequality, which holds for this model through its space-time representation, a description of the quantum model by random configurations in one more dimension, that of imaginary time [BG09, Lemma 4.6]; we derive it instead from a Lee–Yang theorem, which yields more (see Propositions 9.4 and 9.5 below): the partition function has no zeros off the imaginary axis of the complex longitudinal field (the field along the axis of the couplings), and the pressure is real analytic away from zero field.

Fix $d \geq 1$, a coupling $J > 0$, a transverse field $\Gamma \geq 0$ and an inverse temperature $\beta > 0$; the coupling and the transverse field are energies, and the inverse temperature is the reciprocal of the thermal energy $k_B T$, so that βJ and $\beta\Gamma$ are pure numbers. On the torus $\Lambda_N = \{-N, \dots, N\}^d$ of Section 9, now for every $d \geq 1$, with Hilbert space $(\mathbb{C}^2)^{\otimes \Lambda_N}$, the Pauli matrices $\sigma^x = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$ and $\sigma^z = \left(\begin{smallmatrix} 1 & 0 \\ 0 & -1 \end{smallmatrix}\right)$ acting at each site i as σ_i^x and σ_i^z (on the tensor factor of site i, with the identity on the other factors), and $S_N = \sum_i \sigma_i^z$, let

$$H_N = -J \sum_{i \sim j} \sigma_i^z \sigma_j^z - \Gamma \sum_i \sigma_i^x, \qquad Z_N(h) = \operatorname{Tr} \exp\bigl(-\beta H_N + h S_N\bigr),$$

for complex h; as in Section 9, the longitudinal field h includes the factor β. For real h put $p_N(h) = |\Lambda_N|^{-1} \log Z_N(h)$ and $m_N = p_N'$. For $\Gamma = 0$ this is the classical model with βJ in place of β.

(Q1) The function p_N is smooth, convex and even, and $m_N(h)$ is the expectation of $S_N/|\Lambda_N|$ in the Gibbs state, the density operator $e^{-\beta H_N + h S_N}/Z_N(h)$; hence the function m_N is continuous, nondecreasing and odd, with $|m_N| \leq 1$, because $\|S_N\| = |\Lambda_N|$, and $m_N(0) = 0$. For self-adjoint matrices X and S and real h, write $A = X + hS$. The matrix derivative formula

$$\frac{d}{dh} e^{X+hS} = \int_0^1 e^{(1-t)A} S e^{tA}\, dt$$

and cyclicity of the trace give $\frac{d}{dh} \operatorname{Tr} e^{X+hS} = \operatorname{Tr}(S e^{X+hS})$. By the Gibbs variational formula, the function $h \mapsto \log \operatorname{Tr} e^{X+hS} = \sup_\omega \{\operatorname{Tr} \omega(X + hS) - \operatorname{Tr} \omega \log \omega\}$, the supremum over density operators ω, is convex as a supremum of affine functions. Here $-\operatorname{Tr} \omega \log \omega$ is the entropy of the density operator ω, with $0 \log 0 = 0$. The unitary $U = \prod_i \sigma_i^x$ commutes with H_N and satisfies $U S_N U^* = -S_N$, so $Z_N(-h) = Z_N(h)$ for all complex h.

(Q2) The limit $p(h) = \lim_N p_N(h)$ exists for every real h; it is convex and even. For free boundary conditions, in which the bonds that wrap around the torus are omitted, the existence is standard [Sim93], and the periodic Hamiltonian differs in operator norm from the free one by at most J times the number of these bonds, which is at most $d(2N+1)^{d-1}$, so the two pressures differ by at most $\beta J d/(2N+1)$, because $|\log \operatorname{Tr} e^A - \log \operatorname{Tr} e^B| \leq \|A - B\|$ for self-adjoint matrices A and B.

In the proof of Proposition 9.4 below, we show, by the Suzuki–Trotter mapping [Suz76] between quantum spin systems and classical Ising systems in one more dimension, that the partition function of the quantum model is a limit of positive multiples of partition functions

of classical Ising models on a larger graph, in which each site of the torus is repeated M times, for a positive integer M that tends to infinity, one copy for each slice of "imaginary time". All couplings, between neighboring sites of the torus and between consecutive copies of a site, are ferromagnetic, so the classical Lee–Yang theorem applies to each of these classical models: none of their partition functions vanishes off the imaginary axis. A limit of holomorphic functions without zeros has no zeros either, unless it vanishes identically (see Fact 9.3(b)), and so the quantum partition function inherits the zero-free half-planes. The analyticity of the pressure then follows, in Proposition 9.5, from a uniform bound and two classical theorems of complex analysis. The two proofs use only the following standard facts.

Fact 9.3.

(a) Lee–Yang theorem [LY52] *(see* [FV17, Chapter 3]*). Let G be a finite graph with vertex set V and edge set E, let $J_e \geq 0$ for $e \in E$, and let h_v be complex numbers with $\operatorname{Re} h_v > 0$ for $v \in V$. Then $\sum_{s\in\{-1,1\}^V} \exp\bigl(\sum_{e=\{u,w\}\in E} J_e s_u s_w + \sum_{v\in V} h_v s_v\bigr) \neq 0$.*

(b) Hurwitz's theorem [Tit39]. *If holomorphic functions without zeros on a connected open set $U \subseteq \mathbb{C}$ converge uniformly on compact subsets of U to f, then either f has no zeros in U or f vanishes identically.*

(c) Borel–Carathéodory inequality [Tit39]. *If f is holomorphic on a neighborhood of the closed disc $|z - z_0| \leq R$ and $\operatorname{Re} f \leq A$ there, then $|f(z) - f(z_0)| \leq \frac{2r}{R-r}\bigl(A - \operatorname{Re} f(z_0)\bigr)$ for $|z - z_0| \leq r < R$.*

(d) Vitali's theorem [Tit39]. *A sequence of holomorphic functions on a connected open set U that is uniformly bounded on each compact subset of U and converges at every point of a set with an accumulation point in U converges uniformly on compact subsets of U; by Weierstrass's theorem, so do the derivatives.*

Proposition 9.4 (Lee–Yang theorem for the quantum Ising model). *For every N, the function Z_N is entire, $Z_N(h) \neq 0$ whenever $\operatorname{Re} h \neq 0$, and $|Z_N(h)| \leq Z_N(\operatorname{Re} h)$.*

Proof. Write $X = \beta J \sum_{i\sim j} \sigma_i^z \sigma_j^z + h S_N$, which is diagonal in the basis $\{|s\rangle : s \in \{-1,1\}^{\Lambda_N}\}$ of eigenvectors of the operators σ_i^z, and $W = \beta\Gamma \sum_i \sigma_i^x$. The function $Z_N(h) = \operatorname{Tr} e^{X+W}$ is entire, because the exponential series converges locally uniformly in h.

Step 1: the Lie–Trotter formula. For matrices A and B, we have $e^{A/M}e^{B/M} = I + (A+B)/M + O(M^{-2})$, uniformly for A and B in a bounded set; by taking the principal logarithm, we obtain $e^{A/M}e^{B/M} = \exp\bigl((A+B)/M + R_M\bigr)$ with $\|R_M\| = O(M^{-2})$, and hence $(e^{A/M}e^{B/M})^M = \exp(A + B + MR_M) \to e^{A+B}$, uniformly for A and B in a bounded set. Applied to X and W, this yields $Z_{N,M}(h) = \operatorname{Tr}(e^{X/M}e^{W/M})^M \to Z_N(h)$ as $M \to \infty$, uniformly for h in compact subsets of $\mathbb{C}$.

Step 2: a classical ferromagnet. Suppose first that $\Gamma > 0$, and put $\varepsilon = \beta\Gamma/M$. Since $e^{\varepsilon\sigma^x} = \cosh\varepsilon\, I + \sinh\varepsilon\, \sigma^x$, the matrix element $\langle s|e^{\varepsilon\sigma^x}|s'\rangle$ of a single spin is $\cosh\varepsilon$ if $s = s'$ and $\sinh\varepsilon$ if $s \neq s'$, that is, $\langle s|e^{\varepsilon\sigma^x}|s'\rangle = c_\varepsilon e^{J_\varepsilon s s'}$ with $c_\varepsilon = (\sinh\varepsilon\cosh\varepsilon)^{1/2} > 0$ and $J_\varepsilon = \frac{1}{2}\log\coth\varepsilon > 0$. Insertion of the basis between the factors of $(e^{X/M}e^{W/M})^M$ shows that, for $M \geq 3$,

$$Z_{N,M}(h) = c_\varepsilon^{M|\Lambda_N|} \sum_{s^1,\dots,s^M} \exp \sum_{k=1}^{M} \Bigl(\frac{\beta J}{M} \sum_{i\sim j} s_i^k s_j^k + J_\varepsilon \sum_i s_i^k s_i^{k+1} + \frac{h}{M} \sum_i s_i^k \Bigr),$$

where $s^1, \dots, s^M$ range over $\{-1,1\}^{\Lambda_N}$ and $s^{M+1} = s^1$. This is $c_\varepsilon^{M|\Lambda_N|}$ times the partition function of a classical Ising model on the finite graph $\Lambda_N \times \{1,\dots,M\}$, with nonnegative couplings and the same field h/M at every site. By Fact 9.3(a), it does not vanish when

$\operatorname{Re} h > 0$. Moreover, by the triangle inequality applied to the sum, $|Z_{N,M}(h)| \le Z_{N,M}(\operatorname{Re} h)$. If $\Gamma = 0$, then the function Z_N itself is a classical Ising partition function, and the same two statements hold directly.

Step 3: Hurwitz's theorem. The functions $Z_{N,M}$ have no zeros in the connected open set $\{\operatorname{Re} h > 0\}$ and converge to Z_N uniformly on its compact subsets, and Z_N is not identically zero there, since $Z_N(1) > 0$. By Fact 9.3(b), the function Z_N has no zeros in $\{\operatorname{Re} h > 0\}$, and by the evenness in (Q1) none in $\{\operatorname{Re} h < 0\}$. The inequality $|Z_N(h)| \le Z_N(\operatorname{Re} h)$ follows from Step 2 in the limit. □

Proposition 9.5. *The pressure p is real analytic on $\mathbb{R} \setminus \{0\}$, and $m_N(h) \to p'(h)$ for every $h \neq 0$.*

Proof. By evenness it suffices to treat $h > 0$. The half-plane $\{\operatorname{Re} h > 0\}$ is simply connected and Z_N does not vanish there, so the function Z_N has a holomorphic logarithm on it, and we let f_N be $|\Lambda_N|^{-1}$ times the logarithm that is real on $(0,\infty)$; thus $f_N = p_N$ on $(0,\infty)$ and $\operatorname{Re} f_N(h) = |\Lambda_N|^{-1} \log |Z_N(h)| \le p_N(\operatorname{Re} h)$, by Proposition 9.4.

We bound p_N on the real line uniformly in N. The torus has $d|\Lambda_N|$ bonds, whence $\|\beta H_N\| \le \beta(dJ + \Gamma)|\Lambda_N|$. Since the trace of the exponential of a self-adjoint matrix of size $2^{|\Lambda_N|}$ and norm at most c lies between $2^{|\Lambda_N|}e^{-c}$ and $2^{|\Lambda_N|}e^{c}$, it follows that $|p_N(0) - \log 2| \le \beta(dJ + \Gamma) =: b$; as $|p_N'| \le 1$, also $|p_N(x) - \log 2| \le b + |x|$ for real x.

Fix $x_0 > 0$, and put $R = x_0/2$ and $r = x_0/4$. On the disc $|h - x_0| \le R$ we have $\operatorname{Re} f_N \le \log 2 + b + \frac{3}{2}x_0$, and $f_N(x_0) = p_N(x_0) \ge \log 2 - b - x_0$. By the Borel–Carathéodory inequality (see Fact 9.3(c)),

$$|f_N(h) - f_N(x_0)| \le \frac{2r}{R - r}\Big(\sup_{|z - x_0| \le R} \operatorname{Re} f_N(z) - \operatorname{Re} f_N(x_0)\Big) \le 2\big(2b + \tfrac{5}{2}x_0\big)$$

for $|h - x_0| \le r$. Since also $|f_N(x_0)| \le \log 2 + b + x_0$, the functions f_N are uniformly bounded on each disc $\{|h - x_0| < x_0/4\}$, and therefore locally uniformly bounded on the connected open set U that is the union of these discs, which contains $(0,\infty)$. They converge on $(0,\infty)$, by (Q2). By Vitali's theorem (see Fact 9.3(d)), they converge uniformly on compact subsets of U to a holomorphic function F, which equals p on $(0,\infty)$; hence the pressure p is real analytic there, and $m_N = f_N' \to F' = p'$ on $(0,\infty)$. □

Put $m^*(\beta,\Gamma) = p'(0^+)$, the spontaneous magnetization; by evenness, $p'(0^-) = -m^*(\beta,\Gamma)$. With offsets as in Section 9, let $r_{N,a}(h) = m_N(h - a)$ for $N \ge 1$, $a \in [-1,1]$ and $h \in Y = [-2,2]$, and let $\mathcal{R}_\Gamma$ be the family of these functions.

Theorem 9.6. *Let $d \ge 1$, $J > 0$, $\Gamma \ge 0$ and $\beta > 0$ be given, let $m^*(\beta,\Gamma)$ be the spontaneous magnetization of the quantum model, and let $\mathcal{R}_\Gamma$ be its family of responses $r_{N,a}$ with offsets.*

(a) *If $m^*(\beta,\Gamma) = 0$, then every idealized response is continuous: the family $\mathcal{R}_\Gamma$ is stable, and its closure is metrizable.*

(b) *If $m^*(\beta,\Gamma) > 0$, then the family $\mathcal{R}_\Gamma$ is unstable, and its closure is first countable but not hereditarily separable: the family is NIP$_1$ but not NIP$_2$. For each $a \in (-1,1)$, the functions g_{a,m^*} and $g_{a,-m^*}$, defined as before Lemma 9.1 with $m^* = m^*(\beta,\Gamma)$ and with the pressure p of this model, lie in the closure and differ only at a, and their midpoints $g_{a,0}$ form an uncountable discrete set.*

The family $\{m_N\}$ without offsets is stable exactly when $m^*(\beta,\Gamma) = 0$, and it is NIP$_3$ in all cases.

Proof. The discussion preceding Lemma 9.1, the lemma and the proof of Theorem 9.2 use only the following properties: the function m_N is continuous, nondecreasing and odd, with $|m_N| \le 1$ and $m_N(0) = 0$; the limit p exists, is convex and even, and is differentiable at every $h \neq 0$, with $m_N(h) \to p'(h)$; and $p'(0^{\pm}) = \pm m^*$. These hold by (Q1), (Q2) and Proposition 9.5, and the proofs apply verbatim with $m^*(\beta, \Gamma)$ in place of $m^*(\beta)$. □

In physical terms, the values $g_{a,\pm m^*}(a) = \pm m^*$ are the magnetizations of the two pure phases at the transition point $h = a$, and the value $g_{a,0}(a) = 0$ is that of the symmetric state; the classification thus reads off the phase diagram of the quantum model at positive temperature. Both cases of the theorem occur: Björnberg and Grimmett [BG09] write the Hamiltonian as $-\frac{1}{2}\lambda \sum \sigma^z\sigma^z - \delta \sum \sigma^x - \gamma \sum \sigma^z$, in which the parameters are $\lambda = 2J$, $\delta = \Gamma$ and $h = \beta\gamma$, and they define the spontaneous magnetization as the limit, as $\gamma \downarrow 0$, of the infinite-volume magnetization at field γ on the same periodic boxes Λ_N, through the space-time representation. By translation invariance and Proposition 9.5, that magnetization is $p'(\beta\gamma)$, and since the right derivative of a convex function is the right limit of its derivative, their spontaneous magnetization is $p'(0^+) = m^*(\beta, \Gamma)$. For $d = 1$ and $\beta < \infty$ it vanishes, and for $d \ge 2$ and $\Gamma > 0$ there is a critical value of $2J/\Gamma$, depending on $\beta\Gamma$, below which $m^* = 0$ and above which $m^* > 0$ (see [BG09, Section 1 and Theorems 1.1–1.2], where the existence of the critical value is attributed to standard methods). Order at low temperature and small Γ/J goes back to Ginibre [Gin69].

The Lee–Yang theorem [LY52] was extended to general classical ferromagnets by Newman [New74] and by Lieb and Sokal [LS81], and Lee–Yang theorems for quantum spin systems go back to Asano [Asa70] and to Suzuki and Fisher [SF71]. The formulation in Fröhlich and Rodriguez [FR12, Theorem 5], applied to H_N, gives the sufficient zero-free region $\operatorname{Re} h > \beta\Gamma$. In Proposition 9.4 we include a self-contained proof of the whole-half-plane statement for the transverse-field Ising model, by a Suzuki–Trotter approximation by classical ferromagnets followed by Hurwitz's theorem; Proposition 9.5 gives the analytic input needed for the response classification. The argument uses standard quantum Lee–Yang methods, in the spirit of Simon and Griffiths [SG73] and of Suzuki [Suz76].

10. Chiral symmetry breaking and metallic conduction are instabilities

The chiral-condensate and current-response applications distinguish failure of stability from failure of metrizability. For the response families considered below, the limiting response can be discontinuous at one probe value while the full pointwise closure remains metrizable. Under the spectral hypotheses stated below, we compute the discrepancy between the iterated limits from the limiting density of Dirac eigenvalues at zero for the condensate and from the mass at zero of a limiting weighted excitation measure for the current response (see Theorems 10.3 and 10.4). The density describes the near-zero spectrum of the Dirac operator governing quarks in a gluon field; the atom describes the current spectral weight accumulating at zero excitation energy. A common lemma places both response families in NIP_3 and identifies stability with continuity at the exceptional probe value.

Lemma 10.1. *Let Q be a compact metrizable space, q_0 a point of Q that is not isolated, and $q \mapsto g_q$ a map from Q into $\mathbb{R}^{[0,1]}$ that is continuous for the topology of pointwise convergence. Suppose that g_q is continuous for $q \neq q_0$, that $G = g_{q_0}$ is continuous on $(0, 1]$, and that the one-sided limit $G(0^+) = \lim_{s\to 0^+} G(s)$ exists as a finite real number. Then the family $\mathcal{R} = \{g_q : q \neq q_0\}$ is NIP$_3$, and it is stable if and only if G is continuous at 0. For every sequence $(q_k)_{k\in\mathbb{N}}$ in $Q \setminus \{q_0\}$ converging to q_0 and every sequence $(s_l)_{l\in\mathbb{N}}$ in $(0, 1]$ converging*

to 0*, we have*

$$\lim_l \lim_k g_{q_k}(s_l) = G(0^+), \qquad \lim_k \lim_l g_{q_k}(s_l) = G(0).$$

Proof. The image of Q is compact, contains $\mathcal{R}$ and is contained in $\overline{\mathcal{R}}$, because q_0 is not isolated; it is thus $\overline{\mathcal{R}}$. Its members are continuous on $(0,1]$, so two distinct members differ at a rational point of $[0,1]$, and $\overline{\mathcal{R}}$ is metrizable by Proposition 4.5. The function G is of the first Baire class, as the pointwise limit of the continuous functions that equal G on $[1/n,1]$ and are linear on $[0,1/n]$ with value $G(0)$ at 0. For each $s \in [0,1]$, the function $q \mapsto g_q(s)$ is continuous on the compact space Q, so the family $\mathcal{R}$ is pointwise bounded. Every member of its closure is of the first Baire class, and Theorem 3.6 therefore shows that $\mathcal{R}$ is NIP; its closure is compact metrizable and hence separable, so it is NIP_3.

Suppose first that G is continuous at 0, so that every g_q, including g_{q_0}, is continuous on $[0,1]$. To verify stability, let $(p_k)_{k\in\mathbb{N}}$ be a sequence in $Q \setminus \{q_0\}$ and $(t_l)_{l\in\mathbb{N}}$ a sequence in $[0,1]$ for which both iterated limits of $g_{p_k}(t_l)$ exist as real numbers. By compact metrizability, we may pass to subsequences for which $p_k \to p_*$ in Q and $t_l \to t_*$ in $[0,1]$; these passages do not change either existing iterated limit. Pointwise continuity in the parameter gives $\lim_k g_{p_k}(t_l) = g_{p_*}(t_l)$ for each l, while continuity in the probe gives $\lim_l g_{p_k}(t_l) = g_{p_k}(t_*)$ for each k. Consequently,

$$\lim_l \lim_k g_{p_k}(t_l) = \lim_l g_{p_*}(t_l) = g_{p_*}(t_*), \qquad \lim_k \lim_l g_{p_k}(t_l) = \lim_k g_{p_k}(t_*) = g_{p_*}(t_*).$$

Thus the two iterated limits agree, as required.

Now take sequences (q_k) and (s_l) as in the statement. For each l, pointwise continuity in the parameter gives $\lim_k g_{q_k}(s_l) = G(s_l)$; for each k, continuity of g_{q_k} at 0 gives $\lim_l g_{q_k}(s_l) = g_{q_k}(0)$. It follows that

$$\lim_l \lim_k g_{q_k}(s_l) = \lim_l G(s_l) = G(0^+), \qquad \lim_k \lim_l g_{q_k}(s_l) = \lim_k g_{q_k}(0) = G(0).$$

Such a sequence (q_k) exists because q_0 is not isolated and Q is metrizable, and we may take $s_l = 1/(l+1)$. If G is discontinuous at 0, then $G(0^+) \neq G(0)$; the two displayed limits are finite and unequal, so $\mathcal{R}$ is not stable. □

10.1. **The chiral condensate.** We work with a lattice regularization, in which Euclidean spacetime (spacetime with imaginary time) is replaced by a lattice of points in a box of volume V; we use lattice units, with $\hbar = c = 1$ and lattice spacing 1, in which masses and volumes are pure numbers. The quarks are described by an anti-self-adjoint Dirac operator D_V, as for staggered fermions, a standard discretization of quarks on the lattice; the Dirac operator governs the motion of the quarks in a given gauge field, the gluon field with which they interact. For each gauge field in the chosen ensemble, we assume that the spectrum consists of nonzero pairs $\pm i\lambda_n$, indexed by n in a finite set, with $\lambda_n > 0$. This assumption excludes exact zero modes, eigenvectors with eigenvalue 0; if a finite-volume operator has n_0 such modes, they contribute $n_0/(mV)$ to its condensate at mass $m > 0$ before averaging over gauge fields. The spectral density is averaged over gauge fields with a measure that does not depend on the mass m: either the quenched theory, in which the quarks do not act back on the gauge field, or the valence condensate at a fixed sea quark mass, a standard setting for the Banks–Casher relation, whose original form uses a measure that depends on m; here the valence quarks are those of the condensate, whose mass m varies, and the sea quarks are

those that act back on the gauge field. The chiral condensate at quark mass $m > 0$ is

$$\Sigma_V(m) = \Big\langle \frac{1}{V} \operatorname{Tr}(D_V + m)^{-1} \Big\rangle = \int \frac{2m}{m^2 + \lambda^2}\, d\kappa_V(\lambda), \qquad \kappa_V = \Big\langle \frac{1}{V} \sum_n \delta_{\lambda_n} \Big\rangle,$$

where the angle brackets denote the average over gauge fields and δ_{λ_n} is the point mass at λ_n; we put $\Sigma_V(0) = 0$. We assume:

(C1) each κ_V is a positive measure on $(0, \Lambda]$, of total mass at most M_0, against which the function $\lambda \mapsto 1/\lambda$ is integrable, where the cutoff Λ and the bound M_0 do not depend on V;

(C2) as $V \to \infty$ along a given sequence of volumes, the measure κ_V converges weakly on $[0, \Lambda]$ to a measure κ that, on some interval $[0, \varepsilon_0]$, has a density f continuous at 0.

The cutoff and the bound on the total mass in (C1) hold on the lattice; the integrability of $1/\lambda$ holds for each single gauge field, whose spectrum is finite and avoids 0, and for the average over gauge fields it is an assumption on the finite-volume spectrum near 0. Hypothesis (C2) is the physical assumption at issue, the existence of a limiting density of near-zero eigenvalues per unit volume; its value at 0 is $f(0)$. A preparation is a volume V, a probe is a mass $m \in [0, 1]$, and the response is Σ_V. Each Σ_V is continuous on $[0, 1]$, with value 0 at 0, by dominated convergence, because $2m/(m^2 + \lambda^2) \le 1/\lambda$ for $\lambda > 0$.

Lemma 10.2. *Assume* (C1) *and* (C2). *Then* $\Sigma_V(m) \to \Sigma(m) = \int \frac{2m}{m^2+\lambda^2}\, d\kappa(\lambda)$ *for every* $m > 0$, *the function* Σ *is continuous on* $(0, 1]$, *and* $\Sigma(m) \to \pi f(0)$ *as* $m \to 0^+$.

Proof. For fixed $m > 0$, the function $\lambda \mapsto 2m/(m^2 + \lambda^2)$ is continuous and bounded on $[0, \Lambda]$, so the first statement is weak convergence, and the second follows from dominated convergence. For the third, choose $\varepsilon \in (0, \varepsilon_0]$ so small that f is bounded on $[0, \varepsilon]$; since κ has the density f there, it has no atom at 0. On $(\varepsilon, \Lambda]$ the integrand is at most $2m/\varepsilon^2$, and the contribution of this interval therefore tends to 0. On $[0, \varepsilon]$, the substitution $\lambda = mu$ shows that

$$\int_0^{\varepsilon} \frac{2m}{m^2 + \lambda^2} f(\lambda)\, d\lambda = \int_0^{\varepsilon/m} \frac{2}{1 + u^2} f(mu)\, du,$$

and the right side tends to $f(0) \int_0^{\infty} \frac{2\,du}{1+u^2} = \pi f(0)$, by dominated convergence. □

Theorem 10.3. *Assume* (C1) *and* (C2).

(a) *The iterated limits of the condensate are*

$$\lim_{m \to 0} \lim_{V \to \infty} \Sigma_V(m) = \pi f(0), \qquad \lim_{V \to \infty} \lim_{m \to 0} \Sigma_V(m) = 0.$$

(b) *The family* $\{\Sigma_V\}$ *is stable if and only if* $f(0) = 0$, *that is, if and only if the limiting valence condensate in the specified ensemble vanishes.*

(c) *In all cases the family is NIP*$_3$.

Proof. Let Q consist of the numbers $1/V$ along the sequence of volumes, together with $q_0 = 0$, put $g_{1/V} = \Sigma_V$, and let $G(m) = \Sigma(m)$ for $m > 0$ and $G(0) = 0$. By Lemma 10.2 and $\Sigma_V(0) = 0$, the map $q \mapsto g_q$ is continuous, the function G is continuous on $(0, 1]$, and $G(0^+) = \pi f(0)$. The theorem now follows from Lemma 10.1. □

Letting the volume grow before the mass vanishes keeps a density of near-zero modes, and the condensate is $\pi f(0)$; the other order sends the mass to 0 in each finite volume, and the integrability in (C1) makes its condensate tend to 0. When $\Sigma_0 = \pi f(0) > 0$, the microscopic regime with $\mu = mV\Sigma_0$ fixed concerns spectral data at the scale $1/V$. Chiral random-matrix

ensembles provide explicit condensate profiles in this regime, with their dependence on the microscopic spectrum, flavor content and topology [LS92, VW00]. In the Ising model, the intermediate values at the transition are values of idealized responses, obtained by tuning the offset as the volume grows; a natural condensate family with a continuum of transition points is sought in Section 11.

10.2. **The current response.** The Scalapino–White–Zhang criterion concerns the current response of a clean many-body system, one free of disorder, at zero temperature. Let H be its Hamiltonian, with ground state Ω of energy E_0; fix two perpendicular directions in space, and, for a real number $q > 0$, let j_q be the transverse current operator at the wave vector of length q along the first direction, the component along the second direction of the Fourier transform of the current density. For a finite system of volume V, write J_q for the extensive Fourier current and put $j_q = J_q/\sqrt{V}$, so its spectral measure has weights normalized per unit volume; the Hamiltonian and current are operators and the ground state is a vector; for an infinite system, the objects below are the infinite-volume limits of the corresponding finite-volume spectral measures, and Theorem 10.4 below is a statement about these measures. We use units in which $\hbar = 1$, so that frequencies are energies, and fixed units of length and energy, such as the lattice spacing and the hopping amplitude of a lattice model. We consider the case in which the spectral measures of the opposite Fourier currents J_q and $J_{-q} = J_q^*$ agree. The two Lehmann resolvents then combine to give the paramagnetic current response at imaginary frequency $i\omega$ in the form

$$\Pi(q, i\omega) = \int \frac{2\nu}{\nu^2 + \omega^2}\, d\mu_q(\nu),$$

where μ_q is the spectral measure of $H - E_0$ in the vector $j_q\Omega$, restricted to $(0, \infty)$; for a finite system, $\mu_q = \sum_k |\langle \psi_k, j_q\Omega\rangle|^2 \delta_{E_k - E_0}$, the sum running over the k with $E_k > E_0$, where the vectors ψ_k, for k in a countable index set, form an orthonormal basis of eigenvectors of H, with $H\psi_k = E_k\psi_k$. The total response is $\mathcal{K}(q, i\omega) = D_0 - \Pi(q, i\omega)$, where D_0 is the diamagnetic term, the expectation, per unit volume, of minus the kinetic energy of the bonds parallel to the current on a lattice, and, in the continuum, the ratio of the density of the charge carriers to their mass. Following Scalapino, White and Zhang [SWZ93], we define the *superfluid weight* and the *Drude weight*, up to their factor πe^2, where e is the charge of the carriers, by

$$D_s = \lim_{q \to 0} \mathcal{K}(q, 0), \qquad D = \lim_{\omega \to 0} \lim_{q \to 0} \mathcal{K}(q, i\omega).$$

The superfluid weight is the transverse static limit of Scalapino, White and Zhang. Their Drude weight, and that of Kohn [Koh64], is taken at $q = 0$ and continued from imaginary to real frequency; it agrees with D when the spectral measures μ_q depend continuously on q at $q = 0$ away from zero energy and when, in addition, the imaginary-frequency limit agrees with the real-frequency one. In an insulator $D_s = D = 0$; in a clean normal metal $D > 0 = D_s$; positive D_s describes a superconducting response. A gap in the transverse current spectral weight that persists as $q \to 0$ is a sufficient condition for $D = D_s$ [SWZ93]. In a disordered metal the Drude weight vanishes, and at positive temperature the normal fluid, the thermally excited carriers that do not condense, yields $D > D_s$ even in a superconductor; both cases lie outside the setting of this subsection, a clean system at zero temperature.

It is convenient to absorb a factor into the measure: put $d\zeta_q(\nu) = \frac{2}{\nu}\, d\mu_q(\nu)$, whence

$$\Pi(q, i\omega) = \int \frac{\nu^2}{\nu^2 + \omega^2}\, d\zeta_q(\nu), \qquad \Pi(q, 0) = \zeta_q\big((0, \infty)\big).$$

We assume:

(S1) each ζ_q, for $0 < q \leq 1$, is a positive measure on $(0, \Lambda]$ of total mass at most D_0, where the bandwidth Λ does not depend on q;
(S2) there is a measure ζ_0 on $[0, \Lambda]$ such that, for every continuous function h on $[0, \Lambda]$, the map $q \mapsto \int h \, d\zeta_q$, with each ζ_q regarded as a measure on $[0, \Lambda]$, is continuous on $(0, 1]$ and converges to $\int h \, d\zeta_0$ as $q \to 0$.

The bound in (S1) is the statement that $\mathcal{K}(q, 0) \geq 0$, that is, that the ground state is a local minimum of the energy as a function of a static vector potential. This is an assumption: for lattice fermions it can fail, as Lieb's flux-phase theorem shows [Lie94]. The compact support in (S1) is a convenience: since $\nu^2/(\nu^2 + \omega^2)$ extends continuously to $[0, \infty]$, the proof below works for tight families of measures on $(0, \infty)$. The limit ζ_0 may have an atom at 0, which carries the weight of current-carrying excitations whose energy tends to 0 with q. A preparation is the length $q \in (0, 1]$ of the wave vector, a probe is a frequency $\omega \in [0, 1]$, and the response is $g_q(\omega) = \Pi(q, i\omega)$.

Theorem 10.4. *Assume* (S1) *and* (S2).

(a) *We have* $D_s = D_0 - \zeta_0\big([0, \Lambda]\big)$ *and* $D = D_0 - \zeta_0\big((0, \Lambda]\big)$, *whence* $D - D_s = \zeta_0(\{0\})$.
(b) *The family of responses* $\{g_q : 0 < q \leq 1\}$ *is stable if and only if* $D = D_s$. *In particular, insulating responses with* $D = D_s = 0$ *and superconducting responses satisfying* $D = D_s > 0$ *are stable, whereas clean normal-metal responses with* $D > 0 = D_s$ *are unstable.*
(c) *In all cases the family is NIP*$_3$.

Proof. Write $\chi_\omega(\nu) = \nu^2/(\nu^2 + \omega^2)$. For $\omega > 0$ the function χ_ω is continuous on $[0, \Lambda]$ with $\chi_\omega(0) = 0$, while $\chi_0 = 1$ on $(0, \Lambda]$. Each g_q is continuous on $[0, 1]$, by dominated convergence. By (S2), applied to the continuous function 1, we have $g_q(0) = \zeta_q((0, \Lambda]) \to \zeta_0([0, \Lambda])$, and the formula for D_s follows; applied to χ_ω with $\omega > 0$, (S2) shows that $g_q(\omega) \to G(\omega) = \int_{(0,\Lambda]} \chi_\omega \, d\zeta_0$, and, by dominated convergence, $G(\omega) \to \zeta_0((0, \Lambda])$ as $\omega \to 0$, which yields the formula for D. This proves (a). Put $g_0 = G$, with G extended to 0 by $G(0) = \zeta_0([0, \Lambda])$. The map $q \mapsto g_q$ is continuous on $[0, 1]$, by (S2) and the computation just made, and G is continuous on $(0, 1]$ with $G(0) - G(0^+) = \zeta_0(\{0\}) = D - D_s$. Parts (b) and (c) now follow from Lemma 10.1. □

Under (S1) and (S2), the discrepancy $D - D_s$ is exactly the mass at zero of the limiting weighted current spectral measure ζ_0. A positive mass at zero therefore has two equivalent descriptions: the two orders of limits of the current response disagree, and the family of responses is unstable. When this mass vanishes, the limiting response is continuous and the family is stable. In both cases the closure remains metrizable, so the distinction between stable and unstable current responses lies within NIP$_3$.

11. Questions

We close with problems concerning the possible response compacta, the role of noncommutative probe algebras, and extensions of the classification to further physical response families. We also ask for finite combinatorial tests of the finer topological levels and for a proof of the classification that avoids forcing.

(1) *Light cones.* In Theorem 7.10 Bob detects at a fixed time. Locate the causal order in the classification when Bob's detection time is also uncertain, and in spacetimes of dimension $1 + d$, with $d \geq 2$ dimensions of space.
(2) *The minimal families of the box.* By Theorem 8.5, the comparison of positions in a box realizes the minimal families D_3, D_4 and D_7 of the heptachotomy. Does it realize D_6? Which of D_6 and D_7 does Theorem 7.8(b) produce from a given dense sequence?
(3) *Noncommutative probes.* Find a natural observable that is NIP but unstable and whose probe algebra is not commutative, and locate it in the classification. Can noncommutativity produce a phenomenon in the classification that observables with commutative probe algebras cannot produce, for instance in the minimal families of the heptachotomy that arise from a given dense sequence of preparations?
(4) *Orientations.* By Theorem 7.12, the comparison of phases on a circle is NIP but not NIP_1. Locate in the classification the comparison of directions on a sphere, "Bob's direction lies in the open hemisphere centered at Alice's", and relative orientations in higher dimensions.
(5) *Finitary levels.* Is there a finitary, combinatorial characterization of NIP_1, NIP_2 and NIP_3, in the spirit of the fat-shattering dimension for uniform NIP, sharpening Proposition 4.6?
(6) *Quantum phase transitions.* Extend Theorem 9.6 to the ground state of the Ising model in a transverse field, where the transition is driven by the transverse field Γ at zero temperature.
(7) *A continuum of transitions.* Is there a natural family of chiral condensates with a continuum of transition points, obtained for instance from twisted boundary conditions or a chemical potential, that is NIP_1 but not NIP_2?
(8) *Positive temperature.* Extend Theorem 10.4 to positive temperature, where the Matsubara frequencies, the imaginary frequencies at which thermal response functions are evaluated, form a discrete set, and the Drude weight is obtained by extrapolation.
(9) *Unbounded formulas.* In Keisler's setting, classify the energy functional $\rho \mapsto \mathrm{Tr}(\rho H)$ of a Hamiltonian H, taking as shards (the parts of the state space on which every predicate stays within prescribed bounds) the sets of states that satisfy the energy constraints $\mathrm{Tr}(\rho H) \leq E$, for real numbers E.
(10) *Forcing.* Can Todorčević's trichotomy, or its consequences for observables, be proved without forcing?

Acknowledgments

We thank Stevo Todorčević for pointing out to us that Theorem 6.6 is known, in the larger class of locally connected Rosenthal compacta, and for the references that establish it.

Declarations

Competing interests. The authors have no competing interests to declare.
Data availability. No data were generated or analyzed in this work.

References

[Aar07] Scott Aaronson, *The learnability of quantum states*, Proceedings of the Royal Society A **463** (2007), 3089–3114.
[ABDCBH97] Noga Alon, Shai Ben-David, Nicolò Cesa-Bianchi, and David Haussler, *Scale-sensitive dimensions, uniform convergence, and learnability*, Journal of the ACM **44** (1997), no. 4, 615–631.

[ADK08] Spiros A. Argyros, Pandelis Dodos, and Vassilis Kanellopoulos, *A classification of separable Rosenthal compacta and its applications*, Dissertationes Mathematicae **449** (2008), 1–55.

[Ai92] A. V. Arkhangelskiĭ, *Topological function spaces*, Mathematics and its Applications (Soviet Series), vol. 78, Kluwer Academic Publishers Group, Dordrecht, 1992, Translated from the Russian by R. A. M. Hoksbergen. MR 1144519

[Asa70] Taro Asano, *Lee–Yang theorem and the Griffiths inequality for the anisotropic Heisenberg ferromagnet*, Physical Review Letters **24** (1970), 1409–1411.

[Ass83] Patrick Assouad, *Densité et dimension*, Annales de l'Institut Fourier **33** (1983), no. 3, 233–282.

[BC80] Tom Banks and A. Casher, *Chiral symmetry breaking in confining theories*, Nuclear Physics B **169** (1980), 103–125.

[Ben08] Itaï Ben Yaacov, *Continuous first order logic for unbounded metric structures*, Journal of Mathematical Logic **8** (2008), no. 2, 197–223.

[Ben09] ———, *Continuous and random Vapnik–Chervonenkis classes*, Israel Journal of Mathematics **173** (2009), 309–333.

[BFT78] J. Bourgain, D. H. Fremlin, and M. Talagrand, *Pointwise compact sets of Baire-measurable functions*, Amer. J. Math. **100** (1978), no. 4, 845–886. MR 509077

[BG09] Jakob E. Björnberg and Geoffrey R. Grimmett, *The phase transition of the quantum Ising model is sharp*, Journal of Statistical Physics **136** (2009), 231–273, arXiv:0901.0328.

[BK90] Christian Borgs and Roman Kotecký, *A rigorous theory of finite-size scaling at first-order phase transitions*, Journal of Statistical Physics **61** (1990), 79–119.

[BYU10] Itaï Ben Yaacov and Alexander Usvyatsov, *Continuous first order logic and local stability*, Trans. Amer. Math. Soc. **362** (2010), no. 10, 5213–5259. MR 2657678

[CHY15] Hao-Chung Cheng, Min-Hsiu Hsieh, and Ping-Cheng Yeh, *The learnability of unknown quantum measurements*, 2015, arXiv:1501.00559.

[Deb13] Gabriel Debs, *Descriptive aspects of Rosenthal compacta*, Recent Progress in General Topology III, Atlantis Press, 2013, pp. 205–227.

[DI26] Eduardo Dueñez and José Iovino, *Classification of bipartite observables I: stable and uniformly stable observables*, Preprint, arXiv:2610.11002, 2026.

[DIMW+26] E. Dueñez, J. Iovino, T. Matos-Wiederhold, L. Salvetti, and F. D. Tall, *Complexity of deep computations via topology of function spaces*, 2026, arXiv:2601.00528.

[FHS13] Ilijas Farah, Bradd Hart, and David Sherman, *Model theory of operator algebras I: Stability*, Bull. Lond. Math. Soc. **45** (2013), no. 4, 825–838.

[FR12] Jürg Fröhlich and Pierre-François Rodríguez, *Some applications of the Lee–Yang theorem*, Journal of Mathematical Physics **53** (2012), 095218, arXiv:1205.6643.

[FV17] Sacha Friedli and Yvan Velenik, *Statistical mechanics of lattice systems: A concrete mathematical introduction*, Cambridge University Press, 2017.

[Gin69] Jean Ginibre, *Existence of phase transitions for quantum lattice systems*, Communications in Mathematical Physics **14** (1969), 205–234.

[GM22] Eli Glasner and Michael Megrelishvili, *Todorčević' trichotomy and a hierarchy in the class of tame dynamical systems*, Transactions of the American Mathematical Society **375** (2022), no. 7, 4513–4548.

[God80] Gilles Godefroy, *Compacts de Rosenthal*, Pacific J. Math. **91** (1980), no. 2, 293–306. MR 615679

[Gri64] Robert B. Griffiths, *Peierls proof of spontaneous magnetization in a two-dimensional Ising ferromagnet*, Physical Review **136** (1964), A437–A439.

[Gro52] A. Grothendieck, *Critères de compacité dans les espaces fonctionnels généraux*, Amer. J. Math. **74** (1952), 168–186. MR 47313

[Gru88] Gary Gruenhage, *[TITLE TO BE CONFIRMED]*, General Topology and its Relations to Modern Analysis and Algebra VI (Proc. Sixth Prague Topological Symposium, 1986) (Zdeněk Frolík, ed.), Research and Exposition in Mathematics, vol. 16, Heldermann, Berlin, 1988.

[GS84] Yuri Gurevich and Peter H. Schmitt, *The theory of ordered abelian groups does not have the independence property*, Transactions of the American Mathematical Society **284** (1984), no. 1, 171–182.

[HL66] J. D. Halpern and H. Läuchli, *A partition theorem*, Transactions of the American Mathematical Society **124** (1966), 360–367.

[HP11] Ehud Hrushovski and Anand Pillay, *On NIP and invariant measures*, Journal of the European Mathematical Society **13** (2011), no. 4, 1005–1061.

[Kad51] Richard V. Kadison, *A representation theory for commutative topological algebra*, Memoirs of the American Mathematical Society, vol. 7, American Mathematical Society, 1951.

[Kei23] H. Jerome Keisler, *Model theory for real-valued structures*, Beyond First-Order Model Theory, Vol. 2 (José Iovino, ed.), CRC Press, 2023, arXiv:2005.11851, pp. 43–101.

[Koh64] Walter Kohn, *Theory of the insulating state*, Physical Review **133** (1964), A171–A181.

[KPS86] Julia F. Knight, Anand Pillay, and Charles Steinhorn, *Definable sets in ordered structures. II*, Transactions of the American Mathematical Society **295** (1986), no. 2, 593–605.

[KR83] Richard V. Kadison and John R. Ringrose, *Fundamentals of the theory of operator algebras, vol. i*, Academic Press, 1983.

[Lie94] Elliott H. Lieb, *Flux phase of the half-filled band*, Physical Review Letters **73** (1994), 2158–2161.

[LS81] Elliott H. Lieb and Alan D. Sokal, *A general Lee–Yang theorem for one-component and multi-component ferromagnets*, Communications in Mathematical Physics **80** (1981), no. 2, 153–179.

[LS92] H. Leutwyler and A. Smilga, *Spectrum of Dirac operator and role of winding number in QCD*, Physical Review D **46** (1992), 5607–5632.

[LSV93] Alain Louveau, Saharon Shelah, and Boban Veličković, *Borel partitions of infinite subtrees of a perfect tree*, Annals of Pure and Applied Logic **63** (1993), no. 3, 271–281.

[LY52] T. D. Lee and C. N. Yang, *Statistical theory of equations of state and phase transitions. II. Lattice gas and Ising model*, Physical Review **87** (1952), 410–419.

[Mil81] Keith Milliken, *A partition theorem for the infinite subtrees of a tree*, Transactions of the American Mathematical Society **263** (1981), 137–148.

[New74] Charles M. Newman, *Zeros of the partition function for generalized Ising systems*, Communications on Pure and Applied Mathematics **27** (1974), 143–159.

[Pei36] Rudolf Peierls, *On Ising's model of ferromagnetism*, Proceedings of the Cambridge Philosophical Society **32** (1936), 477–481.

[PS86] Anand Pillay and Charles Steinhorn, *Definable sets in ordered structures. I*, Transactions of the American Mathematical Society **295** (1986), no. 2, 565–592.

[Rod21] José Rodríguez, *ε-weakly precompact sets in Banach spaces*, 2021, arXiv:2102.01546.

[Ros74] H. P. Rosenthal, *A characterization of Banach spaces containing l^1*, Proc. Nat. Acad. Sci. U.S.A. **71** (1974), 2411–2413. MR 358307

[Sau72] N. Sauer, *On the density of families of sets*, J. Combinatorial Theory Ser. A **13** (1972), 145–147. MR 307902

[SF71] Masuo Suzuki and Michael E. Fisher, *Zeros of the partition function for the Heisenberg, ferroelectric, and general Ising models*, Journal of Mathematical Physics **12** (1971), 235–246.

[SG73] Barry Simon and Robert B. Griffiths, *The $(\phi^4)_2$ field theory as a classical Ising model*, Communications in Mathematical Physics **33** (1973), 145–164.

[She71] Saharon Shelah, *Stability, the f.c.p., and superstability; model theoretic properties of formulas in first order theory*, Ann. Math. Logic **3** (1971), no. 3, 271–362. MR 0317926

[She72] ———, *A combinatorial problem; stability and order for models and theories in infinitary languages*, Pacific J. Math. **41** (1972), 247–261. MR 307903

[She90] S. Shelah, *Classification theory and the number of nonisomorphic models*, second ed., Studies in Logic and the Foundations of Mathematics, vol. 92, North-Holland Publishing Co., Amsterdam, 1990. MR MR1083551 (91k:03085)

[Sim93] Barry Simon, *The statistical mechanics of lattice gases*, vol. I, Princeton University Press, Princeton, NJ, 1993.

[Sim15a] Pierre Simon, *A guide to NIP theories*, Lecture Notes in Logic, vol. 44, Association for Symbolic Logic, Chicago, IL; Cambridge Scientific Publishers, Cambridge, 2015. MR 3560428

[Sim15b] ———, *Rosenthal compacta and NIP formulas*, Fund. Math. **231** (2015), no. 1, 81–92. MR 3361236

[Suz76] Masuo Suzuki, *Relationship between d-dimensional quantal spin systems and $(d+1)$-dimensional Ising systems: equivalence, critical exponents and systematic approximants of the partition function and spin correlations*, Progress of Theoretical Physics **56** (1976), 1454–1469.

[SWZ93] Douglas J. Scalapino, Steven R. White, and Shoucheng Zhang, *Insulator, metal, or superconductor: The criteria*, Physical Review B **47** (1993), 7995–8007.

[Tit39] E. C. Titchmarsh, *The theory of functions*, second ed., Oxford University Press, Oxford, 1939.

[Tka11] Vladimir V. Tkachuk, *A C_p-theory problem book: Topological and function spaces*, Problem Books in Mathematics, Springer, 2011.

[Tod97] Stevo Todorcevic, *Topics in topology*, Lecture Notes in Mathematics, vol. 1652, Springer Berlin, Heidelberg, 1997.

[Tod99] Stevo Todorčević, *Compact subsets of the first Baire class*, J. Amer. Math. Soc. **12** (1999), no. 4, 1179–1212. MR 1685782

[Val84] L. G. Valiant, *A theory of the learnable*, Commun. ACM **27** (1984), no. 11, 1134–1142.

[VC71] Vladimir N Vapnik and Alexey Ya Chervonenkis, *On the uniform convergence of relative frequencies of events to their probabilities*, Theory of Probability & Its Applications **16** (1971), no. 2, 264–280.

[VC74] ———, *Theory of pattern recognition*, Nauka, Moscow, 1974, German Translation: Theorie der Zeichenerkennung, Akademie-Verlag, Berlin, 1979.

[VW00] J. J. M. Verbaarschot and T. Wettig, *Random matrix theory and chiral symmetry in QCD*, Annual Review of Nuclear and Particle Science **50** (2000), 343–410.

[Wie49] Helmut Wielandt, *Über die Unbeschränktheit der Operatoren der Quantenmechanik*, Mathematische Annalen **121** (1949), 21.

[Wil96] A. J. Wilkie, *Model completeness results for expansions of the ordered field of real numbers by restricted Pfaffian functions and the exponential function*, Journal of the American Mathematical Society **9** (1996), no. 4, 1051–1094.

[Win47] Aurel Wintner, *The unboundedness of quantum-mechanical matrices*, Physical Review **71** (1947), 738–739.

[Zil16] Boris Zilber, *The semantics of the canonical commutation relation*, 2016, arXiv:1604.07745.

Department of Mathematics, The University of Texas at San Antonio, One UTSA Circle, San Antonio, TX 78249, USA

Email address: Eduardo.Duenez@utsa.edu

Department of Mathematics, The University of Texas at San Antonio, One UTSA Circle, San Antonio, TX 78249, USA

Email address: jose.iovino@utsa.edu