COMPLEXITY OF DEEP COMPUTATIONS VIA TOPOLOGY OF FUNCTION SPACES

EDUARDO DUEÑEZ[1] JOSÉ IOVINO[1] TONATIUH MATOS-WIEDERHOLD[2]
LUCIANO SALVETTI[2] FRANKLIN D. TALL[2]

[1]Department of Mathematics, University of Texas at San Antonio
[2]Department of Mathematics, University of Toronto

Abstract. We use topology of function spaces to study the complexity of deep computations, the limits of computations that arise, for instance, in recurrent neural networks. The classification of Rosenthal compacta yields new complexity classes of deep computations, and these classes form a strict hierarchy. Each level of the hierarchy is attained by classical computations, among them the power method, Bresenham's line algorithm and the Euclidean algorithm, while Newton's method on the Riemann sphere lies beyond it. For iterative algorithms, the hierarchy coincides with a hierarchy of tame dynamical systems. The translation between topology and computation is made through the language of model theory, specifically through the *independence property* of Shelah's classification theory. The same methods give conditions under which deep computations are limits of sequences of computations and, in a measure-theoretic form, conditions under which they can be evaluated almost surely by random sampling.

Introduction

In this paper we use topological methods and concepts inspired by model theory to study the asymptotic behavior of computations as a parameter of the computation tends to a limit: for example, as the depth of a neural network tends to infinity, or as the time interval between the layers of a time-series network tends to zero. Particular cases of such limits have recently attracted considerable attention in deep learning research (e.g., Neural Ordinary Differential Equations [CRBD], Physics-Informed Neural Networks [RPK19], and Deep Equilibrium Models [BKK]). Our results concern classical models of computation, in which real-valued quantities are known and manipulated to a finite degree of precision. We work in the framework of compositional computation structures, which was introduced in the first paper of this series [ADIW26] and accommodates all these limits in a single setting.

In that paper, Alva, Dueñez, Iovino and Walton introduced two closely related notions of limit of computations: *ultracomputations*, so called because they arise as ultrafilter limits of standard computations, and *deep computations*, so called after the usage in machine learning [BKK]. The two notions differ in a technical point that plays no role here, and we speak of deep computations throughout. Using

Date: October 25, 2026.

2020 *Mathematics Subject Classification.* Primary 68Q15, 03C45, 54H30; Secondary 03C98, 03D15, 05D10, 26A21, 37B05, 54C35, 68T07.

Key words and phrases. Deep computations, limit computations, complexity classes, deep learning, recurrent neural networks, deep equilibrium models, compositional computation structures, independence property, NIP, model theory, Rosenthal compacta, Baire class 1 functions, Todorčević's trichotomy, Argyros–Dodos–Kanellopoulos heptachotomy, Bourgain–Fremlin–Talagrand theorem, tame dynamical systems, infinite Ramsey theory.

Grothendieck's double limit criterion [Gro52], they proved a dichotomy between *stable* and unstable sets of computations: the deep computations of a stable set are *polynomially approximable*, that is, on every *shard* (a set of states on which each feature obeys a prescribed bound), each feature of their output is a uniform limit of polynomials in the features of the input. Polynomial approximability corresponds to continuous extendibility in topology, and to *stability* and *type definability* in model theory.

In this paper we refine that dichotomy. An unstable set of computations is either *wild*, that is, it has Shelah's *independence property* (IP), or *tame*, that is, it is *NIP*: it does *not* have the independence property. Between stability and NIP we identify three new levels, NIP_3, NIP_2 and NIP_1, which together with NIP form a strict hierarchy of four tame classes beyond the stable ones; we also read the seven minimal families of Baire class 1 functions found by Argyros, Dodos and Kanellopoulos, described below, as primitives of binary search. On the topological side, the theory of *Rosenthal compacta* takes the place of Grothendieck's criterion; on the model-theoretic side, *neostability*, and NIP in particular, takes the place of stability.

The translation between topology and computation goes through model theory. To every intermediate state of a computation we associate its *type*: a "lookup table" of the (real) values of all the features of the computation at that state. The types, together with their limits, form a space of real-valued functions on the set of features, the *space of types*, on which computations act; the embedding of a structure into its space of types is a central device of model theory. This embedding allows us to study topological limits of computations by means of the theory of function spaces with the topology of pointwise convergence, known as $\mathrm{C_p}$*-theory*. The space of types thus becomes a "Rosetta stone" connecting three classification programs: in topology, the classification of Rosenthal compacta pioneered by Todorčević [Tod99]; in logic, the classification of theories developed by Shelah [She71, She90]; and in statistical learning, the theory of VC dimension and PAC learnability [VC71, VC74, Val84]. Under this translation, NIP sets of computations correspond to spaces of deep computations that are, on each shard, Rosenthal compacta (see Theorem 3.1); finiteness of VC dimension implies NIP, and for first-order formulas the two notions coincide (see Section 1.5).

Deep computations arise as topological limits of standard (continuous) computations. Pointwise limits of sequences of continuous functions form the *first Baire class*, one rung above the continuous functions; intuitively, functions of Baire class 1 have "controlled" discontinuities. A *Rosenthal compactum* is a set of Baire class 1 functions on a Polish (i.e., separable and completely metrizable) space, compact in the topology of *pointwise* convergence. Since the late 1970s, Rosenthal compacta have been studied intensively in functional analysis, especially in the theory of Banach spaces; Todorčević's trichotomy for such compacta [Tod99] has been used to organize tame dynamical systems into a hierarchy [GM22]. Todorčević's proof relies on set-theoretic forcing and infinite Ramsey theory, and his methods continue to be analyzed and exploited [Tod23, HT19]. Refining the trichotomy, Argyros, Dodos and Kanellopoulos [ADK08] found a heptachotomy, which we also use: there are seven minimal families of Baire class 1 functions indexed by the binary tree such that every relatively compact family of Baire class 1 functions on a Polish space, indexed by the binary tree, contains, along a dyadic subtree (a copy of the binary

tree inside itself), a family equivalent to one of the seven. Here two families are equivalent when the bijection between them that respects the indexing extends to a homeomorphism of their closures.

The decisive fact for approximation, both in the setting of Grothendieck [Gro52] and in that of Bourgain, Fremlin and Talagrand [BFT78], is that *compactness and sequential compactness coincide*, and that every point in the closure of a relatively compact set of functions is the limit of a sequence from the set. Consequently, when the features are countably many and the set of computations is NIP, every deep computation, in principle a nonconstructive ultrafilter limit over a possibly uncountable set of computations, is, on each shard, the limit of a *sequence* of standard computations.

The results show how the classification invariants used in computation, topology, descriptive set theory and model theory translate into one another. We assume no familiarity with advanced topology, model theory or computing: the prerequisites are undergraduate topology and measure theory, and the background beyond them is given in Section 1.

Our first main result, Theorem 3.1, concerns structures with countably many predicates. It shows that a set of computations has the NIP on every shard, one observable feature at a time, if and only if its deep computations are of the first Baire class on each shard, that is, if and only if, on each shard, the deep computations form a Rosenthal compactum; in that case, every deep computation is, on each shard, the limit of a *sequence* of computations. Guided by Todorčević's trichotomy, we define three levels of tameness (see Definition 3.2) and obtain the hierarchy

$$\text{stable} \Rightarrow \text{NIP}_3 \Rightarrow \text{NIP}_2 \Rightarrow \text{NIP}_1 \Rightarrow \text{NIP},$$

in which every implication is strict.

In Section 5 we realize each level by a classical algorithm:

- a rotation of the circle is stable;
- the power method is NIP_3 but not stable;
- Bresenham's line algorithm is
  - NIP_2 but not NIP_3 in the plane,
  - NIP_1 but not NIP_2 in three-dimensional space;
- the Euclidean algorithm, acting on the projective line by the group that its steps generate, is NIP but not NIP_1, and its deep computations include exact equality tests;
- Newton's method for the roots of $z^2 - 1$ on the extended complex plane is not NIP.

At the level NIP but not NIP_1, the computations of both our examples, the Euclidean algorithm and free reduction in the free group on two generators, form a group with two generators; whether a single iterated map can reach this level is an open question, closely related to a question of Glasner and Megrelishvili (see Question 7.2). For iterative algorithms, the hierarchy coincides with the hierarchy of tame dynamical systems studied by Glasner and Megrelishvili [GM22].

The heptachotomy of Argyros, Dodos and Kanellopoulos refines this picture for families of computations indexed by the binary tree: on each shard and for each feature, every tame family of this kind contains, along a dyadic subtree, a family equivalent to one of the seven minimal families, which we read as primitives of binary search (see Section 3.3). Finally, Theorem 6.7 and Corollary 6.14 relate NIP

and Talagrand stability to the Monte Carlo computability of deep computations, that is, to their evaluation almost surely by random sampling.

Organization of the paper. In Section 1 we present the topological and combinatorial preliminaries, including five canonical Rosenthal compacta (Section 1.4) and the role of NIP in model theory, combinatorics and learning (Section 1.5). In Section 2 we recall compositional computation structures, the source of the structural and model-theoretic viewpoint of the paper, and their stable sets of computations; the section ends with two motivating examples, Newton's method for square roots and Newton's method on the Riemann sphere, whose different behaviors the classification is designed to separate (Section 2.5). Section 3 contains the classification of deep computations and the main theorems. In Section 4 we work out Newton's method in detail, and in Section 5 we place classical algorithms at every level of the hierarchy and beyond it. In Section 6 we replace topological limits by a probabilistic viewpoint, which we intend to develop further in future research. Section 7 contains open questions.

Contents

Introduction 1
Organization of the paper 4
1. Topological preliminaries: From continuity to Baire class 1 5
1.1. From Rosenthal's dichotomy to the Bourgain–Fremlin–Talagrand dichotomy to Shelah's NIP 7
1.2. Rosenthal compacta 9
1.3. The special case $B_1(X, \mathbb{R}^{\mathcal{P}})$ with $\mathcal{P}$ countable 9
1.4. Five canonical Rosenthal compacta 11
1.5. NIP in model theory, combinatorics and learning: polynomial versus exponential growth 12
Historical remarks 13
2. Compositional Computation Structures: A structural approach to arbitrary-precision computations 14
2.1. Computation States Structures 14
2.2. Compositional Computation Structures 15
2.3. Computability of deep computations and the Extendibility Axiom 16
2.4. Stability 17
2.5. Two motivating examples 18
3. Classifying deep computations 19
3.1. NIP, Rosenthal compacta, and deep computations 19
3.2. Todorčević's trichotomy and levels of NIP 19
3.3. The Argyros–Dodos–Kanellopoulos heptachotomy and binary search 21
4. Examples of Compositional Computation Structures 23
4.1. Example: The Newton CCS for the cubic $\rho(z) = z^3 - 2z + 2$ 24
4.2. Example: The Newton CCS for the cubic $\kappa(z) = z^3 - 1$ 24
4.3. Example: Square roots 25
5. Computations across the hierarchy 26
5.1. Iterative algorithms and tame dynamics 26
5.2. Stable and metrizable 28

5.3. Hereditarily separable but not metrizable 29
5.4. First countable but not hereditarily separable 32
5.5. NIP but not first countable: the Euclidean algorithm 32
5.6. Beyond NIP 35
5.7. Summary 37
6. Randomized versions of NIP and Monte Carlo computability of deep computations 37
6.1. NIP and Monte Carlo computability of deep computations 37
6.2. Talagrand stability and Monte Carlo computability of deep computations 40
7. Open questions 43
References 43

1. Topological preliminaries: From continuity to Baire class 1

Deep computations are pointwise limits of continuous maps; hence the two themes of this paper, the approximation of a deep computation by a sequence of computations and the complexity of the set of all deep computations, belong to the theory of spaces of functions with the topology of pointwise convergence. In this section we recall the notions of general topology and of function space theory that these themes require: Polish spaces and Baire category; the first Baire class; the dichotomies of Rosenthal and of Bourgain, Fremlin and Talagrand, from which the NIP emerges; and the Rosenthal compacta, five of which serve as yardsticks for the classification. We close the section with the role of the NIP in model theory, combinatorics and statistical learning. All spaces are assumed to be Hausdorff unless explicitly stated otherwise.

We write $\mathbb{N}$ for the set $\{0, 1, 2, \dots\}$ of natural numbers, so that the semigroups of iterates considered below contain the identity. A subset of a topological space is F_σ if it is a countable union of closed sets, and G_δ if it is a countable intersection of open sets. In a metric space every open set is F_σ, and, equivalently, every closed set is G_δ.

A *Polish space* is a separable topological space whose topology is induced by a complete metric. Countable products of Polish spaces are Polish. Classical examples, besides the real line, are the Cantor space $2^{\mathbb{N}}$ of all infinite binary sequences, the Baire space $\mathbb{N}^{\mathbb{N}}$ of all infinite sequences of natural numbers, and the space $\mathbb{R}^{\mathbb{N}}$ of all sequences of real numbers, each with the product topology.

Fact 1.1. [Eng89, 4.3.24] *A subset of a Polish space is itself Polish in the subspace topology if and only if it is a G_δ set. In particular, closed subsets and open subsets of Polish spaces are also Polish spaces.*

A subset A of a topological space X is *nowhere dense* if its closure has empty interior, and of the *first category*, or *meager*, if it is contained in a countable union of closed nowhere dense sets; otherwise A is of the *second category*. The complements of meager sets are called *comeager*. Since the closure of a nowhere dense set is nowhere dense, the meager subsets of X form the σ-ideal generated by the nowhere dense subsets of X. A topological space X is *Baire* if every comeager subset of X

is dense, or, equivalently, if every countable intersection of dense open sets is dense. The Baire category theorem states that compact Hausdorff spaces and completely metrizable spaces, Polish spaces in particular, are Baire.[1]

For topological spaces X and Y, we denote by $\mathrm{C_p}(X,Y)$ the set of all continuous functions $f\colon X \to Y$ with the topology of pointwise convergence, that is, the topology it inherits as a subspace of the product Y^X, in which X serves merely as an index set. We write $\mathrm{C_p}(X)$ for $\mathrm{C_p}(X,\mathbb{R})$.

A function $f\colon X \to Y$ between topological spaces is of *Baire class 1* if it is the pointwise limit of a sequence $(f_n)_{n\in\mathbb{N}}$ of continuous functions, that is, $f(x) = \lim_{n\to\infty} f_n(x)$ for all $x \in X$. When Y is metrizable, an elementary argument shows that a function of Baire class 1 is continuous at every point outside a set of the first category. We denote by $\mathrm{B_1}(X,Y)$ the set of functions $X \to Y$ of Baire class 1 with the topology of pointwise convergence, and we write $\mathrm{B_1}(X)$ for $\mathrm{B_1}(X,\mathbb{R})$; the inclusions $\mathrm{C_p}(X,Y) \subseteq \mathrm{B_1}(X,Y) \subseteq Y^X$ are topological. Relatively compact subsets of $\mathrm{C_p}(X)$ and of $\mathrm{B_1}(X)$, especially for Polish X, are of interest in analysis and in topological dynamics.

A topological space X is *perfectly normal* if it is normal and every closed subset of X is a G_δ set (equivalently, every open subset of X is an F_σ set). Every metrizable space (hence, every Polish space) is perfectly normal.

The following characterization of Baire class 1 functions is classical; for a proof, see [Tod97, Section 10].

Fact 1.2 (Lebesgue, Hausdorff, Baire[2])**.** *If X is perfectly normal, then the following conditions are equivalent for a function $f : X \to \mathbb{R}$:*

(1) *f is a Baire class 1 function, that is, f is a pointwise limit of continuous functions.*
(2) *$f^{-1}[U]$ is an F_σ subset of X whenever $U \subseteq \mathbb{R}$ is open.*

If X is Polish, then (1) and (2) are equivalent to:

(3) *For every nonempty closed $F \subseteq X$, the restriction $f|_F$ has a point of continuity.*

Moreover, if X is Polish and $f \notin \mathrm{B_1}(X)$, then there exist countable $D_0, D_1 \subseteq X$ and reals $a < b$ such that

$$D_0 \subseteq f^{-1}(-\infty, a], \quad D_1 \subseteq f^{-1}[b,\infty), \quad \overline{D_0} = \overline{D_1}.$$

[1] More generally, any Čech-complete space is Baire [Eng89, Theorem 3.9.3].

[2] The implication (1)⇒(2) holds in every topological space. Indeed, if f is the pointwise limit of a sequence $(f_n)_{n\in\mathbb{N}}$ of continuous functions and a is a real number, then

$$f^{-1}(a,\infty) = \bigcup_{k\geq 1} \bigcup_{m\in\mathbb{N}} \bigcap_{n\geq m} f_n^{-1}[a + 1/k, \infty)$$

is a countable union of closed sets; the same is true of $f^{-1}(-\infty, a)$, and condition (2) follows because every open subset of $\mathbb{R}$ is a countable union of intervals $(a,b) = (a,\infty) \cap (-\infty, b)$. The reverse implication, (2)⇒(1), was established by Lebesgue for real-valued functions of finitely many real variables [Leb04] and extended by Hausdorff to real-valued functions on metric spaces [Kec95, Theorem 24.10]. The argument requires of X only that it be normal and that each of its open subsets be an F_σ set, which are precisely the defining properties of a perfectly normal space. The implication (1)⇒(3) holds whenever X is Polish. If f satisfies (1) and F is a nonempty closed subset of X, the restriction $f|_F$ is of Baire class 1, and its points of discontinuity therefore form a meager subset of F, as noted before Fact 1.2. Since F, being closed in a Polish space, is itself Polish and hence a Baire space, this meager set cannot exhaust F, and so $f|_F$ has a point of continuity. The reverse implication, (3)⇒(1), is due to Baire, who proved it in his thesis

1.1. **From Rosenthal's dichotomy to the Bourgain–Fremlin–Talagrand dichotomy to Shelah's NIP.** A deep computation is accessible to approximation when it is the limit of a *sequence* of standard computations. In a metrizable space every point of the closure of a set is the limit of a sequence of points of the set; in spaces of functions with the topology of pointwise convergence this description of the closure can fail, and the dichotomies of this subsection determine when it fails.

On the Cantor space $X = 2^{\mathbb{N}}$, the n-th coordinate map $p_n \colon X \to \{0,1\}$ is continuous for each $n \in \mathbb{N}$; however, the sequence $(p_n)_{n\in\mathbb{N}}$ has *no* convergent subsequence in $\mathbb{R}^X$ (see [Tod97, Chapter 1.1]). In a sense, this sequence exhibits the most extreme failure of sequential convergence: the closure of $\{p_n : n \in \mathbb{N}\}$ in $\{0,1\}^X$, or in $\mathbb{R}^X$, is homeomorphic to the Stone–Čech compactification $\beta\mathbb{N}$ of the discrete space of natural numbers. The space $\beta\mathbb{N}$ has cardinality $2^{2^{\aleph_0}}$, the largest possible for a separable Hausdorff space, and its points correspond to the ultrafilters on $\mathbb{N}$.[3]

The following theorem, proved by Haskell Rosenthal in 1974, shows that, for pointwise bounded sequences of continuous functions on a Polish space, a copy of $\beta\mathbb{N}$, such as the closure of the coordinate maps, is the only obstruction to the existence of a convergent subsequence.

Theorem 1.3 (Rosenthal's dichotomy [Ros74])**.** *If X is a Polish space and $(f_n)_{n\in\mathbb{N}}$ is a pointwise bounded sequence in $\mathrm{C_p}(X)$, then the sequence (f_n) has either a convergent subsequence or a subsequence whose closure in $\mathbb{R}^X$ is homeomorphic to $\beta\mathbb{N}$.*

The dichotomy grew out of *Rosenthal's ℓ_1-theorem*, which states that every bounded sequence in a Banach space has a subsequence that is either weakly Cauchy or equivalent to the unit vector basis of ℓ_1, the space of absolutely summable sequences; in particular, a Banach space contains no isomorphic copy of ℓ_1 if and only if every bounded sequence in it has a weakly Cauchy subsequence.

When the space $\mathrm{C_p}(X)$ is replaced by the larger space $\mathrm{B}_1(X)$, Bourgain, Fremlin and Talagrand obtained a counterpart of Rosenthal's dichotomy, usually stated as an equivalence rather than as a dichotomy (see Theorem 1.4 below).

To state the theorem, we introduce the following notation. For $f \in \mathbb{R}^X$ and a real number a, define

$$X^f_{\le a} \coloneqq f^{-1}(-\infty, a], \qquad\qquad X^f_{\ge a} \coloneqq f^{-1}[a, +\infty).$$

For a set $A \subseteq \mathbb{R}^X$ and a real number a, define, by a slight abuse of notation,

$$X^A_{\le a} \coloneqq \bigcap_{f\in A} X^f_{\le a} = \{x \in X : f(x) \le a \text{ for all } f \in A\},$$
$$X^A_{\ge a} \coloneqq \bigcap_{f\in A} X^f_{\ge a} = \{x \in X : f(x) \ge a \text{ for all } f \in A\};$$

in particular, $X^\emptyset_{\ge a} = X = X^\emptyset_{\le a}$. For a sequence $(f_n)_{n\in\mathbb{N}}$ in $\mathbb{R}^X$ and a set $I \subseteq \mathbb{N}$, define $I^{\complement} \coloneqq \mathbb{N} \setminus I$ and $f_I \coloneqq \{f_i : i \in I\}$.

---

for real-valued functions of real variables [Bai99]; for Polish spaces, see [Kec95, Theorem 24.15]. Finally, the last assertion of Fact 1.2 follows from the implication (3)⇒(1) together with the Baire Category Theorem.

[3]The proof of the existence of deep equilibria in compositional computation structures given by Alva, Dueñez, Iovino and Walton [ADIW26] relies on a classical result on the existence of idempotent elements of $\beta\mathbb{N}$.

Theorem 1.4 (The Bourgain–Fremlin–Talagrand dichotomy [BFT78])**.** *Let X be a Polish space, and let $A \subseteq \mathrm{C_p}(X)$ be pointwise bounded. The following are equivalent:*

(i) A is relatively compact in $\mathrm{B}_1(X)$.
(ii) For every $(f_n)_{n\in\mathbb{N}} \subseteq A$ and every $a < b$ there is $I \subseteq \mathbb{N}$ such that

$$X^{f_I}_{\leq a} \cap X^{f_{I^{\complement}}}_{\geq b} = \emptyset.$$

(In this form, the Bourgain–Fremlin–Talagrand dichotomy is a special case of the equivalence (ii)⇔(v) of Corollary 4G of Bourgain, Fremlin and Talagrand [BFT78].)

In condition (ii) of Theorem 1.4, the set $X^{f_I}_{\leq a}$ is defined by the inequalities $f_i(x) \leq a$ with $i \in I$, and the set $X^{f_{I^{\complement}}}_{\geq b}$ by the inequalities $f_j(x) \geq b$ with $j \in I^{\complement}$. Since I or $I^{\complement}$ is infinite, at least one of the two sets is defined by countably infinitely many inequalities. For our purposes it is more natural to impose only finitely many inequalities at a time, which leads to the following definitions.

Definition 1.5. A set $A \subseteq \mathbb{R}^X$ of functions has the *finitary No-Independence Property (NIP)* if, for every sequence $(f_n)_{n\in\mathbb{N}}$ in A and all reals $a < b$, there exist finite disjoint sets $E, F \subseteq \mathbb{N}$ such that $X^{f_E}_{\leq a} \cap X^{f_F}_{\geq b} = \emptyset$; we say that such sets E, F *witness finitary NIP for (f_n) and a, b.*

A set $A \subseteq \mathbb{R}^X$ has the *finitary Independence Property (IP)* if it does not have finitary NIP, that is, if there exist a sequence $(f_n)_{n\in\mathbb{N}}$ in A and reals $a < b$ such that $X^{f_E}_{\leq a} \cap X^{f_F}_{\geq b} \neq \emptyset$ for every pair of finite disjoint sets $E, F \subseteq \mathbb{N}$.

If the word "finite" is omitted in these definitions, we obtain *countable* NIP, which is weaker than finitary NIP, and *countable* IP, which is stronger than finitary IP. If, in addition, the witnesses are required to satisfy $F = E^{\complement}$, we obtain the properties that we call *BFT-NIP* and *BFT-IP*. Theorem 1.4 thus states that a pointwise bounded set $A \subseteq \mathrm{C_p}(X)$ is relatively compact in $\mathrm{B}_1(X)$ if and only if it has BFT-NIP.

From now on, IP and NIP mean *finitary* IP and NIP unless otherwise specified.

Theorem 1.6. *Let X be a compact metrizable (hence Polish) space. For every pointwise bounded set $A \subseteq \mathrm{C_p}(X)$, the following are equivalent:*

(i) A is relatively compact in $\mathrm{B}_1(X)$;
(ii) A has BFT-NIP;
(iii) A has countable NIP;
(iv) A has finitary NIP.

(The equivalence *(ii)⇔(iv)* holds for every compact space X, even without the assumption that A is pointwise bounded, and the equivalence *(ii)⇔(iii)* holds for every set $A \subseteq \mathbb{R}^X$ on an arbitrary space X.)

Proof. The equivalence (i)⇔(ii) is Theorem 1.4. The implications (iv)⇒(iii) and (ii)⇒(iii) are trivial, and (iii)⇒(ii) follows by taking $I = E$, since then $F \subseteq I^{\complement}$. To prove (ii)⇒(iv), suppose that (f_n) and $a < b$ witness finitary IP, and let $I \subseteq \mathbb{N}$ be given. The closed sets $X^{f_n}_{\leq a}$ $(n \in I)$ and $X^{f_m}_{\geq b}$ $(m \in I^{\complement})$ have the finite intersection property, so, by compactness, $X^{f_I}_{\leq a} \cap X^{f_{I^{\complement}}}_{\geq b} \neq \emptyset$. Since I is arbitrary, A does not have BFT-NIP, and (ii) fails. □

Theorem 1.6 can be read as a dichotomy: if X is compact metrizable and $A \subseteq \mathrm{C_p}(X)$ is pointwise bounded, then either A is relatively compact in $\mathrm{B}_1(X)$, or A has IP, in the finitary and, equivalently, in the countable sense.

The independence property was isolated by Saharon Shelah [She71, Definition 4.1] as one of the dividing lines of his classification of first-order theories [She90]. We return to it in Section 1.5, where it appears together with its counterparts in combinatorics and in statistical learning.

1.2. **Rosenthal compacta.** The dichotomy of Theorem 1.4 led Godefroy [God80] to the following definition.

Definition 1.7. A *Rosenthal compactum* is a topological space homeomorphic to a compact subset of $\mathrm{B}_1(X)$, with the topology of pointwise convergence, for some Polish space X.

Rosenthal compacta have strong tameness properties; for instance, every point in the closure of a subset of a Rosenthal compactum is the limit of a sequence of points of the subset [BFT78, Theorem 3F]. In this paper they serve as spaces of deep computations: for countably many features, the deep computations of an NIP set of computations form a Rosenthal compactum on each shard (see Theorem 3.1).

1.3. **The special case $\mathrm{B}_1(X, \mathbb{R}^{\mathcal{P}})$ with $\mathcal{P}$ countable.** A state of a computation is described by the values of its features, and a computation is therefore a map with values in a product of copies of the real line indexed by the features; in this subsection we reduce the study of such maps to the study of real-valued functions. Fix a countable set $\mathcal{P}$ (in this paper, "countable" means finite or countably infinite), whose elements we call *predicate symbols* or *formal predicates*. Our goal is to characterize the relatively compact subsets of $\mathrm{B}_1(X, \mathbb{R}^{\mathcal{P}})$, where X is a perfectly normal space, typically a Polish space.

Whenever $\mathcal{P}$ is regarded as a topological space, it carries the discrete topology. The spaces $\mathrm{C_p}(X, \mathbb{R}^{\mathcal{P}}) \subseteq \mathrm{B}_1(X, \mathbb{R}^{\mathcal{P}})$ are subspaces of the ambient space $(\mathbb{R}^{\mathcal{P}})^X$, which can be described in two other ways. The product $X \times \mathcal{P}$ is regarded either as a set or as a topological product, depending on the context. There are natural homeomorphisms

$$(\mathbb{R}^{\mathcal{P}})^X \cong \mathbb{R}^{X \times \mathcal{P}} \cong (\mathbb{R}^X)^{\mathcal{P}},$$

given by

$$\mathbb{R}^{X \times \mathcal{P}} \to (\mathbb{R}^{\mathcal{P}})^X : \varphi \mapsto \hat{\varphi}$$
$$\mathbb{R}^{X \times \mathcal{P}} \to (\mathbb{R}^X)^{\mathcal{P}} : \varphi \mapsto \check{\varphi},$$

where

$$\hat{\varphi}(x) \coloneqq \varphi(x, \cdot) \in \mathbb{R}^{\mathcal{P}}, \qquad \check{\varphi}(P) \coloneqq \varphi(\cdot, P) \in \mathbb{R}^X.$$

These identifications treat X and $\mathcal{P}$ as mere sets; in particular, the topology of X plays no role in them.

For $x \in X$, define the "left projection" map

$$\lambda_x : \mathbb{R}^{X \times \mathcal{P}} \to \mathbb{R}^{\mathcal{P}} : \varphi \mapsto \lambda_x(\varphi) \coloneqq \hat{\varphi}(x);$$

for $P \in \mathcal{P}$, the "right projection" map

$$\rho_P : \mathbb{R}^{X \times \mathcal{P}} \to \mathbb{R}^X : \varphi \mapsto \rho_P(\varphi) \coloneqq \check{\varphi}(P).$$

For fixed $x \in X$ and $P \in \mathcal{P}$, we also have canonical projection maps

$$\lambda_x : \mathbb{R}^X \to \mathbb{R} : f \mapsto f(x), \qquad \rho_P : \mathbb{R}^{\mathcal{P}} \to \mathbb{R} : f \mapsto f(P).$$

When clear from context, rather than using the specific symbols ("λ" for left, "ρ" for right) to denote projections, we may use the generic symbol "π"; thus, π_x may mean λ_x, and π_P may mean ρ_P.

The following proposition reduces the study of continuous functions, and of functions of Baire class 1, from X to $\mathbb{R}^{\mathcal{P}}$ to the case of real-valued functions.

Proposition 1.8. *The identification* $(\mathbb{R}^{\mathcal{P}})^X \cong \mathbb{R}^{X\times\mathcal{P}} \cong (\mathbb{R}^X)^{\mathcal{P}}$ *induces identifications*

$$\mathrm{C_p}(X, \mathbb{R}^{\mathcal{P}}) \cong \mathrm{C_p}(X \times \mathcal{P}) \cong \mathrm{C_p}(X)^{\mathcal{P}}, \quad \mathrm{B}_1(X, \mathbb{R}^{\mathcal{P}}) \cong \mathrm{B}_1(X \times \mathcal{P}) \cong \mathrm{B}_1(X)^{\mathcal{P}}.$$

(The cardinality of $\mathcal{P}$ plays no role here.)

Proof. The identification of the $\mathrm{C_p}$-spaces follows from the definition of the product topology and the discreteness of $\mathcal{P}$: a continuous map $X \to \mathbb{R}^{\mathcal{P}}$ is precisely a $\mathcal{P}$-indexed family of continuous functions $X \to \mathbb{R}$, and such families correspond to continuous functions $X \times \mathcal{P} \to \mathbb{R}$. The identification of the spaces of functions of Baire class 1 follows, since membership in the first Baire class is defined through pointwise limits of sequences of continuous functions, and a sequence of $\mathcal{P}$-indexed families of continuous functions is the same as a $\mathcal{P}$-indexed family of sequences of continuous functions. □

For $A \subseteq Y^X$ and $K \subseteq X$, we write $A|_K \coloneqq \{f|_K : f \in A\}$ for the set of restrictions to K of the functions in A. The following theorem extends Theorem 1.6 to $\mathbb{R}^{\mathcal{P}}$-valued functions.

Theorem 1.9. *Assume that $\mathcal{P}$ is countable, X is a Polish space, and $A \subseteq \mathrm{C_p}(X, \mathbb{R}^{\mathcal{P}})$ is pointwise bounded in the sense that $\pi_P \circ A$ ($\subseteq \mathrm{C_p}(X)$) is pointwise bounded for every $P \in \mathcal{P}$. The following are equivalent for every compact $K \subseteq X$:*

(i) $A|_K$ is relatively compact in $\mathrm{B}_1(K, \mathbb{R}^{\mathcal{P}})$;
(ii) $\pi_P \circ A|_K$ has NIP for every $P \in \mathcal{P}$.

Proof. By Proposition 1.8 and Tychonov's theorem, condition (i) holds if and only if $\pi_P \circ A|_K$ is relatively compact in $\mathrm{B}_1(K)$ for every $P \in \mathcal{P}$. Since K is compact metrizable, Theorem 1.6 shows that this is equivalent to (ii). □

In the applications, the NIP is verified on a set of states, whereas compactness is available only on its closure; the following lemma shows that the NIP passes from a set to its closure.

Lemma 1.10. *Assume that X is Hausdorff and that $A \subseteq \mathrm{C_p}(X)$. The following are equivalent for every $L \subseteq X$:*

(i) $A|_L$ satisfies the NIP;
(ii) $A|_{\overline{L}}$ satisfies the NIP.

Proof. The implication (ii)⇒(i) is trivial. Conversely, suppose that a sequence (f_n) in A and reals $a < b$ witness the IP on $\overline{L}$, and choose a' and b' with $a < a' < b' < b$. For finite disjoint $E, F \subseteq \mathbb{N}$, the open set $\bigcap_{n\in E} f_n^{-1}(-\infty, a') \cap \bigcap_{n\in F} f_n^{-1}(b', \infty)$ contains a point of $\overline{L}$, namely a point at which $f_n \leq a$ for $n \in E$ and $f_n \geq b$ for $n \in F$; hence it meets L. Therefore the sequence (f_n) and the reals $a' < b'$ witness the IP on L. □

1.4. **Five canonical Rosenthal compacta.** Five Rosenthal compacta, all built on the Cantor set, are the canonical members of the classes of Todorčević's trichotomy (see Theorem 3.3 below). By the trichotomy, the space of deep computations of each non-metrizable NIP example of Section 5 contains a copy of one of them; we fix their notation here. They are distinguished by three properties: a topological space is *separable* if it has a countable dense subset, *hereditarily separable* if every subspace is separable, and *first countable* if every point has a countable base of neighborhoods.

Let $2^{\mathbb{N}}$ carry the product topology and the lexicographic order $<$, whose order topology is the product topology. The length of a finite word $t \in 2^{<\mathbb{N}}$ is denoted $|t|$, and, for $a \in 2^{\mathbb{N}}$ and $n \in \mathbb{N}$, we write $a{\restriction}n$ for the word formed by the first n bits of a. Two words are *comparable* if one extends the other, and an *antichain* is a set of pairwise incomparable words. For a finite word $t \in 2^{<\mathbb{N}}$, let $t^\frown 0^\infty$ and $t^\frown 1^\infty$ be the sequences that begin with t and continue with all 0's and with all 1's, respectively; the set of sequences extending t is the clopen interval $[t^\frown 0^\infty, t^\frown 1^\infty]$.

For a statement S, we write $[\![S]\!]$ for its truth value: 1 if S is true and 0 if it is false. We introduce the following functions $2^{\mathbb{N}} \to \{0,1\}$:

- the *prefix test* $v_t \coloneqq [\![\cdot{\restriction}|t| = t]\!]$, the characteristic function of the set of sequences extending a fixed word $t \in 2^{<\mathbb{N}}$;
- the *equality test* $\delta_a \coloneqq [\![\cdot = a]\!]$, the characteristic function of the singleton $\{a\}$ for fixed $a \in 2^{\mathbb{N}}$;
- the *threshold tests* $\chi_{[a} \coloneqq [\![a \le \cdot]\!]$ and $\chi_{(a} \coloneqq [\![a < \cdot]\!]$, the characteristic functions of the final intervals $\{x : a \le x\}$ (closed) and $\{x : a < x\}$ (open), for fixed $a \in 2^{\mathbb{N}}$; the special case $\mathbf{0} \coloneqq \chi_{(1^\infty}$ is the identically zero function.

Prefix tests are continuous, and so are $\chi_{[t^\frown 0^\infty}$ and $\chi_{(t^\frown 1^\infty}$, since each of these functions depends only on the first $|t|$ bits of its argument; equality tests, and threshold tests at all other points, are discontinuous functions of Baire class 1. All closures below are taken in the topology of pointwise convergence.

(1) The *Alexandroff compactification* $A(2^{\mathbb{N}}) = \{\delta_a : a \in 2^{\mathbb{N}}\} \cup \{\mathbf{0}\}$. Each δ_a is isolated, and every neighborhood of $\mathbf{0}$ contains all but finitely many of the functions δ_a. Thus, $A(2^{\mathbb{N}})$ is neither separable nor first countable at $\mathbf{0}$.

(2) The *extended Alexandroff compactification* $\hat{A}(2^{\mathbb{N}}) = \overline{\{v_t : t \in 2^{<\mathbb{N}}\}} = \{v_t : t \in 2^{<\mathbb{N}}\} \cup A(2^{\mathbb{N}})$. As $n \to \infty$, the prefix tests $v_{a{\restriction}n}$ converge to δ_a; along an antichain they converge to $\mathbf{0}$. Thus, $\hat{A}(2^{\mathbb{N}})$ is separable but not first countable.

(3) The *split Cantor set* $S(2^{\mathbb{N}}) = \{\chi_{[a}, \chi_{(a} : a \in 2^{\mathbb{N}}\}$. The continuous thresholds $\chi_{[t^\frown 0^\infty}$, together with the isolated point $\mathbf{0} = \chi_{(1^\infty}$, form a countable dense subset: thresholds approaching a from below converge to $\chi_{[a}$, and from above to $\chi_{(a}$. Thus, $S(2^{\mathbb{N}})$ is separable, first countable and hereditarily separable, but not metrizable.

(4) The *Alexandroff duplicate* $D(2^{\mathbb{N}}) = \{(\mathbf{0}, a), (\delta_a, a) : a \in 2^{\mathbb{N}}\} \subseteq A(2^{\mathbb{N}}) \times 2^{\mathbb{N}}$. The points (δ_a, a) are isolated, and every neighborhood of $(\mathbf{0}, a)$ contains, for all b in some neighborhood of a, the point $(\mathbf{0}, b)$ and, with finitely many exceptions, the point (δ_b, b). Thus, $D(2^{\mathbb{N}})$ is first countable but not separable; it is contained in the separable compactum $\hat{D}(2^{\mathbb{N}}) = \overline{\{(v_t, t^\frown 0^\infty) : t \in 2^{<\mathbb{N}}\}} = D(2^{\mathbb{N}}) \cup \{(v_t, t^\frown 0^\infty) : t \in 2^{<\mathbb{N}}\}$.

(5) The *extended duplicate of the split Cantor set*

$$\hat{D}(S(2^{\mathbb{N}})) = \overline{\{(v_t, \chi_{[t^\frown 0^\infty)}\}_t} \subseteq \hat{A}(2^{\mathbb{N}}) \times S(2^{\mathbb{N}}),$$

which consists of the points $(v_t, \chi_{[t^\frown 0^\infty})$ and, for $a \in 2^{\mathbb{N}}$, the points $(\delta_a, \chi_{[a})$, $(\mathbf{0}, \chi_{[a})$ and $(\mathbf{0}, \chi_{(a)})$ [ADK08, 4.3.7]. Since the sequences $(a{\restriction}n)^\frown 0^\infty$ increase to a, the points $(v_{a{\restriction}n}, \chi_{[(a{\restriction}n)^\frown 0^\infty})$ converge to $(\delta_a, \chi_{[a})$ as $n \to \infty$. Thus, $\hat{D}(S(2^{\mathbb{N}}))$ is separable and first countable; since the points $(\delta_a, \chi_{[a})$ form an uncountable discrete subspace, it is not hereditarily separable.

The constructions of the Alexandroff compactification and of the Alexandroff duplicate make sense with $2^{\mathbb{N}}$ replaced by any compact metric space K; the resulting spaces are denoted $A(K)$ and $D(K)$, respectively. Since $A(K)$ is the one-point compactification of a discrete space of the cardinality of K, it depends only on that cardinality; for instance, $A([0,1])$ is homeomorphic to $A(2^{\mathbb{N}})$.

1.5. **NIP in model theory, combinatorics and learning: polynomial versus exponential growth.** For families of $\{0,1\}$-valued functions, that is, of sets, a combinatorial dividing line closely related to the NIP of Definition 1.5 appeared in 1971–1972, some years before Theorem 1.4, independently and almost simultaneously in three fields: in model theory, in the work of Saharon Shelah [She71, She90]; in combinatorics, in the work of Norbert Sauer [Sau72] and, again, of Shelah [She72, She90]; and in statistical learning, in the work of Vladimir Vapnik and Alexey Chervonenkis [VC71, VC74]. In each of them, the line separates the "tame" families of sets, whose number of traces on an n-element set (the intersections of the members of the family with the set) is bounded by a polynomial in n, from the "wild" ones, for which, for every $n \in \mathbb{N}$, some n-element set has all 2^n possible traces. The precise relation with the NIP is explained at the end of this subsection.

In model theory: Among the dividing lines of Shelah's classification theory [She90], the independence property is detected by counting types over finite sets of parameters. Let T be a complete first-order theory, let $\varphi(x;y)$ be a formula of its language, in which x and y are finite tuples of variables of lengths $|x|$ and $|y|$, and let B be a finite set of $|y|$-tuples of elements of a model M of T. A *φ-type* over B is a maximal set of formulas $\varphi(x;b)$ and $\neg\varphi(x;b)$, with $b \in B$, that is consistent with the theory of M with constants for the elements of M; since B is finite, every φ-type over B is realized by an $|x|$-tuple a of elements of M, and it is determined by the set $\{b \in B : M \models \varphi(a;b)\}$. Hence the set $S_\varphi(B)$ of φ-types over B is in bijection with the set of traces on B of the family of sets $\{b : M \models \varphi(a;b)\}$, with a ranging over $|x|$-tuples of elements of M, and $|S_\varphi(B)| \leq 2^{|B|}$. The formula φ has the *independence property* (IP) if, for every $n \in \mathbb{N}$, some set B of n tuples satisfies $|S_\varphi(B)| = 2^n$; otherwise the formula is *NIP*, or *dependent*, and the theory T is NIP if all its formulas are NIP. Since T is complete, these notions do not depend on the model M.

If φ is NIP, the Sauer–Shelah lemma below bounds $|S_\varphi(B)|$ by a polynomial in $|B|$. Consequently, φ is NIP if and only if the number of φ-types over a finite set A of parameters (that is, with b ranging over the $|y|$-tuples of elements of A) is bounded by a polynomial in $|A|$; if φ has IP, then for every n some set A of at most $n|y|$ parameters satisfies $|S_\varphi(A)| \geq 2^n$ [Sim15, Section 2.3.4]. Over infinite sets of parameters, counting φ-types does not

separate NIP from IP without further set-theoretic hypotheses: if κ is infinite and $2^\kappa = \kappa^+$, every unstable formula, NIP or not, already has 2^κ φ-types over some set of κ parameters [Sim15, Section 2.3.4].

In combinatorics: Sauer [Sau72] and Shelah [She72] proved independently the following dichotomy, now known as the *Sauer–Shelah lemma*. Let $\mathscr{F} = \{S_i : i \in I\}$ be a family of subsets of a set S, indexed by a set I. Then either for every $n \in \mathbb{N}$ there is a set $A \subseteq S$ with $|A| = n$ such that $|\{S_i \cap A : i \in I\}| = 2^n$, in which case $\mathscr{F}$ is said to have "exponential complexity", or there exists $N \in \mathbb{N}$ such that, for every $A \subseteq S$ with $|A| \geq N$,

$$|\{S_i \cap A : i \in I\}| \leq \sum_{j=0}^{N-1} \binom{|A|}{j},$$

in which case $\mathscr{F}$ is said to have "polynomial complexity" (every family of subsets of a *finite* set S has polynomial complexity). The name is justified: for fixed $N > 0$ and for $|A| = m > 0$, we have

$$\sum_{j=0}^{N-1} \binom{m}{j} \leq \sum_{j=0}^{N-1} \frac{m^j}{j!} \leq \left(\sum_{j=0}^{N-1} \frac{1}{j!} \right) \cdot m^{N-1} < e \cdot m^{N-1},$$

a polynomial bound in m.

In machine learning: The Sauer–Shelah lemma is also the dichotomy that Vapnik and Chervonenkis [VC71, VC74] had discovered and proved slightly earlier, in 1971, in their study of uniform convergence in statistics. The largest size of a set $A \subseteq S$ with $2^{|A|}$ traces $S_i \cap A$, that is, the integer $N - 1$ for the least integer N as in the Sauer–Shelah lemma, is called the *VC dimension* of $\mathscr{F}$; if no such integer N exists, the VC dimension of $\mathscr{F}$ is infinite. Through the uniform convergence theorem of Vapnik and Chervonenkis, the lemma bounds the number of data points (the sample size) needed to "learn" a concept class $\mathscr{F}$ of finite VC dimension in a statistical sense, that is, up to a preset error and with a preset confidence. The Fundamental Theorem of Statistical Learning states that a class of $\{0,1\}$-valued functions is PAC-learnable (PAC stands for "Probably Approximately Correct" [Val84]) if and only if its VC dimension is finite, provided that the class satisfies a mild measurability condition [BEHW89]; see also [SSBD14, Theorem 6.7].

For first-order formulas the three dichotomies coincide: by the description of φ-types given above, a formula $\varphi(x;y)$ has IP exactly when the family of sets $\{b : M \models \varphi(a;b)\}$, with a ranging over the $|x|$-tuples of elements of a model M of T, has infinite VC dimension, and for an NIP formula the Sauer–Shelah lemma gives the polynomial bound on the number of φ-types. For an arbitrary family of sets, finiteness of the VC dimension implies that the characteristic functions of its members have the NIP of Definition 1.5; the converse fails, since the finite subsets of $\mathbb{N}$ form a family of infinite VC dimension whose characteristic functions have the NIP.

Historical remarks. The Baire hierarchy of functions was introduced by René-Louis Baire in his 1899 doctoral thesis, *Sur les fonctions de variables réelles*. The field of C_p-theory was pioneered by A. V. Arhangel'skiĭ and his students in the

1970s and 1980s [Ark92], and it has found many applications in model theory and functional analysis. For a recent survey, see [Tka11].

2. Compositional Computation Structures: A structural approach to arbitrary-precision computations

In this section we connect function spaces with arbitrary-precision computations. A computation on real-valued data is observed through its features, each known to finite precision, and a deep computation is a limit of computations in the topology of this observation, the topology of pointwise convergence. We first recall the framework of the first paper of this series [ADIW26]: states are described by their types, the tuples of values of all their features; computations act on states and are composed; and deep computations are the pointwise limits of computations, taken in the space of types. We then recall the stable sets of computations, studied in that paper, and close the section with two examples that motivate the classification of Section 3.

2.1. **Computation States Structures.** A *computation states structure* is a pair $(L, \mathcal{P})$, where L is a set whose elements we call *states* and $\mathcal{P}$ is a collection of real-valued functions on L that we call *predicates* or *features*. For $P \in \mathcal{P}$ and $v \in L$, the value $P(v)$ is the P-th *(observable) feature* of v. Maps $f\colon L \to L$ are called *transitions* of $(L, \mathcal{P})$; the computations of Definition 2.4 below are transitions.

The *type* of a state $v \in L$ is the indexed family[4]

$$\mathrm{tp}(v) = (P(v))_{P\in\mathcal{P}} \in \mathbb{R}^{\mathcal{P}}$$

of its features.

Intuitively, L is the set of states of a computation, and the predicates $P \in \mathcal{P}$ are the primitives that are given and accepted as computable.

We assume that each state $v \in L$ is determined by its type $\mathrm{tp}(v)$; we may then identify L with its image $\mathrm{tp}[L] \subseteq \mathbb{R}^{\mathcal{P}}$ under the type map and give it the subspace topology, that is, the topology of pointwise convergence induced by the product topology of $\mathbb{R}^{\mathcal{P}}$. In particular, for each $P \in \mathcal{P}$, the map $v \mapsto P(v)$ is continuous on L.

Important state spaces include $L = \mathbb{R}^{\mathbb{N}}$, with the predicates $P_i(v) = v_i$ for $i \in \mathbb{N}$, and $L = \mathbb{R}^n$, for a positive integer n, with the predicates $P_i(v) = v_i$ for $0 \le i < n$. Subspaces L of $\mathbb{R}^{\mathbb{N}}$ and of $\mathbb{R}^n$ are of primary interest in this paper; through the identification of each state with its type, every computation states structure with countably many predicates is such a subspace.

Definition 2.1. Let $(L, \mathcal{P})$ be a computation states structure. The type $\mathrm{tp}(v) \in \mathbb{R}^{\mathcal{P}}$ of a state $v \in L$ is called a *realized* (state) type. The closure of the set of realized types in $\mathbb{R}^{\mathcal{P}}$, with the topology of pointwise convergence, is called the *space of (state) types* of $(L, \mathcal{P})$ and is denoted $\mathcal{L}$; its elements $\mathbf{v} \in \mathcal{L}$ are called *state types*, and those in $\mathcal{L} \setminus L$ are called *unrealized* state types.

[4]As in Proposition 1.8, we regard each predicate as named by a formal label P, a *predicate symbol*, whose meaning in the computation states structure is the real function $P(\cdot)\colon v \mapsto P(v)$ on L. The symbol P thus names an observable feature, whose value at a state $v \in L$ is $P(v)$. Whenever $\mathcal{P}$ is used as an index set, as in the product $\mathbb{R}^{\mathcal{P}}$ below, it is regarded as a set of symbols. The distinction between a symbol and its meaning is usually suppressed in mathematics, but the model-theoretic framework requires awareness of it. When necessary, we write $P(\cdot)$ for the function on L and P for the symbol; when no confusion can arise, we also write P for the function.

Intuitively, state types capture a notion of "limit state". For each $P \in \mathcal{P}$, the P-th coordinate map $\mathbb{R}^{\mathcal{P}} \to \mathbb{R}$, $\mathbf{v} = (\mathbf{v}_Q)_{Q\in\mathcal{P}} \mapsto \mathbf{v}_P$, restricts to a continuous extension to $\mathcal{L}$ of the predicate $P(\cdot)$ on L; abusing notation, we still denote this extension by $P(\cdot)$, or just P.

Remark 2.2. One may regard unrealized state types as "hyper-states"; when such a type is obtained as the limit of a sequence (or as an ultralimit) of realized state types, it may be called a "deep state". For readers versed in model theory we note that, at least for the purpose of realizing state types with parameters in the state space L, the type space $(\mathcal{L}, \mathcal{P})$ serves as a "poor man's" ultraproduct of $(L, \mathcal{P})$ in which, by the very definition of $\mathcal{L}$, every type $\mathbf{v}$ is realized.

2.1.1. *Bounds on precision and magnitude. Shards.* Computation states structures are meant to model real-world computing, in which computations with real-valued quantities can only be carried out approximately, to a preset degree of precision. No less important in practice is a bound on the magnitude of the quantities involved: no machine can store, let alone compute with, arbitrarily large numbers. Thus, although the collection $\mathcal{P}$ of predicates may be infinite, even uncountable, an actual computation involves bounds, chosen at the outset, for errors and for the absolute magnitudes of the quantities involved; different parts of a computation may require different error bounds, and may deal with quantities of very different sizes. Focusing on the second of these issues, we consider sets of computations for which a bound for each predicate is fixed in advance. This leads to the notion of *shard*, introduced in the first paper of this series [ADIW26].

Definition 2.3. Let $(L, \mathcal{P})$ be a computation states structure, with L identified with the set $\mathrm{tp}[L] \subseteq \mathbb{R}^{\mathcal{P}}$ of its realized types. A *sizer* is a family $r_\bullet = (r_P)_{P\in\mathcal{P}}$ of positive real numbers indexed by $\mathcal{P}$. For a sizer $r_\bullet$, let $\mathbb{R}[r_\bullet] \coloneqq \prod_{P\in\mathcal{P}}[-r_P, r_P]$, a compact space, and define the $r_\bullet$-*shard* of the state space L by

$$L[r_\bullet] = L \cap \mathbb{R}[r_\bullet].$$

The $r_\bullet$-*type shard* is the closure $\mathcal{L}[r_\bullet] = \overline{L[r_\bullet]}$, a closed and hence compact subset of $\mathbb{R}[r_\bullet]$.

The union of all type shards, as the sizer $r_\bullet$ varies, is denoted $\mathcal{L}_{\mathrm{sh}}$ and called the space of *shard-supported types*.

We have $\mathcal{L}_{\mathrm{sh}} \subseteq \mathcal{L}$, and the inclusion may be proper; equality holds in the important special case when $\mathcal{P}$ is countable (see [ADIW26]).

2.2. **Compositional Computation Structures.**

Definition 2.4. A *compositional computation structure* (CCS) is a triple $(L, \mathcal{P}, \Gamma)$, where

- $(L, \mathcal{P})$ is a computation states structure, and
- $\Gamma \subseteq L^L$ is a semigroup under composition that contains the identity map id of L, that is, a monoid of transformations of L.

The elements of Γ are called the *computations* of $(L, \mathcal{P}, \Gamma)$. The state space L is identified with a subset of the type space $\mathcal{L} \subseteq \mathbb{R}^{\mathcal{P}}$ of $(L, \mathcal{P})$.

We topologize Γ as a subset of the product L^L, in which, as usual, the exponent L serves merely as an index set, while the base $L \subseteq \mathbb{R}^{\mathcal{P}}$ carries the topology of pointwise convergence. The semigroup Γ is thus a subset of the product $\mathcal{L}^L \subseteq$

$(\mathbb{R}^{\mathcal{P}})^L$, and its closure $\overline{\Gamma}$ is contained in $\mathcal{L}^L$. The elements $\xi \in \overline{\Gamma}$ are called (real-valued) *deep computations* or *ultracomputations*.

A collection R of sizers is *exhaustive* if $L = \bigcup_{r_\bullet \in R} L[r_\bullet]$. A computation $\gamma \in \Gamma$ is *R-confined* if it maps $L[r_\bullet]$ into itself for every $r_\bullet \in R$, and a subset $\Delta \subseteq \Gamma$ is *R-confined* if each $\gamma \in \Delta$ is.

Unlike its subspace L^L, which is a semigroup under composition, the space $\mathcal{L}^L$ has no natural semigroup structure; deep computations are therefore not composable in general.

Proposition 2.5. *If $\Delta \subseteq \Gamma$ is confined by an exhaustive collection of sizers, then $\overline{\Delta}$ is a compact subset of $\mathcal{L}^L_{\mathrm{sh}}$.*

Proof. Let R be an exhaustive collection of sizers that confines Δ, and choose, for each $v \in L$, a sizer $r_\bullet^{(v)} \in R$ with $v \in L[r_\bullet^{(v)}]$. Since every $\gamma \in \Delta$ maps v into $L[r_\bullet^{(v)}]$, the set Δ lies in the compact space $\prod_{v \in L} \mathcal{L}[r_\bullet^{(v)}] \subseteq \mathcal{L}^L_{\mathrm{sh}}$, which therefore contains $\overline{\Delta}$. □

For a CCS $(L, \mathcal{P}, \Gamma)$, we regard the elements of Γ as "standard" finitary computations, and the elements of $\overline{\Gamma}$, i.e., deep computations, as possibly infinitary limits of standard computations.

Remarks 2.6. (For readers versed in model theory.)[5]

(1) Compositional computation structures, as defined above, are not literally real-valued structures, because nothing in the structure itself forces Γ to be realized as a subset of the product space L^L. However, as in their original definition [ADIW26], they may be regarded as real-valued structures $(L, \mathcal{P}, \Gamma, \circ, \mathrm{ev})$, where
   - $(L, \mathcal{P})$ is a computation states structure,
   - $(\Gamma, \circ)$ is a semigroup,
   - $\mathrm{ev} : \Gamma \times L \to L : (\gamma, v) \mapsto \mathrm{ev}(\gamma, v)$ is a semigroup action of Γ on L (i.e., $\mathrm{ev}(\gamma \circ \delta, v) = \mathrm{ev}(\gamma, \mathrm{ev}(\delta, v))$ for $\gamma, \delta \in \Gamma$ and $v \in L$).

   The two definitions are reconciled by identifying each $\gamma \in \Gamma$ with the map $v \mapsto \mathrm{ev}(\gamma, v)$.

(2) Regarded as real-valued structures in this sense, the compositional computation structures with a fixed collection $\mathcal{P}$ of predicate symbols form an elementary class. The notion of state type introduced above (essentially, that of a quantifier-free type with parameters of sort L only) may be enriched by further formulas, for instance with parameters in Γ or with quantifiers, and extended to types of sort Γ ("transform-types"). As usual, all types over a structure $\underline{C} = (L, \mathcal{P}, \Gamma, \circ, \mathrm{ev})$ are realized in some ultrapower $\underline{C}^{(\mathcal{U})}$ of $\underline{C}$, constructed as in the references above.[6] In Banach space theory, this approach to spaces of types was introduced by Krivine [Kri74, KM81, Kri72].

2.3. **Computability of deep computations and the Extendibility Axiom.** A machine evaluates only sums and products of finitely many observed features, and only to within a preset error on a preset range of magnitudes; for this reason,

[5] The real-valued structures used here are those of Keisler's model theory for real-valued structures [Kei23]; see also [DnI17, Section 6] for an introduction.

[6] By contrast, no ultrapower is needed to realize state types in a computation states structure (see Remark 2.2).

computability is defined by uniform approximation on shards by polynomials in the features. Throughout this subsection, we fix a computation states structure $(L, \mathcal{P})$ with a *countable* collection $\mathcal{P}$ of predicates.

(1) A *computable predicate* is a function $\varphi\colon L \to \mathbb{R}$ whose restriction to each shard is a uniform limit of polynomials, with real coefficients, in the features $P(\cdot)$.
(2) A transform $f\colon L \to \mathcal{L}$ is *computable* if, for each $Q \in \mathcal{P}$, the output feature $Q \circ f\colon L \to \mathbb{R}$ is a computable predicate.

Alva, Dueñez, Iovino and Walton [ADIW26] showed that a transform $f\colon L \to \mathcal{L}$ is computable, that is, its output features are, on each shard, uniform limits of polynomials in the features, if and only if it extends to a continuous map $\tilde{f}\colon \mathcal{L} \to \mathcal{L}$. This characterization motivates the following definition, in which continuity is required on each type shard.

A computation $\gamma \in \Gamma$ is *extendable* if it has an extension $\tilde{\gamma}\colon \mathcal{L}_{\mathrm{sh}} \to \mathcal{L}_{\mathrm{sh}}$ such that, for every sizer $r_\bullet$, there is a sizer $s_\bullet$ for which the restriction $\tilde{\gamma}|_{\mathcal{L}[r_\bullet]}\colon \mathcal{L}[r_\bullet] \to \mathcal{L}[s_\bullet]$ is continuous. Such an extension is unique, since it is continuous on each type shard $\mathcal{L}[r_\bullet]$, in which $L[r_\bullet]$ is dense. For a set $\Delta \subseteq \Gamma$ of extendable computations, we write $\tilde{\Delta}$ for the set $\{\tilde{\gamma}\colon \gamma \in \Delta\}$.

Extendibility Axiom. *Every computation $\gamma \in \Gamma$ is extendable.*

The Extendibility Axiom is discussed at length in the first paper of this series [ADIW26]. *From now on, all compositional computation structures are assumed to satisfy the Extendibility Axiom.*

The following lemma extends deep computations to the type shards; it is used in Theorems 3.1 and 6.7.

Lemma 2.7. *Let R be an exhaustive collection of sizers and let $\Delta \subseteq \Gamma$ be R-confined. Every deep computation $f \in \overline{\Delta}$ has an extension $\tilde{f} : \mathcal{L}_{\mathrm{sh}} \to \mathcal{L}_{\mathrm{sh}}$ such that, for every $r_\bullet \in R$, $\tilde{f}$ maps $\mathcal{L}[r_\bullet]$ into itself and $\tilde{f}|_{\mathcal{L}[r_\bullet]} \in \overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$.*

Proof. Write $f = \mathcal{U}\lim_i \gamma_i$, where $(\gamma_i)_{i\in I}$ is a family in Δ and $\mathcal{U}$ is an ultrafilter on the index set I. Let $r_\bullet \in R$. Each $\tilde{\gamma}_i$ maps $L[r_\bullet]$ into itself and is continuous on $\mathcal{L}[r_\bullet]$, so it maps the closure $\mathcal{L}[r_\bullet]$ of $L[r_\bullet]$ into itself; since $\mathcal{L}[r_\bullet]$ is compact, the limit $\tilde{f}(v) \coloneqq \mathcal{U}\lim_i \tilde{\gamma}_i(v)$ exists for every $v \in \mathcal{L}[r_\bullet]$ and lies in $\mathcal{L}[r_\bullet]$, and $\tilde{f}|_{\mathcal{L}[r_\bullet]} = \mathcal{U}\lim_i \tilde{\gamma}_i|_{\mathcal{L}[r_\bullet]}$. Since R is exhaustive, this defines $\tilde{f}$ on a set that contains L, and there $\tilde{f}$ agrees with f; at the remaining points of $\mathcal{L}_{\mathrm{sh}}$ we let $\tilde{f}$ be the identity. □

2.4. **Stability.** By the theorem of Alva, Dueñez, Iovino and Walton recalled above, the computable transforms are exactly those that extend to continuous maps on the space of types. The sets of computations all of whose deep computations are computable in this sense sit at the top of the hierarchy studied in this paper.

Definition 2.8. A set $\Delta \subseteq \Gamma$ of computations is *stable* if, for every sizer $r_\bullet$, every deep computation in $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$ is continuous, that is, computable.

Stable sets of computations are the subject of the first paper of this series [ADIW26], where they are characterized by Grothendieck's double limit criterion and identified with stability in the sense of model theory.

2.5. **Two motivating examples.** Before developing the classification, we describe two computations whose deep computations behave very differently. Both are treated in detail in Section 4.

Square roots. Newton's method for $p(x) = x^2 - a$, with $a > 0$, acts on the real projective line $\mathbb{RP}^1 = \mathbb{R} \cup \{\infty\}$ by the map $N_p(x) = (x^2 + a)/(2x)$. Its iterates are continuous, and they converge pointwise to the map $\mathfrak{f}$ that sends every positive number to $\sqrt{a}$, every negative number to $-\sqrt{a}$, and both 0 and ∞ to ∞ (see Figure 1). The deep computations are the iterates together with $\mathfrak{f}$. Since $\mathfrak{f}$ is discontinuous, it is not computable, and the set of iterates is not stable; nevertheless, the map $\mathfrak{f}$ is the limit of a *sequence* of computations, and the space of deep computations is countable.

Newton's method on the Riemann sphere. For $\rho(z) = z^3 - 2z + 2$, Newton's method on the Riemann sphere $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ behaves in three different ways on three parts of the sphere (see Figure 2 in Section 4.1). On the basins of the three roots the iterates converge. On the basin of the cycle $0 \mapsto 1 \mapsto 0$ they do not, although the even iterates and the odd iterates converge there separately, to different limits. On the common boundary of these basins, a fractal set, the iterates are chaotic; for the simpler polynomial $z^2 - 1$, chaos of this kind produces deep computations that are not even limits of sequences of computations (see Proposition 5.10 and Theorem 3.1).

The classification of Section 3 is designed to separate such behaviors. Theorem 3.1 characterizes the sets of computations whose deep computations are limits of sequences of computations, as in the case of square roots, and Todorčević's trichotomy measures the complexity of the space of such deep computations.




FIGURE 1. Newton iterates N_p^n for $p(x) = x^2 - a$ on the real projective line, drawn on the circle with ∞ at both ends of each axis. The continuous maps N_p^n converge pointwise to the deep computation $\mathfrak{f}$, which sends the positive half line to $\sqrt{a}$, the negative half line to $-\sqrt{a}$, and 0 and ∞ to ∞.

3. Classifying deep computations

This section contains the main theorems of the paper. The examples of Section 2.5 show that some natural algorithms have discontinuous deep computations that are nevertheless limits of sequences of computations, while others have deep computations that are not even limits of sequences. We use the topology of function spaces to classify sets of computations by the complexity of their deep computations, and thereby obtain a new hierarchy of complexity classes. The main topological tool is Todorčević's trichotomy for Rosenthal compacta [Tod99], and our model is the work of Glasner and Megrelishvili on tame semigroups in topological dynamics [GM22].

3.1. **NIP, Rosenthal compacta, and deep computations.** Theorem 3.1, on which the classification rests, extends to deep computations of the first Baire class results that the first paper of this series [ADIW26] established for continuous deep computations.

Theorem 3.1. *Let $(L, \mathcal{P}, \Gamma)$ be a compositional computation structure (see Definition 2.4) satisfying the Extendibility Axiom, with $\mathcal{P}$ countable. Let R be an exhaustive collection of sizers, and suppose that $\Delta \subseteq \Gamma$ is R-confined. Then the following conditions are equivalent:*

(i) $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}} \subseteq \mathrm{B}_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$ *for all* $r_\bullet \in R$*;*

(ii) $\pi_P \circ \Delta|_{L[r_\bullet]}$ *satisfies the NIP for all* $P \in \mathcal{P}$ *and* $r_\bullet \in R$*; that is, for all* $P \in \mathcal{P}$*,* $r_\bullet \in R$*, reals* $a < b$ *and sequences* $(\gamma_n)_{n\in\mathbb{N}}$ *in* Δ*, there are finite disjoint sets* $E, F \subseteq \mathbb{N}$ *such that*

$$L[r_\bullet] \cap \bigcap_{n\in E} (\pi_P \circ \gamma_n)^{-1}(-\infty, a] \cap \bigcap_{n\in F} (\pi_P \circ \gamma_n)^{-1}[b, \infty) = \emptyset.$$

Furthermore, if these conditions hold, then every deep computation $f \in \overline{\Delta}$ has an extension $\tilde{f}\colon \mathcal{L}_{sh} \to \mathcal{L}_{sh}$ such that $\tilde{f}|_{\mathcal{L}[r_\bullet]} \in \mathrm{B}_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$ for all $r_\bullet \in R$. In particular, on each shard, every deep computation is the pointwise limit of a sequence of computations.

Proof. Since $\mathcal{P}$ is countable, each type shard $\mathcal{L}[r_\bullet]$ is a compact metrizable, hence Polish, subspace of $\mathbb{R}^{\mathcal{P}}$. By the Extendibility Axiom and confinement, $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ is a pointwise bounded set of continuous functions for all $P \in \mathcal{P}$ and $r_\bullet \in R$. The equivalence of (i) and (ii) therefore follows from Theorem 1.9 and Lemma 1.10. If (i) holds, the extension $\tilde{f}$ of Lemma 2.7 satisfies $\tilde{f}|_{\mathcal{L}[r_\bullet]} \in \overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}} \subseteq \mathrm{B}_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$ for all $r_\bullet \in R$. Finally, for a Polish space X, every compact subset of $\mathrm{B}_1(X)$ is Fréchet–Urysohn, that is, every point in the closure of a subset is the limit of a sequence of points of the subset (see [BFT78, Theorem 3F] or [Deb13, Theorem 4.1]). Applied to $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$, regarded by Proposition 1.8 as a compact subset of $\mathrm{B}_1(\mathcal{L}[r_\bullet] \times \mathcal{P})$, this shows that every deep computation is, on each shard, the pointwise limit of a sequence of computations. □

3.2. **Todorčević's trichotomy and levels of NIP.** By Theorem 3.1, under its hypotheses, the deep computations of a countable NIP set of computations form, on each type shard, a separable Rosenthal compactum. We measure the complexity of such a compactum by three topological properties, metrizability, hereditary separability and first countability, which form a chain: every compact metrizable

space is hereditarily separable, and R. Pol proved that every hereditarily separable Rosenthal compactum is first countable (see [Deb13, Section 10]). These properties define three levels of NIP.

Definition 3.2. Let $(L, \mathcal{P}, \Gamma)$ be a CCS satisfying the Extendibility Axiom, and let R be an exhaustive collection of sizers. Let $\Delta \subseteq \Gamma$ be an R-confined countable set of computations satisfying the NIP on shards and features (condition (ii) in Theorem 3.1). We say that Δ is:


(i) NIP_1 if $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$ is first countable for every $r_\bullet \in R$;
(ii) NIP_2 if $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$ is hereditarily separable for every $r_\bullet \in R$;
(iii) NIP_3 if $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$ is metrizable for every $r_\bullet \in R$.

By the chain of properties above, the levels satisfy $\mathrm{NIP}_3 \Rightarrow \mathrm{NIP}_2 \Rightarrow \mathrm{NIP}_1 \Rightarrow$ NIP. Moreover, a countable stable set of computations (see Definition 2.8), which has the NIP by Theorem 3.1, is NIP_3: on each shard its deep computations form a separable compact subset of a countable product of spaces $\mathrm{C_p}(\mathcal{L}[r_\bullet])$, that is, a separable Eberlein compactum, and separable Eberlein compacta are metrizable [Ark92]. The five compacta of Section 1.4 show that the converse implications fail for Rosenthal compacta: $\hat{A}(2^{\mathbb{N}})$ is separable but not first countable; $\hat{D}(S(2^{\mathbb{N}}))$ and $\hat{D}(2^{\mathbb{N}})$ are separable and first countable but not hereditarily separable; and $S(2^{\mathbb{N}})$ is hereditarily separable but not metrizable. Todorčević proved that these examples are canonical [Tod99], in the following sense.

Theorem 3.3 (Todorčević's trichotomy [Tod99]; see also [ADK08, Theorem 3])**.** *Let K be a separable Rosenthal compactum.*

(i) If K is hereditarily separable but non-metrizable, then $S(2^{\mathbb{N}})$ embeds into K.
(ii) If K is first countable but not hereditarily separable, then either $D(2^{\mathbb{N}})$ or $\hat{D}(S(2^{\mathbb{N}}))$ embeds into K.
(iii) If K is not first countable, then $A(2^{\mathbb{N}})$ embeds into K.

We thus have the following classification for separable Rosenthal compacta:



The three levels can also be described by the number of inputs needed to tell deep computations apart. Let Δ and R be as in Definition 3.2, fix $r_\bullet \in R$, and write $K_{r_\bullet} = \overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$. We identify each $f \in K_{r_\bullet}$ with the real-valued function $(P, x) \mapsto P(f(x))$ on $X_{r_\bullet} = \mathcal{P} \times \mathcal{L}[r_\bullet]$, whose points we call *tests*: the test (P, x) applies the feature P to the output on the input x. The topology of pointwise convergence on $K_{r_\bullet}$ is then the topology of pointwise convergence on tests. A set

D of tests *separates* a set of deep computations if any two distinct members of the set differ on some test in D.

Proposition 3.4. *Let Δ be as in Definition 3.2 and $r_\bullet \in R$.*

(i) $K_{r_\bullet}$ is metrizable if and only if one countable set of tests separates $K_{r_\bullet}$.
(ii) $K_{r_\bullet}$ is first countable if and only if every $f \in K_{r_\bullet}$ is countably determined*: some countable set D_f of tests separates f from every other member of $K_{r_\bullet}$.*
(iii) $K_{r_\bullet}$ is hereditarily separable if and only if there is no set $E \subseteq K_{r_\bullet}$ of cardinality $2^{\aleph_0}$ whose members are finitely separated*: for each $f \in E$ there are finitely many tests and $\varepsilon_f > 0$ such that every $g \in E \setminus \{f\}$ differs from f by at least ε_f on one of them.*

Proof. (i) If a countable set D of tests separates $K_{r_\bullet}$, restriction to D is a continuous injection of the compact space $K_{r_\bullet}$ into the metrizable space $\mathbb{R}^D$, hence a homeomorphism onto its image. Conversely, if $K_{r_\bullet}$ is metrizable, then $K_{r_\bullet} \times K_{r_\bullet}$ is second countable, so the open set $\{(f,g) : f \neq g\}$ is Lindelöf; it is covered by the open sets $\{(f,g) : f(t) \neq g(t)\}$, for tests t, and a countable subcover gives the required set D. (ii) If f is countably determined, the singleton $\{f\}$ is a countable intersection of open sets, and in a compact Hausdorff space such a point has a countable base of neighborhoods. Conversely, if $(U_k)_{k\in\mathbb{N}}$ is a countable base at f and each U_k contains a basic neighborhood of f defined by a finite set F_k of tests, then $D_f = \bigcup_k F_k$ separates f from every other member of $K_{r_\bullet}$. (iii) A set is finitely separated exactly when it is a discrete subspace, and, by Theorem 3.1, $K_{r_\bullet}$ is a separable Rosenthal compactum; for such compacta, hereditary separability is equivalent to the absence of discrete subspaces of cardinality $2^{\aleph_0}$, by a theorem of Todorčević [Tod99] (see also [GM22, Theorem 0.4]). □

Thus the set Δ is NIP_3 when, on each shard, a single countable set of tests separates all its deep computations; it is NIP_2 when no shard carries a finitely separated set of deep computations of the cardinality of the continuum; and it is NIP_1 when, on each shard, each deep computation is separated from all the others by countably many tests of its own.

In Section 5 we exhibit computations at each level of the hierarchy and show that every implication between the classes stable, NIP_3, NIP_2, NIP_1 and NIP is strict. The definitions of NIP_1, NIP_2 and NIP_3 are topological, and the descriptions of Proposition 3.4 still involve infinitely many tests; whether the three levels admit a finitary, combinatorial characterization, like that of NIP in Theorem 3.1, is Question 7.1.

3.3. **The Argyros–Dodos–Kanellopoulos heptachotomy and binary search.** Todorčević's trichotomy describes the closure of a countable set of computations; when the computations are indexed by the binary tree, a finer description is available. The countable dense subsets exhibited in the separable compacta of Section 1.4 are indexed by the binary tree $2^{<\mathbb{N}}$, and this indexing suits our computational emphasis: real numbers are represented by infinite binary sequences, which are the branches of the binary tree, and approximated by finite bitstrings, which are its nodes. A *dyadic subtree* of $2^{<\mathbb{N}}$ is a subset that is isomorphic to $2^{<\mathbb{N}}$ under the order of extension; we write it as $\{s_t : t \in 2^{<\mathbb{N}}\}$, indexed through the canonical isomorphism, which preserves extension and the lexicographic order.

Definition 3.5. Let X be a Polish space.

(1) If I is countable and $\{f_i : i \in I\} \subseteq \mathbb{R}^X$, $\{g_i : i \in I\} \subseteq \mathbb{R}^X$ are two families indexed by I, we say that they are *equivalent* if the map $f_i \mapsto g_i$ extends to a homeomorphism from $\overline{\{f_i : i \in I\}}$ to $\overline{\{g_i : i \in I\}}$.
(2) If $\{f_t : t \in 2^{<\mathbb{N}}\}$ is a pointwise bounded family, we say that it is *minimal* if, for every dyadic subtree $\{s_t : t \in 2^{<\mathbb{N}}\}$ of $2^{<\mathbb{N}}$, the family $\{f_{s_t} : t \in 2^{<\mathbb{N}}\}$ is equivalent to $\{f_t : t \in 2^{<\mathbb{N}}\}$.

Argyros, Dodos and Kanellopoulos proved that, up to equivalence, there are exactly seven minimal families, and that every relatively compact family $\{f_t : t \in 2^{<\mathbb{N}}\} \subseteq \mathrm{B}_1(X)$ contains, along a dyadic subtree, a family equivalent to one of them (see Theorem 3.6 below). We describe the seven minimal families next, in the notation of Argyros, Dodos and Kanellopoulos. A dyadic subtree is *regular* if each of its levels is contained in a level of $2^{<\mathbb{N}}$. Fix a regular dyadic subtree $T = \{s_t : t \in 2^{<\mathbb{N}}\}$ of $2^{<\mathbb{N}}$ such that $s^\frown 0^\infty \neq s'^\frown 0^\infty$ and $s^\frown 1^\infty \neq s'^\frown 1^\infty$ for all distinct $s, s' \in T$. The families are built from the prefix tests v_t and the threshold tests $\chi_{[a}$, $\chi_{(a}$ of Section 1.4; a pair (f, g) of functions on $2^{\mathbb{N}}$ is regarded as a single function on the disjoint union $2^{\mathbb{N}} \sqcup 2^{\mathbb{N}}$.

(1) $D_1 = \{\frac{1}{|t|+1} v_t : t \in 2^{<\mathbb{N}}\}$. Its closure is $D_1 \cup \{\mathbf{0}\}$, the one-point compactification of a countable discrete set.
(2) $D_2 = \{s_t^\frown 0^\infty : t \in 2^{<\mathbb{N}}\}$, a discrete subset of its closure $\overline{D_2} = 2^{\leq\mathbb{N}}$.
(3) $D_3 = \{\chi_{[s_t^\frown 0^\infty} : t \in 2^{<\mathbb{N}}\}$, a discrete subset of its closure $\overline{D_3} = S(2^{\mathbb{N}})$.
(4) $D_4 = \{\chi_{(s_t^\frown 1^\infty} : t \in 2^{<\mathbb{N}}\}$, a discrete subset of its closure $\overline{D_4} = S(2^{\mathbb{N}})$.
(5) $D_5 = \{v_t : t \in 2^{<\mathbb{N}}\}$, a discrete subset of its closure $\overline{D_5} = \hat{A}(2^{\mathbb{N}})$.
(6) $D_6 = \{(v_{s_t}, s_t^\frown 0^\infty) : t \in 2^{<\mathbb{N}}\}$, a discrete subset of its closure $\overline{D_6} = \hat{D}(2^{\mathbb{N}})$.
(7) $D_7 = \{(v_{s_t}, \chi_{[s_t^\frown 0^\infty}) : t \in 2^{<\mathbb{N}}\}$, a discrete subset of its closure $\overline{D_7} = \hat{D}(S(2^{\mathbb{N}}))$.

Theorem 3.6 (Heptachotomy of minimal families [ADK08, Theorem 2])**.** *Let X be a Polish space. For every relatively compact family $\{f_t : t \in 2^{<\mathbb{N}}\} \subseteq \mathrm{B}_1(X)$, there exist $i \in \{1, 2, \ldots, 7\}$ and a regular dyadic subtree $\{s_t : t \in 2^{<\mathbb{N}}\}$ of $2^{<\mathbb{N}}$ such that $\{f_{s_t} : t \in 2^{<\mathbb{N}}\}$ is equivalent to D_i. Moreover, the families D_i are minimal and pairwise non-equivalent.*

For deep computations, the building blocks of the families $D_1, \ldots, D_7$ have a direct reading as primitives of binary search on $2^{\mathbb{N}}$, the space of binary expansions. A node $t \in 2^{<\mathbb{N}}$ determines the dyadic interval $[t^\frown 0^\infty, t^\frown 1^\infty]$, the set of inputs that round to t at precision $|t|$. The prefix test v_t asks whether the input rounds to t; the point $t^\frown 0^\infty$ is the rounded value itself; the comparisons $\chi_{[t^\frown 0^\infty}$ and $\chi_{(t^\frown 1^\infty}$ ask whether the input lies at or to the right of the left endpoint of the interval, or strictly to the right of its right endpoint. Table 1 reads the seven families accordingly.

Let $\{\gamma_t : t \in 2^{<\mathbb{N}}\}$ be a family of computations of a CCS with countably many features, confined by an exhaustive collection R of sizers and satisfying the NIP on shards and features (condition (ii) of Theorem 3.1). By Theorem 3.1, for every feature P and every $r_\bullet \in R$, the functions $P \circ \tilde{\gamma}_t|_{\mathcal{L}[r_\bullet]}$ form a relatively compact subset of $\mathrm{B}_1(\mathcal{L}[r_\bullet])$; by Theorem 3.6, this family contains, along a dyadic subtree, a family equivalent to one of the seven families $D_1, \ldots, D_7$.

Family	Function at node t	Limits
D_1	prefix test weighted by $1/(\|t\|+1)$	the test that never succeeds: confidence vanishes with depth
D_2	return the rounded value $t^\frown 0^\infty$	the exact input: successive approximation
D_3	compare with the left endpoint	comparison with an arbitrary real, ties broken upward
D_4	compare with the right endpoint	comparison with an arbitrary real, ties broken downward
D_5	prefix test	exact equality, or no match
D_6	prefix test and rounded value	a dictionary lookup that returns its key
D_7	prefix test and comparison	a three-way comparison: below, inside or above

TABLE 1. The minimal families of Argyros, Dodos and Kanellopoulos as primitives of binary search.

4. Examples of Compositional Computation Structures

We now turn from the general theory to examples, and compute explicitly the deep computations of three compositional computation structures: Newton's method on the Riemann sphere for two cubic polynomials (Sections 4.1 and 4.2), and Newton's method for square roots on the real projective line (Section 4.3). In Section 5 we place the last of these, together with other classical algorithms, at the levels of the hierarchy of Section 3.

We begin by formalizing the Newton–Raphson root-finding method as a CCS [Gal00], [BFB16, Section 2.3]. Let $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ be the extended complex plane, that is, the Alexandroff (one-point) compactification of $\mathbb{C}$. We identify $\hat{\mathbb{C}}$ with the unit sphere $S^2 \subseteq \mathbb{R}^3$, the Riemann sphere, through the inverse stereographic projection, the homeomorphism $\hat{\mathbb{C}} \to S^2$ given by $z \mapsto (P_1(z), P_2(z), P_3(z))$, where $(P_1(\infty), P_2(\infty), P_3(\infty)) = (0, 0, 1)$ and, for $z \neq \infty$,

$$P_1(z) = \frac{2\,\Re z}{|z|^2+1}, \qquad P_2(z) = \frac{2\,\Im z}{|z|^2+1}, \qquad P_3(z) = \frac{|z|^2-1}{|z|^2+1}.$$

The extended complex plane $\hat{\mathbb{C}}$ thus carries the real predicates P_1, P_2, P_3, with values in $[-1, 1]$, and its topology is metrized by the pullback of the Euclidean metric of S^2.

Newton CCSs: steps and fractals. Let $p = p(z)$ be a nonconstant rational function with complex coefficients, that is, a nonconstant meromorphic map $\hat{\mathbb{C}} \to \hat{\mathbb{C}}$. The *Newton step*, or *Newton map*, of p is the rational map

$$N_p : z \mapsto z - \frac{p(z)}{p'(z)}. \tag{4.1}$$

The iterates of the Newton map, that is, its compositional powers N_p^n, may be regarded as the computations of a CCS with state space $L = \hat{\mathbb{C}}$. The natural predicates are the coordinates P_1, P_2, P_3 of the inverse stereographic projection, since the real and imaginary parts $z \mapsto \Re z$ and $z \mapsto \Im z$ are not defined at $z = \infty$.

Let $\Gamma_p \coloneqq \{N_p^n : n \in \mathbb{N}\}$ be the semigroup of iterates of the Newton map (4.1). The CCS $(\hat{\mathbb{C}}, (P_1, P_2, P_3), \Gamma_p)$ is called *Newton's method for $p(z)$*, or the *Newton CCS for $p(z)$*. Since the state space $\hat{\mathbb{C}}$ is compact and the Newton map N_p is continuous on it, every Newton CCS satisfies the Extendibility Axiom.

Newton's method succeeds at an initial estimate $z_0 \in \hat{\mathbb{C}}$ when the sequence $(z_n)_{n\in\mathbb{N}}$ of Newton steps, where $z_n \coloneqq N_p^n(z_0)$, converges to a root of p.

Fatou and Julia sets. Recall that the *Fatou set* $\mathcal{F}_g$ of a rational map $g\colon \hat{\mathbb{C}} \to \hat{\mathbb{C}}$ consists of the points that have a neighborhood on which every sequence of iterates of g has a subsequence that converges uniformly with respect to the spherical metric; its complement $\mathcal{J}_g \coloneqq \hat{\mathbb{C}} \setminus \mathcal{F}_g$ is the *Julia set* of g, a closed set that is typically a fractal. The Julia set of the Newton map N_p is called the *Newton fractal* of p.

4.1. **Example: The Newton CCS for the cubic $\rho(z) = z^3 - 2z + 2$.** The polynomial $\rho(z) = z^3 - 2z + 2$ has Newton map

$$N_\rho : z \mapsto z - \frac{z^3 - 2z + 2}{3z^2 - 2} = \frac{2z^3 - 2}{3z^2 - 2}.$$

Denote the real root of ρ by r_0, and the roots in the upper and the lower half plane by r_+ and r_-. Starting from $z_0 = 0$, the Newton steps alternate $0, 1, 0, 1, \ldots$, and neither 0 nor 1 is a root of ρ.

Let B_0, B_+, B_- be the basins of the roots r_0, r_+, r_-, and let B_{01} be the basin of the superattracting cycle $\{0, 1\}$. The Fatou set of N_ρ is $B_{01} \cup B_0 \cup B_+ \cup B_-$, and each of these four sets is nonempty and open. Since B_{01} is nonempty, the iterates N_ρ^n do not converge pointwise. Let f_0 and f_1 be the deep computations obtained as limits of the iterates N_ρ^n along nonprincipal ultrafilters on $\mathbb{N}$ that contain the even and the odd numbers, respectively. On the Fatou set the even and the odd iterates converge pointwise, and there the two limits do not depend on the choice of the ultrafilters: both send each basin of a root to that root, while on B_{01} they take only the values 0 and 1 and exchange them, because $f_1 = N_\rho \circ f_0$ on the Fatou set by the continuity of N_ρ. In particular, the two deep computations are distinct.[7] Figure 2 shows f_0 and f_1. On the Newton fractal of ρ, the behavior of the iterates is chaotic.

4.2. **Example: The Newton CCS for the cubic $\kappa(z) = z^3 - 1$.** Figure 3 shows Newton's method for $\kappa(z) = z^3 - 1$, whose roots are the cube roots of unity $r_0 = 1$ and $r_\pm = \exp(\pm 2\pi i/3)$ and whose Newton map is

$$N_\kappa : z \mapsto z - \frac{z^3 - 1}{3z^2} = \frac{2z^3 + 1}{3z^2}.$$

To visualize the iterates themselves, rather than their limits, we color each point z_0 by blending the colors of the three roots with relative weights $|z_n - r|^{-3}$, for $r \in \{r_0, r_+, r_-\}$, where $z_n \coloneqq N_\kappa^n(z_0)$. This coloring is a continuous function of z_0, blurred for small n; as n increases, it sharpens into large regions of nearly constant color, separated by narrow regions where the color varies rapidly, that is, where the coloring is nearly discontinuous. Unlike N_ρ, the map N_κ has no attracting cycle other than its three fixed points, the roots: the Fatou set is the union of the three basins, and on each basin the iterates converge to the corresponding root.

[7] The point ∞ is a repelling fixed point of N_ρ: if $|z|$ is large, then $N_\rho(z) = \frac{2}{3}z + O(1)$. Hence ∞ lies on the Newton fractal of ρ, and $f_0(\infty) = \infty = f_1(\infty)$.




FIGURE 2. The deep computations of Newton's method for $\rho(z) = z^3 - 2z + 2$: the limits of the even iterates N_ρ^{2k} and of the odd iterates N_ρ^{2k+1}, on the whole plane and near the point 1. Points in the basins of the real root and of the roots in the upper and lower half planes (white circles) are purple, orchid and gold, paler where convergence is slower. The remaining open region is the basin of the superattracting cycle $0 \mapsto 1 \mapsto 0$ (white squares). A point there is lilac if the iterate in question is near 0 and plum if it is near 1, so the two colors exchange between even and odd iterates: the two deep computations agree on the basins of the roots and differ on the basin of the cycle. The boundaries of the colored regions form the Newton fractal of ρ.




FIGURE 3. Newton's method for $\kappa(z) = z^3 - 1$ after one and three iterations, and its limit. The coloring blends the colors of the three cube roots of unity (white circles) according to the distance of $N_\kappa^n(z)$ to each of them; the limit is constant on each basin and discontinuous across the Newton fractal; in the right panel, points are paler where convergence is slower.

The chaos of Newton's method on its Julia set is visible to the predicates P_1, P_2, P_3, and it can prevent a Newton CCS on the Riemann sphere from being NIP (see Proposition 5.10).

4.3. **Example: Square roots.** When p has real coefficients, its Newton map N_p maps the real projective line $\mathbb{RP}^1 = \mathbb{R} \cup \{\infty\}$ into itself, and the Newton CCS

for p may be regarded as having state space $\mathbb{RP}^1$. We work out the details in the simplest case, the polynomial $p(x) = x^2 - a$ with real $a > 0$, whose Newton iterates approximate the square roots $\pm\sqrt{a}$.

Let $\mathbb{R} \to S^1$, $x \mapsto (P_1(x), P_2(x))$, be the inverse stereographic projection of the real line onto $S^1 \subseteq \mathbb{R}^2$, given by

$$P_1(x) = \frac{2x}{x^2+1}, \qquad P_2(x) = \frac{x^2-1}{x^2+1},$$

and extended to a homeomorphism of the real projective line $\mathbb{RP}^1 = \mathbb{R} \cup \{\infty\}$ onto S^1 by $\infty \mapsto (P_1(\infty), P_2(\infty)) = (0, 1)$; these predicates are the restrictions to $\mathbb{RP}^1$ of the predicates P_1 and P_3 of the Riemann sphere, and they make $(\mathbb{RP}^1, (P_1, P_2))$ a compact computation states structure. The Newton map N_p is

$$x \mapsto x - \frac{x^2 - a}{2x} = \begin{cases} \frac{x^2+a}{2x}, & x \neq 0, \infty; \\ \infty, & x = 0, \infty. \end{cases}$$

From every initial point $x_0 \neq 0, \infty$, the Newton steps converge to one of the square roots $\pm\sqrt{a}$; more precisely, the iterates of N_p converge pointwise to the map $\mathfrak{f}\colon \mathbb{RP}^1 \to \mathbb{RP}^1$ given by

$$\mathfrak{f}(x) = \lim_{n\to\infty} N_p^n(x) = \begin{cases} \sqrt{a}, & x \in (0, \infty); \\ -\sqrt{a}, & x \in (-\infty, 0); \\ \infty, & x = 0, \infty. \end{cases}$$

The map $\mathfrak{f}$ is therefore discontinuous at 0 and at ∞ (see Figure 1).

5. Computations across the hierarchy

The hierarchy

$$\text{stable} \Rightarrow \text{NIP}_3 \Rightarrow \text{NIP}_2 \Rightarrow \text{NIP}_1 \Rightarrow \text{NIP}$$

of Section 3 was defined through the topology of the space of deep computations. In this section we show that each of its levels is attained by classical algorithms, whence every implication is strict, and that some familiar algorithms lie beyond it. In each example the computations are compositions of the steps of an algorithm. At all levels but one they are the powers of a single step map; at the level NIP but not NIP_1 they form a group generated by two steps, both in the Euclidean algorithm and in free reduction in the free group on two generators. The level of each algorithm is then a theorem about the closure of its computations, and for such computations the hierarchy coincides with a hierarchy studied in topological dynamics by Glasner and Megrelishvili [GM22]. We begin with this coincidence (see Proposition 5.1); Table 2 at the end of the section summarizes the examples.

5.1. **Iterative algorithms and tame dynamics.** The algorithms of this section act on a compact space of states, and each of their steps is continuous. We isolate this situation. A *dynamical CCS* is a compositional computation structure $(L, \mathcal{P}, \Gamma)$ in which L is a compact metrizable space, $\mathcal{P}$ is a countable family of continuous predicates that separates the points of L, and Γ is a countable semigroup of continuous maps $L \to L$ that contains the identity. It is a *cascade CCS* if $\Gamma = \{\gamma^n : n \in \mathbb{N}\}$ is the semigroup of iterates of a single continuous map $\gamma\colon L \to L$. Since $\mathcal{P}$ separates points and L is compact, the map $v \mapsto \text{tp}(v)$ is a homeomorphism of L onto the space of types; hence L is covered by a single shard, every type is realized, and the

Extendibility Axiom holds with $\tilde{\gamma} = \gamma$ for every $\gamma \in \Gamma$. The deep computations of a dynamical CCS form the pointwise closure

$$E(L, \Gamma) = \overline{\Gamma} \subseteq L^L,$$

which is the *enveloping semigroup*, introduced by Ellis, of the dynamical system (L, Γ); for a cascade CCS we write $E(L, \gamma)$. The system (L, Γ) is called *tame* if, for every continuous function $f\colon L \to \mathbb{R}$, the family $\{f \circ \gamma : \gamma \in \Gamma\}$ has the NIP of Definition 1.5 [Köh95, Gla06].

A map $p\colon L \to L$ is *of Baire class* 1 if $f \circ p$ is of Baire class 1 for every continuous $f\colon L \to \mathbb{R}$. It suffices that $P \circ p$ be of Baire class 1 for every $P \in \mathcal{P}$. Indeed, the type map embeds L into $\mathbb{R}^{\mathcal{P}}$, and $\mathcal{P}$ is countable, so every open subset of L is a countable union of finite intersections of sets $P^{-1}(U)$, with $P \in \mathcal{P}$ and $U \subseteq \mathbb{R}$ open; by Fact 1.2, the preimage of every open set under p is then an F_σ set, and so is the preimage of every open set under $f \circ p$, which is therefore of Baire class 1, again by Fact 1.2. In general, this definition is less restrictive than the one adopted in Section 1, according to which a map from L to L is of Baire class 1 when it is the pointwise limit of a sequence of continuous maps from L to L (for example, if $L = [0, 1] \cup \{2\} \subseteq \mathbb{R}$, the map that sends $[0, 1/2]$ to 0 and the rest of L to 2 is of Baire class 1 in the sense just defined, but it is not the pointwise limit of a sequence of continuous maps from L to L, because each such map sends the connected set $[0, 1]$ either into $[0, 1]$ or to the point 2). For the deep computations of a dynamical CCS, however, the two definitions agree: if every deep computation is of Baire class 1 in the sense just defined, then the CCS is NIP, as we show in Proposition 5.1 below, and the last assertion of Theorem 3.1, applied to the single sizer $r_\bullet$ with $L = L[r_\bullet]$, shows that every deep computation is the pointwise limit of a sequence of computations, which are continuous maps from L to L.

Proposition 5.1. *A dynamical CCS is NIP if and only if the dynamical system (L, Γ) is tame, and if and only if every deep computation is of Baire class* 1*. In that case it is NIP$_1$, NIP$_2$ or NIP$_3$ if and only if $E(L, \Gamma)$ is, respectively, first countable, hereditarily separable or metrizable.*

Proof. The single sizer $r_\bullet$ with $L = L[r_\bullet]$ forms an exhaustive collection that confines Γ. If (L, Γ) is tame, the CCS is NIP: take $f = P$ for $P \in \mathcal{P}$. If the CCS is NIP, then, by Theorem 3.1 and Proposition 1.8, $P \circ p$ is of Baire class 1 for every $p \in E(L, \Gamma)$ and $P \in \mathcal{P}$, so every deep computation is of Baire class 1. Finally, suppose that every deep computation is of Baire class 1, and let $f\colon L \to \mathbb{R}$ be continuous. The map $p \mapsto f \circ p$ is continuous on L^L, so the closure of $\{f \circ \gamma : \gamma \in \Gamma\}$ is the compact set $\{f \circ p : p \in E(L, \Gamma)\}$, which lies in $\mathrm{B}_1(L)$. By Theorem 1.6, the family $\{f \circ \gamma : \gamma \in \Gamma\}$ has the NIP, so (L, Γ) is tame. The last statement is Definition 3.2, since $\overline{\tilde{\Gamma}|_L} = E(L, \Gamma)$. □

For metrizable systems, Glasner and Megrelishvili call the tame systems whose enveloping semigroup is first countable, hereditarily separable or metrizable *tame$_1$*, *tame$_2$* and *hereditarily non-sensitive*, respectively. They prove the inclusions between these classes [GM22, Proposition 0.6] and show by examples that the inclusions are proper.

The simplest iterative methods are those whose iterates converge. Their level is decided by the continuity of the limit.

Proposition 5.2. *Let $(L, \mathcal{P}, \Gamma)$ be a cascade CCS whose iterates γ^n converge pointwise to a map $\gamma_\infty \colon L \to L$. Then $E(L,\gamma) = \{\gamma^n : n \in \mathbb{N}\} \cup \{\gamma_\infty\}$, and the CCS is NIP$_3$. It is stable if and only if γ_∞ is continuous.*

Proof. A convergent sequence, together with its limit, is a compact subset of the Hausdorff space L^L, hence closed; so $E(L,\gamma)$ consists of the iterates and γ_∞. For each $P \in \mathcal{P}$, the map $P \circ \gamma_\infty$ is the pointwise limit of the continuous functions $P \circ \gamma^n$, so it is of Baire class 1, and, by Proposition 5.1, the CCS is NIP. The space $E(L,\gamma)$ is a countable compact Hausdorff space, hence metrizable, so the CCS is NIP$_3$. Its members other than γ_∞ are continuous, which gives the statement about stability. □

5.2. **Stable and metrizable.** At the top of the hierarchy, the deep computations form a metrizable space. By Proposition 5.2, every convergent method belongs to this level, and what separates stable from merely NIP$_3$ methods is the continuity of the limit. The iterates of a stable CCS need not converge, however, as the first example shows.

A stable example: rotation. Let $L = \mathbb{T} = \mathbb{R}/\mathbb{Z}$, with the predicates $\cos 2\pi x$ and $\sin 2\pi x$, and let $\gamma(x) = x + \alpha$, where α is irrational. A pointwise limit of rotations is a rotation, and the points $n\alpha$, $n \in \mathbb{N}$, are dense in $\mathbb{T}$; hence the deep computations are all the rotations $x \mapsto x + c$, $c \in \mathbb{T}$. They are continuous, so the rotation CCS is stable, and its space of deep computations is the circle group, a metrizable compactum.

NIP$_3$ but not stable: square roots. The iterates of the Newton map of Section 4.3 converge pointwise to $\mathfrak{f}$, which is discontinuous at 0 and ∞. By Proposition 5.2, the square-root CCS is NIP$_3$ but not stable.

NIP$_3$ but not stable: the power method. The power method computes a dominant eigenvector of a matrix by repeated multiplication and normalization. Let $d \geq 2$ be an integer, let $A = \operatorname{diag}(\lambda_1, \ldots, \lambda_d)$, where $\lambda_1 > \lambda_2 > \cdots > \lambda_d > 0$, let $L = \mathbb{RP}^{d-1}$, and let $\mathcal{P}$ consist of the predicates $p_{ij}([v]) = v_i v_j / \|v\|^2$, for $1 \leq i, j \leq d$, the entries of the orthogonal projection onto the line $[v]$. The step of the algorithm is $\gamma([v]) = [Av]$. For $[v] \in \mathbb{RP}^{d-1}$, let $i(v)$ be the least index i with $v_i \neq 0$, and put $\pi([v]) = [e_{i(v)}]$, where $e_1, \ldots, e_d$ is the standard basis of $\mathbb{R}^d$.

Proposition 5.3. *The deep computations of the power method are $E(L,\gamma) = \{\gamma^n : n \in \mathbb{N}\} \cup \{\pi\}$. Hence the power method is NIP$_3$. Its deep computation π, the only one that is not an iterate, is of Baire class* 1 *and discontinuous, so the power method is not stable.*

Proof. For $i = i(v)$, we have $\lambda_i^{-n} A^n v = v_i e_i + \sum_{j>i} (\lambda_j/\lambda_i)^n v_j e_j \to v_i e_i$, so $\gamma^n \to \pi$ pointwise, and Proposition 5.2 applies. The map π is discontinuous at every $[v]$ with $i(v) > 1$, since $[v + te_1] \to [v]$ as $t \to 0$ while $\pi([v + te_1]) = [e_1] \neq \pi([v])$. □

The deep computation π, which sends each input to the eigenline of the largest eigenvalue whose eigenvector has a nonzero component in the input, is not computable in the sense of Section 2, since it is discontinuous on the space of types L; yet it is the limit of a sequence of computations, and the closure in which it lives is countable and metrizable.

Both levels: the logistic map. The logistic map $f_r(x) = rx(1-x)$, with $0 < r \leq 4$, maps $[0,1]$ into itself. Its CCS has states $L = [0,1]$, the single predicate $x \mapsto x$,

and the semigroup of iterates of f_r. For $0 < r \leq 1$, let ℓ_r be the constant map 0; for $1 < r \leq 3$, let $\ell_r(x) = 1 - 1/r$ for $0 < x < 1$, and $\ell_r(0) = \ell_r(1) = 0$.

Proposition 5.4. *For $0 < r \leq 3$, the iterates of f_r converge pointwise to ℓ_r. Consequently the logistic CCS is stable for $0 < r \leq 1$, and NIP$_3$ but not stable for $1 < r \leq 3$.*

Proof. Write $f = f_r$ and $p = 1 - 1/r$, and note that $f(x)/x = r(1-x)$ for $x > 0$. If $r \leq 1$, then $f(x) < x$ for $0 < x \leq 1$, so every orbit decreases to a fixed point of f, and 0 is the only fixed point in $[0, 1]$.

Suppose that $1 < r \leq 2$. Then $0 < p \leq \frac{1}{2}$, and $f(x) > x$ for $0 < x < p$, while $f(x) < x$ for $p < x \leq 1$. For $0 < x < 1$ we have $f(x) \in (0, \frac{r}{4}] \subseteq (0, \frac{1}{2}]$, and f is increasing on $[0, \frac{1}{2}]$ and fixes p. Hence the orbit of $f(x)$ is increasing and bounded by p if $f(x) < p$, and decreasing and bounded below by p if $f(x) > p$. In both cases, and trivially if $f(x) = p$, it converges to a fixed point of f in $(0, \frac{1}{2}]$, that is, to p.

Suppose that $2 < r \leq 3$, and put $c = \frac{r}{4}$, the maximum of f, and $b = f(c)$. We have $p < c$, since $c - p = (r-2)^2/(4r)$, and therefore $b < f(p) = p$, since f is strictly decreasing on $[\frac{1}{2}, 1]$. We also have $b \geq 1/r$, because $b - 1/r = (r^3(4-r) - 16)/(16r)$ and $r^3(4-r)$ increases from 16 on $[2, 3]$. The concave function f takes the value p at $1/r$ and at p, so $f \geq p$ on $[1/r, p]$. Hence f maps $J_- = [b, p]$ into $J_+ = [p, c]$, and, being decreasing on J_+, it maps J_+ onto J_-. Let $0 < x < 1$. The orbit of $f(x)$ lies in $(0, c]$, and it meets $J_- \cup J_+ = [b, c]$: otherwise all its points would lie in $(0, b)$, where $f(y) > y$ because $b < p$, and the orbit would increase to a fixed point of f in $(0, b]$, which is impossible, since the only positive fixed point of f is p and $p > b$. Since f maps $J_- \cup J_+$ into itself, it remains to be shown that the orbit of every $y \in J_- \cup J_+$ converges to p. The points of period 2 of f that are not fixed are the roots of $r^2y^2 - r(r+1)y + (r+1)$, whose discriminant $r^2(r+1)(r-3)$ is negative for $r < 3$; for $r = 3$ the only root is p. Hence the function $f(f(y)) - y$ vanishes on $[0, 1]$ only at 0 and p, and it is negative on $(p, 1]$, since it is negative at 1. For $y \in J_+$ with $y > p$, the point $f(f(y))$ lies in J_+, so $p \leq f(f(y)) < y$. The sequence $(f^{2n}(y))$ therefore decreases to a fixed point of $f \circ f$ in $[p, y]$, that is, to p, and $f^{2n+1}(y) \to f(p) = p$. Every $y \in J_-$ satisfies $f(y) \in J_+$, so its orbit converges to p as well.

Finally, $f(0) = f(1) = 0$. In all cases the iterates converge pointwise to ℓ_r. The map ℓ_r is continuous for $r \leq 1$, and discontinuous at 0 for $r > 1$, so Proposition 5.2 gives the remaining statements. □

We return to the logistic family at $r = 4$, and to maps of the interval in general, at the end of the section.

5.3. **Hereditarily separable but not metrizable.** Below the metrizable level, no countable set of tests separates the deep computations (see Proposition 3.4). Rounding produces such spaces of deep computations, because an algorithm that rounds at every step makes a fresh decision at every step.

Drawing a line. Fix an irrational number α with $0 < \alpha < 1$. To draw the line $y = \alpha x + \rho$ on a screen, Bresenham's algorithm [Bre65] moves one pixel to the right at each step and decides whether to move one pixel up. The n-th decision is $b_n(\rho) = \lfloor (n+1)\alpha + \rho \rfloor - \lfloor n\alpha + \rho \rfloor = [\![\{n\alpha + \rho\} \in [1-\alpha, 1)]\!]$, where $\{\cdot\}$ denotes the fractional part, and the state of the algorithm is the error term ρ modulo 1.




FIGURE 4. The logistic family $f_r(x) = rx(1 - x)$. Left: the long-run values of the orbit of 0.3 for each r; the CCS is stable for $r \leq 1$, NIP$_3$ but not stable for $1 < r \leq 3$ (see Proposition 5.4), and not NIP at $r = 4$ (see Proposition 5.11); the parameters $3 < r < 4$ are not classified here. Right: for $r = 2.8$, two orbits converge to the fixed point $1 - 1/r$; the deep computation sends 0 and 1 to 0 and every other state to $1 - 1/r$.

The step $\rho \mapsto \rho + \alpha$ is a rotation of the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$, but the decision bit is not a continuous function of ρ. The standard remedy [MH40, Lot02] is to take the decision sequences themselves as states.

Let $D = \{-k\alpha : k \in \mathbb{Z}\} \subseteq \mathbb{T}$, a countable dense set containing both endpoints 0 and $1 - \alpha$ of the decision interval. The *split circle* $\mathbb{T}_D$ is obtained from $\mathbb{T}$ by replacing each $t \in D$ by two points $t^- < t^+$, with the circular order topology; it is a compact metrizable space, homeomorphic to the Cantor set. The rotation by α preserves D and lifts to a homeomorphism R of $\mathbb{T}_D$ that preserves the labels $\pm$. The decision interval becomes clopen in $\mathbb{T}_D$, and the coding map $\mathbb{T}_D \to 2^{\mathbb{Z}}$, $y \mapsto (\llbracket R^n y \in [1 - \alpha, 1) \rrbracket)_{n \in \mathbb{Z}}$, is a homeomorphism onto the *Sturmian shift* X_α that conjugates R to the shift σ. The *Bresenham CCS* has states $L = X_\alpha$, the coordinate predicates, and the semigroup $\Gamma = \{\sigma^n : n \in \mathbb{N}\}$.

For $c \in \mathbb{T}$, let R_c^+ (respectively R_c^-) be the map of $\mathbb{T}_D$ that sends a point over y_0 to $(y_0 + c)^+$ (respectively $(y_0 + c)^-$) when $y_0 + c \in D$, and to $y_0 + c$ otherwise; through the coding map, we regard R_c^+ and R_c^- as maps of X_α. The *double arrow circle* $\mathbb{D}$ is $\mathbb{T} \times \{-, +\}$ with the lexicographic circular order and its order topology; it is compact, first countable, hereditarily separable and not metrizable [Eng89, 3.10.C].

Theorem 5.5. *The Bresenham CCS is NIP. Its deep computations are*

$$E(X_\alpha, \sigma) = \{\sigma^n : n \in \mathbb{N}\} \cup \{R_c^+, R_c^- : c \in \mathbb{T}\},$$

where each σ^n is isolated and $c^\pm \mapsto R_c^\pm$ is a homeomorphism of $\mathbb{D}$ onto the remaining part. In particular, the Bresenham CCS is NIP$_2$ but not NIP$_3$.

Proof. For NIP, note that the predicates $P_j \circ \sigma^n$ are $\{0, 1\}$-valued, and that the j-th coordinate of $\sigma^n y$ is 1 exactly when $R^{n+j} y$ lies in the clopen arc $[1 - \alpha, 1)$ of $\mathbb{T}_D$, that is, when y lies in an arc of $\mathbb{T}_D$. A family of arcs of a circularly ordered

set has VC dimension at most 3, so, by the remark at the end of Section 1.5, the characteristic functions of its members have the NIP.

For the deep computations, let γ be the limit of σ^{n_i} along an ultrafilter, and let $c = \lim n_i\alpha$ in $\mathbb{T}$. If n_i is eventually constant, then γ is a power of σ. Otherwise, since the points $n\alpha$ are distinct, the ultrafilter decides whether $n_i\alpha$ approaches c from the right or from the left. In the first case, for every point y over y_0 the points $R^{n_i}y$ over $y_0 + n_i\alpha$ approach $y_0 + c$ strictly from the right, and they converge in $\mathbb{T}_D$ to $(y_0 + c)^+$ if $y_0 + c \in D$ and to $y_0 + c$ otherwise; so $\gamma = R_c^+$. The second case gives R_c^-. Conversely, every c is approached from both sides by the points $n\alpha$ with $n \geq N$, for every N, so every $R_c^\pm$ is a deep computation, and $R_c^+ \neq R_c^-$, since they differ at the points over $d - c$ with $d \in D$. The same argument shows that $c_i^{\varepsilon_i} \to c^+$ in $\mathbb{D}$ implies $R_{c_i}^{\varepsilon_i} \to R_c^+$ pointwise, and symmetrically for c^-; so $c^\pm \mapsto R_c^\pm$ is a continuous bijection from the compact space $\mathbb{D}$ onto a Hausdorff space, hence a homeomorphism. Finally, fix $d \in D$ and consider the two points $(d - n\alpha)^\pm$ of $\mathbb{T}_D$, which σ^n sends to d^- and d^+. Every map $R_c^\pm$ sends them to the same point, and so does every pointwise limit of σ^{m_i} with $m_i \to \infty$, by the argument above. Hence no net of such maps, and no net of maps σ^m with $m \neq n$ and m bounded, converges to σ^n, so σ^n is isolated. Thus $E(X_\alpha, \sigma)$ is the union of a countable set and a copy of $\mathbb{D}$; it is hereditarily separable and not metrizable. □

The two deep computations R_c^+ and R_c^- are the two ways of drawing the line infinitely far away, where the accumulated error has reached c: they agree except at the pixels where the line passes exactly through a corner, and there they apply the two opposite rounding conventions. Figure 5 shows the two conventions.




FIGURE 5. Left: the pixels drawn by Bresenham's algorithm for $\alpha = (\sqrt{5} - 1)/2$, with the decision bits b_n. Right: where the line passes exactly through a pixel corner, the deep computations R_c^+ and R_c^- break the tie in opposite directions; they agree on every other pixel.

The space $E(X_\alpha, \sigma)$ is the closure of a countable set, so it is a separable Rosenthal compactum; it is hereditarily separable and not metrizable, so, by Todorčević's trichotomy (see Theorem 3.3), it contains a copy of the split Cantor set $S(2^{\mathbb{N}})$. For the action of $\mathbb{Z}$ generated by the shift, Glasner and Megrelishvili proved that Sturmian systems are tame$_2$ but not hereditarily non-sensitive [GM22, Corollary 8.3].

5.4. **First countable but not hereditarily separable.** Running two rounding algorithms side by side moves the computation one level down.

Drawing a line in space. Let α and β be irrational numbers in $(0,1)$ such that $1, \alpha, \beta$ are linearly independent over $\mathbb{Q}$. To draw a line in $\mathbb{Z}^3$ with direction $(1, \alpha, \beta)$, the three-dimensional Bresenham algorithm runs the planar algorithm in two coordinates at once. Its CCS has states $L = X_\alpha \times X_\beta$, the coordinate predicates of both factors, and the semigroup generated by $\sigma \times \sigma$.

Theorem 5.6. *The three-dimensional Bresenham CCS is NIP_1 but not NIP_2.*

Proof. Each predicate is a coordinate of one factor, so the CCS is NIP by Theorem 5.5. The deep computations form a closed subspace of $E(X_\alpha, \sigma) \times E(X_\beta, \sigma)$, which is first countable as a product of two first countable spaces; hence the CCS is NIP_1. By Kronecker's theorem, the points $(n\alpha, n\beta)$ with $n \geq N$ are dense in $\mathbb{T}^2$ for every N, so every point (c, c') is approached by them from inside each of the four open quadrants at (c, c'); as in the proof of Theorem 5.5, it follows that every pair $(R_c^\varepsilon, R_{c'}^{\varepsilon'})$, with $\varepsilon, \varepsilon' \in \{+, -\}$, is a deep computation. Consider the pairs (R_c^+, R_{-c}^+), for $c \in \mathbb{T}$. For $0 < \delta < \frac{1}{2}$, the set $[c^+, (c+\delta)^-) \times [(-c)^+, (-c+\delta)^-)$ is a neighborhood of $(c^+, (-c)^+)$ in $\mathbb{D} \times \mathbb{D}$, and it contains $(s^+, (-s)^+)$ only if $c \leq s < c + \delta$ and $-c \leq -s < -c + \delta$, that is, only if $s = c$. These pairs therefore form an uncountable discrete subspace, and the space of deep computations is not hereditarily separable. □

The uncountable discrete set consists of the deep computations that break ties upward in both coordinates at points where the two accumulated errors are opposite. Each of the two planar algorithms is NIP_2, and drawing in two directions at once leaves the class, just as the Sorgenfrey plane shows that a product of two hereditarily separable spaces need not be hereditarily separable. Glasner and Megrelishvili show similarly that a product of two tame_2 systems need not be tame_2 [GM22, Proposition 4.1]. By Todorčević's trichotomy, the space of deep computations contains a copy of $D(2^{\mathbb{N}})$ or of $\hat{D}(S(2^{\mathbb{N}}))$.

5.5. **NIP but not first countable: the Euclidean algorithm.** At the last tame level, some deep computation is not determined by countably many tests of its own (see Proposition 3.4). An exact equality test has this property: a countable set of inputs cannot tell it apart from the test that always fails. The Euclidean algorithm produces such tests.

Let $L = \mathbb{RP}^1 = \mathbb{R} \cup \{\infty\}$, with the predicates P_1, P_2 of Section 4.3, and let $\Gamma = \mathrm{PGL}(2, \mathbb{Z})$ be the group of Möbius maps $x \mapsto (ax + b)/(cx + d)$ with integer coefficients and $ad - bc = \pm 1$. It is generated by the steps $x \mapsto x + 1$ and $x \mapsto 1/x$ of the Euclidean algorithm, and the maps that the continued fraction algorithm composes are elements of Γ. We call $(\mathbb{RP}^1, (P_1, P_2), \Gamma)$ the *Euclidean CCS*. It is a dynamical CCS whose semigroup is a group, and its deep computations form $E(\mathbb{RP}^1, \Gamma)$.

For an irrational number $\eta = [a_0; a_1, a_2, \dots]$, let p_k/q_k be its convergents, put $p_{-1} = 1$ and $q_{-1} = 0$, and let $\eta_{k+1} = [a_{k+1}; a_{k+2}, \dots]$ be its complete quotients. Let M_k be the Möbius map $t \mapsto (p_k t + p_{k-1})/(q_k t + q_{k-1})$, which belongs to Γ because $p_k q_{k-1} - p_{k-1} q_k = (-1)^{k+1}$. Writing $A_j(t) = a_j + 1/t$, we have $M_k = A_0 \circ A_1 \circ \cdots \circ A_k$, so $M_k^{-1} = A_k^{-1} \circ \cdots \circ A_0^{-1}$ with $A_j^{-1}(t) = 1/(t - a_j)$. Thus M_k^{-1} is the map that the first $k+1$ steps of the continued fraction algorithm for η

apply, and the classical identity $\eta = M_k(\eta_{k+1})$ says that it sends η to its complete quotient η_{k+1}.

Lemma 5.7. *Let η be an irrational number, and let $x \in \mathbb{RP}^1$ be distinct from η. Then $M_k^{-1}(x) + q_{k-1}/q_k \to 0$ as $k \to \infty$, while $M_k^{-1}(\eta) = \eta_{k+1} > 1$.*

Proof. Solving $M_k(t) = x$ gives $M_k^{-1}(\infty) = -q_{k-1}/q_k$ and, for real x,

$$M_k^{-1}(x) = \frac{q_{k-1}x - p_{k-1}}{p_k - q_k x} = -\frac{q_{k-1}}{q_k} \cdot \frac{x - p_{k-1}/q_{k-1}}{x - p_k/q_k}$$

for $k \geq 1$ (for such k we have $q_{k-1} \geq 1$), where the value is ∞ if $x = p_k/q_k$. The convergents of an irrational number are pairwise distinct, so x equals p_k/q_k for at most one k; for all larger k the formula gives a real number. Both convergents tend to $\eta \neq x$, so the last fraction tends to 1, and $0 < q_{k-1}/q_k \leq 1$ for $k \geq 1$; hence $M_k^{-1}(x) + q_{k-1}/q_k \to 0$. The identity $M_k^{-1}(\eta) = \eta_{k+1}$ is the classical one, and $\eta_{k+1} > a_{k+1} \geq 1$ because η_{k+1} is irrational. □

Let $\varphi = (1 + \sqrt{5})/2$. Let S be the set of irrational numbers $\eta = [0; b_1, b_2, \dots]$ whose sequence of partial quotients is a concatenation $1^1 d_1 1^2 d_2 1^3 d_3 \cdots$, where 1^n denotes n consecutive ones and each $d_n \in \{1, 2\}$. The block 1^n occupies the indices $s_n, \dots, s_n + n - 1$ with $s_n = n(n+1)/2$. Distinct choices of (d_n) give distinct numbers, by the uniqueness of continued fraction expansions, so S has the cardinality of the continuum.

Proposition 5.8. *(i) For every $\eta \in S$, the map δ_η that sends η to φ and every other point of $\mathbb{RP}^1$ to $-1/\varphi$ is a deep computation of the Euclidean CCS.*

(ii) The constant map c with value $-1/\varphi$ is a deep computation, and it has no countable base of neighborhoods in $E(\mathbb{RP}^1, \Gamma)$.

Consequently $E(\mathbb{RP}^1, \Gamma)$ is not first countable.

Proof. (i) Let $\eta \in S$, and for each n let $k_n = s_n + \lfloor n/2 \rfloor$ be the middle index of the block 1^n, and $m_n = \lceil n/2 \rceil - 1$. Then $a_{k_n} = a_{k_n - 1} = \cdots = a_{k_n - m_n} = 1$ and $a_{k_n+1} = \cdots = a_{k_n+m_n} = 1$, and $m_n \to \infty$. We have $q_{k-1}/q_k = [0; a_k, a_{k-1}, \dots, a_1]$ and $\eta_{k+1} = [a_{k+1}; a_{k+2}, \dots]$. Suppose that two continued fractions, finite or infinite, $[c_0; c_1, \dots]$ and $[c_0'; c_1', \dots]$ satisfy $c_j = c_j'$ for $j < m$, with $c_1 = \cdots = c_{m-1} = 1$. Then they share the convergents of index at most $m - 1$, and both lie between the last two of them (or equal one of them). It follows that they differ by at most $1/(q_{m-2}q_{m-1}) \leq 1/(F_{m-1}F_m)$, where F_j are the Fibonacci numbers, whatever the remaining partial quotients are. The hypothesis holds with $m = m_n$ for the pair $[0; a_{k_n}, \dots, a_1]$, $[0; 1, 1, \dots]$ and for the pair $[a_{k_n+1}; a_{k_n+2}, \dots]$, $[1; 1, 1, \dots]$. Hence $q_{k_n-1}/q_{k_n} \to [0; 1, 1, \dots] = 1/\varphi$ and $\eta_{k_n+1} \to [1; 1, 1, \dots] = \varphi$. By Lemma 5.7, the maps $M_{k_n}^{-1} \in \Gamma$ converge pointwise to δ_η, at $x = \infty$ as well.

(ii) Let (η_j) be a sequence of distinct elements of S. For each $x \in \mathbb{RP}^1$, at most one j has $\eta_j = x$, so $\delta_{\eta_j}(x) = -1/\varphi$ for all but at most one j. Hence $\delta_{\eta_j} \to c$ pointwise, and $c \in E(\mathbb{RP}^1, \Gamma)$ because this set is closed. Now let $(U_i)_{i \in \mathbb{N}}$ be open neighborhoods of c in $E(\mathbb{RP}^1, \Gamma)$. Each U_i contains a basic neighborhood $\{f : d(f(x), c(x)) < \varepsilon_i \text{ for } x \in F_i\}$ with $F_i \subseteq \mathbb{RP}^1$ finite, $\varepsilon_i > 0$ and d a metric on $\mathbb{RP}^1$. The set $F = \bigcup_i F_i$ is countable, so some $\eta \in S$ lies outside F, and then δ_η agrees with c on F. Hence $\delta_\eta \in \bigcap_i U_i$, while $\delta_\eta \neq c$. Thus no countable family of neighborhoods of c has intersection $\{c\}$. In a Hausdorff space, the intersection of a base of neighborhoods of c is $\{c\}$, so c has no countable base of neighborhoods. □




FIGURE 6. The Euclidean algorithm, drawn on the projective line (each axis runs from ∞ through 0 back to ∞). Left: the maps M_k^{-1} that the continued fraction algorithm applies to a number η whose partial quotients contain longer and longer runs of ones; as k grows along the middles of the runs, every point other than η is sent near $-1/\varphi$, while η is sent near φ. Right: the limits, exact equality tests δ_η, at several points. Every neighborhood of the constant map $-1/\varphi$ contains all but finitely many of the uncountably many tests δ_η, so the space of deep computations is not first countable (see Proposition 5.8).

Corollary 5.9. *The Euclidean CCS is NIP but not NIP$_1$. Its space of deep computations contains the copy $\{\delta_\eta : \eta \in S\} \cup \{c\}$ of the Alexandroff compactification $A(2^{\mathbb{N}})$.*

Proof. Ellis proved that the action of $\mathrm{GL}(2,\mathbb{R})$ on the projective line is tame [Ell93]; see [GM22, Example 5.7]. For every continuous $f\colon \mathbb{RP}^1 \to \mathbb{R}$, the family $\{f \circ \gamma : \gamma \in \Gamma\}$ is contained in the corresponding family for $\mathrm{GL}(2,\mathbb{R})$, so it has the NIP. Hence the action of Γ is tame, and, by Proposition 5.1, the Euclidean CCS is NIP. By Proposition 5.8, it is not NIP$_1$. Each δ_η is isolated in $\{\delta_\eta : \eta \in S\} \cup \{c\}$, since it is the only member that sends η near φ, and every neighborhood of c contains all but finitely many of the δ_η, as in the proof of Proposition 5.8. The set $\{\delta_\eta : \eta \in S\} \cup \{c\}$ is therefore the one-point compactification of a discrete set of cardinality $2^{\aleph_0}$, and hence it is homeomorphic to $A(2^{\mathbb{N}})$ (see Section 1.4). □

The deep computation δ_η is an exact equality test: it takes one value if the input is exactly η and another value otherwise. It is the pointwise limit of maps that the continued fraction algorithm applies to η, when these maps are applied to all inputs. The mechanism is that of convergence groups. By Lemma 5.7, the maps M_k^{-1} draw the images of all points other than η together, near $-q_{k-1}/q_k \in [-1, 0)$, while they send η into $(1, \infty)$; this exhibits every irrational η as a conical limit point of Γ. The choice of S makes the two limits the same for uncountably many η. Glasner and Megrelishvili use the same device to prove that an infinite countable uniform convergence group acting on a compact metric space with infinite limit set is tame but not tame$_1$ [GM22, Proposition 6.1 and Theorem 6.5]. Their theorem does not cover Γ, which contains parabolic elements and is not a uniform convergence group on $\mathbb{RP}^1$. Ellis's example, the projective action of the whole group $\mathrm{GL}(2,\mathbb{R})$, is

also tame but not tame_1 [GM22, Example 5.7]; the Euclidean CCS shows that the countable group generated by the two steps of the algorithm already reaches this level.

A single step cannot produce δ_η. For $g \in \Gamma$, the closure of $\{g^n : n \in \mathbb{Z}\}$ in $(\mathbb{RP}^1)^{\mathbb{RP}^1}$ is countable, hence metrizable, and it does not contain δ_η. Indeed, if g has finite order, the set of its powers is finite. If g is parabolic, with fixed point r, then g^n converges pointwise to the constant map r as $n \to \pm\infty$. If g is hyperbolic, with attracting and repelling fixed points a and r, then g^n converges to the map that fixes r and sends every other point to a as $n \to +\infty$, and to the map that fixes a and sends every other point to r as $n \to -\infty$. If g reverses orientation and has infinite order, then its trace t is nonzero, so g^2 has trace $t^2+2 > 2$ and is hyperbolic, with the same fixed points as g; the odd powers $g^{2m+1} = g \circ g^{2m}$ have the same two limits as the even ones. In every case the closure consists of the powers of g, which are homeomorphisms, and at most two limit maps, each of which is constant or fixes the unique point at which it differs from a constant. The map δ_η is none of these, since it is not injective and $\delta_\eta(\eta) = \varphi \neq \eta$.

Free reduction. A second example at this level is the free group F_2 on two generators a, b, acting on its boundary ∂F_2, the space of infinite reduced words in a, a^{-1}, b, b^{-1}, which is a Cantor set. A generator acts on an infinite word by prepending a letter and cancelling it against the first letter if they are inverse, as a stack does. With the predicates $w \mapsto [\![w_n = x]\!]$, for $n \in \mathbb{N}$ and each letter x, this is a dynamical CCS, and Glasner and Megrelishvili proved that the system is tame but not tame_1 [GM22, Corollary 6.6 and Example 6.7]. By Proposition 5.1, this CCS is NIP but not NIP_1.

An open question. Both examples at this level are actions of groups with two generators. Whether a single map can reach this level, that is, whether some cascade CCS is NIP but not NIP_1, is Question 7.2. For actions of $\mathbb{Z}$, and more generally of abelian groups, the existence of a tame metrizable system whose enveloping semigroup is not first countable is Question 6.8 of Glasner and Megrelishvili [GM22]; to our knowledge, it is open.

5.6. **Beyond NIP.** Beyond the hierarchy lie the computations with the independence property: for some feature, the computations realize every finite pattern of high and low values on suitable inputs. Chaotic iteration produces such patterns.

Newton's method on the Riemann sphere. On its Julia set the Newton map is chaotic, and the chaos is visible to the predicates of the CCS. We exhibit the independence in the simplest case, the square roots of 1.

Proposition 5.10. *Let $p(z) = z^2 - 1$, and let $N(z) = \frac{1}{2}\big(z + \frac{1}{z}\big)$ be its Newton map on $\hat{\mathbb{C}}$. The functions $P_2 \circ N^{3k}$, $k \in \mathbb{N}$, form an independent sequence: for all finite disjoint sets $E, F \subseteq \mathbb{N}$ there is $z \in \hat{\mathbb{C}}$ with $P_2(N^{3k}z) \geq \frac{1}{2}$ for $k \in E$ and $P_2(N^{3k}z) \leq -\frac{1}{2}$ for $k \in F$. In particular, the Newton CCS of p is not NIP.*

Proof. The Möbius transformation $\psi(z) = (z-1)/(z+1)$ conjugates N to the squaring map: $\psi(N(z)) = \psi(z)^2$. It maps the imaginary axis, together with ∞, onto the unit circle, with $\psi^{-1}(e^{i\theta}) = i\cot(\theta/2)$. Hence, for $z = i\cot(\theta/2)$, we have $N^n(z) = i\cot(2^n\theta/2)$ and

$$P_2\big(N^n(z)\big) = \frac{2\cot(2^{n-1}\theta)}{\cot^2(2^{n-1}\theta) + 1} = \sin(2^n\theta).$$

Write $\theta = 2\pi \sum_{j\geq 1} x_j 2^{-j}$ with bits x_j. The fractional part of $2^{3k}\theta/2\pi$ has binary expansion $0.x_{3k+1}x_{3k+2}x_{3k+3}\cdots$. If $(x_{3k+1}, x_{3k+2}, x_{3k+3}) = (0,1,0)$, this fractional part lies in $[\frac{1}{4}, \frac{3}{8}]$, so $\sin(2^{3k}\theta) \geq \sin\frac{3\pi}{4} > \frac{1}{2}$; if the block is $(1,1,0)$, it lies in $[\frac{3}{4}, \frac{7}{8}]$, so $\sin(2^{3k}\theta) < -\frac{1}{2}$. Choosing the block $(0,1,0)$ for $k \in E$ and $(1,1,0)$ for $k \in F$ gives the required z. □

The proof also shows that the doubling map $x \mapsto 2x$ of the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$, with the predicates $\cos 2\pi x$ and $\sin 2\pi x$, is not NIP: with $x = \theta/2\pi$, the functions $\sin(2\pi\, 2^{3k}x)$ form an independent sequence. The doubling map shifts the binary digits of x, and the independence comes from the freedom to choose those digits.

The same algorithm is tame on the real line. The Julia set of N is the imaginary axis together with ∞, which meets $\mathbb{RP}^1$ only at ∞, and on $\mathbb{RP}^1$ the iterates converge to the deep computation of Section 4.3. Whether Newton's method is tame depends on the space of states, not on the formula for the step.

The logistic map at $r = 4$. The doubling map also governs the logistic map $f_4(x) = 4x(1-x)$ of Section 5.2.

Proposition 5.11. *The functions f_4^{3k}, $k \in \mathbb{N}$, form an independent sequence on $[0,1]$: for all finite disjoint sets $E, F \subseteq \mathbb{N}$ there is $x \in [0,1]$ with $f_4^{3k}(x) \geq \frac{3}{4}$ for $k \in E$ and $f_4^{3k}(x) \leq \frac{1}{4}$ for $k \in F$. In particular, the logistic CCS for $r = 4$ is not NIP.*

Proof. Let $h(\theta) = \sin^2(\pi\theta)$ for $\theta \in \mathbb{T}$. Then $f_4(h(\theta)) = 4\sin^2(\pi\theta)\cos^2(\pi\theta) = h(2\theta)$, so $f_4^n(h(\theta)) = \sin^2(\pi\, 2^n\theta)$. Write $\theta = \sum_{j\geq 1} x_j 2^{-j}$ in binary. The fractional part u of $2^{3k}\theta$ is then $0.x_{3k+1}x_{3k+2}\cdots$ in binary. If $(x_{3k+1}, x_{3k+2}, x_{3k+3}) = (0,1,1)$, then $u \in [\frac{3}{8}, \frac{1}{2}]$ and $\sin^2(\pi u) \geq \sin^2\frac{3\pi}{8} > \frac{3}{4}$; if the block is $(0,0,0)$, then $u \in [0, \frac{1}{8}]$ and $\sin^2(\pi u) \leq \sin^2\frac{\pi}{8} < \frac{1}{4}$. Choosing the block $(0,1,1)$ for $k \in E$ and $(0,0,0)$ for $k \in F$, and putting $x = h(\theta)$, gives the required point. □




FIGURE 7. Independence for the doubling map. The values of the predicate $\sin^2(\pi x)$ at the times $0, 3, 6, 9$ follow any prescribed pattern of high (H, at least 3/4) and low (L, at most 1/4): the initial state is chosen with the block 011 of binary digits for H and 000 for L.

Thus the logistic family, which is stable for $0 < r \leq 1$ and NIP_3 but not stable for $1 < r \leq 3$ (see Proposition 5.4), leaves the hierarchy at $r = 4$. A continuous map of a compact interval is tame if and only if it is not chaotic in the sense of Li and Yorke [Li11, Theorems 4.23 and 4.24], and Glasner and Megrelishvili record,

from an unpublished paper, that it is tame if and only if it is hereditarily non-sensitive [GM23, Remark 5.4]. Consequently, for cascade CCSs on $[0,1]$ with the predicate $x \mapsto x$, the levels NIP, NIP_1, NIP_2 and NIP_3 coincide.

The Thue–Morse sequence. Even a sequence produced by a simple substitution rule can be wild. The Thue–Morse sequence t_n is the parity of the number of ones in the binary expansion of n; it is the limit of the words obtained from 0 by the substitution $0 \mapsto 01$, $1 \mapsto 10$. The two-sided sequences all of whose finite blocks occur in (t_n) form the Thue–Morse subshift $X_{TM} \subseteq 2^{\mathbb{Z}}$, on which the shift σ is a homeomorphism. Glasner and Megrelishvili note that the action of $\mathbb{Z}$ that it generates is not tame [GM23, Example 5.3]. Hence the CCS with states X_{TM}, the coordinate predicates, and the group $\{\sigma^n : n \in \mathbb{Z}\}$ is not NIP, by Proposition 5.1. The cascade CCS of σ is not NIP either. Indeed, the words obtained from 0 by an even number of applications of the substitution are palindromes, and every finite block of (t_n) occurs in one of them, so the reversal ρ of two-sided sequences maps X_{TM} onto itself and satisfies $\rho \circ \sigma = \sigma^{-1} \circ \rho$. An independent sequence $f \circ \sigma^{n_k}$, with f continuous and the n_k distinct, has infinitely many indices n_k of the same sign. If they are positive, these terms already form an independent sequence in the cascade; if they are negative, the functions $f \circ \sigma^{n_k} \circ \rho = (f \circ \rho) \circ \sigma^{-n_k}$ form an independent sequence with positive exponents, because ρ is a bijection of X_{TM}. By Proposition 5.1, the cascade CCS is not NIP.

5.7. **Summary.** Table 2 collects the examples of this section. The Euclidean algorithm and free reduction are actions of groups with two generators; every other entry is a single map iterated.

6. Randomized versions of NIP and Monte Carlo computability of deep computations

In Sections 3–5 we studied deep computations as *topological* limits of computations; under the NIP, they are of the first Baire class on the type shards, and on each shard each of them is the limit of a sequence of computations. That theory rests on the type shards being Polish, and hence on the countability of the set $\mathcal{P}$ of predicates, which is used in the proof of Theorem 3.1. In this section we drop that assumption and replace topology by measure: instead of asking that a deep computation be of the first Baire class, we ask that it be *universally measurable*, that is, measurable with respect to every Radon probability measure on the type shards, which allows it to be evaluated almost surely by random sampling. We call such deep computations *universally Monte Carlo computable* (see Definition 6.6). The role of the Bourgain–Fremlin–Talagrand dichotomy is then taken by its measure-theoretic form (see Theorem 6.5), and the role of NIP is shared with Talagrand stability.

The main results of the section are Theorem 6.7, which derives Monte Carlo computability from the NIP, and Corollary 6.14, which derives it from universal Talagrand stability.

6.1. **NIP and Monte Carlo computability of deep computations.** By Fact 1.2, a function $f\colon X \to \mathbb{R}$ on a perfectly normal space X is of Baire class 1 if and only if $f^{-1}[U]$ is an F_σ subset of X for every open $U \subseteq \mathbb{R}$. The following definition replaces this condition by its measure-theoretic counterpart.

Level	Computation	Deep computations; source
stable	rotation of the circle	the circle group
	logistic map, $0 < r \leq 1$	iterates and one continuous limit; Prop. 5.4
NIP_3, not stable	power method	countable; Prop. 5.3
	Newton's method for $x^2 - a$ on $\mathbb{RP}^1$	countable; Prop. 5.2
	logistic map, $1 < r \leq 3$	countable; Prop. 5.4
NIP_2, not NIP_3	Bresenham, plane (Sturmian shift)	isolated iterates and a double arrow circle; Thm. 5.5
NIP_1, not NIP_2	Bresenham, space (two Sturmian shifts)	contains an uncountable discrete set; Thm. 5.6
NIP, not NIP_1	Euclidean algorithm: $\mathrm{PGL}(2,\mathbb{Z})$ on $\mathbb{RP}^1$ (group)	contains $A(2^{\mathbb{N}})$, of exact equality tests; Cor. 5.9
	free reduction: F_2 on ∂F_2 (group)	not first countable; [GM22]
	a single map	open; Question 7.2
not NIP	doubling map of the circle	Prop. 5.10
	Newton's method for $z^2 - 1$ on $\hat{\mathbb{C}}$	Prop. 5.10
	logistic map, $r = 4$	Prop. 5.11
	Thue–Morse shift	[GM23]

TABLE 2. Computations at each level of the hierarchy.

Definition 6.1. Given a Hausdorff space X and a measurable space (Y, Σ), we say that $f : X \to Y$ is *universally measurable* (with respect to Σ) if $f^{-1}(E)$ is μ-measurable for every Radon measure μ on X and every $E \in \Sigma$. When $Y = \mathbb{R}$, we will always take $\Sigma = \mathcal{B}(\mathbb{R})$, the Borel σ-algebra of $\mathbb{R}$.

If X is a compact Hausdorff space, then Radon measures on X are finite, and every nonzero Radon measure on X becomes a probability measure after normalization; hence the following remark holds.

Remark 6.2. If X is compact, then a function $f : X \to \mathbb{R}$ is universally measurable if and only if $f^{-1}(U)$ is μ-measurable for every Radon probability measure μ on X and every open set $U \subseteq \mathbb{R}$.

Intuitively, a function is universally measurable when it is measurable for every reasonable way of measuring sets in its domain. For universally measurable sets and functions, see [Pap02, Chapters 1 and 2].

Notation 6.3. Following Bourgain, Fremlin and Talagrand [BFT78], we denote by $M_r(X)$ the collection of all universally measurable real-valued functions on X; for a Radon measure μ on X, we denote by $\mathscr{M}^0(X, \mu)$ the collection of all μ-measurable real-valued functions on X.

In the study of deep computations we are interested in maps from a state space $L \subseteq \mathbb{R}^{\mathcal{P}}$ into $\mathbb{R}^{\mathcal{P}}$, such as transitions and deep computations themselves. The product space $\mathbb{R}^{\mathcal{P}}$ carries two natural σ-algebras: the Borel σ-algebra, generated by the open subsets of $\mathbb{R}^{\mathcal{P}}$, and the cylinder σ-algebra, generated by the subbasic open sets $\pi_P^{-1}(U)$ with $P \in \mathcal{P}$ and $U \subseteq \mathbb{R}$ open. In general, the cylinder σ-algebra is strictly smaller, but the two coincide when $\mathcal{P}$ is countable. We use the cylinder σ-algebra to define universally measurable maps $f\colon \mathbb{R}^{\mathcal{P}} \to \mathbb{R}^{\mathcal{P}}$, because of the following characterization.

Proposition 6.4. *Let X be a Hausdorff space and $Y = \prod_{i\in I} Y_i$ be any product of measurable spaces (Y_i, Σ_i) for $i \in I$. Let Σ_Y be the cylinder σ-algebra generated by the measurable spaces (Y_i, Σ_i). The following are equivalent for $f : X \to Y$:*

(i) $f : X \to Y$ is universally measurable (with respect to Σ_Y).
(ii) $\pi_i \circ f : X \to Y_i$ is universally measurable (with respect to Σ_i) for all $i \in I$.

Proof. The implication (i)⇒(ii) holds because each projection π_i is measurable and $(\pi_i \circ f)^{-1}(E) = f^{-1}(\pi_i^{-1}(E))$ for every $E \in \Sigma_i$. To prove (ii)⇒(i), fix a Radon measure μ on X, and note that the sets $C \subseteq Y$ for which $f^{-1}(C)$ is μ-measurable form a σ-algebra. This σ-algebra contains every measurable cylinder $C = \prod_{i\in I} C_i$: if J is the finite set of indices $i \in I$ with $C_i \neq Y_i$, then $f^{-1}(C) = \bigcap_{i\in J}(\pi_i \circ f)^{-1}(C_i)$ is μ-measurable by (ii). Since the measurable cylinders generate Σ_Y, the set $f^{-1}(C)$ is μ-measurable for every $C \in \Sigma_Y$. □

By Proposition 6.4, the universal measurability of a map into $\mathbb{R}^{\mathcal{P}}$ can be checked feature by feature, just as membership in the first Baire class can (see Proposition 1.8).

We saw in Section 1 that the NIP corresponds to relative compactness in the space of functions of Baire class 1 (see Theorems 1.4, 1.6 and 1.9), and in Section 3 that, when the set $\mathcal{P}$ of predicates is countable, this correspondence extends to deep computations (see Theorem 3.1). The following theorem, valid on every Hausdorff space, characterizes the NIP in terms of universal measurability; applied to the type shards, it requires no countability of $\mathcal{P}$.

Theorem 6.5 (Bourgain–Fremlin–Talagrand [BFT78, Theorem 2F])**.** *Let X be a Hausdorff space, and let $A \subseteq \mathrm{C_p}(X)$ be pointwise bounded. The following are equivalent:*

(i) $\overline{A} \subseteq M_r(X)$.
(ii) For every compact $K \subseteq X$, $A|_K$ satisfies the NIP.
(iii) For every Radon measure μ on X, A is relatively countably compact in $\mathscr{M}^0(X, \mu)$, i.e., every countable subset of A has a limit point in $\mathscr{M}^0(X, \mu)$.

This theorem allows us to define *Monte Carlo computable* deep computations (see Definition 6.6 below). The definition is motivated by two facts.

(1) Littlewood's second principle, that every Lebesgue measurable function is "nearly continuous", is made precise by Luzin's theorem: if (X, Σ, μ) is a finite Radon measure space and Y is a second-countable topological space (e.g., $Y = \mathbb{R}^{\mathcal{P}}$ with $\mathcal{P}$ countable) with its Borel σ-algebra, then a function $f\colon X \to Y$ is measurable if and only if, for every $E \in \Sigma$ and every $\varepsilon > 0$, there is a closed set $F \subseteq E$ such that the restriction $f|_F$ is continuous and $\mu(E \setminus F) < \varepsilon$.

(2) Computability of a transform is characterized by its continuous extendibility to the space of types (see Section 2); this characterization is at the core of the first paper of this series [ADIW26].

Definition 6.6. Let $(L, \mathcal{P}, \Gamma)$ be a CCS, and let R be a collection of sizers. A transform $f\colon L \to \mathcal{L}_{\mathrm{sh}}$ is *universally Monte Carlo computable on* R if it has an extension $\tilde{f}\colon \mathcal{L}_{\mathrm{sh}} \to \mathcal{L}_{\mathrm{sh}}$ with the following property: for every $r_\bullet \in R$ there is a sizer $s_\bullet$ such that the restriction $\tilde{f}|_{\mathcal{L}[r_\bullet]}\colon \mathcal{L}[r_\bullet] \to \mathcal{L}[s_\bullet]$ is universally measurable, that is, $\pi_P \circ \tilde{f}|_{\mathcal{L}[r_\bullet]}\colon \mathcal{L}[r_\bullet] \to [-s_P, s_P]$ is μ-measurable for every Radon probability measure μ on $\mathcal{L}[r_\bullet]$ and every $P \in \mathcal{P}$.

The deep computations of a set of computations confined by an exhaustive collection of sizers are transforms $L \to \mathcal{L}_{\mathrm{sh}}$ (see Proposition 2.5). Since each type shard $\mathcal{L}[r_\bullet]$ is compact, Remark 6.2 shows that Radon probability measures suffice in Definition 6.6, in place of general Radon measures.

Theorem 6.7. *Let $(L, \mathcal{P}, \Gamma)$ be a CCS satisfying the Extendibility Axiom, let R be an exhaustive collection of sizers, and let $\Delta \subseteq \Gamma$ be R-confined. If $\pi_P \circ \Delta|_{L[r_\bullet]}$ satisfies the NIP for all $P \in \mathcal{P}$ and all $r_\bullet \in R$, then every deep computation in $\overline{\Delta}$ is universally Monte Carlo computable on R.*

Proof. Let $f \in \overline{\Delta}$, let $\tilde{f}$ be the extension of Lemma 2.7, and fix $P \in \mathcal{P}$ and $r_\bullet \in R$. By the Extendibility Axiom and confinement, $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ is a pointwise bounded set of continuous functions on the compact space $\mathcal{L}[r_\bullet]$. Its restriction to $L[r_\bullet]$ is $\pi_P \circ \Delta|_{L[r_\bullet]}$, which has the NIP by hypothesis, and $L[r_\bullet]$ is dense in $\mathcal{L}[r_\bullet]$; hence Lemma 1.10 shows that this set has the NIP, and hence so does its restriction to every compact $K \subseteq \mathcal{L}[r_\bullet]$; Theorem 6.5 then gives $\overline{\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}} \subseteq M_r(\mathcal{L}[r_\bullet])$. By Lemma 2.7, the function $\pi_P \circ \tilde{f}|_{\mathcal{L}[r_\bullet]}$ belongs to this closure, and it is therefore universally measurable. Since $\tilde{f}$ maps $\mathcal{L}[r_\bullet]$ into itself, and since P and $r_\bullet$ are arbitrary, the deep computation f satisfies Definition 6.6 with $s_\bullet = r_\bullet$ for every $r_\bullet \in R$. □

Whether the converse of Theorem 6.7 holds is Question 7.3.

6.2. **Talagrand stability and Monte Carlo computability of deep computations.** Talagrand introduced a notion closely related to the NIP in his study of Pettis integration [Tal84]. Let X be a compact Hausdorff space, let $A \subseteq \mathbb{R}^X$ be a set of functions, and let μ be a Radon probability measure on X. For a μ-measurable set $E \subseteq X$, a positive integer k and real numbers $a < b$, let

$$D_k(A, E, a, b) = \bigcup_{f \in A} \{x \in E^{2k} : f(x_{2i}) \le a,\ f(x_{2i+1}) \ge b \ \text{ for all } i < k\}.$$

Definition 6.8. The set A is *Talagrand μ-stable* if, for every μ-measurable set $E \subseteq X$ of positive measure and all reals $a < b$, there is an integer $k \ge 1$ such that

$$(\mu^{2k})^*(D_k(A, E, a, b)) < (\mu(E))^{2k},$$

where $(\mu^{2k})^*$ denotes the outer measure associated with the product measure μ^{2k} on X^{2k}.

The outer measure is needed in Definition 6.8 because $D_k(A, E, a, b)$ need not be measurable. The set $D_k(A, E, a, b)$ is measurable when A is a countable set of μ-measurable functions, since it is then a countable union of measurable sets.

Remarks 6.9. Let X, μ, $A \subseteq \mathbb{R}^X$ and $E \subseteq X$ be as above, and let $k \geq 1$ be an integer and $a < b$ be reals. The following hold.

(1) If $A \subseteq B \subseteq \mathbb{R}^X$, then
$$D_k(A, E, a, b) \subseteq D_k(B, E, a, b).$$
(2) If $a < a' < b' < b$, then
$$D_k(\overline{A}, E, a, b) \subseteq D_k(A, E, a', b').$$
[*Proof*: If $x \in D_k(\overline{A}, E, a, b)$, then there is $f \in \overline{A}$ such that $f(x_{2i}) \leq a < a'$ and $f(x_{2i+1}) \geq b > b'$ for all $i < k$. By the definition of the topology of pointwise convergence, there is $g \in A$ such that $g(x_{2i}) < a'$ and $g(x_{2i+1}) > b'$ for all $i < k$; hence $x \in D_k(A, E, a', b')$.]
(3) Every subset of a Talagrand μ-stable set is Talagrand μ-stable, by (1).
(4) If A is Talagrand μ-stable, then so is $\overline{A}$, by (2).

The main result of this subsection, Corollary 6.14, states that the deep computations of a confined set of computations are universally Monte Carlo computable when, for every feature and every type shard, the corresponding set of functions is Talagrand μ-stable for every Radon probability measure μ, that is, *universally Talagrand stable* in the sense of Definition 6.12 below. Its proof rests on Lemma 6.11, by which the pointwise closure of a Talagrand μ-stable set consists of μ-measurable functions, and the lemma in turn rests on the following characterization of measurable functions (compare Fact 1.2).

Fact 6.10 ([Fre03, Lemma 413G])**.** *Let (X, Σ, μ) be a measure space, and let $\mathcal{K} \subseteq \Sigma$ be a collection of measurable sets such that the following conditions hold:*

- *(X, Σ, μ) is complete, i.e., for all $E \in \Sigma$ with $\mu(E) = 0$ and $F \subseteq E$ we have $F \in \Sigma$.*
- *(X, Σ, μ) is semi-finite, i.e., for all $E \in \Sigma$ with $\mu(E) = \infty$ there exists $F \subseteq E$ such that $F \in \Sigma$ and $0 < \mu(F) < \infty$.*
- *(X, Σ, μ) is saturated, i.e., $E \in \Sigma$ if and only if $E \cap F \in \Sigma$ for all $F \in \Sigma$ with $\mu(F) < \infty$.*
- *(X, Σ, μ) is inner regular with respect to $\mathcal{K}$, i.e., for all $E \in \Sigma$*
$$\mu(E) = \sup\{\mu(K) : K \in \mathcal{K} \text{ and } K \subseteq E\}.$$

(In particular, if X is compact Hausdorff, $\mathcal{K}$ is the collection of compact subsets of X, μ is a Radon probability measure on X, and Σ is the completion of the Borel σ-algebra by μ, then all these conditions hold.) Then a function $f\colon X \to \mathbb{R}$ is measurable if and only if, for every $K \in \mathcal{K}$ with $0 < \mu(K) < \infty$ and all reals $a < b$, either $\mu^*(K_{\leq a}) < \mu(K)$ or $\mu^*(K_{\geq b}) < \mu(K)$, where $K_{\leq a} = \{x \in K : f(x) \leq a\}$ and $K_{\geq b} = \{x \in K : f(x) \geq b\}$.

We first show that Talagrand stability passes to the pointwise closure and forces measurability.

Lemma 6.11. *If A is Talagrand μ-stable, then $\overline{A}$ is also Talagrand μ-stable and $\overline{A} \subseteq \mathscr{M}^0(X, \mu)$.*

Proof. By Remarks 6.9, the closure $\overline{A}$ is Talagrand μ-stable, and hence so is $\{f\}$ for every $f \in \overline{A}$. It therefore suffices to show that if $\{f\}$ is Talagrand μ-stable, then f is μ-measurable. Suppose that f is not μ-measurable. Then Fact 6.10 gives a compact

set E of positive measure and reals $a < b$ with $\mu^*(E_{\leq a}) = \mu^*(E_{\geq b}) = \mu(E)$, where $E_{\leq a} = \{x \in E : f(x) \leq a\}$ and $E_{\geq b} = \{x \in E : f(x) \geq b\}$. For every $k \geq 1$ we have $(E_{\leq a} \times E_{\geq b})^k \subseteq D_k(\{f\}, E, a, b)$, and this product has outer measure $(\mu(E))^{2k}$ (see [Tal84]); hence $\{f\}$ is not Talagrand μ-stable. □

Definition 6.12. The set A is *universally Talagrand stable* if it is Talagrand μ-stable for every Radon probability measure μ on X.

Universal Talagrand stability implies the NIP.

Proposition 6.13. *Let X be a compact Hausdorff space, and let $A \subseteq \mathrm{C_p}(X)$ be pointwise bounded. If A is universally Talagrand stable, then A satisfies the NIP.*

Proof. By Lemma 6.11, $\overline{A} \subseteq \mathscr{M}^0(X, \mu)$ for every Radon probability measure μ on X, that is, $\overline{A} \subseteq M_r(X)$. By the implication (i)⇒(ii) of Theorem 6.5, applied with $K = X$, the set A satisfies the NIP. □

Corollary 6.14. *Let $(L, \mathcal{P}, \Gamma)$ be a CCS satisfying the Extendibility Axiom, let R be an exhaustive collection of sizers, and let $\Delta \subseteq \Gamma$ be R-confined. If $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ is universally Talagrand stable for all $P \in \mathcal{P}$ and all $r_\bullet \in R$, then every deep computation in $\overline{\Delta}$ is universally Monte Carlo computable on R.*

Proof. Let $P \in \mathcal{P}$ and $r_\bullet \in R$ be given. By the Extendibility Axiom and confinement, $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ is a pointwise bounded set of continuous functions on the compact space $\mathcal{L}[r_\bullet]$, and, by Proposition 6.13, it has the NIP; hence so does its restriction $\pi_P \circ \Delta|_{L[r_\bullet]}$. Theorem 6.7 now completes the proof. □

In the context of deep computations we have thus found two routes to Monte Carlo computability, the NIP and universal Talagrand stability, and the second implies the first (see Proposition 6.13). For countable sets of measurable functions, the following two results show that the converse implication depends on the axioms of set theory. On the one hand, it is consistent with the Zermelo–Fraenkel axioms of set theory with the axiom of choice (ZFC) that, for such sets, the NIP implies universal Talagrand stability.

Theorem 6.15 (Talagrand [Tal84, Theorem 9-3-1(a)])**.** *Let X be a compact Hausdorff space, and let $A \subseteq M_r(X)$ be countable and pointwise bounded. Assume that $[0, 1]$ is not the union of fewer than $\mathfrak{c}$ closed sets of Lebesgue measure zero. If A satisfies the NIP, then A is universally Talagrand stable.*

(The hypothesis on $[0, 1]$ follows, for example, from the Continuum Hypothesis.)

On the other hand, for Lebesgue measure, the failure of this implication is also consistent.

Theorem 6.16 (Fremlin and Shelah [FS93])**.** *It is consistent with ZFC that there exists a countable pointwise bounded set of Lebesgue measurable functions with the NIP that is not Talagrand stable with respect to Lebesgue measure.*

Both results concern countable sets of measurable functions. By the Extendibility Axiom, computations are continuous on the type shards, and continuous functions are measurable; hence, by Theorem 6.15, it is consistent that the NIP implies universal Talagrand stability for countable sets of computations. The example of Fremlin and Shelah, however, is a set of measurable functions, and it does not settle whether some countable set of computations can consistently have the NIP without being universally Talagrand stable; we leave this open (Question 7.4).

7. Open questions

We collect the questions left open by this paper. The first concerns the hierarchy of Section 3.2, whose levels were defined through the topology of the space of deep computations; Proposition 3.4 describes them by the number of tests needed to tell deep computations apart, but these descriptions still involve infinitely many tests.

Question 7.1. Is there a finitary, combinatorial characterization of NIP_1, NIP_2 and NIP_3, in the spirit of the characterization of NIP by finite patterns in Theorem 3.1?

In Section 5 we place a single iterated map at every level of the hierarchy except one: at the level NIP but not NIP_1, both of our examples are actions of groups with two generators (see Section 5.5).

Question 7.2. Is there a cascade CCS that is NIP but not NIP_1? Equivalently, is there a continuous map γ of a compact metrizable space L such that (L, γ) is tame and $E(L, \gamma)$ is not first countable?

For actions of $\mathbb{Z}$, and more generally of abelian groups, the analogous question is Question 6.8 of Glasner and Megrelishvili [GM22].

Theorem 6.7 shows that NIP suffices for universal Monte Carlo computability; whether it is also necessary is the next question.

Question 7.3. Under the assumptions of Theorem 6.7, suppose that every deep computation in $\overline{\Delta}$ is universally Monte Carlo computable on R. Must $\pi_P \circ \Delta|_{L[r_\bullet]}$ have the NIP for all $P \in \mathcal{P}$ and all $r_\bullet \in R$?

Finally, in Section 6 we obtained two routes to Monte Carlo computability, the NIP and universal Talagrand stability; the second implies the first, and the results of Talagrand and of Fremlin and Shelah leave open whether, for computations, the first can hold without the second.

Question 7.4. Is it consistent with ZFC that some countable, pointwise bounded set of continuous functions on a compact Hausdorff space satisfies the NIP but is not universally Talagrand stable? Can this happen, consistently, for the sets $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ arising from a countable set Δ of computations as in Theorem 6.7?

References

[ADIW26] Samson Alva, Eduardo Dueñez, José Iovino, and Claire Walton, *Approximability of deep computations*, Mathematical Structures in Computer Science (2026), To appear; arXiv:2409.06064.

[ADK08] Spiros A. Argyros, Pandelis Dodos, and Vassilis Kanellopoulos, *A classification of separable Rosenthal compacta and its applications*, Dissertationes Mathematicae **449** (2008), 1–55.

[Ark92] A. V. Arkhangel'skii, *Topological function spaces*, Mathematics and its Applications (Soviet Series), vol. 78, Kluwer Academic Publishers Group, Dordrecht, 1992, Translated from the Russian by R. A. M. Hoksbergen. MR 1144519

[Bai99] René Baire, *Sur les fonctions de variables réelles*, Annali di Matematica Pura ed Applicata (3) **3** (1899), 1–123.

[BEHW89] Anselm Blumer, Andrzej Ehrenfeucht, David Haussler, and Manfred K. Warmuth, *Learnability and the Vapnik–Chervonenkis dimension*, J. Assoc. Comput. Mach. **36** (1989), no. 4, 929–965.

[BFB16] Richard L. Burden, J. Douglas Faires, and Annette M. Burden, *Numerical analysis*, 10th ed., Cengage Learning, Boston, MA, 2016.

[BFT78] J. Bourgain, D. H. Fremlin, and M. Talagrand, *Pointwise compact sets of Baire-measurable functions*, American Journal of Mathematics **100** (1978), no. 4, 845–886.

[BKK] Shaojie Bai, J. Zico Kolter, and Vladlen Koltun, *Deep equilibrium models*, Preprint. https://arxiv.org/abs/1909.01377, https://implicit-layers-tutorial.org/.

[Bre65] Jack E. Bresenham, *Algorithm for computer control of a digital plotter*, IBM Systems Journal **4** (1965), no. 1, 25–30.

[CRBD] Ricky T. Q. Chen, Yulia Rubanova, Jesse Bettencourt, and David Duvenaud, *Neural ordinary differential equations*, Preprint. https://arxiv.org/abs/1806.07366.

[Deb13] Gabriel Debs, *Descriptive aspects of Rosenthal compacta*, Recent Progress in General Topology III, Springer, 2013, pp. 205–227.

[DnI17] Eduardo Dueñez and José N. Iovino, *Model theory and metric convergence I: Metastability and dominated convergence*, Beyond First Order Model Theory, Volume I, CRC Press, Boca Raton, FL, 2017, pp. 131–187. MR 3729326

[Ell93] Robert Ellis, *The enveloping semigroup of projective flows*, Ergodic Theory and Dynamical Systems **13** (1993), no. 4, 635–660.

[Eng89] Ryszard Engelking, *General topology*, second ed., Sigma Series in Pure Mathematics, vol. 6, Heldermann Verlag, Berlin, 1989, Translated from the Polish by the author. MR MR1039321 (91c:54001)

[Fre03] David H. Fremlin, *Measure theory, volume 4: Topological measure spaces*, Torres Fremlin, Colchester, UK, 2003.

[FS93] David H. Fremlin and Saharon Shelah, *Pointwise compact and stable sets of measurable functions*, Journal of Symbolic Logic **58** (1993), no. 2, 435–455.

[Gal00] A. Galántai, *The theory of Newton's method*, Journal of Computational and Applied Mathematics **124** (2000), no. 1, 25–44, Numerical Analysis 2000. Vol. IV: Optimization and Nonlinear Equations.

[Gla06] Eli Glasner, *On tame dynamical systems*, Colloquium Mathematicum **105** (2006), no. 2, 283–295.

[GM22] Eli Glasner and Michael Megrelishvili, *Todorčević' trichotomy and a hierarchy in the class of tame dynamical systems*, Transactions of the American Mathematical Society **375** (2022), no. 7, 4513–4548.

[GM23] ———, *More on tame dynamical systems*, arXiv:2302.09571, 2023.

[God80] Gilles Godefroy, *Compacts de Rosenthal*, Pacific J. Math. **91** (1980), no. 2, 293–306. MR 615679

[Gro52] A. Grothendieck, *Critères de compacité dans les espaces fonctionnels généraux*, Amer. J. Math. **74** (1952), 168–186. MR 47313

[HT19] Haim Horowitz and Stevo Todorcevic, *Compact sets of baire class one functions and maximal almost disjoint families*, 2019.

[Kec95] Alexander S. Kechris, *Classical descriptive set theory*, Graduate Texts in Mathematics, vol. 156, Springer-Verlag, 1995.

[Kei23] H. Jerome Keisler, *Model theory for real-valued structures*, Beyond First Order Model Theory, Volume II (José Iovino, ed.), CRC Press, Boca Raton, FL, 2023.

[KM81] Jean-Louis Krivine and Bernard Maurey, *Espaces de Banach stables*, Israel J. Math. **39** (1981), no. 4, 273–295. MR 83a:46030

[Köh95] Angela Köhler, *Enveloping semigroups for flows*, Proceedings of the Royal Irish Academy. Section A **95A** (1995), no. 2, 179–191.

[Kri72] J.-L. Krivine, *Théorie des modèles et espaces L^p*, C. R. Acad. Sci. Paris Sér. A-B **275** (1972), A1207–A1210. MR 46 #7879

[Kri74] ———, *Langages à valeurs réelles et applications*, Fund. Math. **81** (1974), 213–253, Collection of articles dedicated to Andrzej Mostowski on the occasion of his sixtieth birthday, III. MR 50 #1873

[Leb04] Henri Lebesgue, *Une propriété caractéristique des fonctions de classe 1*, Bulletin de la Société Mathématique de France **32** (1904), 229–242.

[Li11] Jian Li, *Chaos and entropy for interval maps*, Journal of Dynamics and Differential Equations **23** (2011), no. 2, 333–352.

[Lot02] M. Lothaire, *Algebraic combinatorics on words*, Encyclopedia of Mathematics and its Applications, vol. 90, Cambridge University Press, Cambridge, 2002.

[MH40] Marston Morse and Gustav A. Hedlund, *Symbolic dynamics II. Sturmian trajectories*, American Journal of Mathematics **62** (1940), 1–42.

[Pap02] E. Pap (ed.), *Handbook of measure theory. Vol. I, II*, North-Holland, Amsterdam, 2002. MR 1953489

[Ros74] Haskell P. Rosenthal, *A characterization of Banach spaces containing l^1*, Proc. Nat. Acad. Sci. U.S.A. **71** (1974), 2411–2413. MR 358307

[RPK19] M. Raissi, P. Perdikaris, and G. E. Karniadakis, *Physics-informed neural networks: a deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations*, J. Comput. Phys. **378** (2019), 686–707. MR 3881695

[Sau72] N. Sauer, *On the density of families of sets*, J. Combinatorial Theory Ser. A **13** (1972), 145–147. MR 307902

[She71] Saharon Shelah, *Stability, the f.c.p., and superstability; model theoretic properties of formulas in first order theory*, Ann. Math. Logic **3** (1971), no. 3, 271–362. MR 317926

[She72] ———, *A combinatorial problem; stability and order for models and theories in infinitary languages*, Pacific J. Math. **41** (1972), 247–261. MR 307903

[She90] ———, *Classification theory and the number of nonisomorphic models*, second ed., Studies in Logic and the Foundations of Mathematics, vol. 92, North-Holland Publishing Co., Amsterdam, 1990.

[Sim15] Pierre Simon, *A guide to nip theories*, Lecture Notes in Logic, vol. 44, Cambridge University Press, 2015.

[SSBD14] Shai Shalev-Shwartz and Shai Ben-David, *Understanding machine learning: From theory to algorithms*, Cambridge University Press, Cambridge, 2014.

[Tal84] Michel Talagrand, *Pettis integral and measure theory*, Memoirs of the American Mathematical Society, vol. 51, American Mathematical Society, Providence, RI, USA, 1984, Includes bibliography (pp. 220–224) and index.

[Tka11] Vladimir V. Tkachuk, *A C_p-theory problem book: Topological and function spaces*, Problem Books in Mathematics, Springer, 2011.

[Tod97] Stevo Todorčević, *Topics in topology*, Lecture Notes in Mathematics, vol. 1652, Springer Berlin, Heidelberg, 1997.

[Tod99] Stevo Todorčević, *Compact subsets of the first Baire class*, Journal of the American Mathematical Society **12** (1999), no. 4, 1179–1212.

[Tod23] ———, *Dense metrizability*, Annals of Pure and Applied Logic **175** (2023), 103327.

[Val84] Leslie G. Valiant, *A theory of the learnable*, Communications of the ACM **27** (1984), no. 11, 1134–1142.

[VC71] Vladimir N Vapnik and Alexey Ya Chervonenkis, *On the uniform convergence of relative frequencies of events to their probabilities*, Theory of Probability & Its Applications **16** (1971), no. 2, 264–280.

[VC74] ———, *Theory of pattern recognition*, Nauka, Moscow, 1974, German Translation: Theorie der Zeichenerkennung, Akademie-Verlag, Berlin, 1979.