

TOPOLOGICAL AND LOGICAL ASPECTS OF THE PROBLEM OF HIDDEN VARIABLES IN QUANTUM MECHANICS

PRELIMINARY NOTES PREPARED BY J. IOVINO

For the University of Toronto–UT San Antonio Mathematics research group

© 2026 José Iovino. All rights reserved.

These notes are a work in progress. Please do not post or redistribute copies;
link instead to the current version at [utsa-quantum-math.github.io](https://github.com/utsa-quantum-math).

Comments are welcome at jose.iovino@utsa.edu.

ABSTRACT. We examine two proofs of the Bell–Kochen–Specker theorem, which rules out non-contextual hidden variables in quantum mechanics. Bell’s proof combines Gleason’s theorem with elementary topology and is infinitary; Kochen and Specker’s proof exhibits a finite configuration of rays and is constructive. The two arguments were found independently in the mid-1960s. We show that the compactness theorem of propositional logic links them: compactness turns Bell’s argument into the existence of a finite configuration of the kind that Kochen and Specker built by hand. We also show that quantum measures have no counterpart of Dirac measures and, unlike classical measures, do not form a lattice. A timeline traces the history of the problem of hidden variables.

0. PRELIMINARIES (ALL THE LINEAR ALGEBRA AND PHYSICS YOU WILL NEED TO READ THESE NOTES)

We assume that the reader is familiar with Hilbert spaces, with eigenvalues and eigenvectors of linear operators, and with orthogonal projections. Everything else we need, from both linear algebra and physics, is recalled in this short section.

Unless stated otherwise (as in Remark 1.6 and Section 5), the symbol H denotes a complex Hilbert space with $3 \leq \dim H \leq \infty$, where ∞ stands for $\aleph_0$: unless stated otherwise, Hilbert spaces are separable. (Separability is not a technicality; Remark 1.7 below explains what large cardinals have to do with it.) Everything below holds verbatim for real and quaternionic Hilbert spaces as well.

We use the physicists’ notation for vectors and functionals, which we now explain, since this notation is not standard in mathematics. The inner product $\langle \xi | \eta \rangle$ of $\xi, \eta \in H$ is linear in the second variable and conjugate-linear in the first. A vector η is also written $|\eta\rangle$ (a *ket*), and $\langle \xi |$ (a *bra*) denotes the linear functional $\eta \mapsto \langle \xi | \eta \rangle$. Juxtaposition is composition: for a bounded operator A , the number $\langle \xi | A | \eta \rangle$ means $\langle \xi | A \eta \rangle$, and $|\xi\rangle \langle \xi |$ is the operator $\eta \mapsto \langle \xi | \eta \rangle \xi$. We write $S(H)$ for the unit sphere of H .

Let $L(H)$ be the lattice of closed subspaces of H — an *orthomodular lattice* in the usual sense — which we identify with the corresponding orthogonal projections. Recall that the orthogonal projection P onto a closed subspace V is defined by writing each $\eta \in H$ uniquely as a sum $\eta = \eta_V + \eta_\perp$ with $\eta_V \in V$ and $\eta_\perp \in V^\perp$, and putting $P\eta = \eta_V$. Two properties follow at once. First, the projection P satisfies $P^2 = P$, because P fixes every vector of V . Second, the projection P satisfies $P^* = P$, because for $\xi, \eta \in H$ we have

$$\langle \xi | P \eta \rangle = \langle \xi_V | \eta_V \rangle = \langle P \xi | \eta \rangle,$$

since the cross terms $\langle \xi_\perp | \eta_V \rangle$ and $\langle \xi_V | \eta_\perp \rangle$ vanish. Conversely, a bounded operator P with $P^2 = P = P^*$ is the orthogonal projection onto the range of P , which is closed. Thus, the

elements of $L(H)$ are exactly the bounded operators P with $P^2 = P = P^*$, and from these two identities

$$\langle \eta | P | \eta \rangle = \langle \eta | P^* P | \eta \rangle = \langle P \eta | P \eta \rangle = \|P \eta\|^2 \quad (\eta \in H). \quad (1)$$

The lattice $L(H)$ carries a partial order $P \leq Q$ (inclusion of ranges), a meet $P \wedge Q$ (intersection of ranges), a join $P \vee Q$ (closed span of the ranges), an orthocomplement $P \mapsto P^\perp = I - P$, and extremal elements 0 and $I = H$. Two projections P and Q are *orthogonal*, written $P \perp Q$, when their ranges are orthogonal, equivalently $P \leq Q^\perp$. A *basis* of $L(H)$ is the set of rank-one projections onto the vectors of some orthonormal basis of H .

For $\psi \in S(H)$, the operator $P_\psi = |\psi\rangle\langle\psi|$ is the orthogonal projection onto the one-dimensional subspace $\mathbb{C}\psi$: the projection P_ψ fixes ψ and kills every vector orthogonal to ψ , and $P_\psi^2 = P_\psi = P_\psi^*$. The rank-one projections are exactly the minimal nonzero elements, the *atoms*, of the lattice $L(H)$.

We now turn to the physical meaning of these objects. In quantum mechanics, a projection $P \in L(H)$ represents a yes/no question about a physical system, that is, an experiment with two possible outcomes, and $P^\perp = I - P$ represents the same question with the answers exchanged. A state of the system assigns to each projection P the probability $\mu(P)$ of obtaining the answer “yes” to P . The question I is always answered “yes”, so $\mu(I) = 1$. If $P_0, P_1, \dots$ are pairwise orthogonal, the corresponding questions are mutually exclusive and can be tested together, and the probability of “yes” to their join $\bigvee_i P_i$ is the sum of the probabilities $\mu(P_i)$. These requirements are the axioms of a quantum measure (see Definition 1.1 below). Which assignments μ actually occur in physics is decided by the Born rule, which leads to the notion of density operator.



Remark 0.1 (The Born rule). The simplest states of a physical system, the *pure states*, are described by unit vectors $\psi \in S(H)$. The *Born rule* says that in the state ψ the probability of the answer “yes” to the question P is

$$\mu(P) = \|P\psi\|^2 = \langle \psi | P | \psi \rangle$$

(the second equality is (1)). An important special case is the question “is the system in the state ξ ?” for another unit vector $\xi \in S(H)$. This question is represented by the rank-one projection $P_\xi = |\xi\rangle\langle\xi|$ onto the line $\mathbb{C}\xi$, which maps a vector η to $\langle \xi | \eta \rangle \xi$. Since $P_\xi \psi = \langle \xi | \psi \rangle \xi$ and $\|\xi\| = 1$, the Born rule gives

$$\mu(P_\xi) = \|P_\xi \psi\|^2 = |\langle \xi | \psi \rangle|^2,$$

the familiar probability that a system in the state ψ is found in the state ξ .

A pure state represents complete knowledge of the system: no further information about how the system was prepared could sharpen the probabilities $\|P\psi\|^2$. Even so, these probabilities are in general strictly between 0 and 1. The randomness they describe is therefore not due to any lack of information on our part; according to quantum mechanics this randomness is intrinsic to the system itself, and the theorems of these notes are precisely about whether this randomness could be explained away as ignorance of some hidden variables.

In practice, however, even this complete knowledge is rarely available. Real physical sources do not produce a single pure state: the light of an ordinary lamp, for instance, consists of photons whose polarizations vary at random from one photon to the next, so no single vector describes the beam. Our information about how a system was prepared is usually incomplete, and the imperfections of the preparing apparatus mean that we know at best which pure states may have been produced, and with what probabilities. This introduces a second kind of randomness, of an entirely different nature from the first: this randomness is classical, of the same kind as in a game of cards, and reflects only our ignorance, that is, the

imperfection of our information about the system, not any property of the system itself. A state that combines the two kinds of randomness is called a *mixed state*, or a *mixture*.¹

Accordingly, suppose one knows only that the system has been prepared in the state ψ_k with probability λ_k , for a family of states $\psi_k \in S(H)$ and weights $\lambda_k \geq 0$ with $\sum_k \lambda_k = 1$. Here the weights λ_k express our ignorance of the preparation, while each $\langle \psi_k | P | \psi_k \rangle$ is a quantum probability given by the Born rule. Combining the two, the probability of “yes” to P is then the weighted average (or expected value)

$$\sum_k \lambda_k \langle \psi_k | P | \psi_k \rangle. \quad (2)$$

We now show that this expression depends on the preparation only through the operator

$$\rho = \sum_k \lambda_k |\psi_k\rangle\langle\psi_k| = \sum_k \lambda_k P_{\psi_k},$$

and we look for a single formula that computes (2) from ρ and P . The trace provides this formula, as we now explain.

When H is finite-dimensional, the *trace* of a linear operator T on H is

$$\text{Tr}(T) = \sum_i \langle e_i | T e_i \rangle,$$

where (e_i) is an orthonormal basis of H . Thus $\text{Tr}(T)$ is the sum of the diagonal entries of the matrix of T in the basis (e_i) , a sum that does not depend on the choice of basis. When H is infinite-dimensional, the same formula defines the trace of the *trace-class* operators, for which the sum converges absolutely and is again independent of the basis. We need only two facts about trace-class operators: every operator of finite rank, in particular every rank-one projection P_ψ , is trace-class; and if T is trace-class and A is bounded, then TA and AT are trace-class. For the definition and these facts, see [RS80, Section VI.6]. The reader who prefers may assume that H is finite-dimensional throughout this section; the infinite-dimensional case differs only in these points of analysis.

The following observation will play an important role in these notes.

Observation 0.2. *Let A be a bounded operator on H and let $\psi \in S(H)$ be given. Then*

$$\text{Tr}(AP_\psi) = \text{Tr}(P_\psi A) = \langle \psi | A | \psi \rangle.$$

In particular, we have $\text{Tr}(P_\psi P) = \langle \psi | P | \psi \rangle = \|P\psi\|^2$ for every $P \in L(H)$, and $\text{Tr}(P_\xi P_\psi) = |\langle \xi | \psi \rangle|^2$ for $\xi \in S(H)$.

Proof. Compute the trace in an orthonormal basis (e_i) of H with $e_0 = \psi$. Then $P_\psi e_0 = \psi$, and $P_\psi e_i = 0$ for $i \geq 1$ because e_i is orthogonal to ψ . So only the first term survives:

$$\text{Tr}(AP_\psi) = \sum_i \langle e_i | AP_\psi e_i \rangle = \langle \psi | A \psi \rangle = \langle \psi | A | \psi \rangle.$$

Similarly, since $P_\psi A e_i = \langle \psi | A e_i \rangle \psi$ and $\langle e_i | \psi \rangle = 0$ for $i \geq 1$, we have

$$\text{Tr}(P_\psi A) = \sum_i \langle e_i | P_\psi A e_i \rangle = \sum_i \langle e_i | \psi \rangle \langle \psi | A e_i \rangle = \langle \psi | A \psi \rangle.$$

¹The same mathematical description arises in a situation where ignorance is not the explanation: a system A that has interacted with another system B , and is therefore entangled with B , has in general no pure state of A alone, even when the combined system is in a pure state, and the statistics of A are again those of a mixture.

If $P \in L(H)$, then by (1) $\langle \psi | P \psi \rangle = \langle P \psi | P \psi \rangle = \|P \psi\|^2$. Finally $P_\psi \xi = \langle \psi | \xi \rangle \psi$, so taking $A = P_\xi$ gives $\text{Tr}(P_\xi P_\psi) = \|P_\psi \xi\|^2 = |\langle \xi | \psi \rangle|^2$. ☺

Returning to the Born rule: by Observation 0.2 and linearity of the trace, the probability (2) is

$$\sum_k \lambda_k \langle \psi_k | P | \psi_k \rangle = \sum_k \lambda_k \text{Tr}(P_{\psi_k} P) = \text{Tr}(\rho P).$$

So the Born rule, for pure and mixed preparations alike, takes the single form $P \mapsto \text{Tr}(\rho P)$. The operator ρ has three properties that do not depend on how ρ was obtained. First, the operator ρ is positive, since $\langle \xi | \rho | \xi \rangle = \sum_k \lambda_k |\langle \psi_k | \xi \rangle|^2 \geq 0$. Second, the operator ρ is trace-class. Third, the trace of ρ is 1; this property is nothing but the requirement that the weights λ_k add up to 1: each P_{ψ_k} has trace $\text{Tr}(P_{\psi_k}) = \langle \psi_k | \psi_k \rangle = 1$ (take $A = I$ in Observation 0.2), so

$$\text{Tr}(\rho) = \sum_k \lambda_k \text{Tr}(P_{\psi_k}) = \sum_k \lambda_k = 1.$$

Equivalently, the number $\text{Tr}(\rho) = \text{Tr}(\rho I)$ is the probability of the answer “yes” to the question I , which is always answered “yes”. These are the properties singled out in the following definition.

Definition 0.3 (Density operator). A *density operator* (or *density matrix*) on H is a trace-class operator ρ that is positive, i.e. $\langle \xi | \rho | \xi \rangle \geq 0$ for all $\xi \in H$, and has $\text{Tr}(\rho) = 1$.

Conversely, every density operator arises from a preparation as in Remark 0.1. Suppose first that H is finite-dimensional. A positive operator ρ on a complex Hilbert space is self-adjoint, so ρ has an orthonormal basis of eigenvectors ψ_k ; the corresponding eigenvalues $\lambda_k = \langle \psi_k | \rho | \psi_k \rangle$ are nonnegative because ρ is positive, and they add up to $\text{Tr}(\rho) = 1$. Hence ρ can be written as

$$\rho = \sum_k \lambda_k |\psi_k\rangle\langle\psi_k|, \quad \lambda_k \geq 0, \quad \sum_k \lambda_k = 1, \quad (3)$$

where ψ_k has eigenvalue λ_k . When H is infinite-dimensional, the same decomposition holds with a countable orthonormal family (ψ_k) of eigenvectors, by the spectral theorem for compact self-adjoint operators [RS80, Section VI.5] (density operators, being trace-class, are compact). So density operators are exactly the probabilistic mixtures of the *pure states* P_ψ with $\psi \in S(H)$.

Finally, taking $A = \rho$ in Observation 0.2, the probability of finding a system in the state ρ in the pure state ψ is

$$\text{Tr}(\rho P_\psi) = \langle \psi | \rho | \psi \rangle.$$

If ψ is an eigenvector of ρ with eigenvalue λ , then $\langle \psi | \rho | \psi \rangle = \lambda \|\psi\|^2 = \lambda$; in general, the number $\langle \psi | \rho | \psi \rangle$ is a weighted average of the eigenvalues of ρ , namely $\sum_k \lambda_k |\langle \psi_k | \psi \rangle|^2$.

This is all the quantum mechanics and linear algebra needed for the rest of these notes.

1. QUANTUM MEASURES AND GLEASON’S THEOREM

In this section we define quantum measures, state the main theorem of these notes, and recall Gleason’s Theorem, which describes all quantum measures when $\dim H \geq 3$.

Definition 1.1 (Quantum measure). A *quantum measure* (or *state*²) on $L(H)$ is a function $\mu : L(H) \rightarrow [0, 1]$ with $\mu(I) = 1$ that is countably additive on orthogonal families: for pairwise orthogonal $\{P_i\}$, we have

$$\mu\left(\bigvee_i P_i\right) = \sum_i \mu(P_i).$$

A *two-valued quantum measure* is a quantum measure $v : L(H) \rightarrow \{0, 1\}$.

What a two-valued quantum measure represents. Definition 1.1 is purely mathematical: the definition concerns functions on a lattice and mentions nothing physical. Read physically, however, a two-valued quantum measure expresses a precise hypothesis about nature. Recall from Section 0 that a projection P represents a yes/no question about a physical system. A two-valued quantum measure v answers every such question in advance: the answer to P is “yes” if $v(P) = 1$ and “no” if $v(P) = 0$, and this answer does not depend on which other compatible questions are asked together with P . The hypothesis that every yes/no question has such a predetermined answer is what physicists call a *non-contextual hidden-variable assignment*, and a two-valued quantum measure is also called a *valuation*.

The “only one winner per basis” lemma. The first mathematical consequence of Definition 1.1 is elementary, but it plays a pivotal role in these notes: the answers that a valuation gives to the questions of an orthonormal basis are consistent, in the sense that exactly one of them is “yes”.

Lemma 1.2. *Let v be a two-valued quantum measure on $L(H)$ and $\{e_i\}_{i \in I}$ an orthonormal basis of H . Exactly one index $i \in I$ has $v(P_{e_i}) = 1$; the rest have $v(P_{e_i}) = 0$.*

Proof. For $i \neq j$ in I , orthogonality gives $v(P_{e_i}) + v(P_{e_j}) = v(P_{e_i} \vee P_{e_j}) \in \{0, 1\}$, so at most one index has value 1. Summing over the basis, we obtain $\sum_i v(P_{e_i}) = v(I) = 1$; since each term is 0 or 1, exactly one equals 1. ☺



Remarks 1.3.

- (1) The proof of Lemma 1.2 does not use $\dim H \geq 3$; the lemma holds for any orthonormal basis of cardinality at least 2, in any dimension. This dimension-independence might suggest that the lemma alone proves Theorem 1.4, which is not the case (see item 2 below).
- (2) The lemma only pins down, separately, one privileged ray in *each* orthonormal basis, and says nothing about how the privileged ray of one basis relates to that of a different, overlapping basis. To prove Theorem 1.4, we must make the choices in different bases interact, and these notes give two ways of doing so. Bell’s argument (see Section 2 below) uses Gleason’s Theorem to force the choices to fit together into one continuous function on the sphere $S(H)$, and then uses the connectedness of $S(H)$ to break that function against itself.³ Kochen and Specker’s argument (see Section 4 below) uses instead a finite family of bases that overlap in a pattern that no choice of privileged rays can respect.

²Gleason called this object a *measure* [Gle57]; the quantum-logic tradition (Mackey [Mac63], Piron [Pir64]) relabeled this object a *state*, and much of the older literature still uses that name; later authors (e.g. [Ham03]) reverted to *quantum measure*, the term used throughout these notes.

³We observe that $S(H)$ is connected; in fact, the sphere $S(H)$ is even path-connected: if $u, w \in S(H)$ and $w \neq -u$, the path $t \mapsto \frac{(1-t)u+tw}{\|(1-t)u+tw\|}$ with $t \in [0, 1]$ joins u to w inside $S(H)$, and the path $\theta \mapsto e^{i\theta}u$ with $\theta \in [0, \pi]$ joins u to $-u$.

Why valuations matter. If valuations existed, the randomness of quantum mechanics could be explained as ignorance: a state would be an average of valuations, just as a classical probability measure is an average of Dirac measures, and quantum probabilities would merely record our uncertainty about which valuation describes the system. The main theorem of these notes rules out this explanation.

Theorem 1.4 (Main Theorem, Bell–Kochen–Specker). *If $3 \leq \dim H \leq \infty$, then no two-valued quantum measure exists on $L(H)$.*

We give two independent proofs of Theorem 1.4: a topological one, due to Bell (see Section 2 below; Bell’s proof is the simpler of the two) and a combinatorial one, due to Kochen and Specker (see Section 4 below). Bell’s proof relies on Gleason’s Theorem, which describes all quantum measures in terms of density operators (see Definition 0.3). For every density operator ρ , the map $P \mapsto \text{Tr}(\rho P)$ is a quantum measure: this map takes values in $[0, 1]$ (by (3) and Observation 0.2, we have $\text{Tr}(\rho P) = \sum_k \lambda_k \|P\psi_k\|^2$), takes the value $\text{Tr}(\rho) = 1$ at I , and is additive on finite orthogonal families, because the join of finitely many pairwise orthogonal projections is their sum and the trace is linear. In finite dimension every orthogonal family of nonzero projections is finite, so finite additivity is all that is needed. In infinite dimension, countable additivity follows in the same way from the continuity of $A \mapsto \text{Tr}(\rho A)$ on bounded increasing sequences of projections [RS80, Section VI.6]. By Remark 0.1, these are exactly the quantum measures given by the Born rule. Gleason’s Theorem (see Theorem 1.5 below) says that, conversely, every quantum measure arises in this way: in dimension at least 3, the Born rule is forced by the lattice structure of $L(H)$ alone.

We can now state Gleason’s Theorem.

Theorem 1.5 (Gleason’s Theorem [Gle57]). *If $3 \leq \dim H \leq \infty$, then every quantum measure μ on $L(H)$ has the form $\mu(P) = \text{Tr}(\rho P)$ for a unique density operator ρ (see Definition 0.3).*

We take Gleason’s Theorem as given, since every known proof (Gleason’s original argument, or any later streamlining) is a substantial piece of real analysis. The following remark, which comes from Gleason’s paper [Gle57, p. 886], shows that the lower bound $\dim H \geq 3$ cannot be weakened to $\dim H \geq 2$. Remark 1.6 is not needed for our proofs and can be skipped without loss of continuity.



Remark 1.6. We show that the lower bound $\dim H \geq 3$ is optimal. On $H = \mathbb{R}^2$, parametrize rays by angle $\theta \in [0, \pi)$. Every orthonormal basis is a pair $\{\theta, \theta + \pi/2 \pmod{\pi}\}$. Define a function $v : [0, \pi) \rightarrow \{0, 1\}$ by

$$v(\theta) = \begin{cases} 1, & \text{if } \theta \in [0, \pi/2), \\ 0, & \text{if } \theta \in [\pi/2, \pi). \end{cases}$$

In every such pair, exactly one of the two angles lands in $[0, \pi/2)$, so v satisfies additivity on every basis; in other words, the function v is a genuine quantum measure in the sense of Definition 1.1, two-valued, on $L(\mathbb{R}^2)$ (with $v(0) = 0$ and $v(I) = 1$). The function v jumps at $\theta = \pi/2$, whereas $\theta \mapsto \langle \theta | \rho | \theta \rangle$ is continuous for every density operator ρ ; so v is not of the form $\text{Tr}(\rho \cdot)$, and the conclusion of Gleason’s Theorem fails when $\dim H = 2$. The same construction works for $H = \mathbb{C}^2$: in each pair of orthogonal rays, assign the value 1 to one ray and 0 to the other, and set $v(0) = 0$ and $v(I) = 1$.

The upper bound on the dimension is just as sharp, but for a reason that has nothing to do with geometry: in nonseparable Hilbert spaces, Gleason's Theorem depends on large cardinals.



Remark 1.7 (Gleason's Theorem and real-valued measurable cardinals). Let H be a Hilbert space with an orthonormal basis $\{e_i\}_{i \in I}$, and suppose that $|I|$ is at least a real-valued measurable cardinal, so that there is a countably additive probability measure ν on the power set of I that vanishes on singletons. Put

$$\mu(P) = \int_I \langle e_i | P | e_i \rangle d\nu(i) \quad (P \in L(H)),$$

which makes sense because ν measures every subset of I . We have $\mu(I) = 1$, and μ is countably additive on orthogonal families: if $P = \bigvee_k P_k$ with the P_k pairwise orthogonal, then $\langle e_i | P | e_i \rangle = \sum_k \langle e_i | P_k | e_i \rangle$ for every i , and the monotone convergence theorem lets us integrate term by term. However, $\mu(P_{e_i}) = \nu(\{i\}) = 0$ for every i , whereas a measure $\text{Tr}(\rho \cdot)$ satisfies $\sum_{i \in I} \text{Tr}(\rho P_{e_i}) = \text{Tr}(\rho) = 1$. So μ is a quantum measure that is not of the form $\text{Tr}(\rho \cdot)$. Dvurečenskij and Solovay proved that this is the only obstruction: a Hilbert space of dimension λ carries a quantum measure that is not of trace form if and only if some real-valued measurable cardinal is at most λ [Dvu93, Sol09]. Gleason's Theorem therefore holds verbatim for every Hilbert space whose dimension lies below the least real-valued measurable cardinal, in particular for every separable one. Theorem 1.4, on the other hand, holds in every dimension at least 3, with no set-theoretic hypothesis (see Remark 1.8 below).

If $|I|$ is measurable, then ν can be taken two-valued, that is, the indicator of a nonprincipal σ -complete ultrafilter $\mathcal{U}$ on I , and then $\mu(P) = \lim_{i, \mathcal{U}} \langle e_i | P | e_i \rangle$. Even so, the measure μ is not two-valued in general, because the numbers $\langle e_i | P | e_i \rangle$ need not tend to 0 or 1 along $\mathcal{U}$. By Ulam's theorem [Ula30], a real-valued measurable cardinal is either at most $2^{\aleph_0}$ or measurable. Solovay showed that adding κ random reals to a model with a measurable cardinal κ makes $2^{\aleph_0} = \kappa$ real-valued measurable [Sol71], while the constructible universe has no real-valued measurable cardinals at all. So, relative to the consistency of a measurable cardinal, Gleason's Theorem for Hilbert spaces of dimension $2^{\aleph_0}$ is independent of ZFC.



Remark 1.8 (Ultrafilters on bases, and the Main Theorem in every dimension). In this remark we drop separability, and we weaken the notion of a two-valued measure as far as possible: let $v : L(H) \rightarrow \{0, 1\}$ satisfy $v(I) = 1$ and $v(P \vee Q) = v(P) + v(Q)$ for orthogonal P, Q , with no countable additivity at all. Let $\{e_i\}_{i \in I}$ be an orthonormal basis of H . The projections onto the closed spans of the sets $\{e_j : j \in J\}$, for $J \subseteq I$, form a Boolean algebra isomorphic to the power set of I , and v restricts to the indicator of an ultrafilter $\mathcal{U}_v$ on I . If v is countably additive and I is countable, then $\mathcal{U}_v$ is a σ -complete ultrafilter on a countable set, hence principal, and its generator is the unique winner of Lemma 1.2: the lemma is the principality of σ -complete ultrafilters on countable sets. On an uncountable basis, the ultrafilter $\mathcal{U}_v$ may be nonprincipal; finite additivity permits any ultrafilter, and countable additivity permits any σ -complete one, which by Ulam's theorem is principal unless $|I|$ is at least a measurable cardinal.

Nevertheless, Theorem 1.4 survives in every dimension at least 3, even for such finitely additive v , by amplifying a finite witness. In finite dimension, finite additivity is all of additivity, and the theorem applies as stated. If $\dim H$ is infinite, write $H = \mathbb{C}^3 \otimes K$ for a Hilbert space K of the same dimension. By Corollary 3.2 below, there are finitely many rays $u_1, \dots, u_m$ of $\mathbb{C}^3$ and orthonormal bases $B_1, \dots, B_n$ drawn from them that admit no assignment taking the value 1 at exactly one ray of each basis. For each ray u , let W_u be the projection onto $u \otimes K$. If $u \perp w$, then $W_u \perp W_w$, and for each basis $\{a, b, c\}$ among

the B_k we have $W_a + W_b + W_c = I$. Finite additivity gives $v(W_a) + v(W_b) + v(W_c) = 1$, so $u \mapsto v(W_u)$ is an assignment that the configuration forbids. So, in dimension at least 3, the lattice $L(H)$ has no points in the strongest sense: it carries no two-valued measure, not even a finitely additive one, although each of its Boolean blocks carries as many ultrafilters as the power set of a basis does.

2. BELL'S 1964 PROOF OF THEOREM 1.4, BASED ON GLEASON'S THEOREM AND ELEMENTARY TOPOLOGY

In this section we present Bell's proof of Theorem 1.4, which combines Gleason's Theorem with the connectedness of the unit sphere. The Kochen–Specker proof, found independently in the same years, is given in Section 4; the history of both is traced in the Timeline (see Section 6 below).

Proof of Theorem 1.4, via Gleason's Theorem and connectedness. Suppose that v is a two-valued quantum measure on $L(H)$. By Gleason's Theorem, we have $v(P) = \text{Tr}(\rho P)$ for some density operator ρ , so $g(\psi) := v(P_\psi) = \langle \psi | \rho | \psi \rangle$ (see Observation 0.2) is continuous on $S(H)$ (a matrix element of the fixed bounded operator ρ) and $\{0, 1\}$ -valued. Since $S(H)$ is connected, a continuous $\{0, 1\}$ -valued function on $S(H)$ is constant; that is, there is $c \in \{0, 1\}$ with $g \equiv c$. But by Lemma 1.2, on any orthonormal basis g equals 1 at exactly one point and 0 at the rest — so g takes *both* values on that basis, contradicting $g \equiv c$. ☹

💡 **Remark 2.1.** In the preceding proof, the hypothesis $\dim H \geq 3$ is used only to invoke Gleason's Theorem; nothing else depends on dimension.

💡 **Remark 2.2 (Sfumato).** Bell's proof has a pictorial reading. Color each unit vector ψ by the value $\mu(P_\psi)$ of a quantum measure μ , from black (0) to white (1). A two-valued quantum measure would paint the sphere $S(H)$ in pure black and white, and by Lemma 1.2 both colors appear on every orthonormal basis, so the picture would have hard contours. When $\dim H \geq 3$, Gleason's Theorem allows only Leonardo's *sfumato*: every quantum measure has the form $\psi \mapsto \langle \psi | \rho | \psi \rangle$, a continuous function on the path-connected space $S(H)$, so between a white vector and a black vector every shade of gray occurs, by the intermediate value theorem. For the pure state $\rho = |u\rangle\langle u|$, the gradation away from u is the Born rule $\psi \mapsto |\langle u | \psi \rangle|^2$ itself. In dimension 2, where Gleason's Theorem fails, hard contours are possible (see Remark 1.6).

🕒 **Historical note.** Gleason proved his theorem in 1957 to answer a question of Mackey's about measures on closed subspaces [Mac57, Gle57], with no thought of hidden variables. In 1963 Jauch pointed out to Bell that the theorem bears on the hidden-variables question, and Bell turned this remark into the argument above; Bell thanks Jauch in a footnote of his paper [Bel66]. Bell wrote the paper in 1964, but the paper appeared only in 1966. The paper also criticizes two earlier impossibility arguments, von Neumann's of 1932 and Jauch and Piron's of 1963: each of them imposes a rule on the answers to questions that cannot be tested together, and nothing in physics justifies such a rule. The Timeline (see Section 6 below) tells the story in more detail.

🔗 **Provenance of the argument.** The proof of Theorem 1.4 given above is, in essential content, due to Bell. His own argument did not use the language of continuity or connectedness: he showed directly, using an auxiliary third orthogonal direction (available exactly because $\dim H \geq 3$), that two rays with v -value 1 and 0 cannot be arbitrarily close; since both values must occur on every basis (see Lemma 1.2), two such close, oppositely valued rays are unavoidable, a contradiction. What we have included here is the now-standard topological restatement of Bell's argument.

3. OBTAINING A FINITE OBSTRUCTION VIA LOGICAL COMPACTNESS

Bell's proof of Theorem 1.4 in Section 2 proves nonexistence over the whole uncountable sphere without ever exhibiting a finite culprit. In this section we show that a finite culprit was *guaranteed* to exist the moment Bell's proof was known (see Corollary 3.2 below). The tool is compactness, which appears here in its logical and its topological form at once.

Throughout this section, we assume that $\dim H = d$ with $3 \leq d < \infty$.

Let $V = S(H)/\sim$ be the set of rays (unit vectors modulo phase factors, where $u \sim e^{i\theta}u$ for $\theta \in \mathbb{R}$), and let $\mathcal{E}$ be the set of all orthonormal bases of H , each basis being a set of exactly d rays. Treat each ray $u \in V$ as a propositional symbol σ_u , and to each basis $e = \{u_1, \dots, u_d\} \in \mathcal{E}$ associate the propositional formula

$$\varphi_e := (\sigma_{u_1} \vee \dots \vee \sigma_{u_d}) \wedge \bigwedge_{1 \leq i < j \leq d} \neg(\sigma_{u_i} \wedge \sigma_{u_j}) \quad (\text{"exactly one } \sigma_{u_i} \text{ is true"}),$$

a formula in the d propositional symbols of e alone. The two-valued quantum measures on $L(H)$ correspond exactly to the truth assignments to $\{\sigma_u : u \in V\}$ that satisfy every φ_e . Indeed, by Lemma 1.2, a two-valued quantum measure v gives such an assignment, namely $\sigma_u \mapsto v(P_u)$. Conversely, let such an assignment be given. For $P \in L(H)$, let $v(P)$ be the number of true σ_u with u in an orthonormal basis of the range of P . This number does not depend on the choice of basis: extending two bases of the range of P by one fixed basis of the range of $P^\perp$ gives two bases of H , each with exactly one true ray. The same argument shows that $v(P) \in \{0, 1\}$, that $v(I) = 1$, and, using bases adapted to a family of pairwise orthogonal projections, that v is additive. So v is a two-valued quantum measure.

The topological picture. A truth assignment to the symbols σ_u is a point of the Cantor cube $2^V = \{0, 1\}^V$, which is compact by Tychonoff's theorem. For each basis $e \in \mathcal{E}$, the set

$$C_e = \{x \in 2^V : x \text{ satisfies } \varphi_e\}$$

depends only on the d coordinates indexed by the rays of e , so C_e is clopen. (In the language of Stone duality, the space 2^V is the Stone space of the free Boolean algebra on the symbols σ_u , and C_e is the clopen set of the formula φ_e .) By the preceding paragraph, the two-valued quantum measures on $L(H)$ correspond to the points of $\bigcap_{e \in \mathcal{E}} C_e$, and Theorem 1.4 says that this intersection is empty. Since 2^V is compact, the finite intersection property then produces a finite subfamily of $\{C_e : e \in \mathcal{E}\}$ with empty intersection. The compactness theorem of propositional logic, for the uncountable set of symbols $\{\sigma_u : u \in V\}$, is exactly this statement about 2^V (see, for example, [End01]). The next proposition and corollary record the argument.

Proposition 3.1. *If every finite subset of bases $\mathcal{E}_0 \subseteq \mathcal{E}$ admits some truth assignment to the finitely many rays occurring in $\mathcal{E}_0$ satisfying φ_e for every $e \in \mathcal{E}_0$, then a two-valued quantum measure exists on all of $L(H)$.*

Proof. Let $\mathcal{E}_0 \subseteq \mathcal{E}$ be finite. An assignment to the rays occurring in $\mathcal{E}_0$ that satisfies every φ_e with $e \in \mathcal{E}_0$ extends arbitrarily to a point of $\bigcap_{e \in \mathcal{E}_0} C_e$, so this intersection is nonempty. The sets C_e are closed in the compact space 2^V , so the finite intersection property gives a point $x \in \bigcap_{e \in \mathcal{E}_0} C_e$. By the correspondence above, the point x is a two-valued quantum measure on $L(H)$. 

Corollary 3.2. *If $3 \leq \dim H < \infty$, then there are a finite set of rays and a finite set of orthonormal bases drawn from them that admit no assignment satisfying "exactly one per basis" on every one of those bases.*

Proof. By Theorem 1.4, proved in Section 2 via Gleason’s Theorem, no two-valued quantum measure exists on $L(H)$. By Proposition 3.1, some finite set of bases $\mathcal{E}_0$, drawn from finitely many rays, admits no assignment satisfying φ_e for every $e \in \mathcal{E}_0$. ☺



Remark 3.3. Corollary 3.2 says that a finite Kochen–Specker configuration exists, and the corollary derives this from Section 2 without constructing one. The derivation is non-constructive in two ways. Compactness of 2^V for uncountable V is not provable in ZF; over ZF, this compactness is equivalent to the Boolean prime ideal theorem, a weak form of the Axiom of Choice. And the argument gives no bound on the size of the finite witness, nor any hint of where to find the witness.

The proof in the next section, in contrast, exhibits an explicit configuration of 18 vectors and uses no choice at all. Finding small explicit configurations took from 1967, when Kochen and Specker used 117 vectors, to 1996, when 18 vectors sufficed in dimension 4; in dimension 3 the smallest size is still unknown (see the [2024](#) entry of the Timeline in Section 6).

The appeal to choice is nevertheless harmless for the conclusion, though not for the proof. The statement of Corollary 3.2 asserts the existence of finitely many complex numbers satisfying finitely many polynomial equations, together with a finite combinatorial condition, so the statement is Σ_1^1 . Since ZFC proves the statement and ZF proves that the constructible universe L satisfies ZFC, the statement holds in L , and Σ_1^1 statements are upward absolute from L to the universe. Hence ZF proves Corollary 3.2. Choice is used by the compactness argument, but never by the theorem that the argument proves.

The topology also hides a trap. One might try to run the compactness argument over a countable dense set of rays, where compactness of 2^ω , which is provable in ZF, would suffice. In dimension 4 this happens to work, because the 18 vectors of Section 4 have coordinates in $\{0, \pm 1\}$. In dimension 3 it fails: Meyer [Mey99] showed that the rays through the rational points of the sphere S^2 admit an assignment that takes the value 1 at exactly one ray of every orthonormal basis made of such rays. So in dimension 3 every finite witness involves irrational directions, and a dense set of rays need not carry any obstruction at all. Density is no substitute for continuity: Bell’s argument uses continuity on the whole sphere, and the assignment of a dense set need not extend continuously.

The same picture, Boolean algebra by Boolean algebra. The lattice $L(H)$ is glued together from Boolean algebras: each family of commuting projections generates a Boolean subalgebra, and an orthonormal basis indexed by a set I generates one isomorphic to the power set of I , whose Stone space is βI . A finitely additive two-valued measure on $L(H)$ (in finite dimension, simply a two-valued quantum measure) is exactly a family of ultrafilters, one on each Boolean subalgebra, that agree on common elements. In the product of the Stone spaces of all Boolean subalgebras, a compact space, each agreement condition defines a clopen set, and the two-valued measures are the points of the intersection of these clopen sets. For a single Boolean algebra, or for a chain of Boolean algebras, such an intersection is never empty. Theorem 1.4 says that for $L(H)$ it is empty, and the finite intersection property says that finitely many agreement conditions already contradict one another, which is Corollary 3.2 once more. In infinite dimension, each Stone space βI has $2^{2^{|I|}}$ points, and all of them together still cannot be made to agree (see Remark 1.8).

4. A FINITE PROOF OF THEOREM 1.4, AFTER KOCHEN AND SPECKER

In this section we give a proof of Theorem 1.4 that uses no measure theory and no continuity. A finite configuration, which can be checked by hand, replaces Bell’s use of Gleason’s Theorem and connectedness, and supplies what both arguments need: a way to force the

separate “one winner per basis” choices of Lemma 1.2 to interact. We use the 18-vector configuration of Cabello, Estebaranz, and García-Alcaine [CEGA96], a descendant of Kochen and Specker’s original 117-vector configuration (see the historical note at the end of this section). One feature should be said at once: the configuration lives in $\mathbb{R}^4$. Kochen and Specker worked in $\mathbb{R}^3$, the smallest dimension in which the theorem holds, and paid for it with 117 vectors; Peres later needed 33 there (see the [1991](#) entry). In $\mathbb{R}^4$, eighteen vectors and a parity count suffice, and an embedding carries the argument to every dimension $d \geq 4$. For dimension 3, these notes rely on Bell’s proof.

Lemma 4.1 (Cabello–Estebaranz–García-Alcaine [CEGA96]). *After normalization, the following 9 lists of 4 pairwise orthogonal vectors are orthonormal bases of $\mathbb{R}^4$. They are built from 18 distinct vectors in total, each of which occurs in exactly 2 of the 9 bases.*

$$\begin{aligned}
B_1 &= \{(0, 0, 0, 1), (0, 0, 1, 0), (1, 1, 0, 0), (1, -1, 0, 0)\} \\
B_2 &= \{(0, 0, 0, 1), (0, 1, 0, 0), (1, 0, 1, 0), (1, 0, -1, 0)\} \\
B_3 &= \{(1, -1, 1, -1), (1, -1, -1, 1), (1, 1, 0, 0), (0, 0, 1, 1)\} \\
B_4 &= \{(1, -1, 1, -1), (1, 1, 1, 1), (1, 0, -1, 0), (0, 1, 0, -1)\} \\
B_5 &= \{(0, 0, 1, 0), (0, 1, 0, 0), (1, 0, 0, 1), (1, 0, 0, -1)\} \\
B_6 &= \{(1, -1, -1, 1), (1, 1, 1, 1), (1, 0, 0, -1), (0, 1, -1, 0)\} \\
B_7 &= \{(1, 1, -1, 1), (1, 1, 1, -1), (1, -1, 0, 0), (0, 0, 1, 1)\} \\
B_8 &= \{(1, 1, -1, 1), (-1, 1, 1, 1), (1, 0, 1, 0), (0, 1, 0, -1)\} \\
B_9 &= \{(1, 1, 1, -1), (-1, 1, 1, 1), (1, 0, 0, 1), (0, 1, -1, 0)\}.
\end{aligned}$$

Verification. Orthogonality within each B_k is a check of $\binom{4}{2} = 6$ dot products; for instance, for B_3 we have $(1, -1, 1, -1) \cdot (1, -1, -1, 1) = 1 + 1 - 1 - 1 = 0$, and the other five products in B_3 vanish in the same way. Table 1 lists, for each of the 18 vectors, the two bases that contain the vector. Since $9 \text{ bases} \times 4 \text{ vectors} = 36 = 18 \times 2$, the table accounts for every entry of every basis. ☺

vector	bases	vector	bases	vector	bases
$(0, 0, 0, 1)$	B_1, B_2	$(1, 0, -1, 0)$	B_2, B_4	$(1, 0, 0, 1)$	B_5, B_9
$(0, 0, 1, 0)$	B_1, B_5	$(1, -1, 1, -1)$	B_3, B_4	$(1, 0, 0, -1)$	B_5, B_6
$(1, 1, 0, 0)$	B_1, B_3	$(1, -1, -1, 1)$	B_3, B_6	$(0, 1, -1, 0)$	B_6, B_9
$(1, -1, 0, 0)$	B_1, B_7	$(0, 0, 1, 1)$	B_3, B_7	$(1, 1, -1, 1)$	B_7, B_8
$(0, 1, 0, 0)$	B_2, B_5	$(1, 1, 1, 1)$	B_4, B_6	$(1, 1, 1, -1)$	B_7, B_9
$(1, 0, 1, 0)$	B_2, B_8	$(0, 1, 0, -1)$	B_4, B_8	$(-1, 1, 1, 1)$	B_8, B_9

TABLE 1. The 18 vectors of Lemma 4.1, each with the two bases that contain the vector.

Table 1 can be read as a graph: the vertices are the 9 bases, and each vector is an edge joining the two bases that contain the vector (see Figure 1). Each vertex has degree 4, because each basis has 4 vectors. A two-valued quantum measure would select, at every vertex, exactly one of the 4 edges, and an edge selected at one endpoint is selected at the other endpoint too, since the edge is a single ray. So the selected edges would form a perfect matching of a graph with 9 vertices, which is impossible, because a perfect matching pairs off the vertices and 9 is odd. The proof below writes this parity argument as a double count.




FIGURE 1. The 18 vectors as the edges of a 4-regular graph on the 9 bases. A two-valued quantum measure would be a perfect matching of this graph, and no graph with an odd number of vertices has one: a maximal matching (pink) always leaves a vertex uncovered (lavender).

Proof of Theorem 1.4 for $H = \mathbb{R}^4$. Suppose that v is a two-valued quantum measure on $L(\mathbb{R}^4)$. Restrict attention to the 18 rays of Lemma 4.1. Since each B_k is an orthonormal basis, additivity gives

$$\sum_{u \in B_k} v(u) = v(I) = 1 \quad \text{for each } k = 1, \dots, 9.$$

Summing over all 9 bases, we have

$$\sum_{k=1}^9 \sum_{u \in B_k} v(u) = 9.$$

We now reorganize the left side by vector instead of by basis. Each of the 18 vectors u occurs in exactly 2 of the bases B_k , so u contributes $2v(u)$ to the double sum. Hence, we have

$$\sum_{k=1}^9 \sum_{u \in B_k} v(u) = \sum_u 2v(u) = 2 \sum_u v(u),$$

which is an even integer. This contradicts the fact that the double sum equals the odd integer 9. 😊

Which dimensions the argument covers. The same argument proves Theorem 1.4 for every real or complex Hilbert space H with $\dim H \geq 4$. Indeed, suppose that v is a two-valued quantum measure on $L(H)$. By Lemma 1.2, some unit vector e has $v(P_e) = 1$. Choose a 4-dimensional subspace K of H containing e , with projection P_K ; then $v(P_K) = 1$, because $P_e \leq P_K$. Fix an orthonormal basis $f_1, \dots, f_4$ of K . The map $(x_1, \dots, x_4) \mapsto \sum_i x_i f_i$ sends orthogonal vectors of $\mathbb{R}^4$ to orthogonal vectors of K , so the 9 bases of Lemma 4.1 become 9 orthonormal bases of K with the same incidence pattern. For each of these bases, the 4 projections add up to P_K , so their values under v add up to $v(P_K) = 1$, and the double count above gives the same contradiction. Dimension 3, as announced at the beginning of the section, requires a configuration in $\mathbb{R}^3$ or Bell's argument.




FIGURE 2. The graph Γ_2 of Kochen and Specker's 1967 paper [KS67, p. 69], whose 117 vertices are the vectors of their original proof; two vertices are joined by an edge when the corresponding vectors are orthogonal. The vertices are the points on the rim, after identifying p_0 with a , q_0 with b , and r_0 with c ; each of the fifteen small gadgets on the rim is a copy of their ten-vertex graph Γ_1 .

✂ **Historical note.** Kochen and Specker's 1967 proof [KS67] uses 117 vectors in $\mathbb{R}^3$ (see Figure 2), and so the proof covers dimension 3 as well.

Both authors were logicians. Ernst Specker (1920–2011), at ETH Zürich, had shown in 1953 that the Axiom of Choice fails in Quine's New Foundations, and he had announced the present result without proof in 1960 [Spe60]. Simon Kochen (b. 1934) had written his Princeton thesis under Alonzo Church on ultraproducts, and was working with James Ax on the papers that became the Ax–Kochen theorem. Their formulation of the problem is accordingly logical. The questions that can be tested together form a Boolean algebra; these Boolean algebras, glued along their common parts, form a *partial Boolean algebra*; and the question is whether the whole structure admits a homomorphism to the two-element Boolean algebra. The 18-vector configuration used above was found in 1996 [CEGA96], after reductions by Peres and by Kernaghan (see the [1991](#), [1994](#), and [1996](#) entries of the Timeline).

5. QUANTUM MEASURES HAVE NO COUNTERPART OF DIRAC MEASURES AND DO NOT FORM A LATTICE

We have now proved the Main Theorem twice, once by continuity and once by counting. This section asks what the theorem says about the structure of quantum events, and the answer, read through ultrafilters, is that the quantum lattice has no points. The answer was already in view at the end of Section 3, where a two-valued measure appeared as a family of ultrafilters, one on each Boolean subalgebra of $L(H)$, that agree on common elements; in this section that picture becomes the starting point.

Classical probability rests on two facts so familiar that they are easy to overlook. The first is that a single outcome decides every event: the events that contain a point ω of the sample space form a principal ultrafilter, and the Dirac measure δ_ω is nothing but the indicator function of that ultrafilter. The second is that measures can always be compared and combined: any two finite measures on the same σ -algebra have a greatest lower bound and a least upper bound, so the measures form a lattice. This section shows that, once $\dim H \geq 3$, quantum measures have neither property. The first failure is the Bell–Kochen–Specker theorem in another guise. The second is of a different nature altogether, since it occurs already in dimension 2, where two-valued measures exist. Throughout the section, H is a Hilbert space with $\dim H \geq 2$, and none of the proofs uses Theorem 1.4.

The classical properties (C1) and (C2). Let Σ be a σ -algebra of subsets of a sample space Ω . The two properties read as follows.

- (C1) Each outcome decides the Boolean operations on events. In terms of the Dirac measures, for every $\omega \in \Omega$ and all $A, B \in \Sigma$, we have

$$\delta_\omega(A \cap B) = \min(\delta_\omega(A), \delta_\omega(B)), \quad \delta_\omega(\Omega \setminus A) = 1 - \delta_\omega(A). \quad (4)$$

(Each side of each equation equals 1 if ω belongs to the event on the left-hand side, and 0 otherwise.)

- (C2) Let $\mathcal{M}(\Sigma)$ be the set of finite positive measures on Σ , ordered by declaring $\mu \leq \nu$ if $\mu(A) \leq \nu(A)$ for every $A \in \Sigma$. Then $(\mathcal{M}(\Sigma), \leq)$ is a lattice (see Proposition 5.10 below).

The quantum questions (Q1) and (Q2). To ask whether (C1) and (C2) survive in quantum mechanics, we replace Σ by the lattice $L(H)$ and the probability measures by the quantum measures of Definition 1.1; for a density operator ρ we write $\mu_\rho(P) = \text{Tr}(\rho P)$. Each classical property then becomes a question.

- (Q1) For a quantum measure μ on $L(H)$, the natural counterpart of the first equation in (4) is

$$\mu(P \wedge Q) = \min(\mu(P), \mu(Q)). \quad (5)$$

Is there a quantum measure μ on $L(H)$ that satisfies (5) whenever $P, Q \in L(H)$ commute?

The second equation in (4) needs no counterpart, since additivity already gives $\mu(P^\perp) = 1 - \mu(P)$. The restriction to commuting pairs is the fair one: commuting projections lie in a common Boolean subalgebra of $L(H)$, and Boolean algebras are where the classical equations (4) live. (If P and Q commute, the projections PQ , $P(I - Q)$, $(I - P)Q$ and $(I - P)(I - Q)$ are pairwise orthogonal with sum I ; the joins of subsets of these four projections form a Boolean subalgebra of $L(H)$ containing $P = PQ + P(I - Q)$ and $Q = PQ + (I - P)Q$.) Without the restriction the question would be too easy, for no quantum measure satisfies (5) for all pairs $P, Q \in L(H)$ (Proposition 5.2).

- (Q2) For the counterpart of (C2), let $\mathcal{C}(H)$ be the set of functions $P \mapsto \text{Tr}(TP)$ on $L(H)$, where T ranges over the positive trace-class operators on H . When $\dim H \geq 3$, Gleason's Theorem identifies $\mathcal{C}(H)$ with the set of nonnegative multiples of quantum measures. As in (C2), we allow all nonnegative multiples and not only the normalized measures, because a greatest lower bound of two probability measures is in general not a probability measure: the greatest lower bound of two distinct Dirac measures is 0. We order $\mathcal{C}(H)$ pointwise:

$$\mu \leq \nu \quad \text{if } \mu(P) \leq \nu(P) \text{ for every } P \in L(H).$$

Is $(\mathcal{C}(H), \leq)$ a lattice?

The main result of this section.

- When $\dim H \geq 3$, the answer to (Q1) is negative (see Theorem 5.6 below); when $\dim H = 2$, the answer is positive (see Remark 1.6).
- When $\dim H \geq 2$, the answer to (Q2) is negative (see Proposition 5.11 below).

Thus, when $\dim H \geq 3$, neither (C1) nor (C2) has a quantum counterpart.

What the answers mean. The negative answer to (Q1) is the Bell–Kochen–Specker theorem in disguise. Lemma 5.1 shows that a quantum measure satisfies (5) for all commuting P, Q exactly when it is two-valued, that is, exactly when it behaves like a Dirac measure. For $\dim H \geq 3$, the negative answer to (Q1) therefore says that there is no two-valued quantum measure on $L(H)$, which is Theorem 1.4, or, in the language of Section 1, that there is no non-contextual hidden-variable assignment. The measures that physics actually uses fare even worse: in every dimension, no measure μ_ρ given by a density operator satisfies (5) for all commuting pairs (Corollary 5.7).

The negative answer to (Q2) says that quantum measures are not ordered like classical ones. Recall that an *order isomorphism* between partially ordered sets $(X, \leq)$ and $(Y, \leq)$ is a bijection $\Phi : X \rightarrow Y$ such that, for all $x, x' \in X$, we have $x \leq x'$ if and only if $\Phi(x) \leq \Phi(x')$. Whatever the σ -algebra Σ , there is no order isomorphism between $(\mathcal{C}(H), \leq)$ and $(\mathcal{M}(\Sigma), \leq)$ (Corollary 5.12). Since this is so already in dimension 2, where two-valued quantum measures exist (Remark 1.6), the failure of (C2) is a phenomenon separate from the Bell–Kochen–Specker theorem.

Ultrafilters, principal and otherwise. The answer to (Q1) is best read in the language of ultrafilters. On a Boolean algebra $\mathcal{B}$, a finitely additive two-valued measure v is the indicator function of the ultrafilter $v^{-1}(1)$, and every ultrafilter arises in this way, so the two-valued measures on $\mathcal{B}$ are the points of the Stone space of $\mathcal{B}$. On the power set of a set Ω , the Dirac measures are exactly the indicators of the principal ultrafilters. (Countable additivity is a strong demand: a countably additive two-valued measure on the power set of Ω that vanishes on singletons is a nonprincipal σ -complete ultrafilter, and such ultrafilters exist only if $|\Omega|$ is at least a measurable cardinal [Ula30]; compare Remark 1.7.)

In $L(H)$, a two-valued quantum measure v defines the set $\mathcal{F}_v = \{P \in L(H) : v(P) = 1\}$. This set is upward closed, contains exactly one of P and $P^\perp$ for each P , and, by Lemma 5.1, is closed under meets of commuting pairs. These are the defining properties of an ultrafilter, read in the only Boolean algebras that $L(H)$ contains, namely those generated by commuting families. So (Q1) asks whether $L(H)$ has ultrafilters, and the Main Theorem says that, when $\dim H \geq 3$, it has none: the quantum lattice has no points. Its atoms do generate principal

filters,

$$\mathcal{F}_\psi = \{P \in L(H) : P \geq P_\psi\} = \{P \in L(H) : \psi \in \text{ran } P\} \quad (\psi \in S(H)),$$

which are even closed under arbitrary meets. But $\mathcal{F}_\psi$ contains P or $P^\perp$ only when ψ is an eigenvector of P , so $\mathcal{F}_\psi$ is not an ultrafilter: the failure of the principal filters to be ultrafilters is the superposition principle in lattice-theoretic dress. What remains of $\mathcal{F}_\psi$ is the pure state $\mu_\psi(P) = \langle \psi | P | \psi \rangle$. We have $\mu_\psi(P) = \|P\psi\|^2 = 1$ if and only if $P\psi = \psi$, that is, if and only if $P \in \mathcal{F}_\psi$, and on the questions that $\mathcal{F}_\psi$ leaves undecided, the pure state takes values strictly between 0 and 1.

The comparison also has a topological side. The Stone space of a Boolean algebra is compact and zero-dimensional, and its clopen sets are the events. The natural space of quantum points, the space of rays, is connected (Proposition 5.3), and so it has no nontrivial clopen sets at all. Section 5.2 turns this contrast into a proof.

The plan of the section. We treat (Q1) first and (Q2) second, and in each case the classical property comes right before its quantum failure. Property (C1) needs no proof beyond the parenthetical remark after (4). Section 5.1 proves Lemma 5.1, which turns (Q1) into a question about two-valued measures, and shows that the restriction to commuting pairs cannot be dropped. Section 5.2 answers (Q1) by playing sliding against jumping: a unit vector can be moved continuously around the unit sphere, Gleason's Theorem forces every quantum measure to follow it continuously, and a two-valued measure can only jump between 0 and 1. In the same section, Remark 5.8 uses the compactness argument of Section 3 to show that finitely many projections in a single 3-dimensional subspace already force the negative answer to (Q1). Section 5.3 then proves (C2) by the Hahn decomposition, and Section 5.4 answers (Q2) with an explicit pair of commuting 2×2 matrices for which the Hahn decomposition argument breaks down.

5.1. Quantum measures that preserve commuting meets. We begin with (Q1), and first show that the condition in (Q1) is far more restrictive than it looks: a quantum measure that satisfies it takes only the values 0 and 1, just as a Dirac measure does. So (Q1) asks exactly whether two-valued quantum measures exist. It is convenient to say that a quantum measure μ on $L(H)$ *preserves commuting meets* if μ satisfies (5) whenever $P, Q \in L(H)$ commute.

Lemma 5.1. *Let μ be a quantum measure on $L(H)$. Then μ preserves commuting meets if and only if μ is two-valued.*

Proof. Additivity gives $\mu(0) = 0$ and $\mu(P^\perp) = 1 - \mu(P)$ for every P . Suppose that μ preserves commuting meets. For every P , the projections P and $P^\perp$ commute and $P \wedge P^\perp = 0$. Applying (5) to the pair $P, P^\perp$ gives

$$\min(\mu(P), 1 - \mu(P)) = \min(\mu(P), \mu(P^\perp)) = \mu(P \wedge P^\perp) = \mu(0) = 0.$$

Since $\min(r, 1 - r) > 0$ for every $r \in (0, 1)$, this forces $\mu(P) \in \{0, 1\}$.

Conversely, suppose that μ is two-valued, and let $P, Q \in L(H)$ be commuting projections. We first recall why $PQ = P \wedge Q$. Since P and Q commute, the operator PQ is self-adjoint and $(PQ)^2 = PQ$, so PQ is a projection. Its range lies in the ranges of both P and Q (because $PQ = QP$), and PQ fixes every vector lying in both ranges. Hence $PQ = P \wedge Q$.

Next, we show that $\mu(P \wedge Q) \leq \min(\mu(P), \mu(Q))$. The projection $P = PQ + P(I - Q)$ is a sum of two orthogonal projections, so additivity gives $\mu(P) = \mu(P \wedge Q) + \mu(P(I - Q)) \geq \mu(P \wedge Q)$. By symmetry, we also have $\mu(Q) \geq \mu(P \wedge Q)$.

Since μ takes only the values 0 and 1, it remains to be shown that $\mu(P \wedge Q) = 1$ whenever $\mu(P) = \mu(Q) = 1$. The projections P and $(I - P)Q$ are orthogonal, so their sum is a projection and additivity gives $\mu(P) + \mu((I - P)Q) \leq 1$. Since $Q = PQ + (I - P)Q$ is also a sum of two orthogonal projections, we have $\mu((I - P)Q) = \mu(Q) - \mu(P \wedge Q)$. Combining these two facts, we have

$$\mu(P) + \mu(Q) - \mu(P \wedge Q) \leq 1.$$

If $\mu(P) = \mu(Q) = 1$, this inequality gives $\mu(P \wedge Q) \geq 1$, and therefore $\mu(P \wedge Q) = 1$. $\odot$

Since the first equation in (4) holds for every pair of events, one might hope to drop the commutativity assumption in (Q1). The next proposition shows that, without this assumption, (Q1) has an immediate negative answer.

Proposition 5.2. *No quantum measure μ on $L(H)$ satisfies $\mu(P \wedge Q) = \min(\mu(P), \mu(Q))$ for all $P, Q \in L(H)$.*

Proof. Suppose that μ is such a quantum measure. In particular, the measure μ preserves commuting meets, so μ is two-valued by Lemma 5.1. Let $\{e_i\}$ be an orthonormal basis of H . Additivity gives $\sum_i \mu(P_{e_i}) = 1$, so $\mu(P_e) = 1$ for some $e \in S(H)$. Let K be a 2-dimensional subspace containing e , and let P_K be the projection onto K . Since $P_K \geq P_e$, we have $\mu(P_K) = 1$. For every orthonormal basis $\{a, b\}$ of K we have $P_a + P_b = P_K$, so exactly one of $\mu(P_a), \mu(P_b)$ equals 1. Fix an orthonormal basis $\{a, b\}$ of K and let $c = (a + b)/\sqrt{2}$ and $d = (a - b)/\sqrt{2}$, so that $\{c, d\}$ is an orthonormal basis of K with no ray in common with $\{a, b\}$. Interchanging a and b , and c and d , if necessary, we may assume that $\mu(P_a) = \mu(P_c) = 1$. Then $P_a \neq P_c$. Two distinct rays meet in 0, so $\mu(P_a \wedge P_c) = \mu(0) = 0$, whereas $\min(\mu(P_a), \mu(P_c)) = 1$. $\odot$

5.2. Topology and the negative answer to (Q1). We can now answer (Q1) when $\dim H \geq 3$. By Lemma 5.1, it suffices to show that no two-valued quantum measure exists, and the argument plays continuity against jumps. Gleason's Theorem makes every quantum measure continuous on rays, whereas a two-valued quantum measure, which is not constant on rays, cannot be continuous on the connected unit sphere.

We give $L(H)$ the operator-norm topology, given by the metric $d(P, Q) = \|P - Q\|$. As in Section 0, for $\psi \in S(H)$ we write $P_\psi = |\psi\rangle\langle\psi|$ for the projection onto $\mathbb{C}\psi$, and we write $\mu_\psi = \mu_{P_\psi}$ for the corresponding pure state, so that $\mu_\psi(P) = \langle\psi|P|\psi\rangle$ (see Observation 0.2). The next proposition collects the three facts that we need: rank-one projections are close in norm exactly when they come from nearly parallel unit vectors, the rank-one projections form a connected space, and the number $\mu_\rho(P)$ depends continuously on both the density operator ρ and the projection P .

Proposition 5.3.

- (1) For $\psi, \varphi \in S(H)$ we have $\|P_\psi - P_\varphi\| = (1 - |\langle\psi|\varphi\rangle|^2)^{1/2}$ and $\|P_\psi - P_\varphi\|_1 = 2(1 - |\langle\psi|\varphi\rangle|^2)^{1/2}$.
- (2) The set $\{P_\psi : \psi \in S(H)\}$ of rank-one projections is a connected subset of $L(H)$.
- (3) For density operators ρ, σ and $P, Q \in L(H)$ we have $|\mu_\rho(P) - \mu_\sigma(Q)| \leq \|\rho - \sigma\|_1 + \|P - Q\|$.

Proof. (1) The operator $X = P_\psi - P_\varphi$ is self-adjoint, has trace 0, and has range in $\text{span}(\psi, \varphi)$. So the nonzero eigenvalues of X are $\pm t$ for some $t \geq 0$, and, by Observation 0.2, we have $2t^2 = \text{Tr } X^2 = \text{Tr } P_\psi + \text{Tr } P_\varphi - 2\text{Tr}(P_\psi P_\varphi) = 2 - 2|\langle\psi|\varphi\rangle|^2$. Hence $t = (1 - |\langle\psi|\varphi\rangle|^2)^{1/2}$, and therefore $\|X\| = t$ and $\|X\|_1 = 2t$.

(2) By (1), we have $\|P_\psi - P_\varphi\| = (1 - |\langle\psi|\varphi\rangle|^2)^{1/2} \leq (2\|\psi - \varphi\|)^{1/2}$, because $1 - |\langle\psi|\varphi\rangle|^2 \leq 2(1 - \operatorname{Re}\langle\psi|\varphi\rangle) = 2\operatorname{Re}\langle\psi|\psi - \varphi\rangle \leq 2\|\psi - \varphi\|$. So the map $\psi \mapsto P_\psi$ is continuous, and the set of rank-one projections is the continuous image of the connected space $S(H)$.

(3) By the triangle inequality, we have $|\operatorname{Tr}(\rho P) - \operatorname{Tr}(\sigma Q)| \leq |\operatorname{Tr}((\rho - \sigma)P)| + |\operatorname{Tr}(\sigma(P - Q))| \leq \|\rho - \sigma\|_1 \|P\| + \|\sigma\|_1 \|P - Q\|$, and moreover $\|P\| \leq 1$ and $\|\sigma\|_1 = 1$. $\odot$

A first contrast with the classical case. Before proving the main theorem of this subsection, we compare the pure states μ_ψ with their classical counterparts, the Dirac measures. Let Ω be a finite set with the discrete topology, and let Σ be the set of all subsets of Ω . For every $A \in \Sigma$, the function $\omega \mapsto \delta_\omega(A)$ takes only the values 0 and 1, and the set $\{\omega : \delta_\omega(A) = 1\} = A$ is both open and closed in Ω . The next proposition shows that both properties fail for the measures μ_ψ on $L(H)$.

Proposition 5.4. *Let $P \in L(H)$ be a projection with $P \neq 0$ and $P \neq I$. The function $\psi \mapsto \mu_\psi(P)$ on $S(H)$ is continuous and takes every value in $[0, 1]$. The set $\{\psi \in S(H) : \mu_\psi(P) = 1\}$ is closed, nonempty, and not open.*

Proof. Let $f(\psi) = \mu_\psi(P) = \langle\psi|P|\psi\rangle$ for $\psi \in S(H)$. The function f is continuous, equal to 1 at the unit vectors in the range of P and to 0 at those in the range of $P^\perp$. By the intermediate value theorem along a path in $S(H)$ joining two such vectors, exactly as in Remark 2.2, the function f takes every value in $[0, 1]$. The set $f^{-1}(1)$ is closed by continuity, nonempty because $P \neq 0$, and proper because $P \neq I$. If $f^{-1}(1)$ were also open, then $S(H)$ would be disconnected. $\odot$

Proposition 5.4 fixes a projection P and lets the measure μ_ψ vary. We now reverse the roles: we fix a measure and let the projection vary. The quantum measures that, like Dirac measures, take only the values 0 and 1 are the two-valued ones, and the next proposition shows that none of them varies continuously with the ray.

Proposition 5.5. *If v is a two-valued quantum measure on $L(H)$, then $\psi \mapsto v(P_\psi)$ is not continuous on $S(H)$.*

Proof. Bell's proof of Theorem 1.4 in Section 2 uses Gleason's Theorem only to show that $\psi \mapsto v(P_\psi)$ is continuous on $S(H)$. If the continuity of $\psi \mapsto v(P_\psi)$ is assumed instead, the rest of Bell's proof applies verbatim and yields a contradiction, using only Lemma 1.2 and the connectedness of $S(H)$; in particular, this part of the argument does not require $\dim H \geq 3$. $\odot$

Gleason's Theorem supplies the continuity that Proposition 5.5 forbids, and (Q1) is answered.

Theorem 5.6. *A quantum measure on $L(H)$ that preserves commuting meets exists if $\dim H = 2$, and does not exist if $\dim H \geq 3$.*

Proof. Suppose that $\dim H \geq 3$, and let μ be a quantum measure that preserves commuting meets. By Lemma 5.1, the measure μ is two-valued. By Gleason's Theorem, we have $\mu = \mu_\rho$ for a density operator ρ , so $\psi \mapsto \mu(P_\psi) = \langle\psi|\rho|\psi\rangle$ is continuous on $S(H)$. This contradicts Proposition 5.5.

If $\dim H = 2$, then, as in Remark 1.6, assign the value 1 to one ray in each pair of orthogonal rays and the value 0 to the other, and set $v(0) = 0$ and $v(I) = 1$. This defines a two-valued quantum measure v , so v preserves commuting meets by Lemma 5.1. $\odot$

The second half of this proof shows what changes in dimension 2. There, Gleason's Theorem fails (see Remark 1.6), and the answer to (Q1) is positive; but by Lemma 5.1 and

Proposition 5.5, every quantum measure that gives this positive answer is two-valued and therefore discontinuous on $S(H)$. The measures μ_ρ used in physics (see Remark 0.1) are continuous, so in every dimension none of them satisfies the condition in (Q1).

Corollary 5.7. *For every density operator ρ , the quantum measure μ_ρ does not preserve commuting meets.*

Proof. If μ_ρ preserved commuting meets, Lemma 5.1 would make μ_ρ two-valued. Since the function $\psi \mapsto \mu_\rho(P_\psi) = \langle \psi | \rho | \psi \rangle$ is continuous, this would contradict Proposition 5.5. $\odot$



Remark 5.8 (A finite witness). The proof of Theorem 5.6 applies Gleason's Theorem to the whole of $L(H)$, an infinite object, but finitely many projections in a single 3-dimensional subspace already force the negative answer to (Q1). By Corollary 3.2 with $d = 3$, there is a finite family $\mathcal{E}_0$ of orthonormal bases of $\mathbb{C}^3$ such that no function from the rays occurring in $\mathcal{E}_0$ to $\{0, 1\}$ takes the value 1 at exactly one ray of each basis, and a unitary map places $\mathcal{E}_0$ in any 3-dimensional space. Now let $3 \leq \dim H \leq \infty$, and let v be a quantum measure on $L(H)$ that preserves commuting meets. By Lemma 5.1, the measure v is two-valued, and by Lemma 1.2 some unit vector e has $v(P_e) = 1$. Let K be a 3-dimensional subspace containing e , so that $v(P_K) = 1$, and place $\mathcal{E}_0$ in K . For each basis $\{u_1, u_2, u_3\}$ in $\mathcal{E}_0$ we have $P_{u_1} + P_{u_2} + P_{u_3} = P_K$, so exactly one of the three rays has value 1, contrary to the choice of $\mathcal{E}_0$. The negative answer to (Q1) is thus witnessed by finitely many commuting triples of rank-one projections in one 3-dimensional subspace, just as the witnesses in Propositions 5.2 and 5.11 (the rays P_a, P_c , and the matrices M, τ) lie in a single 2-dimensional subspace.

So far, the obstruction to (Q1) has come from the continuity of the measures. A second obstruction comes from the meet operation itself. For a Dirac measure δ_ω , the first equation in (4) says that the number $\delta_\omega(A \cap B)$ is computed from $\delta_\omega(A)$ and $\delta_\omega(B)$ by a fixed continuous function, namely $\min$. No measure μ_ρ given by a density operator admits any continuous function of this kind (Proposition 5.9), and the reason can be seen in a picture: the meet is not continuous for the norm topology. Two distinct lines through the origin meet only at 0, however small the angle between them, whereas a line meets itself in the whole line (see Figure 3).




FIGURE 3. Rank-one projections P_n converging to a rank-one projection P , drawn through their ranges. For every n , the meet $P_n \wedge P$ is 0 and the join $P_n \vee P$ is the projection onto the plane spanned by the two lines; in the limit, both the meet and the join of P with itself are P .

Proposition 5.9. *Let P_n and P be rank-one projections with $P_n \neq P$ and $P_n \rightarrow P$ in $L(H)$. Then $P_n \wedge P = 0$ and $\|P_n \vee P - P\| = 1$ for every n , while $P \wedge P = P \vee P = P$. Moreover, for every density operator ρ there is no continuous $m : [0, 1]^2 \rightarrow [0, 1]$ with $\mu_\rho(Q \wedge R) = m(\mu_\rho(Q), \mu_\rho(R))$ for all $Q, R \in L(H)$.*

Proof. Since $P_n \neq P$, the ranges of P_n and P are distinct lines, so $P_n \wedge P = 0$. The join $P_n \vee P$ is the projection onto a 2-dimensional space containing the range of P , so $P_n \vee P - P$ is a rank-one projection and has norm 1.

Now let ρ be a density operator, and suppose that $m : [0, 1]^2 \rightarrow [0, 1]$ is continuous and satisfies $\mu_\rho(Q \wedge R) = m(\mu_\rho(Q), \mu_\rho(R))$ for all $Q, R \in L(H)$. Let $\psi \in S(H)$ be given, choose a unit vector $\varphi \perp \psi$, and put $Q_n = P_{\psi_n}$ with $\psi_n = (\cos \frac{1}{n})\psi + (\sin \frac{1}{n})\varphi$. Then $Q_n \neq P_\psi$ and $Q_n \rightarrow P_\psi$ by Proposition 5.3(1), so $Q_n \wedge P_\psi = 0$ by the first part of this proof, and $\mu_\rho(Q_n) \rightarrow \mu_\rho(P_\psi)$ by Proposition 5.3(3). Hence

$$0 = \mu_\rho(Q_n \wedge P_\psi) = m(\mu_\rho(Q_n), \mu_\rho(P_\psi)) \rightarrow m(\mu_\rho(P_\psi), \mu_\rho(P_\psi)) = \mu_\rho(P_\psi \wedge P_\psi) = \mu_\rho(P_\psi).$$

So μ_ρ vanishes on every rank-one projection. Summing over an orthonormal basis gives $\mu_\rho(I) = 0$, a contradiction. ☹

5.3. Classical measures form a lattice. We now turn to (Q2). Before answering it, we recall why its classical counterpart (C2) holds, since the quantum counterexample of Section 5.4 is built to defeat exactly this argument. Property (C2) holds for a simple reason. Given $\mu, \nu \in \mathcal{M}(\Sigma)$, split Ω into a part where $\mu \geq \nu$ and a part where $\mu \leq \nu$, and take the smaller of the two measures on each part. The Hahn decomposition theorem provides the split, and the resulting measure is the greatest lower bound of μ and ν .

A *finite signed measure* on Σ is a countably additive function $\lambda : \Sigma \rightarrow \mathbb{R}$, for example $\mu - \nu$. Given a finite signed measure λ , a *Hahn set* for λ is a set $S \in \Sigma$ such that $\lambda(B) \geq 0$ for every measurable $B \subseteq S$ and $\lambda(B) \leq 0$ for every measurable $B \subseteq \Omega \setminus S$. The *Hahn decomposition theorem* says that every finite signed measure has a Hahn set (see, e.g., [Rud87, Theorem 6.14]).

Given $\mu, \nu \in \mathcal{M}(\Sigma)$ and a Hahn set S for $\mu - \nu$, define measures $\gamma, \eta \in \mathcal{M}(\Sigma)$ by

$$\gamma(A) = \nu(A \cap S) + \mu(A \setminus S), \quad \eta(A) = \mu(A \cap S) + \nu(A \setminus S) \quad (A \in \Sigma). \quad (6)$$

Proposition 5.10. *Let $\mu, \nu \in \mathcal{M}(\Sigma)$ be given, and let S be a Hahn set for $\mu - \nu$. Then the measures γ and η defined in (6) are, respectively, the greatest lower bound and the least upper bound of μ and ν in $(\mathcal{M}(\Sigma), \leq)$.*

Proof. Since S is a Hahn set for $\mu - \nu$, we have $\mu(B) \geq \nu(B)$ for every measurable $B \subseteq S$ and $\mu(B) \leq \nu(B)$ for every measurable $B \subseteq \Omega \setminus S$. Hence, for every $A \in \Sigma$, we have $\nu(A \cap S) \leq \mu(A \cap S)$ and $\mu(A \setminus S) \leq \nu(A \setminus S)$, so $\gamma(A) \leq \mu(A)$ and $\gamma(A) \leq \nu(A)$; in other words, the measure γ is a lower bound of μ and ν . If $\kappa \in \mathcal{M}(\Sigma)$ is another lower bound, then for every $A \in \Sigma$ we have $\kappa(A) = \kappa(A \cap S) + \kappa(A \setminus S) \leq \nu(A \cap S) + \mu(A \setminus S) = \gamma(A)$. The proof that η is the least upper bound is the same, with the inequalities reversed. ☹

5.4. The negative answer to (Q2). We now show that the answer to (Q2) is negative as soon as $\dim H \geq 2$. The first step translates the order on $\mathcal{C}(H)$ into the familiar order on self-adjoint operators, so that (Q2) becomes a question about positive matrices. Recall that $\mathcal{C}(H) = \{\mu_T : T \geq 0 \text{ trace class}\}$, where $\mu_T(P) = \text{Tr}(TP)$, and that $\mathcal{C}(H)$ is ordered pointwise on $L(H)$. For positive trace-class operators S and T , we have $\mu_S \leq \mu_T$ if and only if $S \leq T$. Indeed, if $\mu_S \leq \mu_T$, then $\langle \psi | T - S | \psi \rangle = \mu_T(P_\psi) - \mu_S(P_\psi) \geq 0$ for every $\psi \in S(H)$, so $T - S \geq 0$; and if $T - S \geq 0$, then $\text{Tr}((T - S)P) = \text{Tr}(P(T - S)P) \geq 0$ for every $P \in L(H)$, so $\mu_S \leq \mu_T$.

To show that $\mathcal{C}(H)$ is not a lattice, it is enough to exhibit two of its elements without a greatest lower bound, and we choose them given by commuting operators, so that the failure

cannot be blamed on noncommutativity. For orthonormal $e_1, e_2 \in H$, let

$$A = P_{e_1} + P_{e_2}, \quad D = 2P_{e_1} + \frac{1}{2}P_{e_2}. \quad (7)$$

Proposition 5.11. *Let $e_1, e_2 \in H$ be orthonormal, and let A and D be the operators defined in (7). Then μ_A and μ_D have no greatest lower bound in $\mathcal{C}(H)$, although A and D commute.*

Proof. Let $K = \text{span}(e_1, e_2)$, and let μ_T be a lower bound of μ_A and μ_D in $\mathcal{C}(H)$, so that $0 \leq T \leq A$ and $T \leq D$. If $\xi \perp K$, then $0 \leq \langle \xi | T | \xi \rangle \leq \langle \xi | A | \xi \rangle = 0$, so $T^{1/2}\xi = 0$ and hence $T\xi = 0$. Since T is self-adjoint, we have $\langle \xi | Tx \rangle = \langle T\xi | x \rangle = 0$ for all $x \in K$ and $\xi \perp K$. Hence T maps K into K and vanishes on $K^\perp$. We may therefore identify the lower bounds of μ_A and μ_D with positive 2×2 matrices in the basis e_1, e_2 . In this basis,

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad D = \begin{pmatrix} 2 & 0 \\ 0 & \frac{1}{2} \end{pmatrix}.$$

Throughout the argument we use the fact that a self-adjoint 2×2 matrix is positive if and only if both diagonal entries and the determinant are nonnegative. In particular, the diagonal matrix

$$M = \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{2} \end{pmatrix}$$

is a lower bound of A and D . Suppose that G were a greatest lower bound. Then $M \leq G \leq A$ forces $G_{11} = 1$. The positive matrix $G - M$ therefore has a zero $(1, 1)$ entry, and the nonnegativity of its determinant forces its off-diagonal entries to vanish. Then $M \leq G \leq D$ forces $G_{22} = \frac{1}{2}$, so $G = M$. To reach a contradiction, consider

$$\tau = \begin{pmatrix} \frac{11}{12} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} \end{pmatrix}.$$

Then $\tau \geq 0$ (since $\det \tau = \frac{1}{6}$), the matrix $A - \tau = \begin{pmatrix} 1/12 & -1/4 \\ -1/4 & 3/4 \end{pmatrix}$ is positive (since $\det(A - \tau) = 0$), and the matrix $D - \tau = \begin{pmatrix} 13/12 & -1/4 \\ -1/4 & 1/4 \end{pmatrix}$ is positive (since $\det(D - \tau) = \frac{5}{24}$), so τ is a lower bound of A and D . However, the matrix $M - \tau = \begin{pmatrix} 1/12 & -1/4 \\ -1/4 & 1/4 \end{pmatrix}$ has determinant $-\frac{1}{24} < 0$, so $\tau \not\leq M = G$, which contradicts the choice of G . Hence no greatest lower bound exists. ☹

The proof shows exactly where the classical argument of Proposition 5.10 breaks down. Among diagonal matrices alone, that is, among operators that commute with P_{e_1} and P_{e_2} , the matrix M would be the greatest lower bound of A and D , just as in the classical case, since we have $\text{diag}(a, b) \leq A$ and $\text{diag}(a, b) \leq D$ if and only if $a \leq 1$ and $b \leq \frac{1}{2}$. The lower bound τ , which shows that M is not the greatest lower bound, is not diagonal.

Proposition 5.11 rules out any identification of the order on quantum measures with the order on classical ones (order isomorphisms were defined at the beginning of this section).

Corollary 5.12. *Let Σ be a σ -algebra on a set Ω . There is no order isomorphism between $(\mathcal{C}(H), \leq)$ and $(\mathcal{M}(\Sigma), \leq)$.*

Proof. Suppose that $\Phi : \mathcal{C}(H) \rightarrow \mathcal{M}(\Sigma)$ is an order isomorphism. Then Φ^{-1} is also an order isomorphism. Let $\mu, \nu \in \mathcal{C}(H)$ be given. By (C2), proved in Proposition 5.10, the measures $\Phi(\mu)$ and $\Phi(\nu)$ have a greatest lower bound γ in $\mathcal{M}(\Sigma)$. We claim that $\Phi^{-1}(\gamma)$ is a greatest lower bound of μ and ν in $\mathcal{C}(H)$. Since γ is a lower bound of $\Phi(\mu)$ and $\Phi(\nu)$, the function $\Phi^{-1}(\gamma)$ is a lower bound of μ and ν . If $\kappa \in \mathcal{C}(H)$ is a lower bound of μ and ν , then $\Phi(\kappa)$ is a lower bound of $\Phi(\mu)$ and $\Phi(\nu)$, so $\Phi(\kappa) \leq \gamma$, and hence $\kappa \leq \Phi^{-1}(\gamma)$. This proves the claim. Thus any two elements of $\mathcal{C}(H)$ have a greatest lower bound, which contradicts Proposition 5.11. ☹



Remark 5.13 (Simplices and the Bloch ball). Two known theorems explain Proposition 5.11 geometrically; they are not used elsewhere in these notes. A compact convex subset of a finite-dimensional real vector space is a *simplex* if every point is a convex combination of extreme points in exactly one way, and a finite-dimensional real vector space ordered by a closed generating cone is a lattice if and only if a base of the cone is a simplex [Alf71]. Classically, the probability measures on a finite set Ω form a simplex whose extreme points are the Dirac measures, that is, the principal ultrafilters once more. For $H = \mathbb{C}^2$, the base of $\mathcal{C}(H)$ is the set of density operators. Every self-adjoint 2×2 matrix of trace 1 can be written uniquely as $\rho = \frac{1}{2}(I + x_1\sigma_1 + x_2\sigma_2 + x_3\sigma_3)$ with $x \in \mathbb{R}^3$, where

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix};$$

the eigenvalues of ρ are $\frac{1}{2}(1 \pm |x|)$, so ρ is a density operator if and only if $|x| \leq 1$. The density operators thus form a closed ball, the *Bloch ball*, whose extreme points, the pure states, fill the whole boundary sphere. The center $\frac{1}{2}I$ is the midpoint of every diameter, since $\frac{1}{2}I = \frac{1}{2}(P_\psi + P_{\psi'})$ for every orthonormal basis $\{\psi, \psi'\}$ of $\mathbb{C}^2$ (see Figure 4). So the Bloch ball is not a simplex, and the order is not a lattice. Kadison [Kad51] proved more: two self-adjoint operators on a Hilbert space have a least upper bound in the self-adjoint operators only if the two operators are comparable.





FIGURE 4. Left: the probability measures on a three-point space form a simplex, and each point is a unique convex combination of the Dirac measures at its vertices. Right: the density operators on $\mathbb{C}^2$ form a ball whose extreme points (pink) are the pure states on the boundary sphere; the center $\frac{1}{2}I$ is the midpoint of every diameter, so the decomposition into extreme points is far from unique.

6. TIMELINE

The story has three acts. In the first, von Neumann's argument against hidden variables is accepted by almost everyone for thirty years, although its key assumption had been questioned as early as 1935. In the second, which lasts only from 1957 to 1967, the theorem of these notes is found twice: once as a corollary of Gleason's Theorem, by Bell, and once as a finite configuration of vectors, by Kochen and Specker. In the third, which is still going on, the configurations are made smaller and smaller, first by hand and then by computer, and the

theorem is pushed beyond projections, beyond exact measurements, and beyond the values 0 and 1.



- 1932

von Neumann. Von Neumann gives the first argument against hidden variables, in his book *Mathematische Grundlagen der Quantenmechanik* [vN32]. The argument assumes that the hidden value of a sum $A + B$ of two observables is the sum of the hidden values of A and B , even when A and B cannot be measured together. The assumption is harmless for averages and unjustified for individual values, and Bell, thirty years later, put his objection bluntly: “the proof of von Neumann is not merely false but foolish!” [MC88].
- 1935

Hermann. Grete Hermann, a mathematician and philosopher who had written her doctoral thesis under Emmy Noether, objects to the same assumption in *Die naturphilosophischen Grundlagen der Quantenmechanik* [Her35]. Her objection went almost unread by physicists until the 1970s [Jam74], and how closely it anticipates Bell’s is still debated [MS18].
- 1957

Gleason. Gleason proves his theorem [Gle57] in answer to a question of Mackey’s about measures on the closed subspaces of a Hilbert space [Mac57, Mac63]. Hidden variables are nowhere mentioned: the theorem that would settle the question sat in plain sight for six years before anyone connected the two.
- 1960

Specker. Specker, meanwhile, approaches the same fact from logic. In *Die Logik nicht gleichzeitig entscheidbarer Aussagen* (“The logic of propositions that cannot be decided simultaneously”) [Spe60], he announces that the closed subspaces of a three-dimensional Hilbert space cannot be mapped into a Boolean algebra in a way that respects the operations on orthogonal subspaces. For a proof he offers only the phrase “an elementary geometric argument” (*ein elementargeometrisches Argument*); the argument itself would take him and Kochen seven more years to publish.
- 1963

Jauch–Piron. In Geneva, Jauch and Piron publish *Can Hidden Variables be Excluded in Quantum Mechanics?* [JP63], an argument against hidden variables in the lattice setting of these notes. Their argument assumes that whenever two questions both have probability 1, so does their meet. Bell’s 1966 paper would object that for questions that cannot be tested together,

the meet cannot be tested by testing the two questions, so nothing in physics forces this assumption either.

1963 • **Jauch.** It is Jauch, nonetheless, who sets Bell on the right track. J. M. Jauch (1914–1974), director of the Institute of Theoretical Physics at the University of Geneva, was then supervising Piron’s doctoral thesis [Pir64], which describes the logic of quantum mechanics by the lattice of closed subspaces, and so Gleason’s Theorem lay at the center of his own research. He pointed the theorem out to Bell, and, as Bell later recalled, he stressed exactly the right feature: Gleason’s Theorem needs additivity only for questions that can be tested together, which is the weakening of von Neumann’s assumption that Bell was looking for [Bel87]. Bell thanks him in a footnote of [Bel66]: “I am much indebted to Professor Jauch for drawing my attention to this work.”

1963–1965 • **Partial Boolean algebras.** Kochen and Specker, for their part, build the framework first. They present *Logical Structures Arising in Quantum Theory* at the 1963 Berkeley symposium on the theory of models, and *The Calculus of Partial Propositional Functions* at the 1964 Jerusalem congress; both appear in 1965 [KS65b, KS65a]. These papers introduce *partial Boolean algebras*: families of Boolean algebras, one for each set of questions that can be tested together, glued along their common parts. The application to hidden variables is held back for the 1967 paper.

1964–1966 • **Bell.** Bell, a particle physicist at CERN who worked on the foundations of quantum mechanics as a side interest, writes *On the Problem of Hidden Variables in Quantum Mechanics* [Bel66] in 1964, during a leave at the Stanford Linear Accelerator Center [Ber14]. The paper disposes of the arguments of von Neumann and of Jauch and Piron, and then derives the nonexistence of non-contextual hidden variables from Gleason’s Theorem, by an argument close to that of Section 2. Publication is delayed until 1966 (Rev. Mod. Phys. **38**, 447–452), so the paper appears between the two stages of Kochen and Specker’s work, and independently of both.

1967 • **Kochen–Specker.** Kochen and Specker publish *The Problem of Hidden Variables in Quantum Mechanics* [KS67]: a self-contained, finite proof with 117 vectors in $\mathbb{R}^3$, written in the language of partial Boolean algebras (see Section 4). Physicists often name the theorem after this paper alone; these notes add Bell’s name, for the reasons given in the subsection after the timeline.

1990 • **Peres, Mermin.** After a long quiet period, the third act opens with a change of viewpoint. Peres and, independently and shortly after, Mermin find the “magic square”: a 3×3 array of observables of a pair of two-dimensional systems, in which the products along the rows and the columns rule out every assignment of predetermined values ± 1 . The square speaks about observables rather than rays, but at heart it is the same kind of

argument as the proof of Section 4: a parity count that cannot come out right.

- 1991 • **Peres.** The shrinking begins. Peres reduces Kochen and Specker’s construction to 33 vectors in $\mathbb{R}^3$, and gives a 24-vector construction in $\mathbb{R}^4$ [Per91]. Conway and Kochen soon go down to 31 vectors in $\mathbb{R}^3$ (unpublished; reported in [Per93]), a record that still stands.
- 1994 • **Kernaghan.** Kernaghan finds 20-vector configurations in $\mathbb{R}^4$ hidden inside Peres’s 24 vectors [Ker94].
- 1996 • **Cabello et al.** Cabello, Estebaranz, and García-Alcaine find the 18-vector configuration in $\mathbb{R}^4$ used in Section 4 [CEGA96]. They started from the configurations of Peres and Kernaghan, and observed that none of their 18-vector configurations lies inside any of the 96 known 20-vector ones, “which probably explains why they were not obtained previously.”
- 2003 • **Cabello.** With the configurations nearly as small as they would get, attention turns to the hypotheses. Cabello extends the argument to measurements that are not given by projections, on a single two-dimensional system [Cab03] (see Appendix A below).
- 2004 • **Barrett–Kent.** Barrett and Kent ask whether the theorem survives when measurements have only finite precision and answers may be random [BK04] (see Appendix A below).
- 2005 • **Pavičić et al.** Pavičić, Merlet, McKay, and Megill close the question in dimension 4: an exhaustive computer search shows that no configuration has fewer than 18 vectors [PMMM05]. The configuration of Section 4 is therefore not merely economical but optimal.
- 2005 • **Spekkens.** In the same year Spekkens proposes a notion of non-contextuality phrased in terms of experimental procedures rather than projections, which applies to imperfect measurements [Spe05] (see Appendix A below).
- 2011 • **Arends et al.** Arends, Ouaknine, and Wampler turn the computer on dimension 3, and prove that every configuration there needs at least 18 vectors [AOW11].
- 2024 • **Li et al.** The gap in dimension 3 is still open: the best configuration known has the 31 vectors of Conway and Kochen, and a computer search has raised the lower bound to 24 [LBG24]. In the same year Ramanathan studies assignments with values in $[0, 1]$ instead of $\{0, 1\}$ [Ram24] (see Appendix A below), so the question that von Neumann asked about averages returns, ninety years later, in a sharper form.

6.1. Why the 1967 paper became the famous one. By Bell’s proof in Section 2, the nonexistence of two-valued quantum measures follows in a few lines from Gleason’s Theorem of 1957. Yet many students of the foundations of physics learn the result under the names of

Kochen and Specker alone. The reason is not that Kochen and Specker found a deeper fact; it lies partly in accidents of publication and mostly in the form of their proof.

The accidents are quickly told. Gleason’s paper never mentions hidden variables: it answers a question of Mackey’s, and nothing in it singles out two-valued measures, so for six years nobody noticed the connection (see the [1963](#) entry). Bell, who did notice, saw his paper delayed (see the [1964–1966](#) entry) until a year before Kochen and Specker’s.

The form of the proof matters more. Kochen and Specker’s argument depends on nothing else: it is finite and can be checked by hand, so the reader need not trust Gleason’s Theorem, a substantial piece of analysis. Finiteness had further consequences. It invited thirty years of shrinking the configuration, from 117 to 33 and then 31 vectors in $\mathbb{R}^3$, and from 24 to 20 to 18 in $\mathbb{R}^4$. It makes sense for structures that do not come from Hilbert spaces, where Gleason’s Theorem says nothing. And it can be taken to the laboratory, since a finite list of measurements can be carried out, whereas “all quantum measures on $L(H)$ at once” cannot.

So Gleason’s Theorem contained the result first, Bell was the first to draw it out, and Kochen and Specker gave it a form that could be checked by hand, generalized, and tested. This is why these notes speak of the Bell–Kochen–Specker theorem.

APPENDIX A. ATTEMPTS BEYOND CLASSICAL PROPOSITIONAL LOGIC

The proofs of Section 2 and Section 4 work throughout with propositional, two-valued ($\{0, 1\}$ -valued) logic. Two natural ways to go beyond that setting are to keep the values binary but enrich the logic, adding bound variables and quantifiers, or to keep the propositional structure but enrich the values, allowing outcomes in $[0, 1]$ rather than $\{0, 1\}$. This appendix records what has been done in each direction.

A.1. Richer logics: quantifiers and first-order quantum logic. The first sustained attempt to give quantum logic quantifiers came from a set theorist. Gaisi Takeuti, a Japanese logician who spent most of his career at the University of Illinois and was already known for Boolean-valued analysis, published *Quantum Set Theory* in 1981 [Tak81]. His idea was to run the Scott–Solovay construction of Boolean-valued models of ZFC with the complete Boolean algebra replaced by the (σ -complete) orthomodular lattice L of projections of a von Neumann algebra. The quantum universe $V^{(L)}$ is built by transfinite recursion, quantifiers receive a literal semantics, namely $\llbracket (\forall x)\varphi(x) \rrbracket = \bigwedge_{x \in V^{(L)}} \llbracket \varphi(x) \rrbracket$, an infimum taken in L , and the real numbers of this universe turn out to be self-adjoint operators. This is genuine first-order, indeed set-theoretic, quantification, evaluated in a truth-value algebra that is not Boolean. Takeuti hoped that a “quantum mathematics” based on quantum logic would grow out of the construction. In 1999 Satoko Titani’s *A Lattice-Valued Set Theory* [Tit99] placed it in a wider family, with arbitrary complete lattices as truth values, alongside the Heyting-valued (intuitionistic) set theories, and proved general completeness results.

Takeuti’s most persistent heir came to the subject from physics. Masanao Ozawa of Nagoya University wanted a rigorous, quantifier-based treatment of Heisenberg’s uncertainty principle, exactly the kind of first-order reasoning that a propositional logic cannot support, and he turned quantum set theory from a suggestive construction into a working apparatus. *Transfer Principle in Quantum Set Theory* [Oza07] proves a rigorous dictionary carrying theorems of ZFC about the reals over to theorems about self-adjoint operators inside $V^{(L)}$, in the spirit of the transfer principle of nonstandard analysis; *Orthomodular-Valued Models for Quantum Set Theory* [Oza09] tightens the model theory; and later work reforms the semantics to recover De Morgan’s laws and extends the framework to quantum conditionals. Ozawa’s interest in uncertainty was not only logical: his own reformulation of the uncertainty principle, proposed

in 2003 [Oza03], was tested with polarized neutrons by Yuji Hasegawa’s group in Vienna in 2012 [ESS⁺12].

The other great tool of modern set theory, Cohen’s method of forcing, was brought to the question by William Boos in 1996, in *Mathematical Quantum Theory I: Random Ultrafilters as Hidden Variables* [Boo96]. Where Takeuti had changed the truth-value algebra, Boos changed the hidden variables: his are *random* ultrafilters, generic objects for forcing with measure algebras, and he reads quantum probabilities through them. The paper remains one of very few attempts to attack the hidden-variable question, and hence the setting of Theorem 1.4, with the methods of set theory rather than those of analysis.

A third line came from physics proper. Between 1998 and 2002 Chris Isham, a quantum-gravity theorist at Imperial College London, and Jeremy Butterfield, a philosopher of physics at Cambridge, joined for one paper by Hamilton, published the four-part series *A Topos Perspective on the Kochen–Specker Theorem* [IB98, BI99, HIB00, IB02]. They took seriously the objection that the orthomodular lattice is too poorly behaved, being non-distributive, to serve as a logic with an ordinary semantics, and they replaced it by the family of its Boolean subalgebras (“contexts”), each of which is classical. In their reformulation, Theorem 1.4 says that there is no family of points, one in the Stone space of each context, that agree whenever one context is contained in another: this is the formulation of Section 3, and the statement of Section 5 that the quantum lattice has no points. Isham and Andreas Döring later extended the approach into a reformulation of physics, *A Topos Foundation for Theories of Physics* [DI08].

A.1.1. Classical first-order logic. Classical first-order logic entered quantum foundations in two quite different ways, and the two are best kept apart. The older tradition, the representation program of Mackey, Piron, Foulis and Randall, Hardy, and Barrett, uses classical logic only as ordinary mathematics is always written: it states physical postulates as axioms about a lattice, a family of tests or a convex set of states, and proves classical representation theorems about the structures that satisfy them. The younger tradition takes first-order logic literally, as an object of study: it treats the orthogonality relation on the closed subspaces of a Hilbert space as a first-order structure and asks the usual model-theoretic questions, namely whether the class of Hilbert lattices is elementary and whether its theory is decidable. Goldblatt and Fritz showed that both answers are negative. We describe the two traditions in this order.

In the early 1960s George Mackey at Harvard [Mac63] and Constantin Piron in Geneva, Jauch’s student [Pir64], chose to state the physical postulates (a set of “questions”, or an atomistic, orthomodular, irreducible lattice satisfying the covering law) as ordinary axioms in ordinary set-theoretic mathematics, with no non-classical logic anywhere, and to *prove*, as a classical representation theorem, that every structure satisfying them is isomorphic to the subspace lattice of a generalized Hilbert space. The quantum content lies in which structures satisfy the axioms, not in the logic used to reason about them. From 1972 on, David Foulis and Charles Randall at the University of Massachusetts continued this tradition with their *manuals* and *test spaces* [FR72], starting from experiments rather than from lattices: outcomes and tests are ordinary sets, and non-Boolean structure emerges only downstream, as a theorem about which propositions can be tested together. Lucien Hardy’s *Quantum Theory from Five Reasonable Axioms* [Har01] of 2001 and Jonathan Barrett’s *Information Processing in Generalized Probabilistic Theories* [Bar07] of 2007 revived the program in a modern form, in which quantum theory appears as one member of a family of probabilistic theories described by simple axioms about mixing states and combining systems.

The direct question, whether the class of Hilbert lattices can be axiomatized by classical first-order sentences in the language of the orthogonality relation, was answered in the negative by Robert Goldblatt, of Victoria University of Wellington, in *Orthomodularity is not elementary* [Gol84] of 1984. An infinite-dimensional pre-Hilbert space and its metric completion are elementarily equivalent as first-order structures, yet only the complete one is orthomodular. Metric completeness, which Theorem 1.4 and Gleason’s Theorem both use (closedness of subspaces, convergence on $S(H)$), is therefore invisible to classical first-order logic here. In 2016 Tobias Fritz, building on William Slofstra’s work on embeddability problems for groups, added a second negative result in *Quantum Logic is Undecidable* [Fri16]: the first-order theory of complex Hilbert lattices is undecidable already at the level of quasi-identities, so that no algorithm decides whether an implication between equations and orthogonality relations holds in every Hilbert space.

A.1.2. The approaches compared. The approaches described above are best read together. Classical first-order logic was not overlooked: tried on the most natural target (the orthogonality relation on subspaces), first-order logic was shown, by the theorems of the field itself, to be structurally too weak to pin down the class of Hilbert lattices at all (Goldblatt) and undecidable where applicable (Fritz). This double failure is a large part of why the two serious efforts at a genuinely quantified quantum logic (Takeuti’s line and the topos line) both abandoned classical semantics rather than pushing harder: Takeuti moved the non-classicality into the truth-value algebra, Isham–Butterfield moved the non-classicality into an intuitionistic internal logic, and both, not coincidentally, gained back exactly the property (extra structure that a bare classical first-order language cannot see) that Goldblatt’s theorem says is missing. The Mackey/Piron/Foulis–Randall/Hardy/Barrett line, meanwhile, never tried to axiomatize Hilbert lattices in first-order logic at all: that line uses classical first-order logic only as ordinary mathematics is always written, aimed at representation theorems about a differently chosen structure (a convex state space, a test space) rather than at characterizing subspace lattices up to elementary equivalence.

A.2. Richer values: $[0, 1]$ -valued and POVM-based assignments. Theorem 1.4 rules out *sharp*, two-valued, non-contextual assignments. A natural physical question is whether an analogous no-go survives once outcomes, or the physical measurements themselves, are allowed to be *unsharp*. The answer depends on which of two things is fuzzed, and pulls in opposite directions.

Unsharp measurements are modeled as follows. An *effect* on H is a self-adjoint operator E with $0 \leq E \leq I$, that is, with spectrum in $[0, 1]$. A *positive-operator-valued measure* (POVM) with finitely many outcomes is a finite family $(E_1, \dots, E_n)$ of effects with

$$E_1 + \dots + E_n = I.$$

By the Born rule, a system in the state given by a unit vector ψ yields outcome i with probability $\langle \psi | E_i | \psi \rangle$; the condition $\sum_i E_i = I$ says exactly that these probabilities sum to 1. A projective measurement, i.e. an orthogonal decomposition of I into projections as in Definition 1.1, is the special case in which every E_i is a projection. Physically, a POVM is what one obtains by coupling the system to an auxiliary system and performing an ordinary projective measurement on the combined system (Naimark’s dilation theorem says every POVM arises this way). The essential new feature is that the effects of a POVM need not be mutually orthogonal, so a POVM may have more outcomes than $\dim H$, and a single effect may belong to many different POVMs.

A.2.1. Fuzzing the measurements, keeping assignments sharp: the no-go strengthens, even at $\dim H = 2$. The analogue of a two-valued quantum measure is now a map v from effects to $\{0, 1\}$ such that, for every POVM $(E_1, \dots, E_n)$, exactly one E_i has $v(E_i) = 1$: each measurement has exactly one outcome. Non-contextuality is built into the requirement that v is a function of the effect alone, not of the POVM in which the effect appears. In $\dim H = 2$ the projections give too few constraints for a contradiction: the only projective measurements are the pairs $\{P, I - P\}$, and a two-valued quantum measure exists (see Remark 1.6). Effects give many more constraints, because the same effect appears in many POVMs, with different companions. Cabello [Cab03] exhibited a finite family of POVMs on the qubit $\mathbb{C}^2$ whose effects admit no such v . Thus the Bell–Kochen–Specker obstruction extends past sharp projections, into the regime $\dim H = 2$, which Theorem 1.4 does not reach.

A.2.2. Fuzzing the assignments, keeping measurements sharp: the naive no-go can be evaded, and needs a better physical definition to survive. Replace the deterministic $v(P) \in \{0, 1\}$ by a probabilistic response function $\xi_P(\lambda) \in [0, 1]$: the probability of outcome 1 given the ontic state λ , a point of a measure space Λ that represents the complete physical state of the system, whether or not anyone knows λ . Barrett and Kent [BK04] observed that once a real experiment resolves outcomes only to finite precision, non-contextual hidden-variable models built from such fuzzy response functions can reproduce quantum statistics to arbitrary accuracy — so naive basis-independence becomes physically toothless once assignments are allowed to be fuzzy. Spekkens [Spe05] closed this gap with a stronger notion of non-contextuality: experimental procedures that no experiment can tell apart must be represented in the same way in the hidden-variable model. Under this notion, new impossibility theorems hold, some of them already in dimension 2.

A.2.3. Extending the outcome alphabet directly. Ramanathan [Ram24] proves a Bell–Kochen–Specker-type theorem for assignments valued in a finite alphabet $\{0, p, 1 - p, 1\} \subseteq [0, 1]$ rather than only $\{0, 1\}$, for finite vector configurations in $\mathbb{C}^3$; the case $p = \frac{1}{2}$ rules out “fundamentally binary” hidden-variable theories.



Remark A.1. Fuzzing the *measurements* strengthens Theorem 1.4 (Cabello, reaching $\dim H = 2$); fuzzing the *assignments* under the naive definition of non-contextuality weakens the theorem (Barrett–Kent), which is why Spekkens’s stronger definition was needed to recover a robust no-go there; extending the outcome set while keeping assignments deterministic recovers a close cousin of the original theorem outright (Ramanathan). In each case the substantive work is a fresh rigidity argument playing the role Gleason’s Theorem played earlier in these notes — never a restatement of the bare fact that local assignments respect the connectives.

REFERENCES

- [Alf71] E. M. Alfsen, *Compact convex sets and boundary integrals*, Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 57, Springer, Berlin, 1971.
- [AOW11] M. Arends, J. Ouaknine, and C. W. Wampler, *On searching for small Kochen–Specker vector systems*, Graph-Theoretic Concepts in Computer Science (WG 2011), Lecture Notes in Computer Science, vol. 6986, Springer, 2011, pp. 23–34.
- [Bar07] J. Barrett, *Information processing in generalized probabilistic theories*, Phys. Rev. A **75** (2007), 032304.
- [Bel66] J. S. Bell, *On the problem of hidden variables in quantum mechanics*, Rev. Mod. Phys. **38** (1966), 447–452.
- [Bel87] ———, *Speakable and unspeakable in quantum mechanics: Collected papers on quantum philosophy*, Cambridge University Press, Cambridge, 1987.

- [Ber14] R. A. Bertlmann, *John Bell and the nature of the quantum world*, J. Phys. A: Math. Theor. **47** (2014), 424007.
- [BI99] J. Butterfield and C. J. Isham, *A topos perspective on the Kochen–Specker theorem: II. Conceptual aspects and classical analogues*, Int. J. Theor. Phys. **38** (1999), 827–859.
- [BK04] J. Barrett and A. Kent, *Non-contextuality, finite precision measurement and the Kochen–Specker theorem*, Stud. Hist. Phil. Mod. Phys. **35** (2004), 151–176, arXiv:quant-ph/0309017.
- [Boo96] William Boos, *Mathematical quantum theory I: Random ultrafilters as hidden variables*, Synthese **107** (1996), no. 1, 83–143.
- [Cab03] A. Cabello, *Kochen–Specker theorem for a single qubit using positive operator-valued measures*, Phys. Rev. Lett. **90** (2003), 190401, arXiv:quant-ph/0210082.
- [CEGA96] A. Cabello, J. M. Estebaranz, and G. García-Alcaine, *Bell–Kochen–Specker theorem: A proof with 18 vectors*, Phys. Lett. A **212** (1996), 183–187.
- [DI08] A. Döring and C. J. Isham, *A topos foundation for theories of physics: I–IV*, J. Math. Phys. **49** (2008), 053515–053518.
- [Dvu93] A. Dvurečenskij, *Gleason’s theorem and its applications*, Kluwer Academic Publishers, Dordrecht, 1993.
- [End01] H. B. Enderton, *A mathematical introduction to logic*, 2nd ed., Academic Press, 2001, Compactness theorem for propositional logic.
- [ESS⁺12] Jacqueline Erhart, Stephan Sponar, Georg Sulyok, Gerald Badurek, Masanao Ozawa, and Yuji Hasegawa, *Experimental demonstration of a universally valid error-disturbance uncertainty relation in spin measurements*, Nature Physics **8** (2012), 185–189.
- [FR72] D. J. Foulis and C. H. Randall, *Operational statistics. I. Basic concepts*, J. Math. Phys. **13** (1972), 1667–1675, and subsequent papers on manuals and test spaces.
- [Fri16] T. Fritz, *Quantum logic is undecidable*, 2016, arXiv:1607.05870.
- [Gle57] A. M. Gleason, *Measures on the closed subspaces of a Hilbert space*, J. Math. Mech. **6** (1957), 885–893.
- [Gol84] R. Goldblatt, *Orthomodularity is not elementary*, J. Symbolic Logic **49** (1984), 401–404.
- [Ham03] J. Hamhalter, *Quantum measure theory*, Fundamental Theories of Physics, vol. 134, Kluwer Academic Publishers, Dordrecht, 2003.
- [Har01] L. Hardy, *Quantum theory from five reasonable axioms*, 2001, arXiv:quant-ph/0101012.
- [Her35] Grete Hermann, *Die naturphilosophischen Grundlagen der Quantenmechanik*, Abhandlungen der Fries’schen Schule, Neue Folge **6** (1935), 75–152.
- [HIB00] J. Hamilton, C. J. Isham, and J. Butterfield, *A topos perspective on the Kochen–Specker theorem: III. Von Neumann algebras as the base category*, Int. J. Theor. Phys. **39** (2000), 1413–1436, arXiv:quant-ph/9911020.
- [IB98] C. J. Isham and J. Butterfield, *A topos perspective on the Kochen–Specker theorem: I. Quantum states as generalised valuations*, Int. J. Theor. Phys. **37** (1998), 2669–2733, arXiv:quant-ph/9803055.
- [IB02] ———, *A topos perspective on the Kochen–Specker theorem: IV. Interval valuations*, Int. J. Theor. Phys. **41** (2002), 613–639.
- [Jam74] Max Jammer, *The philosophy of quantum mechanics*, Wiley, New York, 1974.
- [JP63] J. M. Jauch and C. Piron, *Can hidden variables be excluded in quantum mechanics?*, Helv. Phys. Acta **36** (1963), 827–837.
- [Kad51] R. V. Kadison, *Order properties of bounded self-adjoint operators*, Proc. Amer. Math. Soc. **2** (1951), 505–510.
- [Ker94] M. Kernaghan, *Bell–Kochen–Specker theorem for 20 vectors*, J. Phys. A: Math. Gen. **27** (1994), L829–L830.
- [KS65a] Simon Kochen and Ernst P. Specker, *The calculus of partial propositional functions*, Logic, Methodology and Philosophy of Science (Proceedings of the 1964 International Congress, Jerusalem) (Y. Bar-Hillel, ed.), North-Holland, Amsterdam, 1965, pp. 45–57.
- [KS65b] ———, *Logical structures arising in quantum theory*, The Theory of Models (Proceedings of the 1963 International Symposium at Berkeley) (J. W. Addison, L. Henkin, and A. Tarski, eds.), North-Holland, Amsterdam, 1965, pp. 177–189.
- [KS67] S. Kochen and E. P. Specker, *The problem of hidden variables in quantum mechanics*, J. Math. Mech. **17** (1967), 59–87.

- [LBG24] Z. Li, C. Bright, and V. Ganesh, *A SAT solver and computer algebra attack on the minimum Kochen–Specker problem*, Proc. IJCAI 2024, 2024, arXiv:2306.13319.
- [Mac57] George W. Mackey, *Quantum mechanics and Hilbert space*, Amer. Math. Monthly **64** (1957), no. 8, part II, 45–57.
- [Mac63] G. W. Mackey, *Mathematical foundations of quantum mechanics*, W. A. Benjamin, New York, 1963.
- [MC88] C. Mann and R. Crease, *Interview: John Bell*, Omni **10** (1988), no. 8, 84–86, 88, 90, 92, 121.
- [Mey99] D. A. Meyer, *Finite precision measurement nullifies the Kochen–Specker theorem*, Phys. Rev. Lett. **83** (1999), 3751–3754.
- [MS18] N. David Mermin and Rüdiger Schack, *Homer nodded: von Neumann’s surprising oversight*, Found. Phys. **48** (2018), 1007–1020.
- [Oza03] Masanao Ozawa, *Universally valid reformulation of the Heisenberg uncertainty principle on noise and disturbance in measurement*, Phys. Rev. A **67** (2003), 042105.
- [Oza07] M. Ozawa, *Transfer principle in quantum set theory*, J. Symbolic Logic **72** (2007), 625–648, arXiv:math/0604349.
- [Oza09] ———, *Orthomodular-valued models for quantum set theory*, Rev. Symbolic Logic (2009), arXiv:0908.0367.
- [Per91] A. Peres, *Two simple proofs of the Kochen–Specker theorem*, J. Phys. A: Math. Gen. **24** (1991), L175–L178.
- [Per93] ———, *Quantum theory: Concepts and methods*, Kluwer, Dordrecht, 1993, Reports the unpublished 31-vector Conway–Kochen construction in $\mathbb{R}^3$.
- [Pir64] C. Piron, *Axiomatique quantique*, Helv. Phys. Acta **37** (1964), 439–468.
- [PMMM05] M. Pavičić, J.-P. Merlet, B. McKay, and N. D. Megill, *Kochen–Specker vectors*, J. Phys. A: Math. Gen. **38** (2005), no. 7, 1577–1592, arXiv:quant-ph/0409014.
- [Ram24] R. Ramanathan, *Generalised Kochen–Specker theorem for finite non-deterministic outcome assignments*, npj Quantum Information **10** (2024), 99.
- [RS80] Michael Reed and Barry Simon, *Methods of modern mathematical physics. I: Functional analysis*, revised and enlarged ed., Academic Press, New York, 1980.
- [Rud87] W. Rudin, *Real and complex analysis*, 3rd ed., McGraw-Hill, New York, 1987.
- [Sol71] R. M. Solovay, *Real-valued measurable cardinals*, Axiomatic Set Theory (Proc. Sympos. Pure Math., Vol. XIII, Part I), Amer. Math. Soc., Providence, RI, 1971, pp. 397–428.
- [Sol09] ———, *Gleason’s theorem for non-separable Hilbert spaces: extended abstract*, 2009, Preprint, available at math.berkeley.edu/~solovay/Preprints.
- [Spe60] Ernst Specker, *Die Logik nicht gleichzeitig entscheidbarer Aussagen*, Dialectica **14** (1960), 239–246.
- [Spe05] R. W. Spekkens, *Contextuality for preparations, transformations, and unsharp measurements*, Phys. Rev. A **71** (2005), 052108, arXiv:quant-ph/0406166.
- [Tak81] G. Takeuti, *Quantum set theory*, Current Issues in Quantum Logic (E. G. Beltrametti and B. C. van Fraassen, eds.), Ettore Majorana International Science Series, Plenum Press, New York, 1981, pp. 303–322.
- [Tit99] S. Titani, *A lattice-valued set theory*, Archive for Mathematical Logic **38** (1999), 395–421.
- [Ula30] S. Ulam, *Zur Masstheorie in der allgemeinen Mengenlehre*, Fund. Math. **16** (1930), 140–150.
- [vN32] J. von Neumann, *Mathematische Grundlagen der Quantenmechanik*, Springer, Berlin, 1932.