# § Lecture 21.0

Monday, 2 November 2026 20:22

Limit laws: Let $\sum_{n=m}^{\infty} a_n = x$, $\sum_{n=m}^{\infty} b_n = y$. Then

① $\sum_{n=m}^{\infty} (a_n + b_n) = x + y$

② For any $c \in R$, $\sum_{n=m}^{\infty} c\, a_n = cx$.

③ $\sum_{n=m}^{\infty} a_n$ is convergent iff $\sum_{n=m+k}^{\infty} a_n$ for any $k \geq 0$.

In fact, $\sum_{n=m}^{\infty} a_n = \sum_{n=m}^{m+k-1} a_n + \sum_{n=m+k}^{\infty} a_n$.

[Note that we are not rearranging here]

Claim: An infinite series $\sum_{n=m}^{\infty} a_n$ of nonnegative real numbers is convergent iff there exists a real number $M$ s.t.

$$\sum_{n=m}^{N} a_n \leq M \quad \forall N \geq m.$$

Proof: Recall that an increasing sequence is convergent iff it is bounded from above.

Clearly, $(S_N)_{N\geq m}$, where $S_N = \sum_{n=m}^{N} a_n$ is an increasing sequence as $a_n \geq 0 \ \forall n \geq m$.

Then $(S_N)_{N \geq m}$ is convergent iff it is bounded.

Lemma (Comparison test). Let $\sum_{n=m}^{\infty} a_n$ and $\sum_{n=m}^{\infty} b_n$ be infinite series and $|a_n| \leq b_n \ \forall n \geq m$. Then if $\sum_{n=m}^{\infty} b_n$ is convergent, then $\sum_{n=m}^{\infty} a_n$ is absolutely convergent. Further,

$$\left| \sum_{n=m}^{\infty} a_n \right| \leq \sum_{n=m}^{\infty} |a_n| \leq \sum_{n=m}^{\infty} b_n.$$

Proof: Given: $\sum_{n=m}^{\infty} b_n$ is convergent.

$\Rightarrow (S_N)_{N \geq m}$, where $S_N = \sum_{n=m}^{N} a_n$ is

(i) Increasing as $b_n \geq |a_n| \geq 0$

(ii) Convergent.

This means $(S_N)_{N\geq m}$ is bounded. Let $M \in R$ s.t.

$$S_N \leq M.$$

Let $T_N = \sum_{n=m}^{N} |a_n|$. Then $T_N = \sum_{n=m}^{N} |a_n| \leq \sum_{n=m}^{N} b_n \leq M$.

$\Rightarrow (T_N)_{N \geq m}$ is an increasing sequence, that is bounded. Then $(T_N)_{N\geq m}$ is convergent, thus $\sum_{n=m}^{\infty} |a_n|$ is convergent.

Since $T_N \leq S_N \Rightarrow \lim_{N\to\infty} T_N \leq \lim_{n\to\infty} S_N$

$$\Rightarrow \sum_{n=m}^{\infty} |a_n| \leq \sum_{n=m}^{\infty} b_n.$$

We already have proven $\left|\sum_{n=m}^{\infty} a_n\right| \leq \sum_{n=m}^{\infty} |a_n|$ for absolutely convergent series.

Example 1.

$$\sum_{n=0}^{\infty} x^n = \begin{cases} \text{Divergent if } |x| \geq 1 \\ \text{Absolutely convergent if } |x| < 1. \end{cases}$$

and $= \frac{1}{1-x}$.

Solution: Let $a_n = x^n$. Then $|a_n| = |x|^n \geq 1$. $\forall n \geq 0$.

Thus for $\varepsilon = 1/2$, there is no $N \geq m$ s.t. $|a_n| \leq \frac{1}{2} \ \forall n \geq N$.

(Because $|a_n| \geq 1 \ \forall n \geq m$).

$\Rightarrow \lim_{n\to\infty} a_n \neq 0$

$\Rightarrow \sum_{n=0}^{\infty} x^n$ is not convergent when $|x| \geq 1$.

Case 2: $|x| < 1$,

$$S_N = \sum_{n=0}^{N} x^n = \frac{1-x^{N+1}}{1-x}.$$

$$x S_N = \sum_{n=0}^{N} x^{n+1} = \sum_{n=1}^{N+1} x^n = -1 + S_N + x^{N+1}$$

$\Rightarrow (1-x) S_N = 1 - x^{N+1}$

or $S_N = \frac{1-x^{N+1}}{1-x}$

$$= \frac{1}{1-x} - \frac{1}{1-x} x^{N+1}$$

For $|x| < 1$, $\lim_{N\to\infty} x^N = 0 = \lim_{N\to\infty} x^{N+1}$

$\Rightarrow$

$\lim_{N\to\infty} S_N = 1/1-x$

or $\sum_{n=0}^{\infty} x^n = \frac{1}{1-x} \quad \forall |x| < 1$.

Cauchy criterion: Let $(a_n)_{n\geq 1}$ be a sequence where $a_n \geq 0$ and $a_{n+1} \leq a_n \ \forall n \geq 1$. Then series $\sum_{n=1}^{\infty} a_n$ is convergent iff

$$\sum_{k=0}^{\infty} 2^k a_{2^k} = a_1 + 2a_2 + \cdots \text{ is convergent.}$$

Ex.

$$\sum_{n=1}^{\infty} \frac{1}{n^q} = \begin{cases} \text{Divergent if } 0 < q \leq 1 \\ \text{Convergent if } q > 1. \end{cases}$$

Solution. Note that $\frac{1}{n^q} \geq 0 \quad \forall q > 0$.

Further $(n+1) > n \quad (n \geq 1)$

$\Rightarrow (n+1)^q > n^q \Rightarrow \frac{1}{(n+1)^q} < \frac{1}{n^q}$ (Decreasing)

Thus $\sum_{n=1}^{\infty} \frac{1}{n^q}$ is convergent iff $\sum_{k=0}^{\infty} 2^k \frac{1}{(2^k)^q}$ is convergent.

Let $x = 2^{(1-q)}$, then we need to find $q$ s.t.

$\sum_{k=0}^{\infty} x^k$ is convergent.

We have proven that it is convergent for $|x| < 1$.

i.e. $|2^{1-q}| < 1. \Leftrightarrow 2^{1-q} < 1$ As $2^{1-q} \geq 0$.

$$\Leftrightarrow 1-q < 0$$

$$\Leftrightarrow q > 1.$$

# § Lecture 21.1

Monday, 2 November 2026 21:39

## Rearrangement of series:

$$\sum_{n=m}^{\infty} a_n = \sum_{n=m}^{k} a_n + \sum_{n=k+1}^{\infty} a_n \quad \text{[Not a rearrangement.]}$$

$$\downarrow$$

$$\underbrace{\sum_{n=m}^{\infty} a_{f(n)}}_{\text{Rearrangement of } \sum_{n=m}^{\infty} a_n}$$ , where $f$ is a bijective map from $N \to N$.

**Grouping:** Consider a series $\sum_{n=0}^{\infty} a_n$. Choose indices $n_0 < n_1 < n_2 \cdots$ and write

$$b_0 = \sum_{n=0}^{n_0} a_n, \qquad b_1 = \sum_{n=n_0+1}^{n_1}, \cdots$$

$$\text{or} \quad b_k = \sum_{n=n_{k-1}+1}^{n_k} a_n \qquad (k \geq 1)$$

Then the grouped series $\sum_{k=0}^{\infty} b_k$.

e.g. $\sum_{n=0}^{\infty} (-1)^n$

Choose $n_k = 2k+1$, $n_0 = 1$, $n_1 = 3, \cdots$

$$b_0 = \sum_{n=0}^{n_0} a_n = \sum_{n=0}^{1} a_n = 0$$

$$b_1 = \sum_{n=2}^{3} a_n = 0$$

$$\vdots$$

$$\sum_{n=0}^{\infty} (-1)^n = \overbrace{(1-1)}^{b_0} + \overbrace{(1-1)}^{b_1} + \cdots$$

[Not rearrangement.]

Now let us consider the partial sums:

$$S_N = \sum_{n=0}^{N} a_n, \qquad T_N = \sum_{n=0}^{N} b_n.$$

$(S_N)_{N\geq 0}$ & $(T_N)_{N\geq 0}$. ; $T_0 =$

Notice $S_{n_0} = T_0$

$S_{n_1} = T_1$

$\vdots$

$S_{n_k} = T_k$

That is $(T_k)_{k\geq 0}$ is a subsequence of sequence $(S_N)_{N\geq 0}$.

All the convergence properties of $\sum_{n=0}^{\infty} b_n$ can be obtained by by analyzing the subsequence of the sequence of partial sums of $\sum_{n=0}^{\infty} a_n$.

Clearly, if $\sum_{n=0}^{\infty} a_n$ is convergent, then $\sum_{n=0}^{\infty} b_n$ is convergent for any regrouping.

While a nonconvergent sequence may give rise to a convergent subsequence as happened above.

In the above sense rearrangement is special.