



# Asymptotics of the Chebyshev Polynomials of General Sets

Barry Simon

IBM Professor of Mathematics and Theoretical Physics, Emeritus  
California Institute of Technology  
Pasadena, CA, U.S.A.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Asymptotics of the Chebyshev Polynomials of General Sets

Barry Simon

IBM Professor of Mathematics and Theoretical Physics, Emeritus  
California Institute of Technology  
Pasadena, CA, U.S.A.

Joint Work with Jacob Christiansen and Maxim Zinchenko  
Part 1: Inv. Math., **208** (2017), 217-245

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Asymptotics of the Chebyshev Polynomials of General Sets

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Barry Simon

IBM Professor of Mathematics and Theoretical Physics, Emeritus  
California Institute of Technology  
Pasadena, CA, U.S.A.

Joint Work with Jacob Christiansen and Maxim Zinchenko  
Part 1: Inv. Math, **208** (2017), 217-245  
Part 2: Submitted (also with Peter Yuditskii)



# Asymptotics of the Chebyshev Polynomials of General Sets

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Barry Simon

IBM Professor of Mathematics and Theoretical Physics, Emeritus  
California Institute of Technology  
Pasadena, CA, U.S.A.

Joint Work with Jacob Christiansen and Maxim Zinchenko  
Part 1: Inv. Math, **208** (2017), 217-245  
Part 2: Submitted (also with Peter Yuditskii)  
Part 3: Pavlov Memorial Volume, to appear.



# Chebyshev Polynomials

It is well known that monic orthogonal polynomials minimize the  $L^2(\mathfrak{e}, d\mu)$  norm if  $\mu$  is a measure with compact support,  $\mathfrak{e} \subset \mathbb{C}$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Chebyshev Polynomials

It is well known that monic orthogonal polynomials minimize the  $L^2(\mathfrak{e}, d\mu)$  norm if  $\mu$  is a measure with compact support,  $\mathfrak{e} \subset \mathbb{C}$ . Chebyshev polynomials just replace  $L^2$  by  $L^\infty$  (so only the support matters).

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Chebyshev Polynomials

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

It is well known that monic orthogonal polynomials minimize the  $L^2(\epsilon, d\mu)$  norm if  $\mu$  is a measure with compact support,  $\epsilon \subset \mathbb{C}$ . Chebyshev polynomials just replace  $L^2$  by  $L^\infty$  (so only the support matters).

Specifically, let  $\epsilon \subset \mathbb{C}$  be a compact, infinite, set of points. For any function,  $f$ , define

$$\|f\|_\epsilon = \sup \{|f(z)| \mid z \in \epsilon\}$$



# Chebyshev Polynomials

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

It is well known that monic orthogonal polynomials minimize the  $L^2(\epsilon, d\mu)$  norm if  $\mu$  is a measure with compact support,  $\epsilon \subset \mathbb{C}$ . Chebyshev polynomials just replace  $L^2$  by  $L^\infty$  (so only the support matters).

Specifically, let  $\epsilon \subset \mathbb{C}$  be a compact, infinite, set of points. For any function,  $f$ , define

$$\|f\|_\epsilon = \sup \{|f(z)| \mid z \in \epsilon\}$$

The *Chebyshev polynomial of degree  $n$*  is the monic polynomial,  $T_n$ , with

$$\|T_n\|_\epsilon = \inf \{\|P\|_\epsilon \mid \deg(P) = n \text{ and } P \text{ is monic}\}$$



# Chebyshev Polynomials

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

It is well known that monic orthogonal polynomials minimize the  $L^2(\epsilon, d\mu)$  norm if  $\mu$  is a measure with compact support,  $\epsilon \subset \mathbb{C}$ . Chebyshev polynomials just replace  $L^2$  by  $L^\infty$  (so only the support matters).

Specifically, let  $\epsilon \subset \mathbb{C}$  be a compact, infinite, set of points. For any function,  $f$ , define

$$\|f\|_\epsilon = \sup \{|f(z)| \mid z \in \epsilon\}$$

The *Chebyshev polynomial of degree  $n$*  is the monic polynomial,  $T_n$ , with

$$\|T_n\|_\epsilon = \inf \{\|P\|_\epsilon \mid \deg(P) = n \text{ and } P \text{ is monic}\}$$

The minimizer is unique



# Chebyshev Polynomials

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

It is well known that monic orthogonal polynomials minimize the  $L^2(\epsilon, d\mu)$  norm if  $\mu$  is a measure with compact support,  $\epsilon \subset \mathbb{C}$ . Chebyshev polynomials just replace  $L^2$  by  $L^\infty$  (so only the support matters).

Specifically, let  $\epsilon \subset \mathbb{C}$  be a compact, infinite, set of points. For any function,  $f$ , define

$$\|f\|_\epsilon = \sup \{|f(z)| \mid z \in \epsilon\}$$

The *Chebyshev polynomial of degree n* is the monic polynomial,  $T_n$ , with

$$\|T_n\|_\epsilon = \inf \{\|P\|_\epsilon \mid \deg(P) = n \text{ and } P \text{ is monic}\}$$

The minimizer is unique (as we'll see below in the case that  $\epsilon \subset \mathbb{R}$ ),



# Chebyshev Polynomials

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

It is well known that monic orthogonal polynomials minimize the  $L^2(\epsilon, d\mu)$  norm if  $\mu$  is a measure with compact support,  $\epsilon \subset \mathbb{C}$ . Chebyshev polynomials just replace  $L^2$  by  $L^\infty$  (so only the support matters).

Specifically, let  $\epsilon \subset \mathbb{C}$  be a compact, infinite, set of points. For any function,  $f$ , define

$$\|f\|_\epsilon = \sup \{|f(z)| \mid z \in \epsilon\}$$

The *Chebyshev polynomial of degree n* is the monic polynomial,  $T_n$ , with

$$\|T_n\|_\epsilon = \inf \{\|P\|_\epsilon \mid \deg(P) = n \text{ and } P \text{ is monic}\}$$

The minimizer is unique (as we'll see below in the case that  $\epsilon \subset \mathbb{R}$ ), so it is appropriate to speak of *the* Chebyshev polynomial rather than *a* Chebyshev polynomial.



# Chebyshev Polynomials

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Chebyshev invented his explicit polynomials which obey  $Q_n(\cos(\theta)) = \cos(n\theta)$  not because of their functional relation but because they are the best approximation on  $[-1, 1]$  to  $x^n$  by polynomials of degree  $n - 1$ . In this regard, Sodin and Yuditskii unearthed the following quote from a 1926 report by Lebesgue on the work of S. N. Bernstein.



# Chebyshev Polynomials

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Chebyshev invented his explicit polynomials which obey  $Q_n(\cos(\theta)) = \cos(n\theta)$  not because of their functional relation but because they are the best approximation on  $[-1, 1]$  to  $x^n$  by polynomials of degree  $n - 1$ . In this regard, Sodin and Yuditskii unearthed the following quote from a 1926 report by Lebesgue on the work of S. N. Bernstein.

*I assume that I am not the only one who does not understand the interest in and significance of these strange problems on maxima and minima studied by Chebyshev in memoirs whose titles often begin with, “On functions deviating least from zero . . . ”.*



# Chebyshev Polynomials

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Chebyshev invented his explicit polynomials which obey  $Q_n(\cos(\theta)) = \cos(n\theta)$  not because of their functional relation but because they are the best approximation on  $[-1, 1]$  to  $x^n$  by polynomials of degree  $n - 1$ . In this regard, Sodin and Yuditskii unearthed the following quote from a 1926 report by Lebesgue on the work of S. N. Bernstein.

*I assume that I am not the only one who does not understand the interest in and significance of these strange problems on maxima and minima studied by Chebyshev in memoirs whose titles often begin with, “On functions deviating least from zero . . .”. Could it be that one must have a Slavic soul to understand the great Russian Scholar?*



# Chebyshev Polynomials

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Chebyshev invented his explicit polynomials which obey  $Q_n(\cos(\theta)) = \cos(n\theta)$  not because of their functional relation but because they are the best approximation on  $[-1, 1]$  to  $x^n$  by polynomials of degree  $n - 1$ . In this regard, Sodin and Yuditskii unearthed the following quote from a 1926 report by Lebesgue on the work of S. N. Bernstein.

*I assume that I am not the only one who does not understand the interest in and significance of these strange problems on maxima and minima studied by Chebyshev in memoirs whose titles often begin with, “On functions deviating least from zero . . .”. Could it be that one must have a Slavic soul to understand the great Russian Scholar?*

This quote is a little bizarre given that, as we'll see, Borel (who was Lebesgue's thesis advisor) made important contributions to the subject in 1905!



# The Alternation Theorem

We will focus for most of this talk on the case  $\epsilon \subset \mathbb{R}$ , in which case,  $T_n$  is real, since on  $\mathbb{R}$ ,  $|\operatorname{Re}(T_n)|$  is smaller than  $|T_n|$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# The Alternation Theorem

We will focus for most of this talk on the case  $\epsilon \subset \mathbb{R}$ , in which case,  $T_n$  is real, since on  $\mathbb{R}$ ,  $|\operatorname{Re}(T_n)|$  is smaller than  $|T_n|$ .

We say that  $P_n$ , a degree  $n$  polynomial, has an *alternating set* in  $\epsilon \subset \mathbb{R}$  if there exists  $\{x_j\}_{j=0}^n \subset \epsilon$  with

$$x_0 < x_1 < \dots < x_n$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# The Alternation Theorem

We will focus for most of this talk on the case  $\epsilon \subset \mathbb{R}$ , in which case,  $T_n$  is real, since on  $\mathbb{R}$ ,  $|\operatorname{Re}(T_n)|$  is smaller than  $|T_n|$ .

We say that  $P_n$ , a degree  $n$  polynomial, has an *alternating set* in  $\epsilon \subset \mathbb{R}$  if there exists  $\{x_j\}_{j=0}^n \subset \epsilon$  with

$$x_0 < x_1 < \dots < x_n$$

and so that

$$P_n(x_j) = (-1)^{n-j} \|P_n\|_{\epsilon}$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# The Alternation Theorem

We will focus for most of this talk on the case  $\epsilon \subset \mathbb{R}$ , in which case,  $T_n$  is real, since on  $\mathbb{R}$ ,  $|\operatorname{Re}(T_n)|$  is smaller than  $|T_n|$ .

We say that  $P_n$ , a degree  $n$  polynomial, has an *alternating set* in  $\epsilon \subset \mathbb{R}$  if there exists  $\{x_j\}_{j=0}^n \subset \epsilon$  with

$$x_0 < x_1 < \dots < x_n$$

and so that

$$P_n(x_j) = (-1)^{n-j} \|P_n\|_{\epsilon}$$

While the basic idea of the following theorem goes back to Chebyshev, the result itself is due to Borel and Markov, independently, around 1905.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# The Alternation Theorem

**The Alternation Theorem** The Chebyshev polynomial of degree  $n$  has an alternating set.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# The Alternation Theorem

**The Alternation Theorem** The Chebyshev polynomial of degree  $n$  has an alternating set. Conversely, any monic polynomial with an alternating set is the Chebyshev polynomial.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# The Alternation Theorem

**The Alternation Theorem** The Chebyshev polynomial of degree  $n$  has an alternating set. Conversely, any monic polynomial with an alternating set is the Chebyshev polynomial.

If  $T_n$  is the Chebyshev polynomial, let  $y_0 < y_1 < \dots < y_k$  be the set of all the points in  $\mathbb{C}$  where its takes the value  $\pm \|T_n\|_{\mathbb{C}}$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# The Alternation Theorem

**The Alternation Theorem** The Chebyshev polynomial of degree  $n$  has an alternating set. Conversely, any monic polynomial with an alternating set is the Chebyshev polynomial.

If  $T_n$  is the Chebyshev polynomial, let  $y_0 < y_1 < \dots < y_k$  be the set of all the points in  $\epsilon$  where its takes the value  $\pm \|T_n\|_\epsilon$ . If there are fewer than  $n$  sign changes among these ordered points we can find a degree at most  $n - 1$  polynomial,  $Q$ , non-vanishing at each  $y_j$  and with the same sign as  $T_n$  at those points.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# The Alternation Theorem

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics

**The Alternation Theorem** The Chebyshev polynomial of degree  $n$  has an alternating set. Conversely, any monic polynomial with an alternating set is the Chebyshev polynomial.

If  $T_n$  is the Chebyshev polynomial, let  $y_0 < y_1 < \dots < y_k$  be the set of all the points in  $\epsilon$  where its takes the value  $\pm \|T_n\|_\epsilon$ . If there are fewer than  $n$  sign changes among these ordered points we can find a degree at most  $n - 1$  polynomial,  $Q$ , non-vanishing at each  $y_j$  and with the same sign as  $T_n$  at those points. For  $\epsilon$  small and positive,  $T_n - \epsilon Q$  will be a monic polynomial with smaller  $\|\cdot\|_\epsilon$ .



# The Alternation Theorem

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics

**The Alternation Theorem** The Chebyshev polynomial of degree  $n$  has an alternating set. Conversely, any monic polynomial with an alternating set is the Chebyshev polynomial.

If  $T_n$  is the Chebyshev polynomial, let  $y_0 < y_1 < \dots < y_k$  be the set of all the points in  $\epsilon$  where it takes the value  $\pm \|T_n\|_\epsilon$ . If there are fewer than  $n$  sign changes among these ordered points we can find a degree at most  $n - 1$  polynomial,  $Q$ , non-vanishing at each  $y_j$  and with the same sign as  $T_n$  at those points. For  $\epsilon$  small and positive,  $T_n - \epsilon Q$  will be a monic polynomial with smaller  $\|\cdot\|_\epsilon$ . Thus there must be at least  $n$  sign flips and therefore an alternating set.



# The Alternation Theorem

Conversely, let  $P_n$  be a degree  $n$  monic polynomial with an alternating set and suppose that  $\|T_n\|_{\mathfrak{e}} < \|P_n\|_{\mathfrak{e}}$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# The Alternation Theorem

Conversely, let  $P_n$  be a degree  $n$  monic polynomial with an alternating set and suppose that  $\|T_n\|_{\mathfrak{e}} < \|P_n\|_{\mathfrak{e}}$ . Then at each point,  $x_j$ , in the alternating set for  $P_n$ ,  $Q \equiv P_n - T_n$  has the same sign as  $P_n$ ,

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# The Alternation Theorem

Conversely, let  $P_n$  be a degree  $n$  monic polynomial with an alternating set and suppose that  $\|T_n\|_{\epsilon} < \|P_n\|_{\epsilon}$ . Then at each point,  $x_j$ , in the alternating set for  $P_n$ ,  $Q \equiv P_n - T_n$  has the same sign as  $P_n$ , so  $Q$  has at least  $n$  zeros, which is impossible, since it is of degree at most  $n - 1$ . ■

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# The Alternation Theorem

Conversely, let  $P_n$  be a degree  $n$  monic polynomial with an alternating set and suppose that  $\|T_n\|_{\epsilon} < \|P_n\|_{\epsilon}$ . Then at each point,  $x_j$ , in the alternating set for  $P_n$ ,  $Q \equiv P_n - T_n$  has the same sign as  $P_n$ , so  $Q$  has at least  $n$  zeros, which is impossible, since it is of degree at most  $n - 1$ . ■

The alternation theorem implies uniqueness of the Chebyshev polynomial. For, if  $T_n$  and  $S_n$  are two minimizers, so is

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# The Alternation Theorem

Conversely, let  $P_n$  be a degree  $n$  monic polynomial with an alternating set and suppose that  $\|T_n\|_{\epsilon} < \|P_n\|_{\epsilon}$ . Then at each point,  $x_j$ , in the alternating set for  $P_n$ ,  $Q \equiv P_n - T_n$  has the same sign as  $P_n$ , so  $Q$  has at least  $n$  zeros, which is impossible, since it is of degree at most  $n - 1$ . ■

The alternation theorem implies uniqueness of the Chebyshev polynomial. For, if  $T_n$  and  $S_n$  are two minimizers, so is  $Q \equiv \frac{1}{2}(T_n + S_n)$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# The Alternation Theorem

Conversely, let  $P_n$  be a degree  $n$  monic polynomial with an alternating set and suppose that  $\|T_n\|_{\epsilon} < \|P_n\|_{\epsilon}$ . Then at each point,  $x_j$ , in the alternating set for  $P_n$ ,  $Q \equiv P_n - T_n$  has the same sign as  $P_n$ , so  $Q$  has at least  $n$  zeros, which is impossible, since it is of degree at most  $n - 1$ . ■

The alternation theorem implies uniqueness of the Chebyshev polynomial. For, if  $T_n$  and  $S_n$  are two minimizers, so is  $Q \equiv \frac{1}{2}(T_n + S_n)$ .

At the alternating points for  $Q$ , we must have  $T_n = S_n$ ,

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# The Alternation Theorem

Conversely, let  $P_n$  be a degree  $n$  monic polynomial with an alternating set and suppose that  $\|T_n\|_{\epsilon} < \|P_n\|_{\epsilon}$ . Then at each point,  $x_j$ , in the alternating set for  $P_n$ ,  $Q \equiv P_n - T_n$  has the same sign as  $P_n$ , so  $Q$  has at least  $n$  zeros, which is impossible, since it is of degree at most  $n - 1$ . ■

The alternation theorem implies uniqueness of the Chebyshev polynomial. For, if  $T_n$  and  $S_n$  are two minimizers, so is  $Q \equiv \frac{1}{2}(T_n + S_n)$ .

At the alternating points for  $Q$ , we must have  $T_n = S_n$ , so they must be equal polynomials since there are  $n + 1$  points and their difference has degree at most  $n - 1$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Alternation and Zeros

If  $T_n$  is the Chebyshev polynomial for  $\epsilon \subset \mathbb{R}$  and  $x_0 < x_1 < \dots < x_n$  is an alternating set for  $T_n$ ,

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Alternation and Zeros

If  $T_n$  is the Chebyshev polynomial for  $\epsilon \subset \mathbb{R}$  and  $x_0 < x_1 < \dots < x_n$  is an alternating set for  $T_n$ , there must be at least one zero (in  $\mathbb{R}$ , not necessarily in  $\epsilon$ ) between  $x_{j-1}$  and  $x_j$  because of the sign change.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Alternation and Zeros

If  $T_n$  is the Chebyshev polynomial for  $\epsilon \subset \mathbb{R}$  and  $x_0 < x_1 < \dots < x_n$  is an alternating set for  $T_n$ , there must be at least one zero (in  $\mathbb{R}$ , not necessarily in  $\epsilon$ ) between  $x_{j-1}$  and  $x_j$  because of the sign change. Since this accounts for all  $n$  zeros:

**Fact 1** All the zeros of the Chebyshev polynomials of a set  $\epsilon \subset \mathbb{R}$  lie in  $\mathbb{R}$  and all are simple and lie in  $\text{cvh}(\epsilon)$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Alternation and Zeros

If  $T_n$  is the Chebyshev polynomial for  $\epsilon \subset \mathbb{R}$  and  $x_0 < x_1 < \dots < x_n$  is an alternating set for  $T_n$ , there must be at least one zero (in  $\mathbb{R}$ , not necessarily in  $\epsilon$ ) between  $x_{j-1}$  and  $x_j$  because of the sign change. Since this accounts for all  $n$  zeros:

**Fact 1** All the zeros of the Chebyshev polynomials of a set  $\epsilon \subset \mathbb{R}$  lie in  $\mathbb{R}$  and all are simple and lie in  $\text{cvh}(\epsilon)$ .

Here,  $\text{cvh}(\epsilon)$  is the convex hull of  $\epsilon$  and that result follows from  $x_0, x_n \in \epsilon$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Alternation and Zeros

By a *gap* of  $\epsilon \subset \mathbb{R}$ , we mean a bounded connected component of  $\mathbb{R} \setminus \epsilon$ .

Chebyshev  
Polynomials

**Alternation  
Theorem**

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Alternation and Zeros

By a *gap* of  $\epsilon \subset \mathbb{R}$ , we mean a bounded connected component of  $\mathbb{R} \setminus \epsilon$ . If there are only finitely many gaps and no component of  $\epsilon$  is a single point, we speak of a finite gap set.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Alternation and Zeros

By a *gap* of  $\epsilon \subset \mathbb{R}$ , we mean a bounded connected component of  $\mathbb{R} \setminus \epsilon$ . If there are only finitely many gaps and no component of  $\epsilon$  is a single point, we speak of a finite gap set. Between any two zeros of  $T_n$ , there is a point in the alternating set so

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Alternation and Zeros

By a *gap* of  $\epsilon \subset \mathbb{R}$ , we mean a bounded connected component of  $\mathbb{R} \setminus \epsilon$ . If there are only finitely many gaps and no component of  $\epsilon$  is a single point, we speak of a finite gap set. Between any two zeros of  $T_n$ , there is a point in the alternating set so

**Fact 2** Each gap of  $\epsilon \subset \mathbb{R}$  has at most one zero of  $T_n$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Alternation and Zeros

By a *gap* of  $\epsilon \subset \mathbb{R}$ , we mean a bounded connected component of  $\mathbb{R} \setminus \epsilon$ . If there are only finitely many gaps and no component of  $\epsilon$  is a single point, we speak of a finite gap set. Between any two zeros of  $T_n$ , there is a point in the alternating set so

**Fact 2** Each gap of  $\epsilon \subset \mathbb{R}$  has at most one zero of  $T_n$ .

Above the top zero (resp. below the bottom zero) of  $T_n$ ,  $|T_n(x)|$  is monotone increasing (resp. decreasing). It follows that  $x_n = \sup_{y \in \epsilon} y$  (resp  $x_0 = \inf_{y \in \epsilon} y$ ) so

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Alternation and Zeros

By a *gap* of  $\epsilon \subset \mathbb{R}$ , we mean a bounded connected component of  $\mathbb{R} \setminus \epsilon$ . If there are only finitely many gaps and no component of  $\epsilon$  is a single point, we speak of a finite gap set. Between any two zeros of  $T_n$ , there is a point in the alternating set so

**Fact 2** Each gap of  $\epsilon \subset \mathbb{R}$  has at most one zero of  $T_n$ .

Above the top zero (resp. below the bottom zero) of  $T_n$ ,  $|T_n(x)|$  is monotone increasing (resp. decreasing). It follows that  $x_n = \sup_{y \in \epsilon} y$  (resp  $x_0 = \inf_{y \in \epsilon} y$ ) so

**Fact 3** At the end points of  $\text{cvh}(\epsilon) \subset \mathbb{R}$  we have that  $|T_n(x)| = \|T_n\|_\epsilon$  and

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Alternation and Zeros

By a *gap* of  $\epsilon \subset \mathbb{R}$ , we mean a bounded connected component of  $\mathbb{R} \setminus \epsilon$ . If there are only finitely many gaps and no component of  $\epsilon$  is a single point, we speak of a finite gap set. Between any two zeros of  $T_n$ , there is a point in the alternating set so

**Fact 2** Each gap of  $\epsilon \subset \mathbb{R}$  has at most one zero of  $T_n$ .

Above the top zero (resp. below the bottom zero) of  $T_n$ ,  $|T_n(x)|$  is monotone increasing (resp. decreasing). It follows that  $x_n = \sup_{y \in \epsilon} y$  (resp  $x_0 = \inf_{y \in \epsilon} y$ ) so

**Fact 3** At the end points of  $\text{cvh}(\epsilon) \subset \mathbb{R}$  we have that  $|T_n(x)| = \|T_n\|_\epsilon$  and

$$\epsilon_n \equiv T_n^{-1}([- \|T_n\|_\epsilon, \|T_n\|_\epsilon]) \subset \text{cvh}(\epsilon)$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Coulomb Energies and All That

Szegő realized that Chebyshev polynomials are intimately connected with two dimensional potential theory, so I want to review some of the basics of that subject.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Coulomb Energies and All That

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Szegő realized that Chebyshev polynomials are intimately connected with two dimensional potential theory, so I want to review some of the basics of that subject. Given a probability measure,  $d\mu$ , of compact support on  $\mathbb{C}$ , we define its *Coulomb energy*,  $\mathcal{E}(\mu)$  by

$$\mathcal{E}(\mu) = \int d\mu(x) d\mu(y) \log |x - y|^{-1}$$



# Coulomb Energies and All That

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Szegő realized that Chebyshev polynomials are intimately connected with two dimensional potential theory, so I want to review some of the basics of that subject. Given a probability measure,  $d\mu$ , of compact support on  $\mathbb{C}$ , we define its *Coulomb energy*,  $\mathcal{E}(\mu)$  by

$$\mathcal{E}(\mu) = \int d\mu(x) d\mu(y) \log |x - y|^{-1}$$

and we define the *Robin constant*, of a compact set  $\mathfrak{e} \subset \mathbb{C}$

$$R(\mathfrak{e}) = \inf \{\mathcal{E}(\mu) \mid \text{supp}(\mu) \subset \mathfrak{e} \text{ and } \mu(\mathfrak{e}) = 1\}$$



# Coulomb Energies and All That

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Szegő realized that Chebyshev polynomials are intimately connected with two dimensional potential theory, so I want to review some of the basics of that subject. Given a probability measure,  $d\mu$ , of compact support on  $\mathbb{C}$ , we define its *Coulomb energy*,  $\mathcal{E}(\mu)$  by

$$\mathcal{E}(\mu) = \int d\mu(x) d\mu(y) \log |x - y|^{-1}$$

and we define the *Robin constant*, of a compact set  $\mathfrak{e} \subset \mathbb{C}$

$$R(\mathfrak{e}) = \inf\{\mathcal{E}(\mu) \mid \text{supp}(\mu) \subset \mathfrak{e} \text{ and } \mu(\mathfrak{e}) = 1\}$$

If  $R(\mathfrak{e}) = \infty$ , we say  $\mathfrak{e}$  is a *polar set* or has *capacity zero*.



# Coulomb Energies and All That

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Szegő realized that Chebyshev polynomials are intimately connected with two dimensional potential theory, so I want to review some of the basics of that subject. Given a probability measure,  $d\mu$ , of compact support on  $\mathbb{C}$ , we define its *Coulomb energy*,  $\mathcal{E}(\mu)$  by

$$\mathcal{E}(\mu) = \int d\mu(x) d\mu(y) \log |x - y|^{-1}$$

and we define the *Robin constant*, of a compact set  $\mathfrak{e} \subset \mathbb{C}$

$$R(\mathfrak{e}) = \inf \{\mathcal{E}(\mu) \mid \text{supp}(\mu) \subset \mathfrak{e} \text{ and } \mu(\mathfrak{e}) = 1\}$$

If  $R(\mathfrak{e}) = \infty$ , we say  $\mathfrak{e}$  is a *polar set* or has *capacity zero*. If something holds except for a polar set, we say it holds q.e. (for *quasi-everywhere*).



# Coulomb Energies and All That

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Szegő realized that Chebyshev polynomials are intimately connected with two dimensional potential theory, so I want to review some of the basics of that subject. Given a probability measure,  $d\mu$ , of compact support on  $\mathbb{C}$ , we define its *Coulomb energy*,  $\mathcal{E}(\mu)$  by

$$\mathcal{E}(\mu) = \int d\mu(x) d\mu(y) \log |x - y|^{-1}$$

and we define the *Robin constant*, of a compact set  $\mathfrak{e} \subset \mathbb{C}$

$$R(\mathfrak{e}) = \inf \{\mathcal{E}(\mu) \mid \text{supp}(\mu) \subset \mathfrak{e} \text{ and } \mu(\mathfrak{e}) = 1\}$$

If  $R(\mathfrak{e}) = \infty$ , we say  $\mathfrak{e}$  is a *polar set* or has *capacity zero*. If something holds except for a polar set, we say it holds q.e. (for *quasi-everywhere*). The capacity,  $C(\mathfrak{e})$ , of  $\mathfrak{e}$  is defined by

$$C(\mathfrak{e}) = \exp(-R(\mathfrak{e})) \qquad \qquad R(\mathfrak{e}) = \log(1/C(\mathfrak{e}))$$



# Equilibrium Measures and All That

If  $\epsilon$  is not a polar set, it follows from weak lower semicontinuity of  $\mathcal{E}(\cdot)$  and weak compactness of the family of probability measures that there is a probability measure whose Coulomb energy is  $R(\epsilon)$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Equilibrium Measures and All That

If  $\epsilon$  is not a polar set, it follows from weak lower semicontinuity of  $\mathcal{E}(\cdot)$  and weak compactness of the family of probability measures that there is a probability measure whose Coulomb energy is  $R(\epsilon)$ . Since  $\mathcal{E}(\cdot)$  is strictly convex on the probability measures, this minimizer is unique.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Equilibrium Measures and All That

If  $\epsilon$  is not a polar set, it follows from weak lower semicontinuity of  $\mathcal{E}(\cdot)$  and weak compactness of the family of probability measures that there is a probability measure whose Coulomb energy is  $R(\epsilon)$ . Since  $\mathcal{E}(\cdot)$  is strictly convex on the probability measures, this minimizer is unique. It is called the *equilibrium measure* or *harmonic measure* of  $\epsilon$  and denoted  $d\rho_\epsilon$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Equilibrium Measures and All That

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

If  $\epsilon$  is not a polar set, it follows from weak lower semicontinuity of  $\mathcal{E}(\cdot)$  and weak compactness of the family of probability measures that there is a probability measure whose Coulomb energy is  $R(\epsilon)$ . Since  $\mathcal{E}(\cdot)$  is strictly convex on the probability measures, this minimizer is unique. It is called the *equilibrium measure* or *harmonic measure* of  $\epsilon$  and denoted  $d\rho_\epsilon$ . The second name comes from the fact that if  $f$  is a continuous function on  $\epsilon$ , there is a unique function,  $u_f$ , harmonic on  $(\mathbb{C} \cup \{\infty\}) \setminus \epsilon$ , which approaches  $f(x)$  for q.e.  $x \in \epsilon$  (i.e., solves the Dirichlet problem)



# Equilibrium Measures and All That

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

If  $\epsilon$  is not a polar set, it follows from weak lower semicontinuity of  $\mathcal{E}(\cdot)$  and weak compactness of the family of probability measures that there is a probability measure whose Coulomb energy is  $R(\epsilon)$ . Since  $\mathcal{E}(\cdot)$  is strictly convex on the probability measures, this minimizer is unique. It is called the *equilibrium measure* or *harmonic measure* of  $\epsilon$  and denoted  $d\rho_\epsilon$ . The second name comes from the fact that if  $f$  is a continuous function on  $\epsilon$ , there is a unique function,  $u_f$ , harmonic on  $(\mathbb{C} \cup \{\infty\}) \setminus \epsilon$ , which approaches  $f(x)$  for q.e.  $x \in \epsilon$  (i.e., solves the Dirichlet problem) and

$$u_f(\infty) = \int_{\epsilon} f(x) d\rho_\epsilon(x)$$



# Equilibrium Measures and All That

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

If  $\epsilon$  is not a polar set, it follows from weak lower semicontinuity of  $\mathcal{E}(\cdot)$  and weak compactness of the family of probability measures that there is a probability measure whose Coulomb energy is  $R(\epsilon)$ . Since  $\mathcal{E}(\cdot)$  is strictly convex on the probability measures, this minimizer is unique. It is called the *equilibrium measure* or *harmonic measure* of  $\epsilon$  and denoted  $d\rho_\epsilon$ . The second name comes from the fact that if  $f$  is a continuous function on  $\epsilon$ , there is a unique function,  $u_f$ , harmonic on  $(\mathbb{C} \cup \{\infty\}) \setminus \epsilon$ , which approaches  $f(x)$  for q.e.  $x \in \epsilon$  (i.e., solves the Dirichlet problem) and

$$u_f(\infty) = \int_{\epsilon} f(x) d\rho_\epsilon(x)$$

The function  $\Phi_\epsilon(z) = \int_{\epsilon} d\rho_\epsilon(x) \log |x - z|^{-1}$  is called the *equilibrium potential*.



# Green's Function

The *Green's function*,  $G_{\mathfrak{e}}(z)$ , of a compact subset,  $\mathfrak{e} \subset \mathbb{C}$ , is defined by

$$G_{\mathfrak{e}}(z) = R(\mathfrak{e}) - \Phi_{\mathfrak{e}}(z)$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Green's Function

The *Green's function*,  $G_\epsilon(z)$ , of a compact subset,  $\epsilon \subset \mathbb{C}$ , is defined by

$$G_\epsilon(z) = R(\epsilon) - \Phi_\epsilon(z)$$

It is the unique function harmonic on  $\mathbb{C} \setminus \epsilon$  with q.e. boundary value 0 on  $\epsilon$  and so that  $G_\epsilon(z) - \log |z|$  is harmonic at  $\infty$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Green's Function

The *Green's function*,  $G_\epsilon(z)$ , of a compact subset,  $\epsilon \subset \mathbb{C}$ , is defined by

$$G_\epsilon(z) = R(\epsilon) - \Phi_\epsilon(z)$$

It is the unique function harmonic on  $\mathbb{C} \setminus \epsilon$  with q.e. boundary value 0 on  $\epsilon$  and so that  $G_\epsilon(z) - \log |z|$  is harmonic at  $\infty$ . Moreover,  $G_\epsilon(z) \geq 0$  everywhere and near  $\infty$

$$G_\epsilon(z) = \log |z| + R(\epsilon) + O(1/|z|)$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Green's Function

The *Green's function*,  $G_\epsilon(z)$ , of a compact subset,  $\epsilon \subset \mathbb{C}$ , is defined by

$$G_\epsilon(z) = R(\epsilon) - \Phi_\epsilon(z)$$

It is the unique function harmonic on  $\mathbb{C} \setminus \epsilon$  with q.e. boundary value 0 on  $\epsilon$  and so that  $G_\epsilon(z) - \log |z|$  is harmonic at  $\infty$ . Moreover,  $G_\epsilon(z) \geq 0$  everywhere and near  $\infty$

$$G_\epsilon(z) = \log |z| + R(\epsilon) + O(1/|z|)$$

equivalently,

$$\exp(G_\epsilon(z)) = \frac{|z|}{C(\epsilon)} + O(1)$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Szegő's Lower Bound

Let  $\epsilon \subset \mathbb{C}$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Szegő's Lower Bound

Let  $\epsilon \subset \mathbb{C}$ . Define  $f_n = \{z \mid |T_n(z)| \leq \|T_n\|_\epsilon\}$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Szegő's Lower Bound

Let  $\epsilon \subset \mathbb{C}$ . Define  $f_n = \{z \mid |T_n(z)| \leq \|T_n\|_\epsilon\}$  so that  $\epsilon \subset f_n$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Szegő's Lower Bound

Let  $\epsilon \subset \mathbb{C}$ . Define  $f_n = \{z \mid |T_n(z)| \leq \|T_n\|_\epsilon\}$  so that  $\epsilon \subset f_n$  and thus

$$C(\epsilon) \leq C(f_n)$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Szegő's Lower Bound

Let  $\epsilon \subset \mathbb{C}$ . Define  $\mathfrak{f}_n = \{z \mid |T_n(z)| \leq \|T_n\|_\epsilon\}$  so that  $\epsilon \subset \mathfrak{f}_n$  and thus

$$C(\epsilon) \leq C(\mathfrak{f}_n)$$

The function,  $G(z) = n^{-1} \log (|T_n(z)|/\|T_n\|_\epsilon)$  is the Green's function of  $\mathfrak{f}_n$  (check properties) so

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Szegő's Lower Bound

Let  $\epsilon \subset \mathbb{C}$ . Define  $\mathfrak{f}_n = \{z \mid |T_n(z)| \leq \|T_n\|_\epsilon\}$  so that  $\epsilon \subset \mathfrak{f}_n$  and thus

$$C(\epsilon) \leq C(\mathfrak{f}_n)$$

The function,  $G(z) = n^{-1} \log (|T_n(z)|/\|T_n\|_\epsilon)$  is the Green's function of  $\mathfrak{f}_n$  (check properties) so

$$C(\mathfrak{f}_n) = \|T_n\|_\epsilon^{1/n} \Rightarrow \|T_n\|_\epsilon \geq C(\epsilon)^n$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Szegő's Lower Bound

Let  $\epsilon \subset \mathbb{C}$ . Define  $\mathfrak{f}_n = \{z \mid |T_n(z)| \leq \|T_n\|_\epsilon\}$  so that  $\epsilon \subset \mathfrak{f}_n$  and thus

$$C(\epsilon) \leq C(\mathfrak{f}_n)$$

The function,  $G(z) = n^{-1} \log (|T_n(z)|/\|T_n\|_\epsilon)$  is the Green's function of  $\mathfrak{f}_n$  (check properties) so

$$C(\mathfrak{f}_n) = \|T_n\|_\epsilon^{1/n} \Rightarrow \|T_n\|_\epsilon \geq C(\epsilon)^n$$

an inequality of Szegő with a new proof (not that his proof was complicated).

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Schiefermayr's Theorem

Similarly if  $\epsilon \subset \mathbb{R}$ , we define

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Schiefermayr's Theorem

Similarly if  $\epsilon \subset \mathbb{R}$ , we define

$$\mathfrak{e}_n = \{z \in \mathbb{C} \mid T_n(z) \in [-\|T_n\|_\epsilon, \|T_n\|_\epsilon]\}$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Schiefermayr's Theorem

Similarly if  $\epsilon \subset \mathbb{R}$ , we define

$$\mathfrak{e}_n = \{z \in \mathbb{C} \mid T_n(z) \in [-\|T_n\|_\epsilon, \|T_n\|_\epsilon]\}$$

which the alternation theorem implies is a union of  $n$  intervals in which  $T_n$  is monotone (the intervals can touch).

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Schiefermayr's Theorem

Similarly if  $\epsilon \subset \mathbb{R}$ , we define

$$\mathfrak{e}_n = \{z \in \mathbb{C} \mid T_n(z) \in [-\|T_n\|_\epsilon, \|T_n\|_\epsilon]\}$$

which the alternation theorem implies is a union of  $n$  intervals in which  $T_n$  is monotone (the intervals can touch). As in the complex case, we have that

$$C(\epsilon) \leq C(\mathfrak{e}_n)$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Schiefermayr's Theorem

Similarly if  $\epsilon \subset \mathbb{R}$ , we define

$$\mathfrak{e}_n = \{z \in \mathbb{C} \mid T_n(z) \in [-\|T_n\|_\epsilon, \|T_n\|_\epsilon]\}$$

which the alternation theorem implies is a union of  $n$  intervals in which  $T_n$  is monotone (the intervals can touch). As in the complex case, we have that

$$C(\epsilon) \leq C(\mathfrak{e}_n)$$

but now the Green's function is

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics



# Schiefermayr's Theorem

Similarly if  $\epsilon \subset \mathbb{R}$ , we define

$$\mathfrak{e}_n = \{z \in \mathbb{C} \mid T_n(z) \in [-\|T_n\|_\epsilon, \|T_n\|_\epsilon]\}$$

which the alternation theorem implies is a union of  $n$  intervals in which  $T_n$  is monotone (the intervals can touch). As in the complex case, we have that

$$C(\epsilon) \leq C(\mathfrak{e}_n)$$

but now the Green's function is

$$G_{\mathfrak{e}_n}(z) = \frac{1}{n} \log \left| \left( \frac{T_n(z)}{\|T_n\|_\epsilon} + \sqrt{\frac{T_n(z)^2}{\|T_n\|_\epsilon^2} - 1} \right) \right|$$

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics



# Schiefermayr's Theorem

Similarly if  $\epsilon \subset \mathbb{R}$ , we define

$$\mathfrak{e}_n = \{z \in \mathbb{C} \mid T_n(z) \in [-\|T_n\|_\epsilon, \|T_n\|_\epsilon]\}$$

which the alternation theorem implies is a union of  $n$  intervals in which  $T_n$  is monotone (the intervals can touch). As in the complex case, we have that

$$C(\epsilon) \leq C(\mathfrak{e}_n)$$

but now the Green's function is

$$G_{\mathfrak{e}_n}(z) = \frac{1}{n} \log \left| \left( \frac{T_n(z)}{\|T_n\|_\epsilon} + \sqrt{\frac{T_n(z)^2}{\|T_n\|_\epsilon^2} - 1} \right) \right|$$

For  $z$  near  $\infty$  the argument inside the log is close to  $2z^n/\|T_n\|_\epsilon$  which leads to

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Schiefermayr's Theorem

Similarly if  $\epsilon \subset \mathbb{R}$ , we define

$$\mathfrak{e}_n = \{z \in \mathbb{C} \mid T_n(z) \in [-\|T_n\|_\epsilon, \|T_n\|_\epsilon]\}$$

which the alternation theorem implies is a union of  $n$  intervals in which  $T_n$  is monotone (the intervals can touch). As in the complex case, we have that

$$C(\epsilon) \leq C(\mathfrak{e}_n)$$

but now the Green's function is

$$G_{\mathfrak{e}_n}(z) = \frac{1}{n} \log \left| \left( \frac{T_n(z)}{\|T_n\|_\epsilon} + \sqrt{\frac{T_n(z)^2}{\|T_n\|_\epsilon^2} - 1} \right) \right|$$

For  $z$  near  $\infty$  the argument inside the log is close to  $2z^n/\|T_n\|_\epsilon$  which leads to

$$C(\mathfrak{e}_n) = (\|T_n\|_\epsilon/2)^{1/n} \Rightarrow \|T_n\|_\epsilon \geq 2C(\epsilon)^n$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Schiefermayr's Theorem

Similarly if  $\epsilon \subset \mathbb{R}$ , we define

$$\mathfrak{e}_n = \{z \in \mathbb{C} \mid T_n(z) \in [-\|T_n\|_\epsilon, \|T_n\|_\epsilon]\}$$

which the alternation theorem implies is a union of  $n$  intervals in which  $T_n$  is monotone (the intervals can touch). As in the complex case, we have that

$$C(\epsilon) \leq C(\mathfrak{e}_n)$$

but now the Green's function is

$$G_{\mathfrak{e}_n}(z) = \frac{1}{n} \log \left| \left( \frac{T_n(z)}{\|T_n\|_\epsilon} + \sqrt{\frac{T_n(z)^2}{\|T_n\|_\epsilon^2} - 1} \right) \right|$$

For  $z$  near  $\infty$  the argument inside the log is close to  $2z^n/\|T_n\|_\epsilon$  which leads to

$$C(\mathfrak{e}_n) = (\|T_n\|_\epsilon/2)^{1/n} \Rightarrow \|T_n\|_\epsilon \geq 2C(\epsilon)^n$$

an inequality of Schiefermayr with a new and simpler proof.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Schiefermayr's Theorem

The harmonic measure of a set  $\epsilon \subset \mathbb{R}$  is the boundary value of the harmonic conjugate of the Green's function

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Schiefermayr's Theorem

The harmonic measure of a set  $\epsilon \subset \mathbb{R}$  is the boundary value of the harmonic conjugate of the Green's function (a formula called the Thouless formula by physicists after the recent Nobel Laureate, David Thouless).

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Schiefermayr's Theorem

The harmonic measure of a set  $\epsilon \subset \mathbb{R}$  is the boundary value of the harmonic conjugate of the Green's function (a formula called the Thouless formula by physicists after the recent Nobel Laureate, David Thouless). That shows that each of the sets between two opposite sign extrema of  $T_n$  has  $\epsilon_n$ -harmonic measure  $1/n$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Schiefermayr's Theorem

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

The harmonic measure of a set  $\epsilon \subset \mathbb{R}$  is the boundary value of the harmonic conjugate of the Green's function (a formula called the Thouless formula by physicists after the recent Nobel Laureate, David Thouless). That shows that each of the sets between two opposite sign extrema of  $T_n$  has  $\epsilon_n$ –harmonic measure  $1/n$ . This, in turn implies each connected component of  $\epsilon_n$  has harmonic measure  $k/n$  for some integer  $k$ . Such a set is called a *period-n set*.



# Schiefermayr's Theorem

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics

The harmonic measure of a set  $\epsilon \subset \mathbb{R}$  is the boundary value of the harmonic conjugate of the Green's function (a formula called the Thouless formula by physicists after the recent Nobel Laureate, David Thouless). That shows that each of the sets between two opposite sign extrema of  $T_n$  has  $\epsilon_n$ –harmonic measure  $1/n$ . This, in turn implies each connected component of  $\epsilon_n$  has harmonic measure  $k/n$  for some integer  $k$ . Such a set is called a *period–n set*.

If  $\epsilon$  is a period– $n$  set, one can prove that  $\epsilon_n = \epsilon$  so that all the period– $n$  sets are precisely the possible  $\epsilon_n$ 's.



# Example

**Example ( $\partial\mathbb{D}$ , the unit circle)**

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Example

**Example ( $\partial\mathbb{D}$ , the unit circle)** Its Green's function is  $\log|z|$  so  $R(\mathfrak{e}) = 0$  and  $C(\mathfrak{e}) = 1$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Example

**Example** ( $\partial\mathbb{D}$ , the unit circle) Its Green's function is  $\log|z|$  so  $R(\mathfrak{e}) = 0$  and  $C(\mathfrak{e}) = 1$ . Since  $T_n$  is monic

$$\int_0^{2\pi} \exp(-in\theta) T_n(\exp(i\theta)) d\theta / 2\pi = 1$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Example

**Example** ( $\partial\mathbb{D}$ , the unit circle) Its Green's function is  $\log|z|$  so  $R(\mathfrak{e}) = 0$  and  $C(\mathfrak{e}) = 1$ . Since  $T_n$  is monic

$$\int_0^{2\pi} \exp(-in\theta) T_n(\exp(i\theta)) d\theta / 2\pi = 1$$

we see that  $\|T_n\|_{\mathfrak{e}} \geq 1$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Example

**Example** ( $\partial\mathbb{D}$ , the unit circle) Its Green's function is  $\log|z|$  so  $R(\mathfrak{e}) = 0$  and  $C(\mathfrak{e}) = 1$ . Since  $T_n$  is monic

$$\int_0^{2\pi} \exp(-in\theta) T_n(\exp(i\theta)) d\theta / 2\pi = 1$$

we see that  $\|T_n\|_{\mathfrak{e}} \geq 1$  so that

$$T_n(z) = z^n; \quad \|T_n\|_{\mathfrak{e}} = 1 = C(\mathfrak{e})^n$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Example

**Example** ( $\partial\mathbb{D}$ , the unit circle) Its Green's function is  $\log|z|$  so  $R(\mathfrak{e}) = 0$  and  $C(\mathfrak{e}) = 1$ . Since  $T_n$  is monic

$$\int_0^{2\pi} \exp(-in\theta) T_n(\exp(i\theta)) d\theta / 2\pi = 1$$

we see that  $\|T_n\|_{\mathfrak{e}} \geq 1$  so that

$$T_n(z) = z^n; \quad \|T_n\|_{\mathfrak{e}} = 1 = C(\mathfrak{e})^n$$

**Example** ( $[-1, 1]$ )

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Example

**Example** ( $\partial\mathbb{D}$ , the unit circle) Its Green's function is  $\log|z|$  so  $R(\mathfrak{e}) = 0$  and  $C(\mathfrak{e}) = 1$ . Since  $T_n$  is monic

$$\int_0^{2\pi} \exp(-in\theta) T_n(\exp(i\theta)) d\theta / 2\pi = 1$$

we see that  $\|T_n\|_{\mathfrak{e}} \geq 1$  so that

$$T_n(z) = z^n; \quad \|T_n\|_{\mathfrak{e}} = 1 = C(\mathfrak{e})^n$$

**Example** ( $[-1, 1]$ ) It is known (and follows from results later) that  $C(\mathfrak{e}) = \frac{1}{2}$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Example

**Example** ( $\partial\mathbb{D}$ , the unit circle) Its Green's function is  $\log|z|$  so  $R(\mathfrak{e}) = 0$  and  $C(\mathfrak{e}) = 1$ . Since  $T_n$  is monic

$$\int_0^{2\pi} \exp(-in\theta) T_n(\exp(i\theta)) d\theta / 2\pi = 1$$

we see that  $\|T_n\|_{\mathfrak{e}} \geq 1$  so that

$$T_n(z) = z^n; \quad \|T_n\|_{\mathfrak{e}} = 1 = C(\mathfrak{e})^n$$

**Example** ( $[-1, 1]$ ) It is known (and follows from results later) that  $C(\mathfrak{e}) = \frac{1}{2}$ . By the Alternation Theorem, the polynomials given by  $Q_n(\cos(\theta)) = \cos(n\theta)$  (i.e. “*the Chebyshev polynomials of the first kind*”) are multiples of Chebyshev polynomials as we’ve defined them, so

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom’s Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Example

**Example** ( $\partial\mathbb{D}$ , the unit circle) Its Green's function is  $\log|z|$  so  $R(\mathfrak{e}) = 0$  and  $C(\mathfrak{e}) = 1$ . Since  $T_n$  is monic

$$\int_0^{2\pi} \exp(-in\theta) T_n(\exp(i\theta)) d\theta / 2\pi = 1$$

we see that  $\|T_n\|_{\mathfrak{e}} \geq 1$  so that

$$T_n(z) = z^n; \quad \|T_n\|_{\mathfrak{e}} = 1 = C(\mathfrak{e})^n$$

**Example** ( $[-1, 1]$ ) It is known (and follows from results later) that  $C(\mathfrak{e}) = \frac{1}{2}$ . By the Alternation Theorem, the polynomials given by  $Q_n(\cos(\theta)) = \cos(n\theta)$  (i.e. “*the Chebyshev polynomials of the first kind*”) are multiples of Chebyshev polynomials as we’ve defined them, so

$$T_n(\cos(\theta)) = 2^{-n+1} \cos(n\theta); \quad \|T_n\|_{\mathfrak{e}} = 2^{-n+1} = 2C(\mathfrak{e})^n$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom’s Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Example

**Example** ( $\partial\mathbb{D}$ , the unit circle) Its Green's function is  $\log|z|$  so  $R(\mathfrak{e}) = 0$  and  $C(\mathfrak{e}) = 1$ . Since  $T_n$  is monic

$$\int_0^{2\pi} \exp(-in\theta) T_n(\exp(i\theta)) d\theta / 2\pi = 1$$

we see that  $\|T_n\|_{\mathfrak{e}} \geq 1$  so that

$$T_n(z) = z^n; \quad \|T_n\|_{\mathfrak{e}} = 1 = C(\mathfrak{e})^n$$

**Example** ( $[-1, 1]$ ) It is known (and follows from results later) that  $C(\mathfrak{e}) = \frac{1}{2}$ . By the Alternation Theorem, the polynomials given by  $Q_n(\cos(\theta)) = \cos(n\theta)$  (i.e. “the Chebyshev polynomials of the first kind”) are multiples of Chebyshev polynomials as we’ve defined them, so

$$T_n(\cos(\theta)) = 2^{-n+1} \cos(n\theta); \quad \|T_n\|_{\mathfrak{e}} = 2^{-n+1} = 2C(\mathfrak{e})^n$$

so one can have equality in both lower bounds.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom’s Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# FFS Theorem

**Theorem (Faber–Fekete–Szegő Theorem)** For any compact subset  $\mathfrak{e} \subset \mathbb{C}$ , we have that

$$\lim_{n \rightarrow \infty} \|T_n\|_{\mathfrak{e}}^{1/n} = C(\mathfrak{e})$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# FFS Theorem

**Theorem (Faber–Fekete–Szegő Theorem)** For any compact subset  $\mathfrak{e} \subset \mathbb{C}$ , we have that

$$\lim_{n \rightarrow \infty} \|T_n\|_{\mathfrak{e}}^{1/n} = C(\mathfrak{e})$$

Given Szegő's lower bound, we get a lower bound on the  $\liminf$  by  $C(\mathfrak{e})$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# FFS Theorem

**Theorem (Faber–Fekete–Szegő Theorem)** For any compact subset  $\mathfrak{e} \subset \mathbb{C}$ , we have that

$$\lim_{n \rightarrow \infty} \|T_n\|_{\mathfrak{e}}^{1/n} = C(\mathfrak{e})$$

Given Szegő's lower bound, we get a lower bound on the  $\liminf$  by  $C(\mathfrak{e})$ . One can get an upper bound on  $\|T_n\|_{\mathfrak{e}}^{1/n}$  by

$$\sup_{z_j \in \mathfrak{e}} \prod_{1 \leq j \neq k \leq n+1} |z_j - z_k|^{1/n(n+1)}$$

using suitable trial monic polynomials.

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics



# FFS Theorem

**Theorem (Faber–Fekete–Szegő Theorem)** For any compact subset  $\epsilon \subset \mathbb{C}$ , we have that

$$\lim_{n \rightarrow \infty} \|T_n\|_{\epsilon}^{1/n} = C(\epsilon)$$

Given Szegő's lower bound, we get a lower bound on the  $\liminf$  by  $C(\epsilon)$ . One can get an upper bound on  $\|T_n\|_{\epsilon}^{1/n}$  by

$$\sup_{z_j \in \epsilon} \prod_{1 \leq j \neq k \leq n+1} |z_j - z_k|^{1/n(n+1)}$$

using suitable trial monic polynomials. Fekete proved that as  $n \rightarrow \infty$ , this last quantity had a limit that he called the *transfinite diameter*.

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics



# FFS Theorem

**Theorem (Faber–Fekete–Szegő Theorem)** For any compact subset  $\epsilon \subset \mathbb{C}$ , we have that

$$\lim_{n \rightarrow \infty} \|T_n\|_{\epsilon}^{1/n} = C(\epsilon)$$

Given Szegő's lower bound, we get a lower bound on the  $\liminf$  by  $C(\epsilon)$ . One can get an upper bound on  $\|T_n\|_{\epsilon}^{1/n}$  by

$$\sup_{z_j \in \epsilon} \prod_{1 \leq j \neq k \leq n+1} |z_j - z_k|^{1/n(n+1)}$$

using suitable trial monic polynomials. Fekete proved that as  $n \rightarrow \infty$ , this last quantity had a limit that he called the *transfinite diameter*. One can view this sup as the exponential of the negative of a discrete Coulomb energy of  $n + 1$  point charges, each of charge about  $\frac{1}{n+1}$ , so Szegő's proof that this is  $C(\epsilon)$  is natural from a Coulomb energy point of view.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# History

Faber's name is associated to this theorem because of a 1919 paper in which he proved a result that is both much more restrictive and much stronger than what we call the FFS Theorem.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

**Faber Fekete  
Szegő Theorem**

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# History

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Faber's name is associated to this theorem because of a 1919 paper in which he proved a result that is both much more restrictive and much stronger than what we call the FFS Theorem. It is more restrictive in that he only studied the special case where  $\epsilon$  is a single (closed) analytic Jordan curve.



# History

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Faber's name is associated to this theorem because of a 1919 paper in which he proved a result that is both much more restrictive and much stronger than what we call the FFS Theorem. It is more restrictive in that he only studied the special case where  $\epsilon$  is a single (closed) analytic Jordan curve. But in this case, he proved much more — first he proved that  $\lim_{n \rightarrow \infty} \|T_n\|_\epsilon / C(\epsilon)^n = 1$ .



# History

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Faber's name is associated to this theorem because of a 1919 paper in which he proved a result that is both much more restrictive and much stronger than what we call the FFS Theorem. It is more restrictive in that he only studied the special case where  $\epsilon$  is a single (closed) analytic Jordan curve. But in this case, he proved much more — first he proved that  $\lim_{n \rightarrow \infty} \|T_n\|_\epsilon / C(\epsilon)^n = 1$ .

He also obtained asymptotics for the polynomials themselves. The unbounded component,  $\Omega$ , of  $(\mathbb{C} \cup \{\infty\}) \setminus \epsilon$  is simply connected,



# History

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Faber's name is associated to this theorem because of a 1919 paper in which he proved a result that is both much more restrictive and much stronger than what we call the FFS Theorem. It is more restrictive in that he only studied the special case where  $\epsilon$  is a single (closed) analytic Jordan curve. But in this case, he proved much more — first he proved that  $\lim_{n \rightarrow \infty} \|T_n\|_\epsilon / C(\epsilon)^n = 1$ .

He also obtained asymptotics for the polynomials themselves. The unbounded component,  $\Omega$ , of  $(\mathbb{C} \cup \{\infty\}) \setminus \epsilon$  is simply connected, so  $G_\epsilon(z)$  has a single valued harmonic conjugate



# History

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Faber's name is associated to this theorem because of a 1919 paper in which he proved a result that is both much more restrictive and much stronger than what we call the FFS Theorem. It is more restrictive in that he only studied the special case where  $\epsilon$  is a single (closed) analytic Jordan curve. But in this case, he proved much more — first he proved that  $\lim_{n \rightarrow \infty} \|T_n\|_\epsilon / C(\epsilon)^n = 1$ .

He also obtained asymptotics for the polynomials themselves. The unbounded component,  $\Omega$ , of  $(\mathbb{C} \cup \{\infty\}) \setminus \epsilon$  is simply connected, so  $G_\epsilon(z)$  has a single valued harmonic conjugate and thus, by exponentiating, there is a function,  $B_\epsilon(z)$ , on  $\Omega$  with  $|B_\epsilon(z)| = \exp(-G_\epsilon(z))$  with an overall phase determined by demanding that as  $z \rightarrow \infty$ , we have that  $B_\epsilon(z)^{-1} = \frac{z}{C(\epsilon)} + O(1)$ .



# History

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Faber's name is associated to this theorem because of a 1919 paper in which he proved a result that is both much more restrictive and much stronger than what we call the FFS Theorem. It is more restrictive in that he only studied the special case where  $\epsilon$  is a single (closed) analytic Jordan curve. But in this case, he proved much more — first he proved that  $\lim_{n \rightarrow \infty} \|T_n\|_\epsilon / C(\epsilon)^n = 1$ .

He also obtained asymptotics for the polynomials themselves. The unbounded component,  $\Omega$ , of  $(\mathbb{C} \cup \{\infty\}) \setminus \epsilon$  is simply connected, so  $G_\epsilon(z)$  has a single valued harmonic conjugate and thus, by exponentiating, there is a function,  $B_\epsilon(z)$ , on  $\Omega$  with  $|B_\epsilon(z)| = \exp(-G_\epsilon(z))$  with an overall phase determined by demanding that as  $z \rightarrow \infty$ , we have that  $B_\epsilon(z)^{-1} = \frac{z}{C(\epsilon)} + O(1)$ . Since the curve is analytic,  $B_\epsilon(z)$  has an analytic continuation to a neighborhood of  $\epsilon$ .



# History

Faber proved that uniformly on  $\Omega$  plus a neighborhood of  $\epsilon$ ,  
 $T_n(z)B_\epsilon(z)^n \rightarrow 1$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

**Faber Fekete  
Szegő Theorem**

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# History

Faber proved that uniformly on  $\Omega$  plus a neighborhood of  $\epsilon$ ,  
 $T_n(z)B_\epsilon(z)^n \rightarrow 1$ . Faber didn't mention Green's functions  
or capacities at all!

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

**Faber Fekete  
Szegő Theorem**

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# History

Faber proved that uniformly on  $\Omega$  plus a neighborhood of  $\epsilon$ ,  $T_n(z)B_\epsilon(z)^n \rightarrow 1$ . Faber didn't mention Green's functions or capacities at all! In this case,  $B_\epsilon(z)^{-1}$  can be described as the Riemann map of  $\Omega$  to  $(\mathbb{C} \cup \{\infty\} \setminus \overline{\mathbb{D}})$  (with positive “derivative” at  $\infty$ ) and the capacity appears as inverse of the value of that “derivative”.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

**Faber Fekete  
Szegő Theorem**

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# History

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Faber proved that uniformly on  $\Omega$  plus a neighborhood of  $\epsilon$ ,  $T_n(z)B_\epsilon(z)^n \rightarrow 1$ . Faber didn't mention Green's functions or capacities at all! In this case,  $B_\epsilon(z)^{-1}$  can be described as the Riemann map of  $\Omega$  to  $(\mathbb{C} \cup \{\infty\} \setminus \overline{\mathbb{D}})$  (with positive “derivative” at  $\infty$ ) and the capacity appears as inverse of the value of that “derivative”.

Interestingly enough, for these polynomials, Faber had “Szegő asymptotics” three years before Szegő had his asymptotics (for OPUC, not Chebyshev polynomials).



# History

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Faber proved that uniformly on  $\Omega$  plus a neighborhood of  $\epsilon$ ,  $T_n(z)B_\epsilon(z)^n \rightarrow 1$ . Faber didn't mention Green's functions or capacities at all! In this case,  $B_\epsilon(z)^{-1}$  can be described as the Riemann map of  $\Omega$  to  $(\mathbb{C} \cup \{\infty\} \setminus \overline{\mathbb{D}})$  (with positive “derivative” at  $\infty$ ) and the capacity appears as inverse of the value of that “derivative”.

Interestingly enough, for these polynomials, Faber had “Szegő asymptotics” three years before Szegő had his asymptotics (for OPUC, not Chebyshev polynomials).

Fekete's work on transfinite diameters and its connection to capacity for some special cases is from 1923.



# History

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Faber proved that uniformly on  $\Omega$  plus a neighborhood of  $\epsilon$ ,  $T_n(z)B_\epsilon(z)^n \rightarrow 1$ . Faber didn't mention Green's functions or capacities at all! In this case,  $B_\epsilon(z)^{-1}$  can be described as the Riemann map of  $\Omega$  to  $(\mathbb{C} \cup \{\infty\} \setminus \overline{\mathbb{D}})$  (with positive “derivative” at  $\infty$ ) and the capacity appears as inverse of the value of that “derivative”.

Interestingly enough, for these polynomials, Faber had “Szegő asymptotics” three years before Szegő had his asymptotics (for OPUC, not Chebyshev polynomials).

Fekete's work on transfinite diameters and its connection to capacity for some special cases is from 1923. Szegő had the full theorem in a 1924 paper whose title started “Comments on a paper by Mr. M. Fekete”.



# Character Automorphic Functions

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

In 1969, Widom published a 100+ page brilliant, seminal work on asymptotics of Chebyshev and orthogonal polynomials. In his set up,  $\epsilon$  is a finite union of (closed) analytic Jordan curves and/or (open) Jordan arcs.



# Character Automorphic Functions

In 1969, Widom published a 100+ page brilliant, seminal work on asymptotics of Chebyshev and orthogonal polynomials. In his set up,  $\epsilon$  is a finite union of (closed) analytic Jordan curves and/or (open) Jordan arcs. The cases with  $\epsilon \subset \mathbb{R}$  are exactly the finite gap sets.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Character Automorphic Functions

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

In 1969, Widom published a 100+ page brilliant, seminal work on asymptotics of Chebyshev and orthogonal polynomials. In his set up,  $\epsilon$  is a finite union of (closed) analytic Jordan curves and/or (open) Jordan arcs. The cases with  $\epsilon \subset \mathbb{R}$  are exactly the finite gap sets.

As in the work of Faber, it is natural to look for an analytic function,  $B_\epsilon(z)$ , with  $|B_\epsilon(z)| = \exp(-G_\epsilon(z))$  on  $\Omega$ , the unbounded component of  $(\mathbb{C} \cup \{\infty\}) \setminus \epsilon$ .



# Character Automorphic Functions

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics

In 1969, Widom published a 100+ page brilliant, seminal work on asymptotics of Chebyshev and orthogonal polynomials. In his set up,  $\epsilon$  is a finite union of (closed) analytic Jordan curves and/or (open) Jordan arcs. The cases with  $\epsilon \subset \mathbb{R}$  are exactly the finite gap sets.

As in the work of Faber, it is natural to look for an analytic function,  $B_\epsilon(z)$ , with  $|B_\epsilon(z)| = \exp(-G_\epsilon(z))$  on  $\Omega$ , the unbounded component of  $(\mathbb{C} \cup \{\infty\}) \setminus \epsilon$ . The problem is that  $\Omega$  is no longer simply connected so the magnitude of  $B_\epsilon(z)$  is single valued but its phase is multivalued.



# Character Automorphic Functions

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

In 1969, Widom published a 100+ page brilliant, seminal work on asymptotics of Chebyshev and orthogonal polynomials. In his set up,  $\epsilon$  is a finite union of (closed) analytic Jordan curves and/or (open) Jordan arcs. The cases with  $\epsilon \subset \mathbb{R}$  are exactly the finite gap sets.

As in the work of Faber, it is natural to look for an analytic function,  $B_\epsilon(z)$ , with  $|B_\epsilon(z)| = \exp(-G_\epsilon(z))$  on  $\Omega$ , the unbounded component of  $(\mathbb{C} \cup \{\infty\}) \setminus \epsilon$ . The problem is that  $\Omega$  is no longer simply connected so the magnitude of  $B_\epsilon(z)$  is single valued but its phase is multivalued.

Put differently,  $B_\epsilon(z)$  can be continued along any curve in  $\Omega$  and there is a map from the fundamental group of  $\Omega$  to  $\partial\mathbb{D}$ , which is a character (i.e. group homomorphism), so that after continuation around a closed curve,  $B_\epsilon(z)$  is multiplied by the character applied to that curve.



# Character Automorphic Functions

Indeed, if the curve loops around a subset  $\mathfrak{g} \subset \mathfrak{e}$ , the phase changes by  $-2\pi\rho_{\mathfrak{e}}(\mathfrak{g})$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Character Automorphic Functions

Indeed, if the curve loops around a subset  $\mathfrak{g} \subset \mathfrak{e}$ , the phase changes by  $-2\pi\rho_{\mathfrak{e}}(\mathfrak{g})$ .

If  $T_n(z)B_{\mathfrak{e}}(z)^nC(\mathfrak{e})^{-n}$  had a limit, that limit cannot be  $n$  independent since the character is  $n$  dependent.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Character Automorphic Functions

Indeed, if the curve loops around a subset  $\mathfrak{g} \subset \mathfrak{e}$ , the phase changes by  $-2\pi\rho_{\mathfrak{e}}(\mathfrak{g})$ .

If  $T_n(z)B_{\mathfrak{e}}(z)^nC(\mathfrak{e})^{-n}$  had a limit, that limit cannot be n independent since the character is n dependent. Widom had the idea that there should be functions  $F_{\chi}(z)$  defined for each  $\chi$  in the character group and continuous in  $\chi$  so the limit is the  $F_{\chi}$ , call it  $F_n$ , associated to the character of  $B_{\mathfrak{e}}(z)^n$ . As a function of n, the limit will be almost periodic!

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Widom's Minimizers

He even found a candidate for the functions! Let  $F_\chi(z)$  be that function among all character automorphic functions,  $A(z)$ , on  $\Omega$  with character  $\chi$  and with  $A(\infty) = 1$ , that minimizes  $\sup_{z \in \Omega} \{|A(z)|\}$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Widom's Minimizers

He even found a candidate for the functions! Let  $F_\chi(z)$  be that function among all character automorphic functions,  $A(z)$ , on  $\Omega$  with character  $\chi$  and with  $A(\infty) = 1$ , that minimizes  $\sup_{z \in \Omega} \{|A(z)|\}$ .

Widom proved uniqueness of the minimizer and found a formula for it (in terms of some theta functions and solutions of some implicit equations).

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Widom's Minimizers

He even found a candidate for the functions! Let  $F_\chi(z)$  be that function among all character automorphic functions,  $A(z)$ , on  $\Omega$  with character  $\chi$  and with  $A(\infty) = 1$ , that minimizes  $\sup_{z \in \Omega} \{|A(z)|\}$ .

Widom proved uniqueness of the minimizer and found a formula for it (in terms of some theta functions and solutions of some implicit equations). He also proved that  $\|F_\chi\|_\Omega$  is continuous in  $\chi$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Widom's Minimizers

He even found a candidate for the functions! Let  $F_\chi(z)$  be that function among all character automorphic functions,  $A(z)$ , on  $\Omega$  with character  $\chi$  and with  $A(\infty) = 1$ , that minimizes  $\sup_{z \in \Omega} \{|A(z)|\}$ .

Widom proved uniqueness of the minimizer and found a formula for it (in terms of some theta functions and solutions of some implicit equations). He also proved that  $\|F_\chi\|_\Omega$  is continuous in  $\chi$ . Because of the uniqueness, one can prove that the functions,  $F_\chi(z)$ , defined for  $z \in \Omega$ , are continuous in  $\chi$  on the compact set of characters, uniformly locally in  $z$  (but as functions on the covering space not uniformly in all  $z$ ).

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Widom's Theorems and Conjecture

The Widom minimizers are analogs of the Ahlfors function for which Fisher found a simple elegant proof of uniqueness.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Widom's Theorems and Conjecture

The Widom minimizers are analogs of the Ahlfors function for which Fisher found a simple elegant proof of uniqueness. By modifying his proof, in Part 2 we find a simple proof of uniqueness for any compact set  $\mathfrak{e} \subset \mathbb{C}$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Widom's Theorems and Conjecture

The Widom minimizers are analogs of the Ahlfors function for which Fisher found a simple elegant proof of uniqueness. By modifying his proof, in Part 2 we find a simple proof of uniqueness for any compact set  $\epsilon \subset \mathbb{C}$ .

**Theorem (Widom)** Let  $\epsilon$  be a finite union of disjoint analytic Jordan curves. Let  $F_n(z)$  be as above for the character of  $B_\epsilon(z)^n$ . Then:

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Widom's Theorems and Conjecture

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
  - Totik Widom Bounds
  - Main Results
  - Totik Widom Bounds
  - Szegő–Widom Asymptotics

The Widom minimizers are analogs of the Ahlfors function for which Fisher found a simple elegant proof of uniqueness. By modifying his proof, in Part 2 we find a simple proof of uniqueness for any compact set  $\epsilon \subset \mathbb{C}$ .

**Theorem (Widom)** Let  $\epsilon$  be a finite union of disjoint analytic Jordan curves. Let  $F_n(z)$  be as above for the character of  $B_\epsilon(z)^n$ . Then:

$$\lim_{n \rightarrow \infty} \frac{\|T_n\|_\epsilon}{C(\epsilon)^n \|F_n\|_\Omega} = 1; \quad \lim_{n \rightarrow \infty} \left[ \frac{T_n(z) B_\epsilon(z)^n}{C(\epsilon)^n} - F_n(z) \right] = 0$$

where the limit is uniform on compact subsets of  $\Omega$ .



# Widom's Theorems and Conjecture

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
  - Totik Widom Bounds
  - Main Results
  - Totik Widom Bounds
  - Szegő–Widom Asymptotics

The Widom minimizers are analogs of the Ahlfors function for which Fisher found a simple elegant proof of uniqueness. By modifying his proof, in Part 2 we find a simple proof of uniqueness for any compact set  $\epsilon \subset \mathbb{C}$ .

**Theorem (Widom)** Let  $\epsilon$  be a finite union of disjoint analytic Jordan curves. Let  $F_n(z)$  be as above for the character of  $B_\epsilon(z)^n$ . Then:

$$\lim_{n \rightarrow \infty} \frac{\|T_n\|_\epsilon}{C(\epsilon)^n \|F_n\|_\Omega} = 1; \quad \lim_{n \rightarrow \infty} \left[ \frac{T_n(z) B_\epsilon(z)^n}{C(\epsilon)^n} - F_n(z) \right] = 0$$

where the limit is uniform on compact subsets of  $\Omega$ .

Since  $|B_\epsilon(z)| \rightarrow 1$  and  $\|F_n\|_\Omega$  is taken as  $z \rightarrow \epsilon$ , the  $z$  asymptotics and norm limit fit together.



# Widom's Theorems and Conjecture

**Theorem (Widom)** Let  $\epsilon$  be a finite gap subset of  $\mathbb{R}$ . Let  $F_n(z)$  be as above. Then

$$\lim_{n \rightarrow \infty} \frac{\|T_n\|_\epsilon}{2C(\epsilon)^n \|F_n\|_\Omega} = 1$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Widom's Theorems and Conjecture

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

**Theorem (Widom)** Let  $\epsilon$  be a finite gap subset of  $\mathbb{R}$ . Let  $F_n(z)$  be as above. Then

$$\lim_{n \rightarrow \infty} \frac{\|T_n\|_\epsilon}{2C(\epsilon)^n \|F_n\|_\Omega} = 1$$

**Conjecture (Widom)** Let  $\epsilon$  be a finite gap subset of  $\mathbb{R}$ . Let  $F_n(z)$  be as above. Then:

$$\lim_{n \rightarrow \infty} \left[ \frac{T_n(z)B_\epsilon(z)^n}{C(\epsilon)^n} - F_n(z) \right] = 0$$

uniformly on compact subsets of  $\Omega$ .



# Widom's Theorems and Conjecture

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics

**Theorem (Widom)** Let  $\epsilon$  be a finite gap subset of  $\mathbb{R}$ . Let  $F_n(z)$  be as above. Then

$$\lim_{n \rightarrow \infty} \frac{\|T_n\|_\epsilon}{2C(\epsilon)^n \|F_n\|_\Omega} = 1$$

**Conjecture (Widom)** Let  $\epsilon$  be a finite gap subset of  $\mathbb{R}$ . Let  $F_n(z)$  be as above. Then:

$$\lim_{n \rightarrow \infty} \left[ \frac{T_n(z) B_\epsilon(z)^n}{C(\epsilon)^n} - F_n(z) \right] = 0$$

uniformly on compact subsets of  $\Omega$ .

The norm,  $\|T_n\|_\epsilon$  is twice as large as one might expect!



# Widom's Theorems and Conjecture

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

**Theorem (Widom)** Let  $\epsilon$  be a finite gap subset of  $\mathbb{R}$ . Let  $F_n(z)$  be as above. Then

$$\lim_{n \rightarrow \infty} \frac{\|T_n\|_\epsilon}{2C(\epsilon)^n \|F_n\|_\Omega} = 1$$

**Conjecture (Widom)** Let  $\epsilon$  be a finite gap subset of  $\mathbb{R}$ . Let  $F_n(z)$  be as above. Then:

$$\lim_{n \rightarrow \infty} \left[ \frac{T_n(z)B_\epsilon(z)^n}{C(\epsilon)^n} - F_n(z) \right] = 0$$

uniformly on compact subsets of  $\Omega$ .

The norm,  $\|T_n\|_\epsilon$  is twice as large as one might expect!

Note: This is Widom's conjecture for  $\epsilon \subset \mathbb{R}$ ; he made the conjecture for more general cases of  $\epsilon \subset \mathbb{C}$ .



# Back to $[-1, 1]$

**Example** We return to the case of  $[-1, 1]$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Back to $[-1, 1]$

**Example** We return to the case of  $[-1, 1]$  where  $\Omega$  is simply connected so  $F_n(z) \equiv 1$ . We have that

$$B_{\mathfrak{e}}(z) = z - \sqrt{z^2 - 1}.$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Back to $[-1, 1]$

**Example** We return to the case of  $[-1, 1]$  where  $\Omega$  is simply connected so  $F_n(z) \equiv 1$ . We have that

$$B_{\epsilon}(z) = z - \sqrt{z^2 - 1}. \text{ Notice that } B_{\epsilon}^{-1}(z) = z + \sqrt{z^2 - 1}.$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Back to $[-1, 1]$

**Example** We return to the case of  $[-1, 1]$  where  $\Omega$  is simply connected so  $F_n(z) \equiv 1$ . We have that

$$B_\epsilon(z) = z - \sqrt{z^2 - 1}. \text{ Notice that } B_\epsilon^{-1}(z) = z + \sqrt{z^2 - 1}.$$

On  $[-1, 1]$ , of course,  $B_\epsilon(x)$  has magnitude 1 (since

$$G_\epsilon(x) = 0$$
 so  $B_\epsilon(x) = \exp(i\theta)$  and

$$\cos(\theta) = \frac{1}{2}[B_\epsilon(x) + B_\epsilon^{-1}(x)] = x.$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Back to $[-1, 1]$

**Example** We return to the case of  $[-1, 1]$  where  $\Omega$  is simply connected so  $F_n(z) \equiv 1$ . We have that

$$B_{\epsilon}(z) = z - \sqrt{z^2 - 1}. \text{ Notice that } B_{\epsilon}^{-1}(z) = z + \sqrt{z^2 - 1}.$$

On  $[-1, 1]$ , of course,  $B_{\epsilon}(x)$  has magnitude 1 (since  $G_{\epsilon}(x) = 0$ ) so  $B_{\epsilon}(x) = \exp(i\theta)$  and

$$\cos(\theta) = \frac{1}{2}[B_{\epsilon}(x) + B_{\epsilon}^{-1}(x)] = x.$$

Thus, by  $T_n(\cos(\theta)) = 2^{-n+1} \cos(n\theta)$ , we see that

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Back to $[-1, 1]$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

**Example** We return to the case of  $[-1, 1]$  where  $\Omega$  is simply connected so  $F_n(z) \equiv 1$ . We have that

$B_\epsilon(z) = z - \sqrt{z^2 - 1}$ . Notice that  $B_\epsilon^{-1}(z) = z + \sqrt{z^2 - 1}$ .

On  $[-1, 1]$ , of course,  $B_\epsilon(x)$  has magnitude 1 (since  $G_\epsilon(x) = 0$ ) so  $B_\epsilon(x) = \exp(i\theta)$  and

$$\cos(\theta) = \frac{1}{2}[B_\epsilon(x) + B_\epsilon^{-1}(x)] = x.$$

Thus, by  $T_n(\cos(\theta)) = 2^{-n+1} \cos(n\theta)$ , we see that

$$T_n(z) = 2^{-n}[B_\epsilon^n(z) + B_\epsilon^{-n}(z)].$$



# Back to $[-1, 1]$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

**Example** We return to the case of  $[-1, 1]$  where  $\Omega$  is simply connected so  $F_n(z) \equiv 1$ . We have that

$B_\epsilon(z) = z - \sqrt{z^2 - 1}$ . Notice that  $B_\epsilon^{-1}(z) = z + \sqrt{z^2 - 1}$ .

On  $[-1, 1]$ , of course,  $B_\epsilon(x)$  has magnitude 1 (since  $G_\epsilon(x) = 0$ ) so  $B_\epsilon(x) = \exp(i\theta)$  and

$$\cos(\theta) = \frac{1}{2}[B_\epsilon(x) + B_\epsilon^{-1}(x)] = x.$$

Thus, by  $T_n(\cos(\theta)) = 2^{-n+1} \cos(n\theta)$ , we see that

$T_n(z) = 2^{-n}[B_\epsilon^n(z) + B_\epsilon^{-n}(z)]$ . For  $z \in [-1, 1]$ , both terms contribute and at some points add to 2 and we get

$$\|T_n\|_\epsilon = 2^{-n+1} = 2C(\epsilon)^n.$$



# Back to $[-1, 1]$

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics

**Example** We return to the case of  $[-1, 1]$  where  $\Omega$  is simply connected so  $F_n(z) \equiv 1$ . We have that

$$B_\epsilon(z) = z - \sqrt{z^2 - 1}. \text{ Notice that } B_\epsilon^{-1}(z) = z + \sqrt{z^2 - 1}.$$

On  $[-1, 1]$ , of course,  $B_\epsilon(x)$  has magnitude 1 (since  $G_\epsilon(x) = 0$ ) so  $B_\epsilon(x) = \exp(i\theta)$  and

$$\cos(\theta) = \frac{1}{2}[B_\epsilon(x) + B_\epsilon^{-1}(x)] = x.$$

Thus, by  $T_n(\cos(\theta)) = 2^{-n+1} \cos(n\theta)$ , we see that

$T_n(z) = 2^{-n}[B_\epsilon^n(z) + B_\epsilon^{-n}(z)]$ . For  $z \in [-1, 1]$ , both terms contribute and at some points add to 2 and we get

$\|T_n\|_\epsilon = 2^{-n+1} = 2C(\epsilon)^n$ . On  $\Omega$ ,  $|B_\epsilon(z)| < 1$  so the  $B_\epsilon^n$  term is negligible as  $n \rightarrow \infty$  and we lose the factor of 2.



# Back to $[-1, 1]$

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics

**Example** We return to the case of  $[-1, 1]$  where  $\Omega$  is simply connected so  $F_n(z) \equiv 1$ . We have that

$$B_\epsilon(z) = z - \sqrt{z^2 - 1}. \text{ Notice that } B_\epsilon^{-1}(z) = z + \sqrt{z^2 - 1}.$$

On  $[-1, 1]$ , of course,  $B_\epsilon(x)$  has magnitude 1 (since  $G_\epsilon(x) = 0$ ) so  $B_\epsilon(x) = \exp(i\theta)$  and

$$\cos(\theta) = \frac{1}{2}[B_\epsilon(x) + B_\epsilon^{-1}(x)] = x.$$

Thus, by  $T_n(\cos(\theta)) = 2^{-n+1} \cos(n\theta)$ , we see that

$T_n(z) = 2^{-n}[B_\epsilon^n(z) + B_\epsilon^{-n}(z)]$ . For  $z \in [-1, 1]$ , both terms contribute and at some points add to 2 and we get

$\|T_n\|_\epsilon = 2^{-n+1} = 2C(\epsilon)^n$ . On  $\Omega$ ,  $|B_\epsilon(z)| < 1$  so the  $B_\epsilon^n$  term is negligible as  $n \rightarrow \infty$  and we lose the factor of 2.

It was this example that led Widom to his conjecture.



# Totik–Widom bounds

**Theorem (Totik's  $1/n$  bound)** If  $\epsilon$  is a finite gap set, the period  $n$  sets  $\tilde{\epsilon}_n \supset \epsilon$  can be chosen so that  
$$C(\tilde{\epsilon}_n) \leq C(\epsilon) \left(1 + \frac{E}{n}\right)$$
 for some constant  $E$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

**Totik Widom  
Bounds**

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Totik–Widom bounds

**Theorem** (Totik's  $1/n$  bound) If  $\epsilon$  is a finite gap set, the period  $n$  sets  $\tilde{\epsilon}_n \supset \epsilon$  can be chosen so that  $C(\tilde{\epsilon}_n) \leq C(\epsilon) \left(1 + \frac{E}{n}\right)$  for some constant  $E$ .

Because  $\|T_n\|_{\epsilon} = 2C(\epsilon)^n$ , this bound is equivalent to

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

**Totik Widom  
Bounds**

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Totik–Widom bounds

**Theorem (Totik's  $1/n$  bound)** If  $\epsilon$  is a finite gap set, the period  $n$  sets  $\tilde{\epsilon}_n \supset \epsilon$  can be chosen so that  $C(\tilde{\epsilon}_n) \leq C(\epsilon) \left(1 + \frac{E}{n}\right)$  for some constant  $E$ .

Because  $\|T_n\|_\epsilon = 2C(\epsilon)^n$ , this bound is equivalent to

**Theorem (Totik–Widom bounds in the finite gap case)** If  $\epsilon$  is a finite gap set, then for a constant  $D$  we have that

$$\|T_n\|_\epsilon \leq DC(\epsilon)^n$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Totik–Widom bounds

**Theorem** (Totik's  $1/n$  bound) If  $\epsilon$  is a finite gap set, the period  $n$  sets  $\tilde{\epsilon}_n \supset \epsilon$  can be chosen so that  $C(\tilde{\epsilon}_n) \leq C(\epsilon) \left(1 + \frac{E}{n}\right)$  for some constant  $E$ .

Because  $\|T_n\|_\epsilon = 2C(\epsilon)^n$ , this bound is equivalent to

**Theorem** (*Totik–Widom bounds in the finite gap case*) If  $\epsilon$  is a finite gap set, then for a constant  $D$  we have that

$$\|T_n\|_\epsilon \leq DC(\epsilon)^n$$

This complements the  $2C(\epsilon)^n$  lower bound.

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics



# Totik–Widom bounds

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

**Theorem (Totik's  $1/n$  bound)** If  $\epsilon$  is a finite gap set, the period  $n$  sets  $\tilde{\epsilon}_n \supset \epsilon$  can be chosen so that  $C(\tilde{\epsilon}_n) \leq C(\epsilon) \left(1 + \frac{E}{n}\right)$  for some constant  $E$ .

Because  $\|T_n\|_\epsilon = 2C(\epsilon)^n$ , this bound is equivalent to

**Theorem (Totik–Widom bounds in the finite gap case)** If  $\epsilon$  is a finite gap set, then for a constant  $D$  we have that

$$\|T_n\|_\epsilon \leq DC(\epsilon)^n$$

This complements the  $2C(\epsilon)^n$  lower bound. Because of his asymptotic result, Widom already had this bound in 1969 but Totik's proof was much simpler.



# Totik–Widom bounds

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

**Theorem (Totik's  $1/n$  bound)** If  $\epsilon$  is a finite gap set, the period  $n$  sets  $\tilde{\epsilon}_n \supset \epsilon$  can be chosen so that  $C(\tilde{\epsilon}_n) \leq C(\epsilon) \left(1 + \frac{E}{n}\right)$  for some constant  $E$ .

Because  $\|T_n\|_\epsilon = 2C(\epsilon)^n$ , this bound is equivalent to

**Theorem (Totik–Widom bounds in the finite gap case)** If  $\epsilon$  is a finite gap set, then for a constant  $D$  we have that

$$\|T_n\|_\epsilon \leq DC(\epsilon)^n$$

This complements the  $2C(\epsilon)^n$  lower bound. Because of his asymptotic result, Widom already had this bound in 1969 but Totik's proof was much simpler. Neither proof has very explicit estimates for  $D$ .



# Totik–Widom bounds

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

**Theorem (Totik's  $1/n$  bound)** If  $\epsilon$  is a finite gap set, the period  $n$  sets  $\tilde{\epsilon}_n \supset \epsilon$  can be chosen so that  $C(\tilde{\epsilon}_n) \leq C(\epsilon) \left(1 + \frac{E}{n}\right)$  for some constant  $E$ .

Because  $\|T_n\|_\epsilon = 2C(\epsilon)^n$ , this bound is equivalent to

**Theorem (Totik–Widom bounds in the finite gap case)** If  $\epsilon$  is a finite gap set, then for a constant  $D$  we have that

$$\|T_n\|_\epsilon \leq DC(\epsilon)^n$$

This complements the  $2C(\epsilon)^n$  lower bound. Because of his asymptotic result, Widom already had this bound in 1969 but Totik's proof was much simpler. Neither proof has very explicit estimates for  $D$ . Even though they only had the result for finite gap sets, we will say that a general set  $\epsilon$  has *Totik–Widom bounds*, if there is an upper bound of the above form.



# Canonical Generators

I now want to discuss the case where  $\epsilon$  might have infinitely many components – in the real case, infinitely many gaps.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Canonical Generators

I now want to discuss the case where  $\epsilon$  might have infinitely many components – in the real case, infinitely many gaps. The character group,  $\pi_1^*$ , is in general an infinite dimensional torus.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Canonical Generators

I now want to discuss the case where  $\epsilon$  might have infinitely many components – in the real case, infinitely many gaps. The character group,  $\pi_1^*$ , is in general an infinite dimensional torus. It contains a distinguished element,  $\chi_\epsilon$ , the character of  $B_\epsilon$ . We will say that  $\epsilon \in \mathbb{R}$  has a *canonical generator* if and only if the powers of  $\chi_\epsilon$  are dense in the character group.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Canonical Generators

I now want to discuss the case where  $\epsilon$  might have infinitely many components – in the real case, infinitely many gaps. The character group,  $\pi_1^*$ , is in general an infinite dimensional torus. It contains a distinguished element,  $\chi_\epsilon$ , the character of  $B_\epsilon$ . We will say that  $\epsilon \in \mathbb{R}$  has a *canonical generator* if and only if the powers of  $\chi_\epsilon$  are dense in the character group. This is equivalent to saying that if  $\epsilon$  is decomposed into  $\ell$  closed disjoint sets, the only rational relation among their harmonic masses is that their total sum is 1.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Canonical Generators

I now want to discuss the case where  $\epsilon$  might have infinitely many components – in the real case, infinitely many gaps. The character group,  $\pi_1^*$ , is in general an infinite dimensional torus. It contains a distinguished element,  $\chi_\epsilon$ , the character of  $B_\epsilon$ . We will say that  $\epsilon \in \mathbb{R}$  has a *canonical generator* if and only if the powers of  $\chi_\epsilon$  are dense in the character group. This is equivalent to saying that if  $\epsilon$  is decomposed into  $\ell$  closed disjoint sets, the only rational relation among their harmonic masses is that their total sum is 1.

It seems to us likely that, in some sense, this condition holds generically.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Canonical Generators

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

I now want to discuss the case where  $\epsilon$  might have infinitely many components – in the real case, infinitely many gaps. The character group,  $\pi_1^*$ , is in general an infinite dimensional torus. It contains a distinguished element,  $\chi_\epsilon$ , the character of  $B_\epsilon$ . We will say that  $\epsilon \in \mathbb{R}$  has a *canonical generator* if and only if the powers of  $\chi_\epsilon$  are dense in the character group. This is equivalent to saying that if  $\epsilon$  is decomposed into  $\ell$  closed disjoint sets, the only rational relation among their harmonic masses is that their total sum is 1.

It seems to us likely that, in some sense, this condition holds generically. It follows from results of Totik that among the  $2\nu$  dimensional set of unions of exactly  $\nu$  disjoint unions, the set where the condition fails is a countable union of varieties of dimension  $\nu + 1$  so the set where it fails is both of  $2\nu$  Lebesgue measure zero and a nowhere dense  $F_\sigma$ .



# PW and DCT sets

To state the main new results in part 2, I need to discuss PW (for Parreau–Widom) and DCT (for Direct Cauchy Theorem) sets.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# PW and DCT sets

To state the main new results in part 2, I need to discuss PW (for Parreau–Widom) and DCT (for Direct Cauchy Theorem) sets. We'll act as if these are properties of  $\epsilon$  although they are really properties of  $\Omega \equiv (\mathbb{C} \cup \{\infty\}) \setminus \epsilon$  as an infinitely connected Riemann surface.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# PW and DCT sets

To state the main new results in part 2, I need to discuss PW (for Parreau–Widom) and DCT (for Direct Cauchy Theorem) sets. We'll act as if these are properties of  $\epsilon$  although they are really properties of  $\Omega \equiv (\mathbb{C} \cup \{\infty\}) \setminus \epsilon$  as an infinitely connected Riemann surface. Each has been heavily studied in the literature on such surfaces and each is known to be a family of many different looking equivalent conditions.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# PW and DCT sets

To state the main new results in part 2, I need to discuss PW (for Parreau–Widom) and DCT (for Direct Cauchy Theorem) sets. We'll act as if these are properties of  $\epsilon$  although they are really properties of  $\Omega \equiv (\mathbb{C} \cup \{\infty\}) \setminus \epsilon$  as an infinitely connected Riemann surface. Each has been heavily studied in the literature on such surfaces and each is known to be a family of many different looking equivalent conditions. I'll define the properties via a condition that relates directly to the Widom minimizer problem.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# PW and DCT sets

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

To state the main new results in part 2, I need to discuss PW (for Parreau–Widom) and DCT (for Direct Cauchy Theorem) sets. We'll act as if these are properties of  $\epsilon$  although they are really properties of  $\Omega \equiv (\mathbb{C} \cup \{\infty\}) \setminus \epsilon$  as an infinitely connected Riemann surface. Each has been heavily studied in the literature on such surfaces and each is known to be a family of many different looking equivalent conditions. I'll define the properties via a condition that relates directly to the Widom minimizer problem.

Let  $\chi \in \pi_1^*$  and let  $H^\infty(\Omega, \chi)$  be the family of bounded character automorphic analytic functions on  $\Omega$ .



# PW and DCT sets

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

To state the main new results in part 2, I need to discuss PW (for Parreau–Widom) and DCT (for Direct Cauchy Theorem) sets. We'll act as if these are properties of  $\epsilon$  although they are really properties of  $\Omega \equiv (\mathbb{C} \cup \{\infty\}) \setminus \epsilon$  as an infinitely connected Riemann surface. Each has been heavily studied in the literature on such surfaces and each is known to be a family of many different looking equivalent conditions. I'll define the properties via a condition that relates directly to the Widom minimizer problem.

Let  $\chi \in \pi_1^*$  and let  $H^\infty(\Omega, \chi)$  be the family of bounded character automorphic analytic functions on  $\Omega$ . We say that a compact set,  $\epsilon \in \mathbb{C}$ , has the *PW property* if and only if for every  $\chi \in \pi_1^*$ , we have that  $H^\infty(\Omega, \chi)$  contains non-constant functions.



# PW and DCT sets

Lifted up to the universal cover, for  $\chi \equiv 1$ , this question is equivalent to the existence of automorphic functions, a problem solved in the finitely connected case by Klein and Poincaré.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# PW and DCT sets

Lifted up to the universal cover, for  $\chi \equiv 1$ , this question is equivalent to the existence of automorphic functions, a problem solved in the finitely connected case by Klein and Poincaré. For subsets of  $\mathbb{R}$ , it is often but not always true.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# PW and DCT sets

Lifted up to the universal cover, for  $\chi \equiv 1$ , this question is equivalent to the existence of automorphic functions, a problem solved in the finitely connected case by Klein and Poincaré. For subsets of  $\mathbb{R}$ , it is often but not always true. We'll say more about when it fails later but if  $\epsilon \subset \mathbb{R}$  is perfect, then  $\epsilon$  locally has positive 1D Lebesgue measure,

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# PW and DCT sets

Lifted up to the universal cover, for  $\chi \equiv 1$ , this question is equivalent to the existence of automorphic functions, a problem solved in the finitely connected case by Klein and Poincaré. For subsets of  $\mathbb{R}$ , it is often but not always true. We'll say more about when it fails later but if  $\epsilon \subset \mathbb{R}$  is perfect, then  $\epsilon$  locally has positive 1D Lebesgue measure, so, for example, the classical Cantor set does not have the PW property.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# PW and DCT sets

Lifted up to the universal cover, for  $\chi \equiv 1$ , this question is equivalent to the existence of automorphic functions, a problem solved in the finitely connected case by Klein and Poincaré. For subsets of  $\mathbb{R}$ , it is often but not always true. We'll say more about when it fails later but if  $e \subset \mathbb{R}$  is perfect, then  $e$  locally has positive 1D Lebesgue measure, so, for example, the classical Cantor set does not have the PW property. On the other hand, it is known that homogeneous sets in the sense of Carleson do.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# PW and DCT sets

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Lifted up to the universal cover, for  $\chi \equiv 1$ , this question is equivalent to the existence of automorphic functions, a problem solved in the finitely connected case by Klein and Poincaré. For subsets of  $\mathbb{R}$ , it is often but not always true. We'll say more about when it fails later but if  $\epsilon \subset \mathbb{R}$  is perfect, then  $\epsilon$  locally has positive 1D Lebesgue measure, so, for example, the classical Cantor set does not have the PW property. On the other hand, it is known that homogeneous sets in the sense of Carleson do. It is also known that  $\epsilon \subset \mathbb{C}$  has the PW property iff

$$PW(\epsilon) \equiv \sum_{w \in \mathcal{C}} G_\epsilon(w) < \infty$$

where  $\mathcal{C}$  is the set of critical points of  $G_\epsilon$  (i.e. points in the unbounded component of  $\Omega$  where  $G'_\epsilon(w) = 0$ ).



# PW and DCT sets

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Lifted up to the universal cover, for  $\chi \equiv 1$ , this question is equivalent to the existence of automorphic functions, a problem solved in the finitely connected case by Klein and Poincaré. For subsets of  $\mathbb{R}$ , it is often but not always true. We'll say more about when it fails later but if  $\epsilon \subset \mathbb{R}$  is perfect, then  $\epsilon$  locally has positive 1D Lebesgue measure, so, for example, the classical Cantor set does not have the PW property. On the other hand, it is known that homogeneous sets in the sense of Carleson do. It is also known that  $\epsilon \subset \mathbb{C}$  has the PW property iff

$$PW(\epsilon) \equiv \sum_{w \in \mathcal{C}} G_\epsilon(w) < \infty$$

where  $\mathcal{C}$  is the set of critical points of  $G_\epsilon$  (i.e. points in the unbounded component of  $\Omega$  where  $G'_\epsilon(w) = 0$ ). In particular, any connected set has PW.



# PW and DCT sets

In the case of  $\epsilon \subset \mathbb{R}$  which are regular (i.e.  $G_\epsilon$  continuous on  $\epsilon$ ), there is one critical point in each gap and the value of  $G_\epsilon$  there is the maximum value in the gap.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# PW and DCT sets

In the case of  $\epsilon \subset \mathbb{R}$  which are regular (i.e.  $G_\epsilon$  continuous on  $\epsilon$ ), there is one critical point in each gap and the value of  $G_\epsilon$  there is the maximum value in the gap.

Once one has the PW condition, one can prove there exists a unique Widom minimizer,  $F_\chi$ , which minimizes  $\|A\|_\infty$  among all character automorphic functions with that character and  $A(\infty) = 1$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# PW and DCT sets

In the case of  $\epsilon \subset \mathbb{R}$  which are regular (i.e.  $G_\epsilon$  continuous on  $\epsilon$ ), there is one critical point in each gap and the value of  $G_\epsilon$  there is the maximum value in the gap.

Once one has the PW condition, one can prove there exists a unique Widom minimizer,  $F_\chi$ , which minimizes  $\|A\|_\infty$  among all character automorphic functions with that character and  $A(\infty) = 1$ . One can also consider the *dual Widom maximizer*, the function,  $Q_\chi$ , which is character automorphic with norm 1, non-negative at  $\infty$  that maximizes the value at  $\infty$ . It is easy to see that

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# PW and DCT sets

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

In the case of  $\epsilon \subset \mathbb{R}$  which are regular (i.e.  $G_\epsilon$  continuous on  $\epsilon$ ), there is one critical point in each gap and the value of  $G_\epsilon$  there is the maximum value in the gap.

Once one has the PW condition, one can prove there exists a unique Widom minimizer,  $F_\chi$ , which minimizes  $\|A\|_\infty$  among all character automorphic functions with that character and  $A(\infty) = 1$ . One can also consider the *dual Widom maximizer*, the function,  $Q_\chi$ , which is character automorphic with norm 1, non-negative at  $\infty$  that maximizes the value at  $\infty$ . It is easy to see that

$$Q_\chi = F_\chi / \|F_\chi\|_\infty, \quad F_\chi = Q_\chi / Q_\chi(\infty),$$

$$Q_\chi(\infty) = 1 / \|F_\chi\|_\infty$$



# PW and DCT sets

As the name implies the basic DCT condition has something to do with validity of a Cauchy formula involving boundary values of  $H^\infty$  functions but for us a more useful definition is an equivalent condition:

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# PW and DCT sets

As the name implies the basic DCT condition has something to do with validity of a Cauchy formula involving boundary values of  $H^\infty$  functions but for us a more useful definition is an equivalent condition: we say that  $\epsilon$  obeys the DCT condition if and only if it is a PW set and the map  $\chi \mapsto Q_\chi(\infty)$  is a continuous function of  $\pi_1^*$  to  $(0, 1)$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# PW and DCT sets

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

As the name implies the basic DCT condition has something to do with validity of a Cauchy formula involving boundary values of  $H^\infty$  functions but for us a more useful definition is an equivalent condition: we say that  $\epsilon$  obeys the DCT condition if and only if it is a PW set and the map  $\chi \mapsto Q_\chi(\infty)$  is a continuous function of  $\pi_1^*$  to  $(0, 1)$ . This implies various functions like  $n \mapsto F_{\chi_\epsilon^n}(z)$  are almost periodic.



# Totik Widom Bounds

Recall that we say that  $\mathfrak{e} \subset \mathbb{R}$  obeys a Totik-Widom bound if there is a  $D$  with  $\|T_n\|_{\mathfrak{e}} \leq DC(\mathfrak{e})^n$  and that this was only known for finite gap sets.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Totik Widom Bounds

Recall that we say that  $\epsilon \subset \mathbb{R}$  obeys a Totik-Widom bound if there is a  $D$  with  $\|T_n\|_\epsilon \leq DC(\epsilon)^n$  and that this was only known for finite gap sets. One of the (two) main results in Part 1 is

**Theorem** If  $\epsilon \subset \mathbb{R}$  is a regular Parreau-Widom set, then

$$\|T_n\|_\epsilon \leq 2 \exp(PW(\epsilon)) C(\epsilon)^n$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Totik Widom Bounds

Recall that we say that  $\epsilon \subset \mathbb{R}$  obeys a Totik-Widom bound if there is a  $D$  with  $\|T_n\|_\epsilon \leq DC(\epsilon)^n$  and that this was only known for finite gap sets. One of the (two) main results in Part 1 is

**Theorem** If  $\epsilon \subset \mathbb{R}$  is a regular Parreau-Widom set, then

$$\|T_n\|_\epsilon \leq 2 \exp(PW(\epsilon)) C(\epsilon)^n$$

Homogeneous sets are regular and obey a Parreau Widom condition (a theorem of Jones and Marshall).

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Totik Widom Bounds

Recall that we say that  $\epsilon \subset \mathbb{R}$  obeys a Totik-Widom bound if there is a  $D$  with  $\|T_n\|_\epsilon \leq DC(\epsilon)^n$  and that this was only known for finite gap sets. One of the (two) main results in Part 1 is

**Theorem** If  $\epsilon \subset \mathbb{R}$  is a regular Parreau-Widom set, then

$$\|T_n\|_\epsilon \leq 2 \exp(PW(\epsilon)) C(\epsilon)^n$$

Homogeneous sets are regular and obey a Parreau Widom condition (a theorem of Jones and Marshall). This explicit constant is interesting even for the finite gap case.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Totik Widom Bounds

Recall that we say that  $\epsilon \subset \mathbb{R}$  obeys a Totik-Widom bound if there is a  $D$  with  $\|T_n\|_\epsilon \leq DC(\epsilon)^n$  and that this was only known for finite gap sets. One of the (two) main results in Part 1 is

**Theorem** If  $\epsilon \subset \mathbb{R}$  is a regular Parreau-Widom set, then

$$\|T_n\|_\epsilon \leq 2 \exp(PW(\epsilon)) C(\epsilon)^n$$

Homogeneous sets are regular and obey a Parreau Widom condition (a theorem of Jones and Marshall). This explicit constant is interesting even for the finite gap case. We also proved a weak converse:

**Theorem** If  $\epsilon \subset \mathbb{C}$  is a regular set for which a TW bound holds and  $\epsilon$  has a canonical generator, then  $\epsilon$  is a PW set.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Totik Widom Bounds

**Interesting Open Question** Does potential theory regularity + Parreau-Widom  $\Rightarrow$  Totik-Widom bound for general  $e \subset \mathbb{C}$  (our proof is only for  $e \subset \mathbb{R}$ ).

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Totik Widom Bounds

**Interesting Open Question** Does potential theory regularity + Parreau-Widom  $\Rightarrow$  Totik-Widom bound for general  $e \subset \mathbb{C}$  (our proof is only for  $e \subset \mathbb{R}$ ).

For a time I suspected the answer was yes, then no, and now I'm unsure, but I lean towards yes.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Totik Widom Bounds

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

**Interesting Open Question** Does potential theory regularity + Parreau-Widom  $\Rightarrow$  Totik-Widom bound for general  $e \subset \mathbb{C}$  (our proof is only for  $e \subset \mathbb{R}$ ).

For a time I suspected the answer was yes, then no, and now I'm unsure, but I lean towards yes. A key example is the solid Koch snowflake.



# Totik Widom Bounds

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

**Interesting Open Question** Does potential theory regularity + Parreau-Widom  $\Rightarrow$  Totik-Widom bound for general  $e \subset \mathbb{C}$  (our proof is only for  $e \subset \mathbb{R}$ ).

For a time I suspected the answer was yes, then no, and now I'm unsure, but I lean towards yes. A key example is the solid Koch snowflake. Since it is simply connected, it is a PW set.



# Totik Widom Bounds

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

**Interesting Open Question** Does potential theory regularity + Parreau-Widom  $\Rightarrow$  Totik-Widom bound for general  $e \subset \mathbb{C}$  (our proof is only for  $e \subset \mathbb{R}$ ).

For a time I suspected the answer was yes, then no, and now I'm unsure, but I lean towards yes. A key example is the solid Koch snowflake. Since it is simply connected, it is a PW set. On the other hand the fact that its boundary has dimension greater than 1 makes it a candidate for failure of TW bounds if the theorem does not extend to the general complex case so much so I suspect it was false.



# Totik Widom Bounds

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

**Interesting Open Question** Does potential theory regularity + Parreau-Widom  $\Rightarrow$  Totik-Widom bound for general  $e \subset \mathbb{C}$  (our proof is only for  $e \subset \mathbb{R}$ ).

For a time I suspected the answer was yes, then no, and now I'm unsure, but I lean towards yes. A key example is the solid Koch snowflake. Since it is simply connected, it is a PW set. On the other hand the fact that its boundary has dimension greater than 1 makes it a candidate for failure of TW bounds if the theorem does not extend to the general complex case so much so I suspect it was false. But, recently, Andrievskii proved a TW bound holds for a class of sets that includes the Koch snowflake, so I've oscillated back.



# Limit Points of the ratio

Paper 3 has the following result about the quantity  
 $r_n = \|T_n\|_{\mathfrak{e}} / C(\mathfrak{e})^n$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Limit Points of the ratio

Paper 3 has the following result about the quantity

$$r_n = \|T_n\|_{\mathfrak{e}} / C(\mathfrak{e})^n$$

**Theorem** Let  $\mathfrak{e} \subset \mathbb{R}$  be a DCT set and have canonical generators. Then the set of limit points of the ratio  $r_n$  is exactly the closed interval  $[2, 2 \exp(PW(\mathfrak{e}))]$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Limit Points of the ratio

Paper 3 has the following result about the quantity

$$r_n = \|T_n\|_{\epsilon}/C(\epsilon)^n$$

**Theorem** Let  $\epsilon \subset \mathbb{R}$  be a DCT set and have canonical generators. Then the set of limit points of the ratio  $r_n$  is exactly the closed interval  $[2, 2 \exp(PW(\epsilon))]$ .

Thus the upper and lower bounds are asymptotically exact and cannot be improved.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Limit Points of the ratio

Paper 3 has the following result about the quantity

$$r_n = \|T_n\|_{\epsilon}/C(\epsilon)^n$$

**Theorem** Let  $\epsilon \subset \mathbb{R}$  be a DCT set and have canonical generators. Then the set of limit points of the ratio  $r_n$  is exactly the closed interval  $[2, 2 \exp(PW(\epsilon))]$ .

Thus the upper and lower bounds are asymptotically exact and cannot be improved. For example, in the finite gap case, this result holds if the harmonic measures of the bands are rational independent.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Limit Points of the ratio

Paper 3 has the following result about the quantity

$$r_n = \|T_n\|_{\epsilon}/C(\epsilon)^n$$

**Theorem** Let  $\epsilon \subset \mathbb{R}$  be a DCT set and have canonical generators. Then the set of limit points of the ratio  $r_n$  is exactly the closed interval  $[2, 2 \exp(PW(\epsilon))]$ .

Thus the upper and lower bounds are asymptotically exact and cannot be improved. For example, in the finite gap case, this result holds if the harmonic measures of the bands are rational independent. But the limits can be less than the whole interval in the non-generic case.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Limit Points of the ratio

Paper 3 has the following result about the quantity  
 $r_n = \|T_n\|_{\epsilon}/C(\epsilon)^n$

**Theorem** Let  $\epsilon \subset \mathbb{R}$  be a DCT set and have canonical generators. Then the set of limit points of the ratio  $r_n$  is exactly the closed interval  $[2, 2 \exp(PW(\epsilon))]$ .

Thus the upper and lower bounds are asymptotically exact and cannot be improved. For example, in the finite gap case, this result holds if the harmonic measures of the bands are rational independent. But the limits can be less than the whole interval in the non-generic case. For example, if  $\epsilon$  has two bands of equal size, the the set limit points is the two point set  $\{2, 2 \exp(PW(\epsilon))\}$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Widom's Conjecture

The other main result of Part 1 settled a 45 year old conjecture:

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Widom's Conjecture

The other main result of Part 1 settled a 45 year old conjecture:

**Theorem** Widom's conjecture on the almost periodic Szegő asymptotics outside  $\epsilon$  for the Chebyshev polynomials of finite gap sets is true.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Widom's Conjecture

The other main result of Part 1 settled a 45 year old conjecture:

**Theorem** Widom's conjecture on the almost periodic Szegő asymptotics outside  $\epsilon$  for the Chebyshev polynomials of finite gap sets is true.

In Part 2, we extended this:

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Widom's Conjecture

The other main result of Part 1 settled a 45 year old conjecture:

**Theorem** Widom's conjecture on the almost periodic Szegő asymptotics outside  $\epsilon$  for the Chebyshev polynomials of finite gap sets is true.

In Part 2, we extended this:

**Theorem** Szegő–Widom asymptotics outside  $\epsilon$  holds for the Chebyshev polynomials of any  $\epsilon \subset \mathbb{R}$  for any  $\epsilon$  that is both PW and DCT.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Widom's Conjecture

The other main result of Part 1 settled a 45 year old conjecture:

**Theorem** Widom's conjecture on the almost periodic Szegő asymptotics outside  $\epsilon$  for the Chebyshev polynomials of finite gap sets is true.

In Part 2, we extended this:

**Theorem** Szegő–Widom asymptotics outside  $\epsilon$  holds for the Chebyshev polynomials of any  $\epsilon \subset \mathbb{R}$  for any  $\epsilon$  that is both PW and DCT.

The proof of this in Part 2 is simpler than the proof of Part 1 (and doesn't require Widom's result on the norm apriori but rather proves it).

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Szegő–Widom Asymptotics

We also proved that:

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Szegő–Widom Asymptotics

We also proved that:

**Theorem** If  $\epsilon \subset \mathbb{R}$  is a regular set PW set for which  $n \mapsto A_{\chi_\epsilon^n}$  is almost periodic in  $n$  and which has a canonical generator, then  $\epsilon$  is a DCT set.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

**Main Results**

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Szegő–Widom Asymptotics

We also proved that:

**Theorem** If  $\epsilon \subset \mathbb{R}$  is a regular set PW set for which  $n \mapsto A_{\chi_\epsilon^n}$  is almost periodic in  $n$  and which has a canonical generator, then  $\epsilon$  is a DCT set.

Since almost periodicity of the limit is part of Szegő–Widom asymptotics, this is a kind of converse to

DCT  $\Rightarrow$  Szegő–Widom asymptotics.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Proof modulo Lemma

We turn to the proof of TW bounds from Part 1:

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Proof modulo Lemma

We turn to the proof of TW bounds from Part 1:

**Lemma** Let  $\epsilon \subset \epsilon_n \subset \mathbb{R}$  be a compact subset and its canonical period  $n$  superset.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Proof modulo Lemma

We turn to the proof of TW bounds from Part 1:

**Lemma** Let  $\epsilon \subset \epsilon_n \subset \mathbb{R}$  be a compact subset and its canonical period  $n$  superset. Let  $K$  be a gap of  $\epsilon$  and  $d\rho_n$ , the equilibrium measure of  $\epsilon_n$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Proof modulo Lemma

We turn to the proof of TW bounds from Part 1:

**Lemma** Let  $\epsilon \subset \epsilon_n \subset \mathbb{R}$  be a compact subset and its canonical period  $n$  superset. Let  $K$  be a gap of  $\epsilon$  and  $d\rho_n$ , the equilibrium measure of  $\epsilon_n$ . Then  $\rho_n(K) \leq 1/n$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Proof modulo Lemma

We turn to the proof of TW bounds from Part 1:

**Lemma** Let  $\epsilon \subset \epsilon_n \subset \mathbb{R}$  be a compact subset and its canonical period  $n$  superset. Let  $K$  be a gap of  $\epsilon$  and  $d\rho_n$ , the equilibrium measure of  $\epsilon_n$ . Then  $\rho_n(K) \leq 1/n$ .

Accepting this for the moment, let  $h(z) \equiv G_\epsilon(z) - G_{\epsilon_n}(z)$  which is harmonic at infinity with

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Proof modulo Lemma

We turn to the proof of TW bounds from Part 1:

**Lemma** Let  $\epsilon \subset \epsilon_n \subset \mathbb{R}$  be a compact subset and its canonical period  $n$  superset. Let  $K$  be a gap of  $\epsilon$  and  $d\rho_n$ , the equilibrium measure of  $\epsilon_n$ . Then  $\rho_n(K) \leq 1/n$ .

Accepting this for the moment, let  $h(z) \equiv G_\epsilon(z) - G_{\epsilon_n}(z)$  which is harmonic at infinity with

$$h(\infty) = R(\epsilon) - R(\epsilon_n) = \log \left[ \frac{C(\epsilon_n)}{C(\epsilon)} \right]$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Proof modulo Lemma

We turn to the proof of TW bounds from Part 1:

**Lemma** Let  $\epsilon \subset \epsilon_n \subset \mathbb{R}$  be a compact subset and its canonical period  $n$  superset. Let  $K$  be a gap of  $\epsilon$  and  $d\rho_n$ , the equilibrium measure of  $\epsilon_n$ . Then  $\rho_n(K) \leq 1/n$ .

Accepting this for the moment, let  $h(z) \equiv G_\epsilon(z) - G_{\epsilon_n}(z)$  which is harmonic at infinity with

$$h(\infty) = R(\epsilon) - R(\epsilon_n) = \log \left[ \frac{C(\epsilon_n)}{C(\epsilon)} \right]$$

Since  $d\rho_n$  is harmonic measure and  $h(x) = G_\epsilon(x)$  on  $\epsilon_n$ , if  $\{K_j\}_{j=1}^M$  are the gaps for  $\epsilon$ , then, using the Lemma

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Proof modulo Lemma

We turn to the proof of TW bounds from Part 1:

**Lemma** Let  $\epsilon \subset \epsilon_n \subset \mathbb{R}$  be a compact subset and its canonical period  $n$  superset. Let  $K$  be a gap of  $\epsilon$  and  $d\rho_n$ , the equilibrium measure of  $\epsilon_n$ . Then  $\rho_n(K) \leq 1/n$ .

Accepting this for the moment, let  $h(z) \equiv G_\epsilon(z) - G_{\epsilon_n}(z)$  which is harmonic at infinity with

$$h(\infty) = R(\epsilon) - R(\epsilon_n) = \log \left[ \frac{C(\epsilon_n)}{C(\epsilon)} \right]$$

Since  $d\rho_n$  is harmonic measure and  $h(x) = G_\epsilon(x)$  on  $\epsilon_n$ , if  $\{K_j\}_{j=1}^M$  are the gaps for  $\epsilon$ , then, using the Lemma

$$h(\infty) \leq \sum_{j=1}^M \rho_n(K_j) \max_{x \in K_j} (G_\epsilon(x)) \leq \frac{1}{n} \sum_{j=1}^M G_\epsilon(w_j)$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Proof modulo Lemma

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics

We turn to the proof of TW bounds from Part 1:

**Lemma** Let  $\epsilon \subset \epsilon_n \subset \mathbb{R}$  be a compact subset and its canonical period  $n$  superset. Let  $K$  be a gap of  $\epsilon$  and  $d\rho_n$ , the equilibrium measure of  $\epsilon_n$ . Then  $\rho_n(K) \leq 1/n$ .

Accepting this for the moment, let  $h(z) \equiv G_\epsilon(z) - G_{\epsilon_n}(z)$  which is harmonic at infinity with

$$h(\infty) = R(\epsilon) - R(\epsilon_n) = \log \left[ \frac{C(\epsilon_n)}{C(\epsilon)} \right]$$

Since  $d\rho_n$  is harmonic measure and  $h(x) = G_\epsilon(x)$  on  $\epsilon_n$ , if  $\{K_j\}_{j=1}^M$  are the gaps for  $\epsilon$ , then, using the Lemma

$$h(\infty) \leq \sum_{j=1}^M \rho_n(K_j) \max_{x \in K_j} (G_\epsilon(x)) \leq \frac{1}{n} \sum_{j=1}^M G_\epsilon(w_j)$$
 since regularity of  $\epsilon$  implies  $G_\epsilon$  vanishes at the ends of each gap so the maximum is taken a critical point  $w_j$ . Exponentiating and using  $\|T_n\|_\epsilon \leq 2C(\epsilon_n)^n$  we get the result. ■



# Proof of the Lemma

Because the integrated equilibrium measure of  $\epsilon_n$  is  $\frac{1}{\pi n} \arccos \left( \frac{T_n(x)}{\|T_n\|_{\epsilon}} \right)$ , each band of  $\epsilon_n$  has  $\rho_{\epsilon_n}$  measure  $\frac{1}{n}$  and the part of a band from a zero of  $T_n$  to a nearby band edge has  $\rho_{\epsilon_n}$  measure  $\frac{1}{2n}$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

**Totik Widom  
Bounds**

Szegő–Widom  
Asymptotics



# Proof of the Lemma

Because the integrated equilibrium measure of  $\epsilon_n$  is  $\frac{1}{\pi n} \arccos \left( \frac{T_n(x)}{\|T_n\|_{\epsilon}} \right)$ , each band of  $\epsilon_n$  has  $\rho_{\epsilon_n}$  measure  $\frac{1}{n}$  and the part of a band from a zero of  $T_n$  to a nearby band edge has  $\rho_{\epsilon_n}$  measure  $\frac{1}{2n}$ . Recall that any gap  $K$  has at most one zero.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

**Totik Widom  
Bounds**

Szegő–Widom  
Asymptotics



# Proof of the Lemma

Because the integrated equilibrium measure of  $\epsilon_n$  is  $\frac{1}{\pi n} \arccos \left( \frac{T_n(x)}{\|T_n\|_{\epsilon}} \right)$ , each band of  $\epsilon_n$  has  $\rho_{\epsilon_n}$  measure  $\frac{1}{n}$  and the part of a band from a zero of  $T_n$  to a nearby band edge has  $\rho_{\epsilon_n}$  measure  $\frac{1}{2n}$ . Recall that any gap  $K$  has at most one zero.

**Case 1** ( $T_n$  has no zero in  $K$ ) Then there are zeros above and below  $K$  not in  $K$ . Thus  $K$  contains at most two half bands.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Proof of the Lemma

Because the integrated equilibrium measure of  $\epsilon_n$  is  $\frac{1}{\pi n} \arccos \left( \frac{T_n(x)}{\|T_n\|_{\epsilon}} \right)$ , each band of  $\epsilon_n$  has  $\rho_{\epsilon_n}$  measure  $\frac{1}{n}$  and the part of a band from a zero of  $T_n$  to a nearby band edge has  $\rho_{\epsilon_n}$  measure  $\frac{1}{2n}$ . Recall that any gap  $K$  has at most one zero.

**Case 1** ( $T_n$  has no zero in  $K$ ) Then there are zeros above and below  $K$  not in  $K$ . Thus  $K$  contains at most two half bands. (In fact, using the Alternation Theorem, one can show at most one half band).

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Proof of the Lemma

Chebyshev  
Polynomials  
  
Alternation  
Theorem  
  
Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Because the integrated equilibrium measure of  $\epsilon_n$  is  $\frac{1}{\pi n} \arccos \left( \frac{T_n(x)}{\|T_n\|_{\epsilon}} \right)$ , each band of  $\epsilon_n$  has  $\rho_{\epsilon_n}$  measure  $\frac{1}{n}$  and the part of a band from a zero of  $T_n$  to a nearby band edge has  $\rho_{\epsilon_n}$  measure  $\frac{1}{2n}$ . Recall that any gap  $K$  has at most one zero.

**Case 1** ( $T_n$  has no zero in  $K$ ) Then there are zeros above and below  $K$  not in  $K$ . Thus  $K$  contains at most two half bands. (In fact, using the Alternation Theorem, one can show at most one half band).

**Case 2** ( $T_n$  has a zero in  $K$ ) By the Alternation Theorem, one of the two extreme points immediately below the zero must lie in  $\epsilon$ , so there is at most a half band below the zero. Similarly, at most a half band above, so no more than a full band. ■



# Size of $\epsilon_n \setminus \epsilon$

In fact, one can prove if there is a zero not too close to a gap edge and  $n$  is large, then there is exactly a full exponentially small (in Lebesgue measure) band of  $\epsilon_n$  totally inside  $K$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Size of $\epsilon_n \setminus \epsilon$

In fact, one can prove if there is a zero not too close to a gap edge and  $n$  is large, then there is exactly a full exponentially small (in Lebesgue measure) band of  $\epsilon_n$  totally inside  $K$ .

This implies that if  $K$  is a gap and  $n_j$  is such as  $j \rightarrow \infty$  and any zeros of  $T_{n_j}$  in  $K$  go to the edges,

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Size of $\epsilon_n \setminus \epsilon$

In fact, one can prove if there is a zero not too close to a gap edge and  $n$  is large, then there is exactly a full exponentially small (in Lebesgue measure) band of  $\epsilon_n$  totally inside  $K$ .

This implies that if  $K$  is a gap and  $n_j$  is such as  $j \rightarrow \infty$  and any zeros of  $T_{n_j}$  in  $K$  go to the edges, then asymptotically there is none of  $\epsilon_{n_j}$  in  $K$  in the sense that  $\bigcap_{k=1}^{\infty} \bigcup_{j=k}^{\infty} (K \cap \epsilon_{n_j}) = \emptyset$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Size of $\epsilon_n \setminus \epsilon$

In fact, one can prove if there is a zero not too close to a gap edge and  $n$  is large, then there is exactly a full exponentially small (in Lebesgue measure) band of  $\epsilon_n$  totally inside  $K$ .

This implies that if  $K$  is a gap and  $n_j$  is such as  $j \rightarrow \infty$  and any zeros of  $T_{n_j}$  in  $K$  go to the edges, then asymptotically there is none of  $\epsilon_{n_j}$  in  $K$  in the sense that  $\bigcap_{k=1}^{\infty} \bigcup_{j=k}^{\infty} (K \cap \epsilon_{n_j}) = \emptyset$ .

Similarly if, for  $j$  large  $T_{n_j}$  has a zero,  $x_j$ , in  $K$  and  $x_j \rightarrow x_{\infty} \in K$ , then only  $x_{\infty}$  is asymptotically in  $\epsilon_{n_j} \cap K$  in the sense that  $\bigcap_{k=1}^{\infty} \bigcup_{j=k}^{\infty} (K \cap \epsilon_{n_j}) = \{x_{\infty}\}$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Overall Strategy

Finally, some remarks on the proof of SW asymptotics. For any  $x \in \Omega$ , we can look at the lifts of  $x$  in the universal cover, thought of as the unit disk

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Overall Strategy

Finally, some remarks on the proof of SW asymptotics. For any  $x \in \Omega$ , we can look at the lifts of  $x$  in the universal cover, thought of as the unit disk and form the product of Blaschke factors with those zeros.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Overall Strategy

Finally, some remarks on the proof of SW asymptotics. For any  $x \in \Omega$ , we can look at the lifts of  $x$  in the universal cover, thought of as the unit disk and form the product of Blaschke factors with those zeros. If  $\epsilon$  is PW, then one can prove this Blaschke product converges to a function vanishing precisely at the those lifts.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Overall Strategy

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Finally, some remarks on the proof of SW asymptotics. For any  $x \in \Omega$ , we can look at the lifts of  $x$  in the universal cover, thought of as the unit disk and form the product of Blaschke factors with those zeros. If  $e$  is PW, then one can prove this Blaschke product converges to a function vanishing precisely at the those lifts. This defines a character automorphic function,  $B(z, x)$ , which is the unique character automorphic function with infinity norm 1, positive at  $\infty$  with zero's precisely at  $x$  (and its images under the deck transformations).



# Overall Strategy

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Finally, some remarks on the proof of SW asymptotics. For any  $x \in \Omega$ , we can look at the lifts of  $x$  in the universal cover, thought of as the unit disk and form the product of Blaschke factors with those zeros. If  $\epsilon$  is PW, then one can prove this Blaschke product converges to a function vanishing precisely at the those lifts. This defines a character automorphic function,  $B(z, x)$ , which is the unique character automorphic function with infinity norm 1, positive at  $\infty$  with zero's precisely at  $x$  (and its images under the deck transformations).

$\mathbb{R} \setminus \epsilon$  is a disjoint union of bounded open components (plus two unbounded components),  $K \in \mathcal{G}$ . We'll call these the gaps and  $\mathcal{G}$  the set of gaps.



# Overall Strategy

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Finally, some remarks on the proof of SW asymptotics. For any  $x \in \Omega$ , we can look at the lifts of  $x$  in the universal cover, thought of as the unit disk and form the product of Blaschke factors with those zeros. If  $\epsilon$  is PW, then one can prove this Blaschke product converges to a function vanishing precisely at the those lifts. This defines a character automorphic function,  $B(z, x)$ , which is the unique character automorphic function with infinity norm 1, positive at  $\infty$  with zero's precisely at  $x$  (and its images under the deck transformations).

$\mathbb{R} \setminus \epsilon$  is a disjoint union of bounded open components (plus two unbounded components),  $K \in \mathcal{G}$ . We'll call these the gaps and  $\mathcal{G}$  the set of gaps. A *gap collection* is a subset  $\mathcal{G}_0 \subset \mathcal{G}$ .



# Overall Strategy

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

Finally, some remarks on the proof of SW asymptotics. For any  $x \in \Omega$ , we can look at the lifts of  $x$  in the universal cover, thought of as the unit disk and form the product of Blaschke factors with those zeros. If  $\epsilon$  is PW, then one can prove this Blaschke product converges to a function vanishing precisely at the those lifts. This defines a character automorphic function,  $B(z, x)$ , which is the unique character automorphic function with infinity norm 1, positive at  $\infty$  with zero's precisely at  $x$  (and its images under the deck transformations).

$\mathbb{R} \setminus \epsilon$  is a disjoint union of bounded open components (plus two unbounded components),  $K \in \mathcal{G}$ . We'll call these the gaps and  $\mathcal{G}$  the set of gaps. A *gap collection* is a subset  $\mathcal{G}_0 \subset \mathcal{G}$ . A *gap set* is a gap collection,  $\mathcal{G}_0$ , and for each  $K_k \in \mathcal{G}_0$  a point  $x_k \in K_k$ .



# Overall Strategy

For any gap set,  $S$ , we define the associated Blaschke product

$$B_S(z) = \prod_{K_k \in \mathcal{G}_0} B_{\epsilon}(z, x_k)$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Overall Strategy

For any gap set,  $S$ , we define the associated Blaschke product

$$B_S(z) = \prod_{K_k \in \mathcal{G}_0} B_{\epsilon}(z, x_k)$$

To prove Szegő–Widom asymptotics, it suffices, by Montel's theorem and uniqueness of minimizers, to show that any limit point of the  $L_n(z) \equiv T_n(z)B_{\epsilon}(z)^n/C(\epsilon)^n$  is a Widom minimizer.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Overall Strategy

For any gap set,  $S$ , we define the associated Blaschke product

$$B_S(z) = \prod_{K_k \in \mathcal{G}_0} B_{\epsilon}(z, x_k)$$

To prove Szegő–Widom asymptotics, it suffices, by Montel's theorem and uniqueness of minimizers, to show that any limit point of the  $L_n(z) \equiv T_n(z)B_{\epsilon}(z)^n/C(\epsilon)^n$  is a Widom minimizer. We'll instead look at

$$M_n(z) = B(z)^n/B_n(z)^n$$

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics



# Overall Strategy

For any gap set,  $S$ , we define the associated Blaschke product

$$B_S(z) = \prod_{K_k \in \mathcal{G}_0} B_{\mathfrak{e}}(z, x_k)$$

To prove Szegő–Widom asymptotics, it suffices, by Montel's theorem and uniqueness of minimizers, to show that any limit point of the  $L_n(z) \equiv T_n(z)B_{\mathfrak{e}}(z)^n/C(\mathfrak{e})^n$  is a Widom minimizer. We'll instead look at

$$M_n(z) = B(z)^n / B_n(z)^n$$

which obeys

$$|M_n(z)| = \exp(-h_n(z)); \quad h_n(z) \equiv G_{\mathfrak{e}}(z) - G_{\mathfrak{e}_n}(z)$$

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics



# Overall Strategy

For any gap set,  $S$ , we define the associated Blaschke product

$$B_S(z) = \prod_{K_k \in \mathcal{G}_0} B_{\mathfrak{e}}(z, x_k)$$

To prove Szegő–Widom asymptotics, it suffices, by Montel's theorem and uniqueness of minimizers, to show that any limit point of the  $L_n(z) \equiv T_n(z)B_{\mathfrak{e}}(z)^n/C(\mathfrak{e})^n$  is a Widom minimizer. We'll instead look at

$$M_n(z) = B(z)^n / B_n(z)^n$$

which obeys

$$|M_n(z)| = \exp(-h_n(z)); \quad h_n(z) \equiv G_{\mathfrak{e}}(z) - G_{\mathfrak{e}_n}(z)$$

so that one can write  $L_n$  in terms of  $M_n$

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics



# Overall Strategy

$$L_n(z) = (1 + B_n(z)^{2n})H_n(z)$$

$$H_n(z) = \frac{C(\mathfrak{e}_n)^n}{C(\mathfrak{e})^n} \frac{B(z)^n}{B_n(z)^n} = \frac{M_n(z)}{M_n(\infty)}$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Overall Strategy

$$L_n(z) = (1 + B_n(z)^{2n})H_n(z)$$

$$H_n(z) = \frac{C(\epsilon_n)^n}{C(\epsilon)^n} \frac{B(z)^n}{B_n(z)^n} = \frac{M_n(z)}{M_n(\infty)}$$

This means it suffices to prove that any limit point of the  $M_n$  is a dual Widom maximizer.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Overall Strategy

$$L_n(z) = (1 + B_n(z)^{2n}) H_n(z)$$

$$H_n(z) = \frac{C(\epsilon_n)^n}{C(\epsilon)^n} \frac{B(z)^n}{B_n(z)^n} = \frac{M_n(z)}{M_n(\infty)}$$

This means it suffices to prove that any limit point of the  $M_n$  is a dual Widom maximizer. Moreover, we can pass to subsequences and suppose that for the limit point is also true that the zeros in each gap have a limit or else leave the gap (through the edges).

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics



# Overall Strategy

$$L_n(z) = (1 + B_n(z)^{2n}) H_n(z)$$

$$H_n(z) = \frac{C(\epsilon_n)^n}{C(\epsilon)^n} \frac{B(z)^n}{B_n(z)^n} = \frac{M_n(z)}{M_n(\infty)}$$

This means it suffices to prove that any limit point of the  $M_n$  is a dual Widom maximizer. Moreover, we can pass to subsequences and suppose that for the limit point is also true that the zeros in each gap have a limit or else leave the gap (through the edges). The limit points of zeros then define a gap set.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Overall Strategy

$$L_n(z) = (1 + B_n(z)^{2n})H_n(z)$$

$$H_n(z) = \frac{C(\epsilon_n)^n}{C(\epsilon)^n} \frac{B(z)^n}{B_n(z)^n} = \frac{M_n(z)}{M_n(\infty)}$$

This means it suffices to prove that any limit point of the  $M_n$  is a dual Widom maximizer. Moreover, we can pass to subsequences and suppose that for the limit point is also true that the zeros in each gap have a limit or else leave the gap (through the edges). The limit points of zeros then define a gap set.

The proof has two steps: first, prove that the limit is the Blaschke product of this gap set

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Overall Strategy

$$L_n(z) = (1 + B_n(z)^{2n})H_n(z)$$

$$H_n(z) = \frac{C(\epsilon_n)^n}{C(\epsilon)^n} \frac{B(z)^n}{B_n(z)^n} = \frac{M_n(z)}{M_n(\infty)}$$

This means it suffices to prove that any limit point of the  $M_n$  is a dual Widom maximizer. Moreover, we can pass to subsequences and suppose that for the limit point is also true that the zeros in each gap have a limit or else leave the gap (through the edges). The limit points of zeros then define a gap set.

The proof has two steps: first, prove that the limit is the Blaschke product of this gap set and, secondly, prove that any such product is a dual Widom maximizer.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Overall Strategy

$$L_n(z) = (1 + B_n(z)^{2n}) H_n(z)$$

$$H_n(z) = \frac{C(\epsilon_n)^n}{C(\epsilon)^n} \frac{B(z)^n}{B_n(z)^n} = \frac{M_n(z)}{M_n(\infty)}$$

This means it suffices to prove that any limit point of the  $M_n$  is a dual Widom maximizer. Moreover, we can pass to subsequences and suppose that for the limit point is also true that the zeros in each gap have a limit or else leave the gap (through the edges). The limit points of zeros then define a gap set.

The proof has two steps: first, prove that the limit is the Blaschke product of this gap set and, secondly, prove that any such product is a dual Widom maximizer. The proof of the second half follows, in part, ideas of Volberg–Yuditskii, who considered a related problem and uses some deep 1997 results of Sodin–Yuditskii on the Abel map in this setting.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Limits of $M_n$ are Gap Blaschke Products

Rather than control  $M_n$  globally, it suffices, by a little complex analysis, to prove convergence of the absolute values and only for  $z$  in a small neighborhood of  $\infty$ . Taking into account the formula we had for  $|M_n(z)|$  and taking logs, it suffices to prove that, for  $z$  near  $\infty$ , we have that

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Limits of $M_n$ are Gap Blaschke Products

Rather than control  $M_n$  globally, it suffices, by a little complex analysis, to prove convergence of the absolute values and only for  $z$  in a small neighborhood of  $\infty$ . Taking into account the formula we had for  $|M_n(z)|$  and taking logs, it suffices to prove that, for  $z$  near  $\infty$ , we have that

$$nh_n(z) \rightarrow \sum_{K_k \in \mathcal{G}_0} G_{\epsilon}(x_k, z)$$

where  $G_{\epsilon}(x, z)$  is the Green's function with pole at  $x$  (so that  $G_{\epsilon}(\infty, z)$  is what we called  $G_{\epsilon}(z)$ ).

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Limits of $M_n$ are Gap Blaschke Products

Rather than control  $M_n$  globally, it suffices, by a little complex analysis, to prove convergence of the absolute values and only for  $z$  in a small neighborhood of  $\infty$ . Taking into account the formula we had for  $|M_n(z)|$  and taking logs, it suffices to prove that, for  $z$  near  $\infty$ , we have that

$$nh_n(z) \rightarrow \sum_{K_k \in \mathcal{G}_0} G_{\epsilon}(x_k, z)$$

where  $G_{\epsilon}(x, z)$  is the Green's function with pole at  $x$  (so that  $G_{\epsilon}(\infty, z)$  is what we called  $G_{\epsilon}(z)$ ).

Part 1, we proved Totik–Widom bounds for PW sets,  $\epsilon \subset \mathbb{R}$  by using that when  $z = \infty$ , we have that

$$h_n(\infty) = \int_{\bigcup_{K_j \in \mathcal{G}} K_j} G_{\epsilon}(x) d\rho_n(x)$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# A Poisson Formula

We proved this by thinking of  $d\rho_n$  as harmonic measure at  $\infty$ , i.e. if  $H$  is harmonic on  $(\mathbb{C} \cup \{\infty\}) \setminus \mathfrak{e}_n$  with boundary values  $H(x)$  on  $\mathfrak{e}_n$ , then  $H(\infty) = \int_{\mathfrak{e}_n} H(x)d\rho_n(x)$ .

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# A Poisson Formula

We proved this by thinking of  $d\rho_n$  as harmonic measure at  $\infty$ , i.e. if  $H$  is harmonic on  $(\mathbb{C} \cup \{\infty\}) \setminus \mathfrak{e}_n$  with boundary values  $H(x)$  on  $\mathfrak{e}_n$ , then  $H(\infty) = \int_{\mathfrak{e}_n} H(x) d\rho_n(x)$ . If we wrote the analog of this for general  $z$ , we'd get

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# A Poisson Formula

We proved this by thinking of  $d\rho_n$  as harmonic measure at  $\infty$ , i.e. if  $H$  is harmonic on  $(\mathbb{C} \cup \{\infty\}) \setminus \mathfrak{e}_n$  with boundary values  $H(x)$  on  $\mathfrak{e}_n$ , then  $H(\infty) = \int_{\mathfrak{e}_n} H(x)d\rho_n(x)$ . If we wrote the analog of this for general  $z$ , we'd get

$$H(z) = \int_{\mathfrak{e}_n} H(x)d\rho_n(x, z)$$

varying the harmonic measure.

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# A Poisson Formula

We proved this by thinking of  $d\rho_n$  as harmonic measure at  $\infty$ , i.e. if  $H$  is harmonic on  $(\mathbb{C} \cup \{\infty\}) \setminus \mathfrak{e}_n$  with boundary values  $H(x)$  on  $\mathfrak{e}_n$ , then  $H(\infty) = \int_{\mathfrak{e}_n} H(x)d\rho_n(x)$ . If we wrote the analog of this for general  $z$ , we'd get

$$H(z) = \int_{\mathfrak{e}_n} H(x)d\rho_n(x, z)$$

varying the harmonic measure. Instead we use

$$h_n(z) = \int_{\bigcup_{K_j \in \mathcal{G}} K_j} G_{\mathfrak{e}}(x, z)d\rho_n(x)$$

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics



# A Poisson Formula

We proved this by thinking of  $d\rho_n$  as harmonic measure at  $\infty$ , i.e. if  $H$  is harmonic on  $(\mathbb{C} \cup \{\infty\}) \setminus \epsilon_n$  with boundary values  $H(x)$  on  $\epsilon_n$ , then  $H(\infty) = \int_{\epsilon_n} H(x)d\rho_n(x)$ . If we wrote the analog of this for general  $z$ , we'd get

$$H(z) = \int_{\epsilon_n} H(x)d\rho_n(x, z)$$

varying the harmonic measure. Instead we use

$$h_n(z) = \int_{\bigcup_{K_j \in \mathcal{G}} K_j} G_\epsilon(x, z)d\rho_n(x)$$

which follows from noting that  $h_n$  vanishes on  $\epsilon$  and, on  $(\mathbb{C} \cup \{\infty\}) \setminus \epsilon$  obeys,  $\Delta h_n = d\rho_n \upharpoonright (\epsilon_n \setminus \epsilon)$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# A Poisson Formula

We proved this by thinking of  $d\rho_n$  as harmonic measure at  $\infty$ , i.e. if  $H$  is harmonic on  $(\mathbb{C} \cup \{\infty\}) \setminus \epsilon_n$  with boundary values  $H(x)$  on  $\epsilon_n$ , then  $H(\infty) = \int_{\epsilon_n} H(x) d\rho_n(x)$ . If we wrote the analog of this for general  $z$ , we'd get

$$H(z) = \int_{\epsilon_n} H(x) d\rho_n(x, z)$$

varying the harmonic measure. Instead we use

$$h_n(z) = \int_{\bigcup_{K_j \in \mathcal{G}} K_j} G_\epsilon(x, z) d\rho_n(x)$$

which follows from noting that  $h_n$  vanishes on  $\epsilon$  and, on  $(\mathbb{C} \cup \{\infty\}) \setminus \epsilon$  obeys,  $\Delta h_n = d\rho_n \upharpoonright (\epsilon_n \setminus \epsilon)$  so this is just the Poisson formula (proven by checking that the difference is harmonic on  $\Omega$  and vanishes on  $\epsilon$ ).

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Limits of $M_n$ are Gap Blaschke Products

By using the PW bound and the fact that

$\int G_{\epsilon}(x, z)\delta(x - x_k) = G_{\epsilon}(x_k, z)$ , it suffices to prove that, for all  $K \in \mathcal{G}$ , we have that

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Limits of $M_n$ are Gap Blaschke Products

By using the PW bound and the fact that

$\int G_{\epsilon}(x, z) \delta(x - x_k) = G_{\epsilon}(x_k, z)$ , it suffices to prove that, for all  $K \in \mathcal{G}$ , we have that

$$nd\rho \upharpoonright K \rightarrow \begin{cases} \delta(x - x_k), & \text{if } K \in \mathcal{G}_0 \\ 0, & \text{if } K \notin \mathcal{G}_0 \end{cases}$$

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# Limits of $M_n$ are Gap Blaschke Products

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

By using the PW bound and the fact that

$\int G_{\epsilon}(x, z) \delta(x - x_k) = G_{\epsilon}(x_k, z)$ , it suffices to prove that, for all  $K \in \mathcal{G}$ , we have that

$$nd\rho \upharpoonright K \rightarrow \begin{cases} \delta(x - x_k), & \text{if } K \in \mathcal{G}_0 \\ 0, & \text{if } K \notin \mathcal{G}_0 \end{cases}$$

If the zeros in a gap have a limit,  $x_k$ , in the gap, there is a single narrow band of  $\rho_n$ -weight  $1/n$  near the point so the first case is handled.



# Limits of $M_n$ are Gap Blaschke Products

- Chebyshev Polynomials
- Alternation Theorem
- Potential Theory
- Lower Bounds and Special Sets
- Faber Fekete Szegő Theorem
- Widom's Work
- Totik Widom Bounds
- Main Results
- Totik Widom Bounds
- Szegő–Widom Asymptotics

By using the PW bound and the fact that

$\int G_\epsilon(x, z) \delta(x - x_k) = G_\epsilon(x_k, z)$ , it suffices to prove that, for all  $K \in \mathcal{G}$ , we have that

$$nd\rho \upharpoonright K \rightarrow \begin{cases} \delta(x - x_k), & \text{if } K \in \mathcal{G}_0 \\ 0, & \text{if } K \notin \mathcal{G}_0 \end{cases}$$

If the zeros in a gap have a limit,  $x_k$ , in the gap, there is a single narrow band of  $\rho_n$ -weight  $1/n$  near the point so the first case is handled. If there is a zero that approaches an edge, the limit is 0 since regularity implies the Green's function  $G_\epsilon(x, z) = G_\epsilon(z, x)$  vanishes as  $x$  approaches  $\epsilon$ . ■



# And Now a Word from Our Sponsor

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics



# And Now a Word from Our Sponsor

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

*A Comprehensive Course in Analysis* by Poincaré Prize winner Barry Simon is a five-volume set that can serve as a graduate-level analysis textbook with a lot of additional bonus information, including hundreds of problems and numerous notes that extend the text and provide important historical background. Depth and breadth of exposition make this set a valuable reference source for almost all areas of classical analysis.

Part I is devoted to real analysis. From one point of view, it presents the infinitesimal calculus of the twentieth century with the ultimate integral calculus (measure theory) and the ultimate differential calculus (distribution theory). From another, it shows the triumph of abstract spaces: topological spaces, Banach and Hilbert spaces, measure spaces, Riesz spaces, Polish spaces, locally convex spaces, Fréchet spaces, Schwartz space, and  $L^p$  spaces. Finally it is the study of big techniques, including the Fourier series and transform, dual spaces, the Baire category, fixed point theorems, probability ideas, and Hausdorff dimension. Applications include the constructions of nowhere differentiable functions, Brownian motion, space-filling curves, solutions of the moment problem, Haar measure, and equilibrium measures in potential theory.



For additional information  
and updates on this book, visit  
[www.ams.org/bookpages/simon](http://www.ams.org/bookpages/simon)  
AMS on the Web  
[www.ams.org](http://www.ams.org)



Real Analysis

ANALYSIS  
Part  
1

Simon

Real Analysis  
A Comprehensive Course in Analysis, Part 1

Barry Simon

$$xy \leq \frac{x^p}{p} + \frac{y^q}{q}$$



$$\hat{f}(\mathbf{k}) = (2\pi)^{-\nu/2} \int \exp(-i\mathbf{k} \cdot \mathbf{x}) f(\mathbf{x}) d^\nu x$$



Google *simon comprehensive course preview*



# And Now a Word from Our Sponsor

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

*A Comprehensive Course in Analysis* by Poincaré Prize winner Barry Simon is a five-volume set that can serve as a graduate-level analysis textbook with a lot of additional bonus information, including hundreds of problems and numerous notes that extend the text and provide important historical background. Depth and breadth of exposition make this set a valuable reference source for almost all areas of classical analysis.

Part 2A is devoted to basic complex analysis. It interweaves three analytic threads associated with Cauchy, Riemann, and Weierstrass, respectively. Cauchy's view focuses on the differential and integral calculus of functions of a complex variable, with the key topics being the Cauchy integral formula and contour integration. For Riemann, the geometry of the complex plane is central, with key topics being fractional linear transformations and conformal mapping. For Weierstrass, the power series is King, with key topics being spaces of analytic functions, the product formulas of Weierstrass and Hadamard, and the Weierstrass theory of elliptic functions. Subjects in this volume that are often missing in other texts include the Cauchy integral theorem when the contour is the boundary of a Jordan region, continued fractions, two proofs of the big Picard theorem, the uniformization theorem, Ahlfors's function, the sheaf of analytic germs, and Jacobi, as well as Weierstrass, elliptic functions.




## Basic Complex Analysis

**Basic Complex Analysis**  
*A Comprehensive Course in Analysis, Part 2A*

Barry Simon

$$f(z_0) = \frac{1}{2\pi i} \int_{|z|=1} \frac{f(z)}{z - z_0} dz$$



ANALYSIS  
Part  
2A

Simon



1584-9705-1-4774-1100-8  
9 781470 411008  
SIMON/2.I

For additional information  
and updates on this book, visit  
[www.ams.org/bookpages/simon](http://www.ams.org/bookpages/simon)

AMS on the Web  
[www.ams.org](http://www.ams.org)



AMERICAN MATHEMATICAL SOCIETY

Google *simon comprehensive course preview*



# And Now a Word from Our Sponsor

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

*A Comprehensive Course in Analysis* by Poincaré Prize winner Barry Simon is a five-volume set that can serve as a graduate-level analysis textbook with a lot of additional bonus information, including hundreds of problems and numerous notes that extend the text and provide important historical background. Depth and breadth of exposition make this set a valuable reference source for almost all areas of classical analysis.

Part 2B provides a comprehensive look at a number of subjects of complex analysis not included as Part 2A. Presented in this volume are the theory of conformal metrics (including the Poincaré metric, the Ahlfors-Robinson proof of Picard's theorem, and Bell's proof of the Painlevé smoothness theorem), topics in analytic number theory (including Jacobi's two- and four-square theorems, the Dirichlet prime progression theorem, the prime number theorem, and the Hardy-Littlewood asymptotics for the number of partitions), the theory of Fuchsian differential equations, asymptotic methods (including Euler's method, stationary phase, the saddle-point method, and the WKB method), univalent functions (including an introduction to SLE), and Nevanlinna theory. The chapters on Fuchsian differential equations and on asymptotic methods can be viewed as a minicourse on the theory of special functions.



For additional information  
and updates on this book, visit  
[www.ams.org/bookpages/simon](http://www.ams.org/bookpages/simon)

AMS on the Web  
[www.ams.org](http://www.ams.org)



AMS

Advanced Complex Analysis

ANALYSIS  
Part  
2B

Simon

Advanced Complex Analysis  
A Comprehensive Course in Analysis, Part 2B

Barry Simon

$$\frac{\pi(x)}{(x/\log x)} \rightarrow 1$$



$$J_\alpha(x) \sim \sqrt{\frac{2}{\pi x}} \cos\left(x - \frac{\alpha\pi}{2} - \frac{\pi}{4}\right) + o(x^{-1/2})$$

AMS  
AMERICAN MATHEMATICAL SOCIETY

Google *simon comprehensive course preview*



# And Now a Word from Our Sponsor

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

*A Comprehensive Course in Analysis* by Poincaré Prize winner Barry Simon is a five-volume set that can serve as a graduate-level analysis textbook with a lot of additional bonus information, including hundreds of problems and numerous notes that extend the text and provide important historical background. Depth and breadth of exposition make this set a valuable reference source for almost all areas of classical analysis.



Part 3 returns to the themes of Part 1 by discussing pointwise limits (going beyond the usual focus on the Hardy-Littlewood maximal function by including ergodic theorems and martingale convergence), harmonic functions and potential theory, frames and wavelets,  $H^p$  spaces (including bounded mean oscillation (BMO) and, in the final chapter, lots of inequalities, including Sobolev spaces, Calderon-Zygmund estimates, and hypercontractive semigroups).



For additional information  
and updates on this book, visit  
[www.ams.org/bookpages/simon](http://www.ams.org/bookpages/simon)  
AMS on the Web  
[www.ams.org](http://www.ams.org)



AMS

Harmonic Analysis


ANALYSIS  
Part  
3

Simon

Harmonic Analysis

*A Comprehensive Course in Analysis, Part 3*

Barry Simon



$$|f - f_Q|_Q = \frac{1}{|Q|} \int_Q |f(x) - f_Q| d^d x$$

$$|\{x \mid M_{\text{HL}} f(x) > \alpha\}| \leq \frac{3^d}{\alpha} \|f\|_{L^1(\mathbb{R}^d, d^d x)}$$



AMERICAN MATHEMATICAL SOCIETY

Google *simon comprehensive course preview*



# And Now a Word from Our Sponsor

Chebyshev  
Polynomials

Alternation  
Theorem

Potential Theory

Lower Bounds  
and Special Sets

Faber Fekete  
Szegő Theorem

Widom's Work

Totik Widom  
Bounds

Main Results

Totik Widom  
Bounds

Szegő–Widom  
Asymptotics

A Comprehensive Course in Analysis by Poincaré Prize winner Barry Simon is a five-volume set that can serve as a graduate-level analysis textbook with a lot of additional bonus information, including hundreds of problems and numerous notes that extend the text and provide important historical background. Depth and breadth of exposition make this set a valuable reference source for almost all areas of classical analysis.

Part 4 focuses on operator theory, especially on a Hilbert space. Central topics are the spectral theorem, the theory of trace class and Fredholm determinants, and the study of unbounded self-adjoint operators. There is also an introduction to the theory of orthogonal polynomials and a long chapter on Banach algebras, including the commutative and non-commutative Gel'fand–Naimark theorems and Fourier analysis on general locally compact abelian groups.

ISBN: 978-1-4704-1103-9  
9 781470 411039  
SIMON/4

For additional information and updates on this book, visit [www.ams.org/bookpages/simon](http://www.ams.org/bookpages/simon)

AMS on the Web [www.ams.org](http://www.ams.org)



Operator Theory

ANALYSIS  
Part  
4

Simon

Operator Theory

A Comprehensive Course in Analysis, Part 4

Barry Simon



$$A = \int t \, dE_t$$

$$\det(1 + zA) = \prod_{k=1}^{N(A)} (1 + z\lambda_k(A))$$



Google *simon comprehensive course preview*