

HIERARCHICAL COMPLEXITY OF FINITE GROUPS

CHRYSTOPHER L. NEHANIV

In any field of mathematics, complexity is first level of sophistication beyond knowing the building blocks. –John Rhodes

ABSTRACT. What are simplest ways to construct a finite group from its atomic constituents? To understand part-whole relations between finite simple groups (“atoms”) and the global structure of finite groups, we axiomatize complexity measures on finite groups. From the Jordan-Hölder theorem and Frobenius-Lagrange embedding in an iterated wreath product, any finite group G can be constructed from a unique collection of simple groups, its Jordan-Hölder factors, each with well-defined multiplicities through iterated extension by simple groups. What is the least number of levels needed in such a hierarchical construction if a level is allowed to include several of these atomic pieces? To pose and answer this question rigorously, we give a natural set of hierarchical complexity axioms for finite groups relating to constructability, extension, quotients, and products, and prove these axioms are satisfied by a unique maximal complexity function \mathbf{cx} . We prove this function is the same as the minimal number of “spans of gems” (direct products of simple groups) in a subnormal series with all factors of this type. Hierarchical complexity \mathbf{cx} is thus effectively computable, and bounded below by all other complexity measures satisfying the axioms, including generalizations of derived length, Fitting height and solvability. Also, the hierarchical complexity of a normal subgroup is bounded above by the complexity of the whole group, although this is not assumed in the axioms and does not follow from them.

For solvable groups, the unique maximal group complexity measure defined satisfying the axioms agrees with the restriction of the one for all finite groups, and in addition satisfies an embedding axiom - which decidedly cannot be applied in the general case of all finite groups. In both cases, the complexity of a group is bounded above and below by various natural functions. In particular, hierarchical complexity is sharply bounded above by socle length, which yields a canonical decomposition and satisfies all the axioms except the extension axiom. Examples illustrate applications of the bounds and axiomatic methods in determining complexity of groups. We show also that minimal decompositions need not be unique in terms of what components occur nor their ordering.

The complexity axioms are also shown to be independent.

This work was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC), funding reference number RGPIN-2019-04669. Cette recherche a été financée par le Conseil de recherches en sciences naturelles et en génie du Canada (CRSNG), numéro de référence RGPIN-2019-04669.

The author also is grateful to Mr. Thomas George, recipient of a University of Waterloo President’s Award as undergraduate research assistant, for assistance with constructing software tools with which the counterexample groups in this paper were initially found and studied by the author.

CONTENTS

1. Introduction	3
1.1. Motivation and Brief Overview	3
1.2. Complexity Axioms for Finite Groups	3
2. The Unique Maximal Complexity Function	6
2.1. The Jordan-Hölder Function	6
2.2. Existence of a Maximal Group Complexity Function	7
3. The Subgroup Axiom (or “Embedding Axiom”)	8
4. Spans of Gems and Hierarchical Complexity	9
4.1. Spans of Gems: Splitting Lemmas I	10
4.2. Characterization of the Maximal Complexity Function	12
4.3. Computing Hierarchical Complexity	15
4.4. Wreath Product of Complexity 1 Groups	18
5. Minimal Normal Subgroups, Socle Length and Hierarchical Complexity	18
5.1. Minimal Normal Subgroups	19
5.2. Fragmentation and Splitting Lemma II	20
5.3. A Characteristic Series and Complexity Upper Bound	22
5.4. Socle Length and the Complexity Axioms	24
6. Other Complexity Functions: Lower and Upper Bounds	27
6.1. Uncountably Many Complexity Functions	27
6.2. Solvable Groups and the Embedding Axiom	29
6.3. Hierarchical Complexity of an Iterated Wreath Product	31
6.4. Hierarchical Complexity of p -torsion and other Groups	32
6.5. Derived Complexity, Fitting Complexity and Solvability Complexity	32
6.6. Wreath Product of Abelian Groups and Hierarchical Complexity of Iterated Wreath Products	36
7. Mutual Independence of the Complexity Axioms	37
7.1. Normal Subgroup Property is Independent from the Other Complexity Axioms	37
7.2. Independence of the Quotient Axiom	39
7.3. Independence of Each Complexity Axiom from the Rest	39
References	40

1. INTRODUCTION

1.1. Motivation and Brief Overview. This article follows the viewpoint of Rhodes [15] on complexity of mathematical objects, developing it for finite groups using a global, axiomatic approach. Such an axiomatic approach complements the study of extensions of finite groups initiated by Otto Hölder in the 19th century which attends to the local process of putting together groups from simpler parts. Such complexity theories develop the study of part-whole relations and provide a global hierarchical picture for understanding how simple building block components of objects may be put together to constituent the whole (albeit perhaps non-uniquely). Previously such an axiomatic complexity theory has been developed by Krohn, Rhodes, Tilson, Eilenberg, Steinberg and many others for finite semigroups and finite automata ([12], [7], [19], [14],[16, Ch. 4]) and for finite aperiodic semigroups [13]. Remarkably, although these areas are close to finite group theory and their axioms are abstracted from the research practices of group theory [15, Ch. 2-3], such a complexity theory for all finite groups has not been developed.¹ An appropriate formulation of the complexity axioms in the context of finite group theory allows us to do so here, at least some initial steps.

We axiomatize hierarchical complexity of finite group and study the functions satisfy the axioms and subsets of them. The axioms are shown to be independent and there is a unique maximal complexity function \mathbf{cx} on finite groups satisfying them. Certain well-known functions are also complexity functions on finite or finite solvable groups (Fitting height, derived length and others), which are lower bounds to \mathbf{cx} . The socle length is closely related to \mathbf{cx} , but shown to be a sharp upper bound and fails one of the axioms.

1.2. Complexity Axioms for Finite Groups. We consider functions \mathbf{c} from the class of finite groups to the natural numbers satisfying the following natural complexity axioms² :

- (1) Extension. If N is normal in G , then $\mathbf{c}(G) \leq \mathbf{c}(N) + \mathbf{c}(G/N)$.
- (2) Quotient. If a homomorphism maps G onto H , then $\mathbf{c}(G) \geq \mathbf{c}(H)$.
- (3) Normal. If N is normal in G , $\mathbf{c}(N) \leq \mathbf{c}(G)$.
- (4) Constructability from building blocks. Each group G either has complexity at most 1 or can be constructed by iterated extension from groups with complexity at most 1.
- (5) Product. If $G = H \times K$, then $\mathbf{c}(G) = \max(\mathbf{c}(H), \mathbf{c}(K))$.
- (6) Initial Condition. The complexity of the trivial group $\mathbf{1}$ is 0, i.e. $\mathbf{c}(\mathbf{1}) = 0$

¹The treatment of subgroups (or equivalently embedding) presents a subtle point (cf. Section 3 for this “Michelangelo Problem”), that is not an issue for solvable groups. It turns out we can take an agnostic approach that avoids any axiom mentioning the complexity of subgroups or complexity of normal subgroups and still develop an axiomatic theory that reveals how to handle this issue. As a consequence of this approach, it naturally follows that maximal complexity function and other measures satisfy the Normal property, i.e., that a normal subgroup has complexity no more than the complexity of the whole – although, as we show, this statement too is independent of the other axioms!

²All groups considered here are finite.

That “the constituent parts are not more complex than the whole” is captured by the quotient and normal properties.

We say a group G is an *extension of Q by N* if N is isomorphic to a normal subgroup of G and G/N is isomorphic to Q . The extension axiom expresses that the complexity of group is bounded by a simple function of the complexity of component parts of the group.

Since isomorphic groups map onto each other, any \mathbf{c} satisfying the quotient axiom must assign them the same value. So we may regard such a \mathbf{c} as a function on the set of isomorphism classes of finite groups. The quotient axioms says that a homomorphic image of a group is no more complex than the group - this is also called “the mapping axiom”.

Another immediate consequence of the quotient axiom is:

(2'). If N is normal in G , $\mathbf{c}(G/N) \leq \mathbf{c}(G)$.

For functions that are well-defined on isomorphism classes of finite groups, (2') is equivalent to the quotient axiom (2) since for $\varphi : G \twoheadrightarrow H$, by the first isomorphism theorem $H \cong G/N$ for $N = \ker \varphi$. Hence, (2') plus well-definedness on isomorphism classes may be used interchangeably with (2).

The product axiom captures the fact that direct product of factors does not combine them in a hierarchical way, but in parallel, which is not more complex than the independent factors.

Precisely, the constructability axiom says each G is *constructable from groups of complexity at most 1*, i.e., either $\mathbf{c}(G) \leq 1$ or G is an extension of Q by N where Q and N are constructable from groups of complexity at most 1.

Recall that if G is an extension of Q by N then G is isomorphic to a subgroup of the wreath product $N \wr Q$ which acts on the set $N \times Q$ (e.g., Frobenius-Lagrange coordinates [1, pp. 29–33], [4, Theorem 1.17], Kalužnin-Krasner Universal Embedding Theorem [11], [3, Theorem 2.6A]).³

Therefore, the constructability axiom may be restated as follows:

(4') Each group G has a subnormal series

$$1 = G_n \triangleleft G_{n-1} \triangleleft \cdots \triangleleft G_0 = G, \text{ with } \mathbf{c}(G_i/G_{i+1}) \leq 1,$$

for each $0 \leq i < n$ and some $n \geq 0$.

³Note that we follow [15, 16, 4] for basic results and notation for wreath products: Recall the (permutational) wreath product $W = N \wr Q$ is the group of mappings on $N \wr Q$ of the form $w = (w_2, w_1)$ where $(n, q)w = (nw_2(q), qw_1)$ for some $w_2 : Q \rightarrow N$ and $w_1 \in Q$. Then W is a semidirect product $N^Q \rtimes Q$. Similarly, the n -fold wreath product of G_n, \dots, G_1 is $W = G_n \wr \cdots \wr G_1$, consisting of functions w on the set $G_n \times \cdots \times G_1$ of the form $w = (w_n, \dots, w_1)$ with $w_i : G_{i-1} \times \cdots \times G_1 \rightarrow G_i$ ($1 \leq i \leq n$), so $w_1 \in G_1$, with $(g_n, \dots, g_1)w = (g_n w_n(g_{n-1}, \dots, g_2 w_2(g_1), g_1 w_1), g_i \in G_i)$. More generally, if $G_i \leq \text{Sym}(X_i)$, the symmetric group on set X_i , we write $(X_n, G) \wr \cdots \wr (X_1, G_1)$ for the permutation group W acting on $X_n \times \cdots \times X_1$ consisting of all $w = (w_n, \dots, w_1)$ with $w_i : X_{i-1} \times \cdots \times X_1 \rightarrow G_i$ ($1 \leq i \leq n$), $w_1 \in G_1$, with

$$(x_n, \dots, x_1)w = (x_n w_n(x_{n-1}, \dots, x_2 w_2(g_1), x_1 w_1),$$

$x_i \in X_i, g_i \in G_i$. Thus $G_n \wr \cdots \wr G_1$ denotes $(G_n, G_n) \wr \cdots \wr (G_1, G_1)$, where each G_i acts on itself by right multiplication.

Remarks and Questions on Global Structure: As the iterated (permutational) wreath product is associative, we may ‘parenthesize’ the quotients G_{i+1}/G_i ($0 \leq i < n$), corresponding to a subnormal series for a given group G arbitrarily in constructing the embedding of G in the iterated wreath product of these quotients:

$$G \leq G_{n-1}/G_n \wr \cdots \wr G_0/G_1.$$

Observe that the Jordan-Hölder factors of G are the union of the Jordan-Hölder factors for G_i/G_{i+1} with the same multiplicities. If each quotient G_i/G_{i+1} has complexity at most 1, then n bounds the complexity of G above (by the extension axiom). *Therefore, determining complexity of G is solved if we determine how to ‘pack’ the simple pieces of G (its Jordan-Hölder factors) into these quotient groups with a shortest hierarchical decomposition using the wreath product, giving the global structure of how the parts can be put together to build the whole.*⁴

Is such a minimal decomposition of G into constituent components canonical? Are the factors G_i/G_{i+1} constituting G for a minimal decomposition unique (up to isomorphism)? Is their order and manner of putting them unique? We shall see that for all finite groups there is a canonical decomposition via the socle series that sharply bounds complexity above, but that for hierarchical complexity itself neither the constituents nor their ordering are unique, as we show by examples. On the other hand, for solvable groups, both the Fitting series and derived series give lower bounds with canonical constituents.

Remarks on Axioms, Independence and Bounds: Hierarchical complexity will be constructed and shown to be the unique maximal complexity function satisfying the axioms. Therefore by constructing many complexity functions and functions that fail a single axiom or property, we shall be able to use them to show lower and upper bounds, respectively, for the hierarchical complexity on all groups and particular classes of groups as well as in concrete cases of computing complexity. This will also allow us to show that the complexity axioms themselves together with the normal subgroup property are logically independent of one another.

Moreover, the study and systematization of the properties of many well-known functions (such as nilpotency class, derived length, Fitting height, socle length, minimal number of generators, exponent, etc., - sometimes called *arithmetic functions*) on groups and subclasses of groups is achieved by placing them in the context of the complexity axioms allowing to be related and compared in a larger framework.

Lemma 1.1. *Let S be a finite simple group and \mathbf{c} a function on finite groups that satisfies the constructability axiom. Then $\mathbf{c}(S) \leq 1$.*

Proof. This follows from (4'), since each simple group has a unique subnormal series of length 1. □

⁴The embedding a group using iterated extensions into the wreath product constitutes a so-called a ‘cascade product’ of the factors synthesizing the group from these constituents [6] and is supported by computer algebraic tools of the SGPDEC package for GAP [5].

Conversely,

Lemma 1.2. *If a function on finite groups \mathbf{c} satisfies $\mathbf{c}(S) \leq 1$ for every simple group S , then \mathbf{c} satisfies the constructability axiom.*

Proof. Since every finite group G is either simple or has a maximal normal subgroup, G can be constructed by iterated extension from finite simple groups. In particular, G is isomorphic to a subgroup of the iterated wreath product of its Jordan-Hölder factors (e.g. [4, Theorem 1.17]). Alternatively, since each of the quotients S in a composition series for G is a simple group, constructability in the form (4') follows from the hypothesis that $\mathbf{c}(S) \leq 1$ for each of these. \square

It is now immediate that the constructability axiom is equivalent to the complexity of simple groups being bounded above by 1:

Lemma 1.3. *Let \mathbf{c} be a function from finite groups to \mathbb{N} . Then: \mathbf{c} satisfies the constructability axiom if and only if $\mathbf{c}(S) \leq 1$ for all S simple.*

Important Note: *In the sequel, we shall **not** assume the Normal property (3) hold for our complexity functions. Nevertheless, it will be shown to hold for the unique maximal complexity function \mathbf{cx} on finite groups and related measures.*

2. THE UNIQUE MAXIMAL COMPLEXITY FUNCTION

By the constructability axiom, each simple group has complexity at most one. The following is clear.

Lemma 2.1. (1) *The trivial complexity function z that assigns to each group the number zero satisfies the complexity axioms.*

(2) *The function δ with $\delta(1) = 0$ for the trivial group, and $\delta(G) = 1$ for G a nontrivial finite group satisfies the complexity axioms.*

So complexity functions exist.

A complexity function \mathbf{c} is *maximal* if it pointwise dominates any other one \mathbf{c}' satisfying the axioms, i.e., $\mathbf{c}(G) \geq \mathbf{c}'(G)$ for all finite groups G . If \mathbf{c} and \mathbf{c}' are both maximal, then $\mathbf{c}(G) \geq \mathbf{c}'(G)$ and $\mathbf{c}'(G) \geq \mathbf{c}(G)$, whence $\mathbf{c} = \mathbf{c}'$. Thus, if there is a maximal complexity function, it is unique.

2.1. The Jordan-Hölder Function. Let $\mathbf{JH}(G)$ be the number of Jordan-Hölder factors of a finite group G counting multiplicities.

Lemma 2.2. (1) *\mathbf{JH} satisfies the constructability, quotient, normal subgroup, initial condition, and extension axioms, but not the product axiom.*

(2) *If \mathbf{c} satisfies the axioms except possibly the product axiom, then $\mathbf{c}(G) \leq \mathbf{JH}(G)$ for any finite group.*

(3) *\mathbf{JH} is the unique maximal function on groups satisfying the axioms except possibly the product axiom.*

Proof. (1) Since the number of Jordan-Hölder factors of a simple group is 1 and every finite group can be constructed from simple groups by iterated extension, constructability holds for \mathbf{JH} . If G is an extension of Q by N ,

then since a composition series for G can be constructed from the one from Q by multiply each subgroup in the series by N , and followed by a composition series for N down to the trivial group. The quotients of this series are exactly the Jordan-Hölder factors of Q together with those of N counting multiplicities. Therefore $\mathbf{JH}(G) = \mathbf{JH}(N) + \mathbf{JH}(Q)$, whence extension holds for \mathbf{JH} . The quotient axiom $\mathbf{JH}(G/N) \leq \mathbf{JH}(G)$ holds since quotienting removes the occurrences of Jordan-Hölder factors of N from those of G . The product axiom does not hold since, e.g. $\mathbf{JH}(K \times H) = 2$ but $\max(\mathbf{JH}(K), \mathbf{JH}(H)) = 1$ for any simple groups K and H . Since the trivial group has no Jordan-Hölder factors $\mathbf{JH}(1) = 0$.

(2) Consider functions \mathbf{c} on all finite groups which satisfy all axioms except possibly the direct product axiom. Let

$$\epsilon(G) = \sup\{\mathbf{c}(G) : \mathbf{c} \text{ satisfies the complexity axioms, except possibly the product axiom}\}.$$

Suppose \mathbf{c} satisfies the axioms except possibly the direct product axiom. We claim $\mathbf{c}(G) \leq \mathbf{JH}(G)$ for every finite group G , showing this by induction on the length of a composition series for G , i.e., on $\mathbf{JH}(G)$. The case $\mathbf{JH}(G) = 0$ is trivial. If $\mathbf{JH}(G) = 1$, then G is simple, and G cannot be constructed by iterated extension from other groups, so $\mathbf{c}(G) \leq 1$ by constructability, and the claim holds. If $\mathbf{JH}(G) > 1$, then G is not simple and G has a maximal normal subgroup N with G/N is simple, so $\mathbf{JH}(G/N) = 1$. By the extension axiom, $\mathbf{c}(G) \leq \mathbf{c}(N) + \mathbf{c}(G/N)$. Now $\mathbf{c}(N) \leq \mathbf{JH}(N)$ by induction hypothesis, so $\mathbf{c}(G) \leq \mathbf{JH}(N) + 1 = \mathbf{JH}(G)$. We conclude that $\mathbf{c}(G) \leq \mathbf{JH}(G)$ for all G by induction. Since \mathbf{c} was arbitrary, $\epsilon(G) \leq \mathbf{JH}(G)$. Since \mathbf{JH} itself satisfies the axioms (except the product axiom), it follows that $\epsilon(G) = \mathbf{JH}(G)$ for all G . This proves (3). \square

2.2. Existence of a Maximal Group Complexity Function.

Theorem 2.3. *There is exists a unique maximal hierarchical complexity function \mathbf{cx} on finite groups satisfying the axioms, where*

$$\mathbf{cx}(G) = \sup\{\mathbf{c}(G) : \mathbf{c} \text{ satisfies the complexity axioms}\}.$$

Proof. By Lemma 2.1, the supremum is over a non-empty set of natural numbers. By Lemma 2.2(2), $\mathbf{JH}(G)$ is an upper bound on $\mathbf{c}(G)$ for each complexity function \mathbf{c} . Therefore $\mathbf{cx}(G)$ is natural number no larger than $\mathbf{JH}(G)$.

We claim \mathbf{cx} satisfies the complexity axioms.

Given G choose \mathbf{c} satisfying the axioms such that $\mathbf{c}(G)$ is maximal, then for any N normal in G , we have

$$\mathbf{cx}(G) = \mathbf{c}(G) \leq \mathbf{c}(N) + \mathbf{c}(G/N) \leq \mathbf{cx}(N) + \mathbf{cx}(G/N).$$

The first inequality holds since the axiom extension holds for \mathbf{c} . The second holds by definition of \mathbf{cx} . Therefore \mathbf{cx} satisfies extension.

For the quotient axiom, choose a complexity function \mathbf{c}_Q for which $\mathbf{c}_Q(G/N)$ is greatest, then by the quotient axiom for \mathbf{c}_Q and definition of \mathbf{cx}

$$\mathbf{cx}(G) \geq \mathbf{c}_Q(G) \geq \mathbf{c}_Q(G/N) = \mathbf{cx}(G/N).$$

Clearly, $\mathbf{cx}(1) = 0$ since $\mathbf{c}(1) = 0$ for each \mathbf{c} satisfying the axioms. For each finite simple group K , we have $\mathbf{cx}(K) \leq 1$ since for each complexity function $\mathbf{c}(K) \leq 1$. Hence, each group G can be constructed by iterated

extension from groups K for which $\mathbf{cx}(K) \leq 1$. Thus constructability axiom holds for \mathbf{cx} . (In fact, $\mathbf{cx}(K) \geq \delta(K) = 1$ for simple K shows $\mathbf{cx}(K) = 1$.)

For the product axiom, suppose $G = H \times K$, and choose a complexity function \mathbf{c} for which $\mathbf{c}(G)$ is greatest, \mathbf{c}' for which $\mathbf{c}'(H)$ is greatest, and \mathbf{c}'' for which $\mathbf{c}''(K)$ is greatest. Then we have

$$\begin{aligned} \mathbf{cx}(G) &= \mathbf{c}(G) \text{ by choice of } \mathbf{c} \\ &= \max\{\mathbf{c}(H), \mathbf{c}(K)\} \text{ by product axiom for } \mathbf{c} \\ &\leq \max\{\mathbf{c}'(H), \mathbf{c}''(K)\} \text{ by choice of } \mathbf{c}' \text{ and } \mathbf{c}'' \\ &= \max\{\mathbf{cx}(H), \mathbf{cx}(K)\} \text{ by choice of } \mathbf{c} \text{ and } \mathbf{c}'' \text{ and definition of } \mathbf{cx} \end{aligned}$$

On the other hand,

$$\begin{aligned} \mathbf{cx}(G) &\geq \max\{\mathbf{c}'(G), \mathbf{c}''(G)\} \text{ since } \mathbf{cx} \text{ is maximal at } G \\ &= \max\{\max\{\mathbf{c}'(H), \mathbf{c}'(K)\}, \max\{\mathbf{c}''(H), \mathbf{c}''(K)\}\} \text{ by the product axiom for } \mathbf{c}' \text{ and } \mathbf{c}'' \\ &\geq \max\{\mathbf{c}'(H), \mathbf{c}''(K)\} \\ &= \max\{\mathbf{cx}(H), \mathbf{cx}(K)\} \text{ by choice of } \mathbf{c}' \text{ and } \mathbf{c}'' \text{ and definition of } \mathbf{cx} \text{ at } H \text{ and at } K. \end{aligned}$$

This proves $\mathbf{cx}(H \times K) = \max\{\mathbf{cx}(H), \mathbf{cx}(K)\}$ as required.

By its definition $\mathbf{cx}(G) \geq \mathbf{c}(G)$ for any \mathbf{c} that also satisfies the axioms. It follows \mathbf{cx} is unique, since if \mathbf{c} were also pointwise maximal then $\mathbf{cx}(G) \geq \mathbf{c}'(G) \geq \mathbf{cx}(G)$, and so $\mathbf{cx}(G) = \mathbf{c}(G)$ for every group G . \square

From the argument of the proof, the following is clear.

Corollary 2.4. *The function given by the pointwise maximum of any set of complexity functions is complexity function. That is, let I index a set of complexity functions, then the function*

$$\mathbf{c}_I(G) = \max_{i \in I} \mathbf{c}_i(G),$$

for G a finite group, satisfies the complexity axioms.

3. THE SUBGROUP AXIOM (OR “EMBEDDING AXIOM”)

The sculpture is already complete within the marble block, before I start my work. It is already there, I just have to chisel away the superfluous material. —Michelangelo

It is possible to consider another axiom, namely that the complexity of a subgroup should be no more than the complexity of a group containing it:

Subgroup Axiom. If H is a subgroup of G , then $\mathbf{c}(H) \leq \mathbf{c}(G)$.

While this axiom sounds reasonable, it leads to mainly uninteresting complexity functions reminiscent of detecting only the marble block in which a beautiful sculpture might be found.

Proposition 3.1 (“The Michelangelo Problem”). *If a complexity measure \mathbf{c} satisfies the subgroup axiom, its value is bounded above by 1.*

Proof. By the next lemma, any finite group G embeds into a finite simple group S . This simple group has complexity at most 1 by Lemma 1.1, whence the subgroup axiom implies $\mathbf{c}(G) \leq \mathbf{c}(S) \leq 1$. \square

Lemma 3.2. *Every finite group G embeds in a finite simple alternating group.*

Proof. If $|G| \leq 3$ then G embeds in the smallest simple alternating group A_5 . If $|G|$ is at least 3, then consider the right regular representation of G acting on itself. This gives an embedding of G in the symmetric group on $|G|$ points. From this we can construct an embedding of G into an alternating group on $n = 2|G|$ points, which is a simple group since $n \geq 5$ as shown by Galois. Taking two disjoint copies of G as a set, let G act on each one as in the right regular representation within each copy by right multiplication. Whether $g \in G$ acts on the itself by an odd or even permutation, the action of g on the $2|G|$ points is always even since we have two disjoint copies of each cycle in g action on $|G|$. This gives an embedding of G in the simple group. \square

We shall see in section 6.2 that for complexity measures on solvable groups the subgroup axiom does not lead to the triviality seen in “Michelangelo problem” but is useful (and actually a consequence for the hierarchical complexity function and related functions on solvable groups).

4. SPANS OF GEMS AND HIERARCHICAL COMPLEXITY

Viewing finite simple groups as the rare ‘gems’ among finite groups, from which all finite groups are built, call a finite group G a *span of gems* if it is simple or the direct product of simple groups, i.e.,

$$G \cong K_1 \times \cdots \times K_n \text{ (with each } K_i \text{ simple, } 0 \leq i \leq n).$$

(Note: $n = 0$, G is the trivial group $\mathbf{1}$.) We also refer to such a group as a *mālā* (or *necklace*) of simple groups⁵ in line with terminology introduced for other complexity functions (cf. Section 6.5).

Indeed, from the constructability and product axioms, it follows that a span of gems can have complexity at most one. We shall see that for the maximal complexity function \mathbf{cx} on finite groups, the nontrivial spans of gems are exactly the groups of complexity of 1, and layering them leads to all higher complexity groups.

We first investigate spans of gems in Section 4.1, showing in detail how these groups are split as internal direct products by their normal subgroups, and also by their quotients. In Section 4.2, Theorem 4.6 characterizes the maximal complexity function on groups \mathbf{cx} as the hierarchical complexity measure μ where spans of gems comprise the ‘layers’ in a hierarchy in a minimal length decomposition of group G , with $\mathbf{cx}(G)$ -many layers in the recursive construction of finite groups from simpler ones by iterated extensions.

Notation: For a collection of groups \mathcal{S} , let $\text{Span}(\mathcal{S})$ denote all groups isomorphic to direct products $G_1 \times \cdots \times G_n$ with each $G_i \in \mathcal{S}$ ($0 \leq i \leq n$). We say G is a *span of groups from \mathcal{S}* (or a *mālā of \mathcal{S} groups*) if $G \in \text{Span}(\mathcal{S})$.

⁵This is a Sanskrit word referring to a circular chain of beads in the form of a rosary.

4.1. Spans of Gems: Splitting Lemmas I. In this subsection we prove two variants of “the span of gems splitting lemma”. Namely, we shall see that they are split by any normal subgroup in complementary spans of gems in a particular manner that is “rigid” for their simple non-abelian group (SNAG) factors but “fluid” for their maximal elementary abelian factors (direct powers of prime-order cyclic groups). These results reformulating classical facts (see references in the first paragraph of Section 5) are needed in the sequel.

We begin with this observation:

Lemma 4.1 (SNAG Factor Rigidity). *Let H be a simple non-abelian group (SNAG). Consider any direct product $H \times K$ with another group K . Let $\pi : H \times K \rightarrow K$ be the projection morphism. For $N \triangleleft H \times K$, if $(h_0, k_0) \in N$ with $h_0 \neq 1$ then $H \times \mathbf{1} \triangleleft N$, and $N = H \times \pi(N)$. Otherwise $N = \mathbf{1} \times \pi(N)$. In either case, $\pi(N) \triangleleft K$.*

Proof. If there is no (h_0, k_0) with h_0 a non-trivial element of H , then clearly $N = \mathbf{1} \times \pi(N)$, and also $\pi(N) \triangleleft K$ since $\mathbf{1} \times \pi(N) = N$ is normal in $H \times K$. Thus, the assertion holds.

Otherwise there is a $(h_0, k_0) \in N$ with $h_0 \neq 1$, so also $(h_0^{-1}, k_0^{-1}) \in N$. Since N is normal, for all $(h, k) \in H \times K$, $(h^{-1}h_0^{-1}h, k^{-1}k_0^{-1}k) \in N$. So taking $k = k_0$, and multiplying the following two members of N yields

$$(h^{-1}h_0^{-1}h, k_0^{-1}k_0^{-1}k_0)(h_0, k_0) = (h^{-1}h_0^{-1}hh_0, 1) \in N.$$

Since H is non-abelian, there is an $h \in H$ so $h^{-1}h_0^{-1}hh_0$ is not 1. Write $h_1 = h^{-1}h_0^{-1}hh_0$. Since $h_1 \neq 1$ and H is simple, the normal closure in $H \times K$ of the group generated by $(h_1, 1) \in N$ contains $H \times \mathbf{1}$. Thus, $H \times \mathbf{1} \leq N$. Of course $H \times \mathbf{1}$ is normal in N since it is normal in $H \times K$. Now for any $(h, k) \in N$, also $(1, k) \in N$ since $(h^{-1}, 1) \in H \times \mathbf{1} \leq N$ and $(1, k) = (h, k)(h^{-1}, 1)$. That is $\mathbf{1} \times \pi(N) \leq N$. Hence we can write each (h, k) in N as a member of $H \times \mathbf{1}$ times a member of $\mathbf{1} \times \pi(N)$. By normality of N , any conjugate of $(1, k)$ lies in N , so $\pi(N) \triangleleft K$ (since the H -coordinate of such a conjugate is 1). Thus N is the (internal) direct product of $H \times \mathbf{1}$ and $\mathbf{1} \times \pi(N) \triangleleft N$. \square

Thus if a normal subgroup intersects a SNAG factor of a direct product nontrivially, it contains the whole SNAG. By iterated the Lemma, we have,

Corollary 4.2. *A normal subgroup N of a direct product of SNAGs and another group K is the direct product of a subset of these SNAGs and the projection $\pi(K)$ of N to K , and, moreover, $\pi(N) \triangleleft K$.*

The ‘span of gems’ groups play a special role in the hierarchical complexity theory of finite groups and have special properties. Such a group can be “split along any normal subgroup” into a direct product:

Lemma 4.3 (Span of Gems - Splitting Lemma I). *If a finite group G is a direct product of simple groups, and $A \triangleleft G$, then there exists $B \triangleleft G$, such that $G = A \times B$ as an internal direct product.*

Proof. We may write G as $S_1 \times \cdots \times S_k$, with $k \geq 1$ and each factor S_i a simple group. We proceed by induction on $|G/A|$. If $|G/A| = 1$, then we

may choose the trivial subgroup of G as B and the assertion holds trivially. If $|G/A| > 1$, there exists must some S_i with $S_i \not\leq A$ (otherwise A would be all of G). Now S_i and A are normal in G properly contained $S_i \cap A$, which is normal in S_i , hence trivial. Since the commutator of any $s \in S_i$ and $a \in A$ lies in $S_i \cap A$ it follows that elements of S_i commute with elements of A . So $SA = AS \cong S \times A$ is an internal direct and clearly normal in G . Now $|G/SA| < |G/A|$, so by induction hypothesis applied to SA we have $G = (A \times S) \times C$, an internal direct product, for some $C \triangleleft G$. Letting $B = S \times C$, we have $G = A \times B$ as an internal direct product. \square

We call B as in Splitting Lemma I (Lemma 4.3) a *complement* to A in G .

Remarks on Splitting Lemma I: The following are not hard to see :

1. Note complements are not generally unique as can be seen by considering $\mathbb{Z}_p \times \mathbb{Z}_p$, and $\mathbb{Z}_p \times \mathbf{1}$, which has complements any the other $p^2 - p$ subgroups of size p .
2. The hypothesis of normality of A cannot dropped in if G has a nonabelian simple group as a factor, as, e.g., any proper subgroup of simple group as no complement in G .
3. Normality of A can be dropped if G has no simple nonabelian factor.
4. If all the factor S_i , then A consists of the factors it intersects has a unique complement consisting of the factors it does not intersect.

The proof of a variant of Lemma 4.3 gives more detail on this structure.

Lemma 4.4 (Span of Gems - Splitting Lemma I (Detailed Variant)). *Let G be a finite group which is the direct product of simple groups, and let N be a normal subgroup of G . Then,*

- (1) N is isomorphic to the direct product of simple groups.
- (2) G/N is isomorphic to a direct product of simple groups.
- (3) $G \cong N \times G/N$ as an internal direct product.

Proof. Let $N \triangleleft G = K_1 \times \cdots \times K_n$ with the K_i simple. Let $\pi_i : G \rightarrow K_i$ be the projection onto the i th factor. Let $I = \{i : \pi_i(N) \neq \mathbf{1}, 1 \leq i \leq n\}$. We may assume the first $m \leq n$ factors have $\pi_i(N) \neq \mathbf{1}$. Then $N = K_1 \times \cdots \times K_m \times \mathbf{1} \times \cdots \times \mathbf{1}$. the first ℓ factors are SNAGS and the rest are prime order cyclic groups ($0 \leq \ell \leq m$). By the SNAG rigidity applied successively for each SNAG factor we have that

$$N = K_1 \times \cdots \times K_\ell \times \pi(K_{\ell+1} \times \cdots \times K_m \times \mathbf{1} \times \cdots \times \mathbf{1}),$$

where $\pi : G \rightarrow K_{\ell+1} \times \cdots \times K_n$ is projection morphism and

$$\pi(N) \triangleleft K_{\ell+1} \times \cdots \times K_m \times \mathbf{1} \times \cdots \times \mathbf{1}.$$

Now $K_{\ell+1} \times \cdots \times K_m$ is a product of elementary abelian groups (direct products of prime power order groups), so we may assume

$$K_{\ell+1} \times \cdots \times K_m = (\mathbb{Z}_{p_1})^{n_1} \times \cdots \times (\mathbb{Z}_{p_k})^{n_k},$$

where the p_i are pairwise distinct primes and $n_i > 0$. It follows from the structure theory of finitely generated abelian groups that

$$\pi(N) \cong (\mathbb{Z}_{p_1})^{m_1} \times \cdots \times (\mathbb{Z}_{p_k})^{m_k},$$

with $0 \leq m_i \leq n_i$. Thus,

$$N = K_1 \times \cdots \times K_\ell \times (\mathbb{Z}_{p_1})^{m_1} \times \cdots \times (\mathbb{Z}_{p_k})^{m_k} \times \mathbf{1} \times \cdots \times \mathbf{1}.$$

This proves (1). Each elementary abelian group $(\mathbb{Z}_{p_i})^{n_i}$ is a n_i -dimensional \mathbb{Z}_p vector space and is an internal direct product of its m_i dimensional subspace $(\mathbb{Z}_{p_i})^{m_i}$ and a (non-unique) $n_i - m_i$ dimensional complement $(\mathbb{Z}_{p_i})^{n_i - m_i}$. It follows that

$$G/N \cong \mathbf{1} \times \cdots \times \mathbf{1} \times (\mathbb{Z}_{p_1})^{n_1 - m_1} \times \cdots \times (\mathbb{Z}_{p_k})^{n_k - m_k} \times K_{m+1} \times \cdots \times K_n,$$

where the first ℓ factors are $\mathbf{1}$ and the last $n - m$ are the factors for which $\pi_i(N) = \mathbf{1}$. This shows (2), and now (3) that

$$G = K_1 \times \cdots \times K_\ell \times (\mathbb{Z}_{p_1})^{n_1} \times \cdots \times (\mathbb{Z}_{p_k})^{n_k} \times K_m \times \cdots \times K_n$$

is the internal direct product of N and G/N follows. \square

Remark: Lemma 4.4(1) is not true if the word “normal” is removed, since by Lemma 3.2 any finite group is a subgroup of a (simple) alternating group.

4.2. Characterization of the Maximal Complexity Function. For each finite group G , define the *hierarchical complexity* $\mu(G)$ to be the shortest length ℓ for which G has a subnormal series

$$(\star) \quad \mathbf{1} = V_0 \triangleleft V_1 \triangleleft \cdots \triangleleft V_\ell = G$$

such that V_{i+1}/V_i is isomorphic to a span of gems (i.e., the direct product of simple groups) for $0 \leq i < \ell$. We say a series of length $\mu(G)$ of this kind gives *decomposition of G at its hierarchical complexity*.

We show μ is bounded by **JH** and satisfies the complexity axioms.

Proposition 4.5. *Let G be a finite group.*

- (1) $\mu(G) \leq \mathbf{JH}(G)$
- (2) (*Initial Condition*). $\mu(\mathbf{1}) = 0$,
- (3) $\mu(K) = 1$ if K is simple or the product of one or more simple groups.
- (4) (*Constructability Axiom*). G can be constructed by iterated extension from groups on which μ is no more than 1.
- (5) (*Product Axiom*). If $G = H \times K$, then $\mu(G) = \max(\mu(H), \mu(K))$.
For N normal in G ,
- (6) (*Quotient Axiom*). $\mu(G/N) \leq \mu(G)$
- (7) (*Extension Axiom*). $\mu(G) \leq \mu(N) + \mu(G/N)$.
- (8) (*Normality Property*). For N normal in G , $\mu(N) \leq \mu(G)$.

Proof. (1) holds since a composition series for G has simple groups as its quotients the Jordan-Hölder factors of G , each of which is trivially a 1-fold product of simple groups. (2) is trivial, while (3) is immediate from the definition of μ .

(4) The values of μ on $\mathbf{1}$, simple groups and their products follow immediately from the definition of μ . This entails constructability by iterated extension from groups on which μ takes value 1 (cf. Lemma 1.3).

(5) Suppose $\mu(H) = a$ and $\mu(K) = b$. Choose minimal length subnormal series for H and K such that each has quotients that are spans of gems, i.e., simple or the direct products of simple groups

$$1 = H_0 \triangleleft H_1 \triangleleft \cdots \triangleleft H_a = H,$$

$$1 = K_0 \triangleleft K_1 \triangleleft \cdots \triangleleft K_b = K,$$

We may assume $a \leq b$. If $a < b$, define $H_{a+1} = \cdots = H_b = H$. Now consider the series:

$$1 = H_0 \times K_0 \triangleleft H_1 \times K_1 \triangleleft \cdots \triangleleft H_b \times K_b = H \times K.$$

We have

$$(H_{i+1} \times K_{i+1})/(H_i \times K_i) \cong H_{i+1}/H_i \times K_{i+1}/K_i.$$

By choice of the series for H and K it follows that these factors are products of finite simple groups. This shows $\mu(H \times K) \leq \max\{\mu(H), \mu(K)\}$. On the other hand, if $G = H \times K$ had a shorter series with direct products of simple groups as its quotients G_{i+1}/G_i ,

$$1 = G_0 \triangleleft \cdots \triangleleft G_\ell = H \times K,$$

then we could project on the first coordinate with $\pi_H : H \times K \twoheadrightarrow H$ to obtain a subnormal series with groups $\pi_H(G_i) \leq H$: Since $\pi_H(gG_i) = \pi_H(g)\pi_H(G_i)$ for all $g \in G_{i+1}$, the projection π_H induces a morphism from G_{i+1}/G_i onto $\pi_H(G_{i+1})/\pi_H(G_i)$. By Lemma 4.3 (Splitting Lemma I), since G_{i+1}/G_i is a direct product simple groups so is $\pi_H(G_{i+1})/\pi_H(G_i)$. Now $\pi(G_{i+1})/\pi(G_i)$ are the quotients of a subnormal series for H of length less than $\max\{\mu(H), \mu(K)\}$. Similarly, for K . This would contradict the minimality of the chosen of series for H or K . Therefore, $\mu(H \times K) = \max\{\mu(H), \mu(K)\}$.

(6) On the other hand, applying the natural morphism from G to G/N to the series for G yields a series for G/N in which the successive quotients are homomorphic images of the quotients from the original series [18, pp. 102-3, first part of proof of Theorem 5.16]. Since homomorphic images of direct products of simple groups are also direct products of simple groups by Lemma 4.3, this series shows G/N has $\mu(G/N) \leq \mu(G)$.

(7) To establish the extension axiom we use strong induction on $\mu(G)$. Let $N \triangleleft G$. If $\mu(G) = 0$, then $G = 1$ and the result is trivial. If $\mu(G) = 1$, then G is the direct product of simple groups. By Lemma 4.4 (Splitting Lemma I - Detailed Variant), N and G/N are each also direct products of simple groups with $G \cong N \times G/N$. So $\mu(N)$ and $\mu(G/N)$ are less than or equal to 1. By (5) the direct product axiom for $\mu(G) = \max\{\mu(N), \mu(G/N)\} = 1$, and also $\mu(G) \leq \mu(N) + \mu(G/N)$ holds. Next suppose the extension property holds for all groups on which with μ is less than ℓ . Let G be such that $\mu(G) = \ell$, and take a subnormal minimal series of G whose quotients are products of simple groups.

Let $N \triangleleft G$ be a proper normal subgroup. We then have a series for N obtained from the series for G of length ℓ as in (\star) above.

$$1 = V_0 \cap N \triangleleft V_1 \cap N \triangleleft \cdots \triangleleft V_\ell \cap N = N,$$

Then each factor is

$$W_i = (V_{i+1} \cap N)/(V_i \cap N).$$

Now by Noether's isomorphism theorem that $HK/K \cong H/(H \cap K)$ for K normalized by H , we can take $H = V_{i+1} \cap N$ and $K = V_i$ (which is normal in V_{i+1} hence normalized by $V_{i+1} \cap N$), to conclude that

$$(V_{i+1} \cap N)V_i/V_i \cong (V_{i+1} \cap N)/(V_i \cap (V_{i+1} \cap N)) = (V_{i+1} \cap N)/(V_i \cap N) = W_i.$$

But note that

$$(V_{i+1} \cap N)V_i/V_i \leq V_{i+1}/V_i,$$

since $V_{i+1} \cap N$ and V_i are subgroups of V_{i+1} . Moreover, since $(V_{i+1} \cap N)V_i$ is normal in V_{i+1} ,

$$(V_{i+1} \cap N)V_i/V_i \triangleleft V_{i+1}/V_i,$$

by the Correspondence Theorem (e.g. [18, Theorem 2.28]). By choice of the series for G , V_{i+1}/V_i is the direct product of simple groups, so it follows by Lemma 4.4 (Splitting Lemma I) that $(V_{i+1} \cap N)V_i/V_i \cong W_i$ is the direct product of simple groups. Thus we have obtained series for N witnessing the fact that $\mu(N) \leq \ell = \mu(G)$.

Here is an alternative proof of (7). Now we show $\mu(G) \leq \mu(N) + \mu(G/N)$: Taking minimal series of whose whose groups are direct products of simple groups for N and G/N

$$\begin{aligned} \mathbf{1} &= A_0 \triangleleft A_1 \triangleleft \cdots \triangleleft A_a = N, \\ \mathbf{1} &= B_0 \triangleleft B_1 \triangleleft \cdots \triangleleft B_b = G/N, \end{aligned}$$

and letting $\varphi : G \twoheadrightarrow G/N$ be the natural quotient map, to obtain

$$N = \varphi^{-1}(B_0) \triangleleft \varphi^{-1}(B_1) \triangleleft \cdots \triangleleft \varphi^{-1}(B_b) = G,$$

with quotients $\varphi^{-1}(B_{i+1})/\varphi^{-1}(B_i)$ isomorphic to B_{i+1}/B_i by the Correspondence Theorem (e.g. [18, Theorem 2.28]). Concatenating the first and third series above yields a subnormal series for G , with all quotients direct products of simple groups, establishing $\mu(G) \leq a + b = \mu(N) + \mu(G/N)$. \square

Theorem 4.6. *The hierarchical complexity measure μ satisfies the complexity axioms and dominates any other complexity function \mathbf{c} . Hence, μ is \mathbf{cx} , the unique maximal complexity measure on groups.*

Proof. By Proposition 4.5, μ satisfies the complexity axioms. Let \mathbf{c} be any complexity measure satisfying the axioms. Then $\mu(\mathbf{1}) = \mathbf{c}(\mathbf{1}) = 0$. By definition of μ , if $\mu(G) = 1$ then G is a non-trivial direct product of simple groups. For simple groups K , $\mu(K) = 1 \geq \mathbf{c}(\mathbf{1})$ by Lemma 1.1 for \mathbf{c} . Thus, $\mathbf{c}(G)$ is the max of \mathbf{c} on the simple factors of G by the product axiom, hence at most one, i.e. $\mu(G) = 1 \geq \mathbf{c}(G)$.

We now continue by induction on $\mu(G) = \ell > 1$:

Take a shortest series for G with quotients isomorphic to direct products of simple groups :

$$\mathbf{1} = G_0 \triangleleft G_1 \triangleleft \cdots \triangleleft G_{\ell-1} \triangleleft G_\ell = G = G,$$

This series must a shortest one for $G_{\ell-1}$ of this type, lest G had a shorter such series. Therefore

$$\mu(G_{\ell-1}) = \ell - 1$$

and by induction hypothesis $\mathbf{c}(G_{\ell-1}) \leq \mu(G_{\ell-1})$, and so by the extension axiom for \mathbf{c}

$$\mathbf{c}(G) \leq \mathbf{c}(G_{\ell-1}) + \mathbf{c}(G/G_{\ell-1}) \leq (\ell - 1) + 1 = \mu(G).$$

We conclude by induction that μ dominates an arbitrary complexity function \mathbf{c} . Thus, by Theorem 2.3, $\mu = \mathbf{cx}$ \square

Corollary 4.7. *A nontrivial group has hierarchical complexity 1 if and only if it is a direct product of simple groups. That is, for a finite group $G \neq 1$,*

$$G \in \text{Span}(\text{SJMPL}\mathcal{E}) \Leftrightarrow \mathbf{cx}(G) = 1.$$

We observe that \mathbf{cx} satisfies the normal property though it was not assumed:

Theorem 4.8 (Normal Property of Hierarchical Complexity). *If N is a normal subgroup of G , then $\mu(N) \leq \mu(G)$. Hence, $\mathbf{cx}(N) \leq \mathbf{cx}(G)$.*

Proof. If N is normal in G , a minimal subnormal series for N with quotients all spans of gems can be extended to one for G . Thus $\mu(N) \leq \mu(G)$. By Theorem 4.6, $\mu = \mathbf{cx}$. \square

Do the complexity axioms, which do not include the normal property, imply it for an arbitrary \mathbf{c} satisfying them? No, the normal subgroup property is independent of the other axioms (as shown in Section 7.1). However, it holds for $\mu = \mathbf{cx}$ and many other important complexity functions (see Sections 6.2 and 6.5).

4.3. Computing Hierarchical Complexity. Let G be a finite group. By Theorem 4.6, $\mathbf{cx}(G) = \mu(G)$ can be computed by examining all subnormal chains whose quotients are spans of gems to find one of minimal length. To make determination of $\mathbf{cx}(G)$ easier, we can often compute or bound complexity of groups already from the axioms. Moreover, the normal property, as well as upper and lower bounds on \mathbf{cx} can simplify this further (Theorems 4.8, 6.20 & 5.12, and other results).

Example 1 (Span of Gems). By definition of μ , if $\mu(G) = 1$ then $G \neq 1$ is a direct product $G_1 \times \cdots \times G_n$ of finite simple groups G_i ($1 \leq i \leq n$, $n \geq 1$). So by Theorem 4.6, $\mathbf{cx}(G) = 1$ if and only if G is of this form, i.e., G is a “span of gems”.

Example 2 (Two Jordan-Hölder Factors). Let G be an extension of Q by N with Q and N simple. We have $\mathbf{cx}(Q) = \mathbf{cx}(N) = 1$. By the extension axiom $\mathbf{cx}(G) \leq \mathbf{cx}(Q) + \mathbf{cx}(N) = 2$. If G is the direct product $Q \times N$, $\mathbf{cx}(G) = 1$ by the product axiom. Otherwise, $\mathbf{cx}(G) = 2$ since the unique subnormal series $1 \triangleleft N \triangleleft G$ has length two, and quotients are spans of gems since they are simple.

Example 3 (Symmetric Groups). Let S_n be the symmetric group of all permutations on $\{1, \dots, n\}$, and let A_n be the alternating group, the subgroup of even permutations. If $n = 2$, $S_2 \cong \mathbb{Z}_2$ is simple and has complexity 1. Now $S_n/A_n \cong \mathbb{Z}_2$ and for $2 < n \neq 4$, A_n and \mathbb{Z}_2 are simple, so $\mathbf{cx}(S_n) = 2$ by example 2. For $n = 4$, $\mathbf{cx}(S_4) \leq \mathbf{cx}(A_4) + \mathbf{cx}(\mathbb{Z}_2)$. A_4 has a normal subgroup $\mathbb{Z}_2 \times \mathbb{Z}_2 \cong \langle (1, 2)(3, 4), (1, 3)(2, 4) \rangle$, with quotient \mathbb{Z}_3 , so $\mathbf{cx}(A_4) \leq \mathbf{cx}(\mathbb{Z}_2 \times \mathbb{Z}_2) + \mathbf{cx}(\mathbb{Z}_3) = 2$ by the extension and product axioms,

and the simplicity of \mathbb{Z}_2 and \mathbb{Z}_3 . It follows $\mathbf{cx}(S_4) \leq 3$. However, inspection shows $1 \triangleleft \langle \mathbb{Z}_2 \times \mathbb{Z}_2 \rangle \triangleleft A_4 \triangleleft S_4$ is a unique minimal subnormal series for S_4 with quotients all spans of gems, so $\mathbf{cx}(S_4) = 3$.

Example 4. Let \mathbb{Z}_{p^n} denote the cyclic group generated by x of order p^n for p prime, $n \geq 1$. Then \mathbb{Z}_{p^n} has a normal subgroup $\langle x^p \rangle$ isomorphic to $\mathbb{Z}_{p^{n-1}}$ with quotient $\mathbb{Z}_{p^n}/\mathbb{Z}_{p^{n-1}} \cong \mathbb{Z}_p$, so by the extension axiom, $\mathbf{cx}(\mathbb{Z}_{p^n}) \leq \mathbf{cx}(\mathbb{Z}_{p^{n-1}}) + \mathbf{cx}(\mathbb{Z}_p)$. Since $\mathbf{cx}(\mathbb{Z}_p) = 1$, it follows by induction that $\mathbf{cx}(\mathbb{Z}_{p^n}) \leq n$. (We shall see that n is also a lower bound in example 11.)

The number of factor groups in any minimal decomposition at complexity is $\mu(G)$ but neither the order nor the isomorphism classes of the quotients need to be the same:

Counterexample 5 (Order of Factors Not Unique at Complexity). The group of symmetries of the square, $D_4 \cong \mathbb{Z}_2 \wr \mathbb{Z}_2$ has $\mu(D_4) = 2$ but the quotients \mathbb{Z}_2 and $\mathbb{Z}_2 \times \mathbb{Z}_2$ can appear in either order in decompositions at complexity.

Group	Subnormal Chain	Group	Subnormal Chain
D_4	$= \langle (1, 2), (3, 4), (1, 3)(2, 4) \rangle$	D_4	$= \langle (1, 2), (3, 4), (1, 3)(2, 4) \rangle$
\mid	quotient $\mathbb{Z}_2 \times \mathbb{Z}_2$	\mid	quotient \mathbb{Z}_2
\mathbb{Z}_2	$= \langle (1, 2)(3, 4) \rangle$	$\mathbb{Z}_2 \times \mathbb{Z}_2$	$= \langle (1, 2)(3, 4), (1, 2) \rangle$
\mid	quotient \mathbb{Z}_2	\mid	quotient $\mathbb{Z}_2 \times \mathbb{Z}_2$
$\mathbf{1}$	$= \mathbf{1}$	$\mathbf{1}$	$= \mathbf{1}$

Counterexample 6 (Isomorphism Types of Factors Not Unique at Complexity). Also the isomorphism types of the quotients are not unique for decompositions at complexity, for $\mathbb{Z}_2 \wr \mathbb{Z}_4$. Inspection of all chains of subnormal subgroups shows complexity $\mu(\mathbb{Z}_2 \wr \mathbb{Z}_4) = 3$, where all three quotients may be isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$, or the non-isomorphic \mathbb{Z}_2 , $\mathbb{Z}_2 \times \mathbb{Z}_2$, $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ (with order not unique) for different subnormal chains, or even $(\mathbb{Z}_2)^4$, \mathbb{Z}_2 , \mathbb{Z}_2 for the subnormal series

$$\mathbf{1} \triangleleft (\mathbb{Z}_2)^4 \cong (\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2) \rtimes \mathbf{1} \triangleleft (\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2) \rtimes \mathbb{Z}_2 \triangleleft \mathbb{Z}_2 \wr \mathbb{Z}_4,$$

Either \mathbb{Z}_2 or $\mathbb{Z}_2 \times \mathbb{Z}_2$ must occur as the topmost quotient of $\mathbb{Z}_2 \wr \mathbb{Z}_4$, but with this constraint all possible orderings of direct products of the six \mathbb{Z}_2 Jordan-Hölder factors of $\mathbb{Z}_2 \wr \mathbb{Z}_4$ occur in some decomposition at complexity. Here are some more examples:

$$\begin{array}{lcl} \mathbb{Z}_2 \wr \mathbb{Z}_4 = \langle (1, 2), (3, 4), (5, 6), (7, 8), (1, 3, 5, 7)(2, 4, 6, 8) \rangle & & \\ \mid & & \text{quotient } \mathbb{Z}_2 \\ (\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2) \rtimes \mathbb{Z}_4 \cong \langle (5, 6)(7, 8), (3, 4)(5, 6), (1, 2)(3, 4), (1, 3, 5, 7)(2, 4, 6, 8) \rangle & & \\ \mid & & \text{quotient } \mathbb{Z}_2 \times \mathbb{Z}_2 \\ \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 = \langle (1, 2)(5, 6), (3, 4)(7, 8), (1, 5)(2, 6)(3, 7)(4, 8) \rangle & & \end{array}$$

Also, the subnormal series

$$\begin{array}{lcl} \mathbb{Z}_2 \wr \mathbb{Z}_4 = \langle (1, 2), (3, 4), (5, 6), (7, 8), (1, 3, 5, 7)(2, 4, 6, 8) \rangle & & \\ \mid & & \text{quotient } \mathbb{Z}_2 \times \mathbb{Z}_2 \\ \mathbb{Z}_2 \times D_4 \cong \langle (5, 6)(7, 8), (3, 4)(5, 6), (1, 2)(3, 4), (1, 5)(2, 6)(3, 7)(4, 8) \rangle & & \end{array}$$

may be continued to a subnormal series at complexity 3 by various normal subgroups of $\mathbb{Z}_2 \times D_4$ yielding non-isomorphic quotients in different orders:

$$\begin{array}{ccc}
 & | & \text{quotient } \mathbb{Z}_2 \\
 \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \cong & \langle (1, 2)(3, 4)(5, 6)(7, 8), (1, 5)(2, 6)(3, 7)(4, 8), (3, 4)(7, 8) \rangle & \\
 & \text{or} & \\
 & | & \text{quotient } \mathbb{Z}_2 \times \mathbb{Z}_2 \\
 \mathbb{Z}_2 \times \mathbb{Z}_2 \cong & \langle (1, 2)(3, 4)(5, 6)(7, 8), (1, 5)(2, 6)(3, 8)(4, 7) \rangle & \\
 & \text{or} & \\
 & | & \text{quotient } \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \\
 \mathbb{Z}_2 \cong & \langle (1, 2)(3, 4)(5, 6)(7, 8) \rangle &
 \end{array}$$

In fact, there are 15 different subnormal subseries at complexity for $\mathbb{Z}_2 \wr \mathbb{Z}_4$.

Remark 4.9 (Diversity of Constructions at Complexity).

- (1) There may be multiple ways to build a group by a *minimal number* of extensions from simple groups and their direct products (as Counterexamples 5 and 6 show). In other words, ‘the’ manner of constructing a group G by iterated extension even *in a minimal way* (i.e., at complexity) is not always unique in terms of type of ‘pieces’ and the order of extension to constitute the ‘whole’. There can be multiple dissimilar solutions to building G using $\mathbf{cx}(G)$ pieces of complexity 1 in terms of how Jordan-Hölder factors are distributed amongst the ‘pieces’ (i.e., constituents of the iterated extension).
- (2) Iterated extensions using isomorphic pieces in the same order may correspond to essentially different minimal constructions of G . For example, in the first 3-mālā minimal decomposition of $\mathbb{Z}_2 \wr \mathbb{Z}_4$ in Counterexample 6, the normal group of $(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2) \rtimes \mathbb{Z}_2 = \langle (1, 5)(2, 6)(3, 7)(4, 8), (7, 8), (5, 6), (3, 4), (1, 2) \rangle$ can be replaced by $\langle (3, 4)(7, 8), (1, 2)(5, 6), (5, 6)(7, 8) \rangle \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$, to yield a minimal construction of $\mathbb{Z}_2 \wr \mathbb{Z}_4$ at complexity with the same constituents \mathbb{Z}_2 , $(\mathbb{Z}_2)^2$, $(\mathbb{Z}_2)^3$ in the same order as in the second decomposition in Counterexample 6 above. Although built from the same pieces $(\mathbb{Z}_2)^2$ and $(\mathbb{Z}_2)^3$ in the same order, the middle groups are of the form $(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2) \rtimes \mathbb{Z}_4$ and $(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2) \rtimes \mathbb{Z}_2$ but not isomorphic, since as can be shown they have centers of size 2 and 4, respectively. Nevertheless, \mathbb{Z}_2 can be extended by either of these middle groups to yield a minimal construction of $\mathbb{Z}_2 \wr \mathbb{Z}_4$.

Example 7 (Nilpotent Groups). Since a nilpotent group N is a direct product $N = P_1 \times \cdots \times P_k$ of its Sylow p -groups (e.g., [10, Theorem 10.3.4]), by the product axiom, $\mathbf{cx}(N)$ is the maximum complexity $\mathbf{cx}(P_i)$. This reduces the computation of the complexity of nilpotent groups to computing complexity of p -groups.

Example 8 (Solvable Group Complexity Bounds). If G is a solvable group, G has a minimal length subnormal series in which quotients N_i are nilpotent. The length of such a series is the Fitting height of G (see also Section 6.5, Rhodes’ Lemma 6.23). Refining this series using minimal length series for the N_i whose quotients are spans of gems from example 7 yields a subnormal

series whose length is an upper bound for $\mathbf{cx}(G)$. By the extension property of complexity,

$$\mathbf{cx}(G) \leq \sum_{i=1}^n \mathbf{cx}(N_i) \leq \sum_{i=1}^n \sum_{j=1}^{k_j} \max\{\mathbf{cx}(P_1^{(i)}), \dots, \mathbf{cx}(P_{k_j}^{(i)})\},$$

where $P_1^{(i)}, \dots, P_{k_j}^{(i)}$ ($k_j \geq 1$) denote the Sylow p -subgroups of the nilpotent group N_i as in example 7.

4.4. Wreath Product of Complexity 1 Groups. We can apply some of what we have just seen to compute the (hierarchical) complexity of the wreath product of two complexity 1 groups.

Proposition 4.10. *If N and Q have hierarchical complexity 1, then the wreath product $N \wr Q$ has hierarchical complexity 2. In particular, if N and Q are simple groups, their wreath product has complexity 2.*

Proof. The wreath product $W = N \wr Q$ is an extension of Q by the $|Q|$ -fold direct product $N^Q = N \times \dots \times N$, and is a semidirect product $N^Q \rtimes Q$. By the extension and product axioms $\mathbf{cx}(W) \leq \mathbf{cx}(Q) + \mathbf{cx}(N^Q) = \mathbf{cx}(Q) + \mathbf{cx}(N)$.

If N and Q are spans of gems (complexity 1), then $0 \neq \mathbf{cx}(W) \leq 2$. We show $\mathbf{cx}(W) = 2$. Suppose for a contradiction that $\mathbf{cx}(W) = 1$ then we would have $N \wr Q$ a span of gems. Consider the natural morphism $\pi : W \rightarrow Q$. The kernel of π is $N^Q \rtimes \mathbf{1} \triangleleft W$ and $\ker(\pi) \cong N^Q$. By Splitting Lemma I (Lemma 4.3), W is an internal direct product $(N^Q \rtimes \mathbf{1}) \times B$, for B a normal complement to $(N^Q \rtimes \mathbf{1})$ in W , $B \triangleleft W$ with $B \cap (N^Q \rtimes \mathbf{1})$ is trivial. We have $B \cong W/(N^Q \rtimes \mathbf{1}) \cong Q$. Consider the subgroup $\mathbf{1}^Q \times Q$ of W . Choose any $(c_1, q) \in \mathbf{1}^Q \times Q$ where $c_1 : Q \rightarrow \mathbf{1} \leq N$ is the constant function, with $q \neq 1$. Since $W = (N^Q \rtimes \mathbf{1})B$, we may write (c_1, q) as

$$(c_1, q) = (w_2, 1)(b_2, b_1) \text{ with } (w_2, 1) \in N^Q \rtimes \mathbf{1} \text{ and } (b_2, b_1) \in B$$

Here $w_2, b_2 \in N^Q N$, and $1, b_1 \in Q$. Now $(c_1, q) = (w_2, 1)(b_2, b_1) = (w_2 b_2, b_1)$. Hence $b_1 = q$ and $c_1 = w_2 b_2$, so $b_2 = w_2^{-1}$. Now since $B \triangleleft W$, we have $(c_1, q)^{-1}(b_2, b_1)(c_1, q) = (c_1, q^{-1})(w_2^{-1}, q)(c_1, q) = (w_2^{-1}, q^{-1}qq) = (w_2^{-1}, 1) \in B$. So, $(w_2^{-1}, 1) \in (N^Q \rtimes \mathbf{1}) \cap B = \mathbf{1}$. This shows (b_2, b_1) is conjugate to the identity, hence $(b_2, b_1) = 1$. Hence $(c_1, q) = (w_2, 1)(b_2, b_1) = (w_2, 1)$. This contradicts $q \neq 1$. Hence $\mathbf{cx}(W) \neq 1$, but rather $\mathbf{cx}(W) = 2$. \square

Remark: One can easily check $\mathbf{1}^Q \rtimes N$ is not normal in $W = N \wr Q$, but this is implied by the above argument, for otherwise $\mathbf{1}^Q \rtimes Q$ would be a normal complement to $(N^Q \rtimes \mathbf{1})$ in W .

5. MINIMAL NORMAL SUBGROUPS, SOCLE LENGTH AND HIERARCHICAL COMPLEXITY

To begin to develop the ‘semi-local’ theory relating finite groups of complexity 1 (spans of gems) occurring as hierarchical layers within the structure of higher complexity groups that contain them, this section develops material related on minimal normal subgroups (e.g., [10], [3], [18]) and the socle of a finite group (e.g., [3], [2]). While much of this is classical and well-known, there are also some new formulations and results that we shall need

subsequently for the complexity theory of finite groups. These include more detailed splitting and fragmentation lemmas of Sections 4.1 and 5.2, and lead to proofs of original results that socle length satisfies the normal and quotient axioms, but *not* the extension axiom, and is a sharp upper bound for hierarchical complexity.

5.1. Minimal Normal Subgroups. A normal subgroup $M \neq \mathbf{1}$ of a group G is a *minimal normal subgroup* of G if it contains no other non-trivial normal subgroup of G . That is, $\mathbf{1} \neq K \triangleleft G$ with $K \leq M$ implies $K = M$. We make some observations on minimal normal subgroups. By the Correspondence Theorem (e.g., [18, Theorem 2.28]), the following is clear.

Lemma 5.1. *The image of a minimal normal subgroup under a surjective map is either a minimal normal subgroup of the image or is trivial.*

We shall make use of the following well-known elementary Fact.

Fact 5.2. *If A and B are normal subgroups of a group G and $A \cap B = \mathbf{1}$, then $ab = ba$ for all $a \in A, b \in B$. Moreover, the group $\langle A, B \rangle$ generated by A and B equals $AB \triangleleft G$, and AB is the internal direct product isomorphic to $A \times B$*

Proof. $(b^{-1}a^{-1}b)a \in A$ since $b^{-1}a^{-1}b \in A$, and similarly $b^{-1}(a^{-1}ba) \in B$ since $a^{-1}ba \in B$. So $b^{-1}a^{-1}ba = 1$, i.e., a and b commute. In any product in $G = \langle A, B \rangle$, we can move all a 's to the left and all b 's to the right, so $\langle A, B \rangle = AB$. To multiply in AB , we have $(ab)(a'b') = a(ba')b' = aa'bb'$ ($a, a' \in A, b, b' \in B$). Also for $g \in G, ab \in AB$, we have $g^{-1}abg = (g^{-1}ag)(g^{-1}bg) \in AB$, so AB is normal in G . Each element of AB can be uniquely written in the form ab : for if $ab = a'b'$ then it follows $a^{-1}a' = bb'^{-1} \in A \cap B = \mathbf{1}$, whence $a = a'$ and $b = b'$. Thus the map $(a, b) \mapsto ab$ is a surjective and injective morphism, showing $A \times B \cong AB$. \square

Fact 5.3. (1) *Let N_1, \dots, N_n be minimal normal subgroups of M . Then the subgroup they generate is an internal direct product and normal in M . Furthermore, $\langle N_1, \dots, N_n \rangle \cong N_{i_1} \times \dots \times N_{i_m}$, for some $1 \leq m \leq n$ and $1 = i_1 < \dots < i_m \leq n$.*

(2) *Let M be a minimal normal subgroup of G , and N be a minimal normal subgroup of M . Then M is the internal direct product of N and (zero or more) conjugates of N .*

Proof. (1) Inductively define $M_1 = N_1$, and for $0 \leq i < n$,

$$M_{i+1} := \begin{cases} \langle M_i, N_{i+1} \rangle & \text{if } \mathbf{1} = M_i \cap N_{i+1} \\ M_i & \text{otherwise.} \end{cases}$$

We proceed by induction from 1 up to n , to show that $M_i = \langle N_j : 1 \leq j \leq i \rangle$, that M_i is the internal direct product of a subset of $\{N_1, \dots, N_i\}$, and that $M_i \triangleleft M$: This is trivially for $i = 1$, since $M_1 = N_1 \triangleleft M$. Now suppose this assertion is true for i ($1 \leq i < n$). On the one hand, if $\mathbf{1} = M_i \cap N_{i+1}$, by Fact 5.2, $M_{i+1} = \langle M_i, N_{i+1} \rangle = M_i N_{i+1} \cong M_i \times N_{i+1}$, and $M_{i+1} \triangleleft M$. By induction hypothesis $M_i = \langle N_j : 1 \leq j \leq i \rangle$ is the internal direct product a subset of $\{N_1, \dots, N_i\}$, so it follows M_{i+1} is the internal direct product of a subset of $\{N_1, \dots, N_{i+1}\}$. On the other hand if $\mathbf{1} \neq M_i \cap N_{i+1}$, then

$M_i \cap N_{i+1} \triangleleft M$ since $M_i \triangleleft M$ (by induction hypothesis) and $N_{i+1} \triangleleft M$. So minimal normality of N_{i+1} in M entails $M_i \cap N_{i+1} = N_{i+1}$, so $N_{i+1} \leq M_i$. In either case, $N_{i+1} \leq M_{i+1}$, and so $M_{i+1} = \langle M_i, N_{i+1} \rangle = \langle N_1, \dots, N_i, N_{i+1} \rangle$. This proves the assertion for $i + 1$. By induction, the assertion holds $i = n$ and (1) is proved.

(2) Let $g_1 (= 1), \dots, g_n$ be elements of G , such that the conjugates $N_i = g_i^{-1}Ng_i$ are pairwise distinct for $i \neq j$ ($1 \leq i, j \leq n$), and each conjugate of $g^{-1}Ng$ in G occurs as $g_i^{-1}Ng_i$ for some i . Since conjugation is an automorphism of G , we have $g^{-1}Ng$ is minimal normal in $g^{-1}Mg = M$ (since $M \triangleleft G$). Applying (1) yields that

$$M' := \langle g^{-1}Ng : g \in G \rangle = \langle g_i^{-1}Ng_i : 1 \leq i \leq n \rangle = N_{i_1} \dots N_{i_n} \triangleleft M, \text{ and}$$

$$M' \cong N_{i_1} \times \dots \times N_{i_n},$$

where $i_1 = 1$. Clearly, M' is the normal closure of N in G , hence by minimal normality of M in G , $M' = M$. \square

Fact 5.4. *A minimal normal subgroup M of a finite group G is simple or the (internal) direct product of isomorphic simple groups.*

Proof. Let S be a minimal normal subgroup of M . By Fact 5.3(2), M is the internal direct product of conjugates of S (including S itself). Suppose $\mathbf{1} < K \triangleleft S$, then S normalizes K and each internal direct product factor different from S commutes with S , hence with K . Thus $K \triangleleft M$. Minimal normality of S in M now implies $K = S$. Therefore S is simple. \square

In particular, a minimal normal subgroup is a span of gems.

5.2. Fragmentation and Splitting Lemma II. The *socle* $\text{soc}(G)$ of a finite group is the subgroup of G generated by the minimal normal subgroups of G .

Theorem 5.5. *For a finite group $G \neq \mathbf{1}$, $\text{cx}(\text{soc}(G)) = 1$.*

Proof. It is the internal direct product of minimal normal subgroups and normal in G by Lemma 5.3(1). By Fact 5.4, each of these minimal normal subgroups is a direct product of simple groups, so $\text{soc}(G)$ is too. It follows $\text{cx}(G) = \mu(G) = 1$. \square

If $\text{soc}(G)$ intersects the normal subgroup N of G non-trivially, a minimal normal subgroup M of G completely “fragments” as a product of isomorphic minimal normal subgroups of N , and lies in $\text{soc}(N)$:

Fact 5.6 (Fragmentation Lemma). *Let M be a minimal normal subgroup of G and $N \triangleleft G$. Suppose $M \cap N \neq \mathbf{1}$. Then*

- (1) *M is internal direct product of isomorphic minimal normal subgroups of N .*
- (2) *$M = M \cap N$.*
- (3) *(Absorption 1). $M \triangleleft N$.*
- (4) *(Absorption 2). $M \triangleleft \text{soc}(N)$.*

Proof. (2) Since $N \triangleleft G$, $\mathbf{1} \neq M \cap N$ is normal in G . Since M is minimal normal in G , $M \cap N = M$. (3) follows from (2), since $M \cap N \triangleleft N$. (1) We have $\mathbf{1} \neq M \cap N \triangleleft N$. Let $K \neq \mathbf{1}$ be a minimal normal a subgroup of N

contained in $M \cap N$. For any $g \in G$, $g^{-1}Kg$ lies in N since $N \triangleleft G$. Since conjugation is an automorphism, $g^{-1}Kg$ is a minimal normal subgroup of $N = g^{-1}Ng$. By minimal normality in N , either $K \cap g^{-1}Kg = 1$ or $K = g^{-1}Kg$. Choose representatives $g_1, \dots, g_k \in G$ so that $g_1^{-1}Kg_1, \dots, g_k^{-1}Kg_k$ comprise the collection of all the distinct conjugates of K . Since $K \leq M \cap N = M$ (by (2)) and each $g^{-1}Kg \leq g^{-1}Mg = M$, therefore, $M' = \langle g^{-1}Kg : g \in G \rangle \leq M$, but this is the normal closure of K in G so $M' \triangleleft M$. By minimal normality of M in G , $M' = M$. By Fact 5.3(1), the group M' is normal in N and is isomorphic to an internal direct product copies of K . Therefore M is the internal direct product of K and conjugates of K in G . (4) By (1), M is the product of minimal normal subgroups of N , hence $M \triangleleft \text{soc}(N)$. \square

Lemma 5.7 (Span of Gems - Splitting Lemma II). *Let N be a normal subgroup of a finite group G , then*

- (1) $\text{soc}(G) \cap N = M_1 \dots M_k = N_1 \dots N_m$, is the internal direct product of some minimal normal subgroups M_i of G ($m \geq 0, 1 \leq i \leq m$), and also the internal direct product of some minimal normal subgroups N_j of N ($m \geq 0, 1 \leq j \leq m \leq n$).
- (2) In particular, $\text{soc}(G) \cap N \leq \text{soc}(N)$, and $\text{soc}(G) \cap N \triangleleft \text{soc}(N)$.
- (3) The complement of $N_1 \dots N_m$ in $\text{soc}(G)$ is an internal direct product of minimal normal subgroups M' of G that intersect N trivially. That is, for some M'_1, \dots, M'_s minimal normal in G

$$\text{soc}(G) = (\text{soc}(G) \cap N) \times M'_1 \times \dots \times M'_s,$$

is an internal direct product, with each $M'_i \cap \text{soc}(N) = 1$, $s \geq 0$ ($1 \leq i \leq s$).

- (4) The complement of $N_1 \dots N_m$ in $\text{soc}(N)$ is the internal direct product of minimal normal subgroups of N' that intersect $\text{soc}(G)$ trivially. That is, for some N'_1, \dots, N'_t minimal normal in N

$$\text{soc}(N) = (\text{soc}(G) \cap N) \times N'_1 \times \dots \times N'_t,$$

is an internal direct product, with each $N'_i \cap \text{soc}(G) = 1$, $t \geq 0$ ($1 \leq i \leq t$).

Proof. (1) Let $P = \text{soc}(G) \cap N$. Then $P \triangleleft G$, since $\text{soc}(N)$ and P are normal in G , and so $P \triangleleft \text{soc}(G)$. By Lemma 4.4, P is a direct product of (zero or more) simple groups since $P \triangleleft \text{soc}(G)$. Hence $P = \langle S : S \text{ simple, } S \triangleleft P \rangle$. Note for simple $S \triangleleft P$, we have $P = S \times Q$ and $\text{soc}(G) = P \times B$ by Lemma 4.4, so $\text{soc}(G) = S \times Q \times B$ and $S \triangleleft \text{soc}(G)$. Let $M_S = \langle g^{-1}Sg : g \in G \rangle$ which is a minimal normal subgroup of G . Every conjugate $g^{-1}Sg$ of S is simple and normal in $g^{-1}Pg = P$. So $M_S \leq P$. Thus

$$P = \langle S : S \text{ simple, } S \triangleleft P \rangle \leq \langle M_S : S \text{ simple, } S \triangleleft P \rangle \leq P.$$

So by Fact 5.3(1) applied to the set of these minimal normal subgroups M_S of G , we have that $P = \langle M_S : S \text{ simple, } S \triangleleft P \rangle = M_{S_1} \dots M_{S_n}$ is the internal direct product for some simple $S_1, \dots, S_n \triangleleft P$. Then, since, $1 \neq S \leq M_S \cap N$, the Fragmentation Lemma (Lemma 5.61) shows that each M_S is the internal direct product of some minimal normal subgroups of N . We have that P is the internal direct product of some minimal normal

subgroups of N . (2) Since $\text{soc}(N)$ is the generated by all the minimal normal subgroups of N , this follows from (1). Also $\text{soc}(G) \cap N = P$ is normal in G , hence in N . (3) follows from Lemma 5.3(1) applied the set consisting of M_1, \dots, M_n and all the minimal normal subgroups of G that intersect N trivially, since these together generate $\text{soc}(G)$. Similarly, (4) follows from Lemma 5.3(1) applied to the set consisting of N_1, \dots, N_m and all N' minimal normal in N that intersect $\text{soc}(G)$ trivially, since these together generate $\text{soc}(N)$. \square

5.3. A Characteristic Series and Complexity Upper Bound.

Definition 5.8 (Socle Characteristic Series and \mathbf{sx}). Let G be a finite group. Let $V_0 = \mathbf{1}$, define $V_1 = \text{soc}(G)$, and inductively for $i \geq 1$, define the natural surjective homomorphism $h_i : G \twoheadrightarrow G/V_i$, $\text{soc}_{i+1}(G) = \text{soc}(G/V_i)$, $V_{i+1} = h_i^{-1}(\text{soc}_i(G))$. Since $\text{soc}(G/V_i)$ is normal in G/V_i , its inverse image V_{i+1} is normal in G . Observe that if $V_i \neq G$, then V_{i+1} properly contains V_i . Therefore, $V_\ell = G$ for some $\ell \geq 0$. Let $\mathbf{sx}(G)$, the *socle length* (or *socle 'complexity'*) of G , be the least integer $\ell \geq 0$ such that $V_\ell = G$.

Fact 5.9. Let $G \neq \mathbf{1}$ be a finite group.

- (1) The socle $\text{soc}(G) = \langle N_1 \dots N_v \rangle$ is generated by the distinct minimal normal subgroups N_i of G , is the internal direct product of a subset $\{N_1, \dots, N_v\}$, and is normal in G .
(Note that by minimality $N_i \cap N_j = \mathbf{1}$ (the trivial subgroup) of G for $i \neq j$.)
- (2) $\text{soc}(G)$ is isomorphic to $S_1^{k_1} \times \dots \times S_m^{k_m}$ where $m \geq 1$, S_i is simple, $1 \leq k_i$ for $1 \leq i \leq m$.
- (3) $\text{soc}(G)$ is a characteristic normal subgroup of G .
- (4) $\mathbf{1} = V_0 \triangleleft V_1 \triangleleft V_2 \triangleleft \dots \triangleleft V_\ell = G$ is a characteristic series with each V_i is characteristic in G .
- (5) $\text{soc}(\text{soc}(G)) = \text{soc}(G)$
- (6) $\text{soc}(G/V_i) = V_{i+1}/V_i = \text{soc}(V_{i+1}/V_i)$.
- (7) $\mathbf{sx}(G/V_i) = \ell - i$ for $0 \leq i \leq \ell$.
- (8) If G is a simple or a direct product of simple groups, $\mathbf{sx}(G) = 1$.

Proof. (1) holds by definition of socle and Lemma 5.3(1). (2) follows from (1) and Lemma 4.3. (3): An automorphism α of G send a minimal normal subgroup M of G to a minimal normal subgroup $\alpha(M)$ of G . Hence, α permutes the minimal normal subgroups of G . Thus, the group the minimal normal subgroups generate, namely $\text{soc}(G)$, is mapped onto itself under automorphisms. (4): We show by induction on i that V_i is characteristic in G , i.e., $\alpha(V_i) = V_i$. (In particular V_i is normal since conjugation is an automorphism.) For $i = 0$, $\alpha(V_0) = V_0$ is trivial. Suppose the assertion is true for i . Let $w \in V_{i+1}$, we have $h_i(w) \in \text{soc}(G/V_i)$, i.e. $wV_i \in \text{soc}(G/V_i)$. By (1) we may write

$$wV_i = w_1V_i \dots w_kV_i,$$

for some $w_i \in G$, where the each w_jV_i lies in a minimal normal subgroup of G/V_i . Note: w and w_j lie V_{i+1} , since $wV_i, w_jV_i \in \text{soc}(G/V_i)$. Now $\alpha(w_jV_i)$ lies in a minimal normal subgroup of $\alpha(G)/\alpha(V_i) = G/V_i$ by the induction hypothesis that $\alpha(V_i) = V_i$. Therefore $\alpha(wV_i) = \alpha(w)V_i$ is a

product of elements of minimal normal subgroups of G/V_i . This shows $\alpha(w)V_i \in \text{soc}(G/V_i)$, so $\alpha(w) \in V_{i+1}$. Since w was an arbitrary element of V_{i+1} , $\alpha(V_{i+1}) \leq V_{i+1}$. Now α is injective and V_{i+1} finite, so $\alpha(V_{i+1}) = V_{i+1}$, i.e., as α was arbitrary, V_{i+1} is characteristic in G . So the result follows by induction.

(5): The $\text{soc}(G)$ is the direct product of simple groups $S_1 \times \cdots \times S_k$. Hence each factor is a minimal normal subgroup of $\text{soc}(G)$, hence contained in $\text{soc}(\text{soc}(G))$. This implies the group these S_i generate, namely, $\text{soc}(G)$ is a subgroup of $\text{soc}(\text{soc}(G))$; but the latter is a subgroup of $\text{soc}(G)$.

(6): By definition V_{i+1} is the inverse image in G of $\text{soc}(G/V_i)$ under the quotient map, and so $\text{soc}(G/V_i) = V_{i+1}/V_i$. So $\text{soc}(\text{soc}(G/V_i)) = \text{soc}(V_{i+1}/V_i)$, whence by (5), $\text{soc}(G/V_i) = \text{soc}(V_{i+1}/V_i)$ too.

(7) follows directly from the definition of socle length.

(8) follows since a direct product of simple groups is its own socle by (5). \square

A finite simple group has complexity at most 1. Socle length \mathbf{sx} assigns it complexity 1.

Theorem 5.10. *Socle length \mathbf{sx} dominates any hierarchical complexity function on finite groups.*

Proof. Let \mathbf{c} be any complexity function on finite groups. We prove for all finite groups $\mathbf{c}(G) \leq \mathbf{sx}(G)$ by strong induction on $|G|$. If $G = 1$, $\mathbf{sx}(G) = \mathbf{c}(G) = 0$. We have $\text{soc}(G)$ embeds in G with quotient $G/\text{soc}(G)$, therefore by the extension axiom $\mathbf{c}(G) \leq \mathbf{c}(\text{soc}(G)) + \mathbf{c}(G/\text{soc}(G))$. Now $\text{soc}(G)$ is a direct product of simple groups S_1, \dots, S_k , so by the product axiom $\mathbf{c}(\text{soc}(G)) = \max_{i=1}^k \mathbf{c}(S_i) \leq 1 = \mathbf{sx}(\text{soc}(G))$ (by Fact 5.9(8)) since simple groups S_i have $\mathbf{c}(S_i) \leq 1$ (by Lemma 1.1). By induction hypothesis, $\mathbf{c}(G/\text{soc}(G)) \leq \mathbf{sx}(G/\text{soc}(G))$. Now the socle length of G is one more than the socle length of $G/\text{soc}(G)$ by definition of socle length (Fact 5.9(7)), hence

$$\mathbf{c}(G) \leq \mathbf{sx}(\text{soc}(G)) + \mathbf{sx}(G/\text{soc}(G)) = 1 + \mathbf{sx}(G/\text{soc}(G)) = \mathbf{sx}(G).$$

\square

In particular

Corollary 5.11. $\mathbf{cx}(G) \leq \mathbf{sx}(G)$ for all finite groups G .

Theorem 5.12 (Complexity Upper Bounds). *For every finite group G ,*

$$\mu(G) = \mathbf{cx}(G) \leq \mathbf{sx}(G) \leq \mathbf{chief}(G) \leq \mathbf{JH}(G),$$

where \mathbf{chief} is the length of a chief series for G (which can include one type of simple group in the quotients).

Proof. We have seen $\mu(G) = \mathbf{cx}(G)$ in Theorem 4.6 and that $\mathbf{cx}(G) \leq \mathbf{sx}(G)$ in Corollary 5.11. One can obtain a chief series for G by refining the socle series of G : the socle series has a quotients V_{i+1}/V_i , each a product minimal normal subgroups of V_{i+1} , by adding appropriate subgroups one at a time between V_i and V_{i+1} one obtains a refined normal series with quotients isomorphic to these minimal normal subgroup. This is a chief series, hence $\mathbf{sx}(G) \leq \mathbf{chief}(G)$. Clearly $\mathbf{chief}(G) \leq \mathbf{JH}(G)$ since a single quotient in a chief series has one or more Jordan-Hölder factors. \square

But does $\mathbf{sx} = \mathbf{cx}$? If \mathbf{sx} satisfied the complexity axioms, it would follow by maximality of \mathbf{cx} that the two are equal. We'll now see this is not the case.

5.4. Socle Length and the Complexity Axioms.

Lemma 5.13. *For a direct product of finite groups $H \times K$, we have*

$$\text{soc}(H \times K) = \text{soc}(H) \times \text{soc}(K).$$

Proof. If M is minimal normal in H then $M \times 1$ is minimal normal in $H \times K$; similarly, if M' minimal normal in K , then $1 \times M'$ is minimal normal in $H \times K$. It follows that $\text{soc}(H) \times \text{soc}(K)$ is contained in $\text{soc}(H \times K)$. Conversely, if M is minimal normal in $H \times K$, then M projects to a minimal normal subgroup or 1 in each factor (Lemma 5.1), hence $M \leq \pi_H(M) \times \pi_K(M)$, where π_H and π_K are the projections. Since $M \neq 1$, either one or both of these projections is minimal normal in H resp. K . Thus $M \leq \text{soc}(H) \times \text{soc}(K)$, whence $\text{soc}(H \times K) \leq \text{soc}(H) \times \text{soc}(K)$. This proves the lemma. \square

Proposition 5.14. *Socle length \mathbf{sx} satisfies the constructability, product, and initial condition axioms on finite groups.*

Proof. Socle length satisfies these axioms:

Initial condition. We have $\mathbf{sx}(1) = 0$ by the definition of socle length.

Constructability axiom. Every finite group can be constructed from simple groups by iterated extension from simple groups S , namely its Jordan-Hölder factors, which have $\mathbf{sx}(S) = 1$ by Lemma 8.

Product axiom. Let $G = H \times K$. We now use strong induction on $|G|$ to show $\mathbf{sx}(G) = \max(\mathbf{sx}(H), \mathbf{sx}(K))$. If $G = 1$, then $\mathbf{sx}(G) = 0$ and the assertion holds. Otherwise, by Lemma 5.13,

$$\text{soc}(G) = \text{soc}(H \times K) = \text{soc}(H) \times \text{soc}(K),$$

so

$$G/\text{soc}(G) \cong (H \times K)/(\text{soc}(H) \times \text{soc}(K)) \cong H/\text{soc}(H) \times K/\text{soc}(K),$$

where the last isomorphism holds since the socles are normal in the respective factors (e.g. [18, Theorem 2.30]). By induction hypothesis,

$$\mathbf{sx}(G/\text{soc}(G)) = \max(\mathbf{sx}(H/\text{soc}(H)), \mathbf{sx}(K/\text{soc}(K))).$$

Therefore, by Fact 5.9(7) applied to G , H and K ,

$$\begin{aligned} \mathbf{sx}(G) &= 1 + \mathbf{sx}(G/\text{soc}(G)) = 1 + \max(\mathbf{sx}(H/\text{soc}(H)), \mathbf{sx}(K/\text{soc}(K))) \\ &= \max(1 + \mathbf{sx}(H/\text{soc}(H)), 1 + \mathbf{sx}(K/\text{soc}(K))) = \max(\mathbf{sx}(H), \mathbf{sx}(K)), \end{aligned}$$

completing the induction. \square

Counterexample 9 (Extension Axiom Fails for Socle Length). Consider

$$G = \langle (1, 2), (3, 4), (5, 6), (7, 8), (1, 3, 5, 7)(2, 4, 6, 8) \rangle.$$

It is not hard to see that $|G| = 64$ and $G \cong \mathbb{Z}_2 \wr \mathbb{Z}_4$, the wreath product of cyclic groups of order 4 and 2, projecting onto the \mathbb{Z}_4 factor.⁶

⁶To see the isomorphism, using depth-preserving maps of a two-level tree branches 4 ways from the root to depth 1, and 2 ways to 8 nodes at depth 2. Then the first 4 generators each fix the nodes at depth 1 and permute the two nodes under a node at

The center $Z(G)$ of G is a two-element group

$$Z(G) = \langle (1, 2)(3, 4)(5, 6)(7, 8) \rangle$$

and intersects any normal subgroup $N \neq 1$ of G nontrivially since G is a p -group (as is well-known $Z(G) \cap N \neq 1$ for G a p -group since N is a union of conjugacy classes, but of order a power of p , hence there must be conjugacy classes other than 1 of size 1, i.e., other central elements in N , since the sum of the sizes of conjugacy classes is $|N| \equiv 0$ modulo p). It follows that this two-element $Z(G)$ is the unique minimal normal subgroup of G . Thus $\text{soc}(G) = Z(G)$. Direct computation shows $\text{sx}(G) = 4$, with socle series $1 \triangleleft V_1 \triangleleft V_2 \triangleleft V_3 \triangleleft V_4 = G$, where $V_1 = Z(G)$, $V_2 = \langle (3, 4)(7, 8), (1, 2)(3, 4)(5, 6)(7, 8) \rangle$, $V_3 = \langle (1, 5)(2, 6)(3, 7)(4, 8), (5, 6)(7, 8), (3, 4)(7, 8) \rangle$, with successive quotients isomorphic to $\mathbb{Z}_2, \mathbb{Z}_2, \mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_2 \times \mathbb{Z}_2$.

On the other hand consider $N = \langle (3, 4)(7, 8), (1, 2)(5, 6) \rangle$. One easily checks N is normal in G and is a Klein 4-group. Since it is a direct product of simple groups, $\text{sx}(N) = 1$. Direct computations show $Q = G/N$ is isomorphic to a semidirect product $(\mathbb{Z}_4 \times \mathbb{Z}_2) \rtimes \mathbb{Z}_2 \cong \langle (2, 6)(5, 8), (1, 2, 3, 5)(4, 6, 7, 8) \rangle$ with socle $Q_1 = \langle (1, 4)(2, 6)(3, 7)(5, 8), (1, 3)(2, 5)(4, 7)(6, 8) \rangle \cong \mathbb{Z}_2 \times \mathbb{Z}_2$, and $Q/Q_1 \cong \mathbb{Z}_2 \times \mathbb{Z}_2$, so $\text{sx}(G/N) = 2$. Now

$$\text{sx}(G) = 4 \not\leq 1 + 2 = \text{sx}(N) + \text{sx}(G/N),$$

so this is a counterexample to the extension axiom for sx .

Counterexample 10. The smallest counterexample to extension for sx is the 32-element group $H = \langle (1, 2)(3, 5)(4, 6)(7, 8), (2, 5, 6, 8)(3, 7), (2, 6)(5, 8) \rangle$, whose socle length is 3 with socle series is $1 \triangleleft V_1 \triangleleft V_2 \triangleleft V_3 = H$. The center of H is a unique minimal normal subgroup

$$V_1 = \text{soc}(H) = Z(H) = \langle (1, 2)(3, 5)(4, 6)(7, 8) \rangle \cong \mathbb{Z}_2,$$

$V_2 = \langle (2, 6)(5, 8), (1, 7)(2, 8)(3, 4)(5, 6), (1, 4)(2, 6)(3, 7)(5, 8) \rangle \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$, whose quotient by $\text{soc}(H)$ is $\mathbb{Z}_2 \times \mathbb{Z}_2$, with $H/V_2 \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. So $\text{sx}(H) = 3$. However, since $V_2 \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ and $H/V_2 \cong \mathbb{Z}_2 \times \mathbb{Z}_2$, shows $\text{cx}(H) \leq 2$, but $\text{cx}(H) \neq 1$ since H is not the direct product of \mathbb{Z}_2 's as it has an element of order 4, whence $\text{cx}(H) = 2$. Now $\text{cx}(H) = 2 < 3 = \text{sx}(H)$. Note also that the $\text{sx}(H)$ is *not* bounded above by $\text{sx}(V_2) + \text{sx}(H/V_2) = 1 + 1 = 2$.

Since $\mu = \text{cx}$ satisfies the extension axiom by Prop. 4.5, we have this

Corollary 5.15. *sx does not satisfy the extension axiom. Hence, $\text{cx} \neq \text{sx}$.*

Proposition 5.16. *sx satisfies the quotient and normal subgroup axioms.*

Lemma 5.17. *Let G be a finite group and N a normal subgroup of G . Then, the socle characteristic series of N and of G/N are each no longer than that of G . That is, (1) $\text{sx}(N) \leq \text{sx}(G)$ and (2) $\text{sx}(G/N) \leq \text{sx}(G)$.*

Proof. (1) Normal Subgroup Property. Let G be finite group with socle characteristic series as defined in Section 5.3

$$1 = V_0 \triangleleft V_1 \triangleleft \cdots \triangleleft V_\ell \triangleleft V_{\ell+1} = G.$$

depth one. The final generator cyclically permutes the depth 1 nodes but maps the nodes at depth two trivially.

For N a normal subgroup of G , let

$$1 = W_0 \triangleleft W_1 \triangleleft \cdots \triangleleft W_m = N$$

be the socle characteristic series for N . We claim

Claim 1. *For all natural numbers i , $V_i \cap N \leq W_i$.*

(By definition of the socle characteristic series, $W_i = N$ for $i \geq m$ and $V_i = G$ for $i \geq \ell$.)

Since $V_\ell = G$, from the claim it follows that $N = V_\ell \cap N \leq W_\ell$. Thus $N = W_\ell$, but by definition of the socle characteristic series for N , m is the least natural number such that $W_m = N$, so $m \leq \ell$, i.e., $\mathbf{sx}(N) \leq \mathbf{sx}(G)$.

So to prove (1), it suffices to prove Claim 1.

Proof of Claim 1 by Induction: The assertion is trivial for $i = 0$.

Suppose it is true up to i

$$V_i \cap N \subseteq W_i.$$

Then $V_i \cap N \triangleleft W_i$ since $V_i \triangleleft$ and $N \triangleleft G$. We must show $V_{i+1} \cap N \triangleleft W_{i+1}$. Consider the natural map $\varphi : G \twoheadrightarrow G/V_i = \varphi(G)$. Then $\varphi(N) \triangleleft G/V_i$, since φ is surjective. We shall denote $\varphi(N) = NV_i/V_i$ by N/V_i . By Splitting Lemma II (Lemma 5.7(1)),

$$\text{soc}(G/V) \cap N/V_i = \tilde{N}_1 \cdots \tilde{N}_k$$

for direct factors \tilde{N}_i minimal normal in $\varphi(N)$. By induction hypothesis, $V_i \cap N \triangleleft W_i$, so we have a natural surjective morphism:

$$\psi : \frac{N}{V_i \cap N} \twoheadrightarrow \frac{N}{W_i}.$$

Each \tilde{N}_i maps to $\mathbf{1}$ or a minimal normal subgroup of N/W_i under ψ by Lemma 5.1. Hence \tilde{N}_i maps to $\text{soc}(N/W_i) = W_{i+1}/W_i$. Therefore all of $\text{soc}(G) \cap N/V_i$ maps to W_{i+1}/W_i . Now $\text{soc}(N/V_i)$ is generated by minimal normal subgroups of N/V_i , and by Splitting Lemma II (Lemma 5.7(1)), so is $\text{soc}(G/V_i) \cap N/V_i$. Now $\text{soc}(G/V_i) = V_{i+1}/V_i$. Suppose $x \in V_{i+1} \cap N$. Then $xV_i \in NV_i$, and $xV_i = n_1V_i \cdots n_kV_i$ where each $n_jV_i \in \tilde{N}_j$ ($1 \leq j \leq k$). Applying ψ we have

$$xV_i = n_1V_i \cdots n_kV_i \mapsto n_1W_i \cdots n_kW_i \in \text{soc}(N/W_i),$$

whence

$$xW_i = n_1 \cdots n_kW_i \in W_{i+1}/W_i.$$

with $n_1 \cdots n_k \in W_{i+1}$. Therefore, $(n_1 \cdots n_k)^{-1}x \in W_i \triangleleft W_{i+1}$. It follows that $x \in W_{i+1}$. Since x was an arbitrary element of $V_{i+1} \cap N$, this completes the induction step. Therefore the assertion holds for all $i \geq 0$. This proves Claim 1, hence (1) is proved.

(2) Quotient Axiom. Similarly, let

$$\mathbf{1} = W_0 \triangleleft W_1 \triangleleft \cdots \triangleleft W_n = H$$

be the socle characteristic series for $H = G/N$. Using the notation V_i/N to denote the homomorphic image of V_i under the quotient map $G \twoheadrightarrow G/N$, we claim

Claim 2. *with each $M'_i \cap \text{soc}(N) = \mathbf{1}$, $s \geq 0$ Let $\varphi : G \twoheadrightarrow H$ be a surjective homomorphism. For all $i \geq 0$, $\varphi(V_i) \leq W_i$.*

(By definition, $W_i = G/N$ for $i \geq n$ and $V_i = G$ for $i \geq \ell$.)

Similarly to be before, since $V_\ell = G$, from the claim it follows that $\varphi(V_\ell) \leq W_\ell$. Thus $G/N = W_\ell$, but by definition of the socle characteristic series for G/N , n is the natural number such that $W_n = G/N$, so $n \leq \ell$, i.e., $\text{sx}(H) \leq \text{sx}(G)$.

So to prove (2), it suffices to prove Claim 2.

Proof of Claim 2. We proceed by induction. The assertion is trivial for $i = 0$, since $\varphi(V_0) = \mathbf{1} \leq \mathbf{1} = W_0$. Suppose $i \geq 0$ and $\varphi(V_i) \leq W_i$. Then φ is surjective so $\varphi(V_i) \triangleleft H$ and so $\varphi(V_i) \triangleleft W_i$. We must show $\varphi(V_{i+1}) \leq W_{i+1}$.

Consider a minimal normal subgroup M of G/V_i . By Lemma 5.1, M is either trivial or a minimal normal subgroup of G/V_i . Since $\varphi(V_i) \leq W_i$ by induction hypothesis, there is well-defined surjective morphism from H/V_i to H/W_i ; and by Lemma 5.1, $\varphi(M)$ is either trivial or a minimal normal subgroup in H/W_i . Hence M maps to a $\text{soc}(W_{i+1}/W_i) = \text{soc}(H/W_i)$. Since M is an arbitrary minimal normal subgroup of G/V_i , and $\text{soc}(H/W_i)$ is generated by minimal normal subgroups of H/W_i , it follows (using Fact 5.9(6)) that $\text{soc}(G/V_i) = \text{soc}(V_{i+1}/V_i) = V_{i+1}/V_i$ maps homomorphically to $\text{soc}(H/W_i) = \text{soc}(W_{i+1}/W_i) = W_{i+1}/W_i$.

Now let v be arbitrary element of V_{i+1} . We have that

$$vV_i \mapsto \varphi(v)W_i$$

under the composite morphism from G/V_i to H/W_i . Therefore $\varphi(v)W_{i+1}$ lies in the socle of H/W_i . By definition of W_{i+1} , $\varphi(v) \in W_{i+1}$. Therefore $\varphi(V_{i+1}) \leq W_{i+1}$, and the result of Claim 2 follows by induction. This proves (2). \square

6. OTHER COMPLEXITY FUNCTIONS: LOWER AND UPPER BOUNDS

6.1. Uncountably Many Complexity Functions. Here we show for any collection of finite simple groups we obtain a distinct maximal complexity function assigning 1 to these simple groups and 0 to all others.

Definition 6.1. Let \mathcal{S} be a collection of finite simple groups. Define the *characteristic complexity function with respect to \mathcal{S}* by

$$\chi_{\mathcal{S}}(G) = \begin{cases} 1 & \text{if some Jordan-Hölder factor of } G \text{ lies in } \mathcal{S} \\ 0 & \text{otherwise.} \end{cases}$$

It is immediate to check

Lemma 6.2. *The function $\chi_{\mathcal{S}}$ satisfies the complexity axioms.*

Examples 1. We have the following complexity functions:

- (1) $z = \chi_{\emptyset}$
- (2) Let $\mathcal{SJMPL}\mathcal{E}$ be the finite simple groups. $\delta = \chi_{\mathcal{SJMPL}\mathcal{E}}$.
- (3) Let $\mathcal{SNA}\mathcal{G}$ be the collection of all finite simple non-abelian groups.
Then $\chi_{\mathcal{SNA}\mathcal{G}}(G) = 0$ if and only if G is solvable.
- (4) Let \mathcal{Z} be all simple groups \mathbb{Z}_p of prime order. Then $\delta = \max(\chi_{\mathcal{SNA}\mathcal{G}}, \chi_{\mathcal{Z}})$.

Definition 6.3. If \mathcal{S} is collection of finite simple groups. $\chi_{\mathcal{S}}^*$ is the maximal function pointwise dominating all complexity functions which assign zero to all finite simple groups not in \mathcal{S} . That is,

$$\chi_{\mathcal{S}}^*(G) = \max\{\mathbf{c}(G) : \mathbf{c} \text{ is a complexity function with } \mathbf{c}(K) \leq \chi_{\mathcal{S}}(K) \text{ for all } K \text{ simple}\}.$$

As for $\mathbf{c}\mathbf{x}$, which is $\chi_{\mathcal{S}\text{JMPLE}}^*$, one shows that $\chi_{\mathcal{S}}^*$ exists and is the unique complexity function assigning 1 to members of \mathcal{S} and 0 to other finite simple groups.

For a subnormal series for a group G in which each quotient G_{i+1}/G_i is the direct product of simple groups, we say *the i th level of the series has members in \mathcal{S}* if the i th quotient group has at least one factor in \mathcal{S} . Let $\mu_{\mathcal{S}}(G)$ be the least number of levels over all such series for which the series has members of \mathcal{S} .

Theorem 6.4. *For any set \mathcal{S} of simple groups, $\mu_{\mathcal{S}} = \chi_{\mathcal{S}}^*$.*

Proof. Just as in the proofs the $\mu = \mathbf{c}\mathbf{x}$ by Prop. 4.5 and Theorem 2.3, one proves $\mu_{\mathcal{S}}$ satisfies the axioms and dominates all complexity functions \mathbf{c} with $\mathbf{c}(K) \leq \chi_{\mathcal{S}}(K)$ for all K simple. The only subtle point is the product axiom, which is verified in the Lemma 6.5. \square

Lemma 6.5 (Alignment Lemma). *The product axiom holds for $\mu_{\mathcal{S}}$.*

Proof. Let H have a subnormal series with quotients direct products of simple groups

$$1 = H_0 \triangleleft H_1 \triangleleft \cdots \triangleleft H_a = H,$$

with $h \leq a$ levels having factors in \mathcal{S} . Similarly, let K have subnormal series of this type with $k \leq b$ levels having factors in \mathcal{S}

$$1 = K_0 \triangleleft K_1 \triangleleft \cdots \triangleleft K_b = K.$$

Suppose these series have least possible h and k , respectively. That is, $\mu_{\mathcal{S}}(H) = h$ and $\mu_{\mathcal{S}}(K) = k$. If $H \times K$ had $\mu_{\mathcal{S}}(H \times K) < \max\{h, k\}$, then it would have a series projecting onto series for H and K with less than h and k levels with factors in \mathcal{S} , respectively, a contradiction (just as in the proof of Prop. 4.5), so $\mu_{\mathcal{S}}(H \times K) \geq \max\{\mu_{\mathcal{S}}(H), \mu_{\mathcal{S}}(K)\}$.

Next we claim $H \times K$ has subnormal series with quotients direct products of simple groups having exactly $\max\{h, k\}$ levels with factors in \mathcal{S} .

Let $i_1 < \cdots < i_h$ be the indices in $[0, a-1]$ such that $H_{i_{\ell}+1}/H_{i_{\ell}}$ has factors in \mathcal{S} . Similarly, let $j_1 < \cdots < j_k$ be the indices in $[0, b-1]$ such that $H_{j_{\ell}+1}/H_{j_{\ell}}$ has factors in \mathcal{S} . We have $H_{i_{\ell}} \times K_{j_{\ell}} \triangleleft H_{i_{\ell}+1} \times K_{j_{\ell}+1}$, where if $h < k$ we take $H_{i_{\ell}} = H_{i_h}$ for all $\ell > h$, and similarly if $h > k$ we take $K_{j_{\ell}} = K_{j_k}$ for all $\ell > k$. Then the groups

$$(H_{i_{\ell}+1} \times K_{j_{\ell}+1})/(H_{i_{\ell}} \times K_{j_{\ell}}) \cong (H_{i_{\ell}+1}/H_{i_{\ell}}) \times (K_{j_{\ell}+1}/K_{j_{\ell}})$$

are each a direct product of simple groups with factors coming factors in \mathcal{S} . It is easy to see that the normal inclusions

$$H_{i_{\ell}} \times K_{j_{\ell}} \triangleleft H_{i_{\ell}+1} \times K_{j_{\ell}+1}$$

for $1 \leq \ell \leq \max\{h, k\}$ can be extended to a subnormal series for $H \times K$ in which these are the only quotients with factors in \mathcal{S} : One simply inserts products of factors from the above series for H and K , e.g., include the

subgroups $H_0 \times K_j$ for $j \leq j_1$, followed by $H_i \times K_{j_1}$ for $i < i_1$, and so on inserting product groups between $H_{i_{\ell+1}} \times K_{j_{\ell+1}}$ and $H_{i_{\ell+1}} \times K_{j_{\ell+1}}$ by first increasing the factors from the H_i and then the ones from the K_j . Finally insert $H_{i_{h+1}} \times K_j$ for all j with $b \geq j > j_k + 1$, if any, followed by $H_i \times K$ for all i with $a \geq i > i_h + 1$, if any. This yields a subnormal series from 1 to $H \times K$ having $\max\{h, k\}$ levels with factors in \mathcal{S} . This shows $\mu_{\mathcal{S}}(H \times K) \leq \max\{\mu_{\mathcal{S}}(H), \mu_{\mathcal{S}}(K)\}$ and completes the proof. \square

Corollary 6.6. *There are uncountably many pairwise distinct complexity functions on finite groups. In particular, if $\mathcal{S}, \mathcal{S}' \subseteq \text{SJMPLE}$ are distinct then $\mu_{\mathcal{S}} \neq \mu_{\mathcal{S}'}$.*

Proof. $\mu_{\mathcal{S}}$ and $\mu_{\mathcal{S}'}$ disagree on each member of the symmetric difference $(\mathcal{S} \cup \mathcal{S}') \setminus (\mathcal{S} \cap \mathcal{S}')$ which is nonempty since for $\mathcal{S}, \mathcal{S}' \subseteq \text{SJMPLE}$ with $\mathcal{S} \neq \mathcal{S}'$. Hence $\mu_{\mathcal{S}} \neq \mu_{\mathcal{S}'}$. Since there are countably many isomorphism classes of finite simple groups comprising SJMPLE , this yields a distinct complexity function for each of the uncountably many subsets \mathcal{S} of SJMPLE . \square

Since $\mu_{\mathcal{S}}$ satisfies the complexity axioms and $\mathbf{cx} = \mu = \mu_{\text{SJMPLE}}$ is the maximal complexity function on finite groups, we conclude:

Theorem 6.7. *Let $\mathcal{S} \subseteq \text{SJMPLE}$, the $\mu_{\mathcal{S}}$ is a lower bound for hierarchical complexity \mathbf{cx} .*

6.2. Solvable Groups and the Embedding Axiom. Next we consider complexity functions on finite solvable groups.

Theorem 6.8. *There exists a unique maximal complexity function $\mathbf{cx}_{\text{solv}}$ on finite solvable groups satisfying the axioms.*

Proof. By the same reasoning as in Theorem 2.3, there exists a unique maximal complexity function on solvable groups. \square

Lemma 6.9. *For a finite solvable group G and prime p , let $\log_p(G)$ be the greatest natural number n such that G has an element of order p^n . Then \log_p is a complexity function on solvable groups.*

Caveat: \log_p is not a complexity function on all finite groups since it can assign value greater than one to simple non-abelian groups. Indeed, by Lemma 3.2, we can embed \mathbb{Z}_{p^n} into a simple group K_n results in $\log_p(K_n) \geq n$ for any positive n .

Proof. Product. If $G = H \times K$ and $g = (h, k)$, then the order of g is the least common multiple the orders of h and k . Therefore, if $g \in G$ has order p^n , then h and k both have order dividing p^n , so they have orders p^a and p^b , with $\max(a, b) = n$. This entails that $\log_p(G) = \max\{\log_p(H), \log_p(K)\}$.

Extension. Suppose $a = \log_p(Q)$, $b = \log_p(N)$, and G is an extension of Q by N . Then G embeds in a wreath product $N \wr Q$. Let g in G be an element whose order is a power of p , then in the embedding, $g = (\mathbf{n}, q)$, $q \in Q$, $\mathbf{n} \in N \times \cdots \times N$ (the $|Q|$ -fold direct power of N). We have $\log_p(N) = \log_p(N^Q)$ (since the product axiom holds). Now

$$g^{p^a} = (\mathbf{n}', q^{p^a}) = (\mathbf{n}', 1)$$

for some $\mathbf{n}' \in N^Q$, whence

$$g^{p^{(a+b)}} = (g^{p^a})^{p^b} = (\mathbf{n}', 1)^{p^b} = (\mathbf{n}'^{p^b}, 1) = (1, 1) = 1 \in G.$$

This shows the order of g is at most p^{a+b} . This shows

$$\log_p(G) \leq \log_p(N) + \log_p(Q),$$

i.e., the extension axiom holds.

Quotient. If g has order p^n in G , its image in any quotient Q has order dividing p^n , whence $\log_p(G) \geq \log_p(Q)$.

Normal Subgroup Axiom / Subgroup Axiom. If N is a subgroup of G , whether normal or not, and $n \in N$ has order p^n , so G also contains this element, hence $\log_p(G) \geq \log_p(N)$.

Constructability. Every finite solvable group can be constructed by iterated extension from simple cyclic groups \mathbb{Z}_q of prime order q . Since $\log_p(\mathbb{Z}_q) = 1$ for $p = q$ but is 0 otherwise, this shows the constructability axiom holds.

Initial Condition. Finally, the element of largest prime power in the trivial group has order $1 = p^0$, so \log_p of the trivial group is 0. \square

Remark 6.10. Note that $\log_p(S)$ can be greater than 1 if S is a simple non-abelian group. Therefore \log_p is *not* a complexity function on all finite groups as it fails to satisfy the constructability axiom.

Corollary 6.11 (Some Complex Bounds). *The following bounds hold related the complexity $\mathbf{cx}_{\text{solv}}$ of solvable groups.*

- (1) \log_p is unbounded since $\log_p(\mathbb{Z}_{p^n}) = n$ for cycle groups of order p^n .
- (2) For a finite solvable group G ,

$$\mathbf{cx}_{\text{solv}}(G) \geq \sup_{p \text{ prime}} \log_p(G) = \max_{p \text{ prime divisor of } |G|} \log_p(G).$$

- (3) The complexity function $\mathbf{cx}_{\text{solv}}$ is unbounded.
- (4) $\mathbf{sx} \geq \mathbf{cx}_{\text{solv}}$.

Proof. (1) follows from the definition of \log_p . (2) follows since $\mathbf{cx}_{\text{solv}}$ is a maximal complexity function on solvable groups so dominates each complexity function \log_p . (3) follows from (2) and (1). (4) follows since \mathbf{sx} bounds any complexity function on solvable groups from above (same proof as for \mathbf{sx} bounding \mathbf{cx} on all finite groups). \square

Theorem 6.12. (1) If G is a solvable group, then $\mathbf{cx}(G) = \mathbf{cx}_{\text{solv}}(G)$.
 (2) $\mathbf{cx}_{\text{solv}}$ satisfies the embedding axiom: $\mathbf{cx}_{\text{solv}}(H) \leq \mathbf{cx}_{\text{solv}}(G)$ for H a subgroup of G .
 (3) \mathbf{sx} restricted to solvable groups satisfies the embedding axiom. If G is solvable and H a subgroup, then $\mathbf{sx}(H) \leq \mathbf{sx}(G)$.

Proof. (1) Recall $\mathbf{cx}(G) = \mu(G)$ the minimal length of a composition series for G with all quotients direct products of simple groups. The same proof that $\mu(G)$ is a maximal complexity function on finite groups works to prove $\mu(G)$ restricted to solvable groups is a maximal complexity function on solvable groups. Since μ restricted to solvable groups satisfies the complexity axioms (recall the embedding axiom was not required) and $\mathbf{cx}_{\text{solv}}$ is a maximal complexity function on solvable groups,

$$\mathbf{cx}_{\text{solv}}(G) = \mu(G).$$

(2) Consider a subnormal series of length $\mu(G)$ for G ,

$$1 = V_0 \triangleleft V_1 \triangleleft \dots \triangleleft V_n = G,$$

witnessing $\mu(G) = n$ with V_{i+1}/V_i a direct product of simple abelian groups. Intersection with subgroup H yields a subnormal series for H :

$$1 = V_0 \cap H \triangleleft V_1 \cap H \triangleleft \dots \triangleleft V_m \cap H = G \cap H = H.$$

By the 2nd of Noether's isomorphism theorems, this has quotients

$$(V_{i+1} \cap H)/(V_i \cap H) \cong (V_{i+1} \cap H)V_i/V_i \leq V_{i+1}/V_i,$$

where the inclusion holds since $(V_{i+1} \cap H)V_i \leq V_{i+1}$ for each i . Since V_{i+1}/V_i is a direct product of abelian simple groups, so is its subgroup the quotient from the series for H . This shows $\mu(H) \leq \mu(G)$. The proof for $\mathbf{sx}(H) \leq \mathbf{sx}(G)$ is the same, but starts with the socle characteristic series of G . \square

By Theorem 6.12, when restricted to solvable groups, the hierarchical group complexity and socle length satisfy the

Subgroup Property/ Embedding Axiom:

If H is a subgroup of G , then $\mathbf{c}(H) \leq \mathbf{c}(G)$.

The importance of the theorem is that when working with solvable groups, we can use the embedding axiom to compute their complexity and socle length.

6.3. Hierarchical Complexity of an Iterated Wreath Product.

Example 11. We compute, for p prime, and any n , the complexity of the n -fold wreath product of k_i -fold direct products of a simple abelian group with itself,

$$W = \wr_{i=1}^n \left(\prod_{i=1}^{k_i} \mathbb{Z}_p \right) = (\mathbb{Z}_p)^{k_n} \wr \dots \wr (\mathbb{Z}_p)^{k_1},$$

where $k_i > 0$ ($1 \leq i \leq n$). Namely, $\mathbf{cx}(W) = n$. Also, $\mathbf{cx}(\mathbb{Z}_{p^n}) = n$.

Proof. From example 4, we know $\mathbf{cx}(\mathbb{Z}_{p^n}) \leq n$. Since \mathbb{Z}_{p^n} is solvable, \log_p gives a lower bound for \mathbf{cDer} ,

$$n = \log_p(\mathbb{Z}_{p^n}) \leq \mathbf{cx}_{\text{solv}}(\mathbb{Z}_{p^n}) = \mathbf{cx}(\mathbb{Z}_{p^n})$$

by Theorem 6.12. Therefore $\mathbf{cx}(\mathbb{Z}_{p^n}) = n$. The iterated wreath product W of direct products of simple groups \mathbb{Z}_p contains a cyclic group of order p^n , but not p^{n+1} since \log_p satisfies the extension and product axioms (applied iteratively to W). Since the embedding axiom holds for \mathbf{cx} on solvable groups,

$$n = \mathbf{cx}(\mathbb{Z}_p^n) \leq \mathbf{cx}(W).$$

On the other hand, W is an n -fold iterated extension by products of simple groups, so $\mathbf{cx}(W) \leq n$ by the extension axiom. Hence $\mathbf{cx}(W) = n$. \square

6.4. Hierarchical Complexity of p -torsion and other Groups.

Example 12. Let p be prime and G be a finite p -torsion group, i.e. $x^p = 1$ for all $x \in G$. Then $\mathbf{cx}(G) \leq \mathbf{sx}(G) = \text{length of the ascending central series of } G$.

Proof. $\text{soc}(G) = N_1 \cdots N_k$, N_i a minimal normal subgroup of G . Each $N_i \cap Z(G) \neq 1$, since G is a p -group.⁷ By minimality N_i , $N_i \cap Z(G) = N_i$. Thus every minimal normal subgroup is contained the center of G . Therefore $\text{soc}(G) \leq Z(G)$. Conversely if $1 \neq x \in Z(G)$, then $\langle x \rangle \cong \mathbb{Z}_p$ is minimal normal in G . Therefore $Z(G) = \text{soc}(G)$. Therefore $Z(G/V_i) = \text{soc}(G/V_i) = V_{i+1}/V_i$ in the socle series, so it coincides with the ascending central series. The conclusion then follows from Theorem 5.10. \square

Open Question 1 (Complexity of Finite Prime-Exponent Burnside Groups). What is the hierarchical complexity of $B = B(k, p)$ a largest finite k -generated group of exponent p prime?

From example 12 and Theorem 6.20 on lower bounds proved below, for any p -torsion group B ,

$$\mathbf{Der}(B) \leq \mathbf{cx}(B) \leq \mathbf{sx}(B) = \text{length of ascending central series of } B.$$

Since the derived series for B has abelian quotients Q_i and each of these quotients is a homomorphic image of B , it follows that Q_i is an abelian p -torsion group, hence necessarily of the form $\mathbb{Z}_p \times \cdots \times \mathbb{Z}_p$, a span of gems, whence $\mathbf{cx}(B) \leq \mathbf{Der}(B)$. Since $\mathbf{Der}(B) \leq \mathbf{cx}(B)$, it follows that the derived series of B gives a decomposition of B at its hierarchical complexity.⁸ Therefore $\mathbf{Der}(B) = \mathbf{cx}(B)$, for any p -torsion group B .

The same argument shows that if the factors of n contain no square of a prime, then any finite n -torsion solvable group has $\mathbf{Der}(B) = \mathbf{cx}(B)$: since the quotients of the derived series are abelian and n -torsion, they must therefore be direct products of simple cyclic groups by the fundamental structure theorem for finite abelian groups.⁹

Open Question 2. For which finite groups G , does $\mathbf{sx}(G) = \mathbf{cx}(G)$?

From Theorem 5.11, $\mathbf{cx}(x) \leq \mathbf{sx}(G)$, and this bound is sharp since, e.g., $\mathbf{sx}(\mathbb{Z}_{p^n}) = \mathbf{cx}(\mathbb{Z}_{p^n})$: For $\mathbb{Z}_{p^n} = \langle x \rangle$, we have $\text{soc}(\mathbb{Z}_{p^n}) = \langle x^{p^{n-1}} \rangle \cong \mathbb{Z}_p$, whence by induction $\mathbf{sx}(\mathbb{Z}_{p^n}) = n$. By example 11, $\mathbf{cx}(\mathbb{Z}_{p^n}) = n$ too.

6.5. Derived Complexity, Fitting Complexity and Solvability Complexity. Here we recover the concepts of derived length and Fitting height within the framework of group complexity axioms. We show that the usual

⁷This well-known fact is shown in Counterexample 9.

⁸Note it does *not* necessarily follow that there is not another subnormal series showing $\mathbf{cx}(B)$ has this value, since the quotient factors at complexity need not be unique (cf. Counterexample 6).

⁹Similarly, the assertion $\mathbf{Der}(G) = \mathbf{cx}(G)$ holds for any finite group G with square-free order, even without the assumption of solvability of G : By a classical result of Frobenius [8, Sec. 4], a finite group whose order is the product of distinct primes is necessarily solvable, so the derived series terminates at 1 and has abelian n -torsion quotients which again must be spans of gems. (See exposition of Frobenius's result in [9].)

derived length of solvable groups and the the usual Fitting height of solvable groups are restrictions of more general complexity functions on all finite groups, and give lower bounds on hierarchical complexity.

To obtain derived length we can add to the complexity axioms the axiom:

(Com) Commutative groups have complexity at most 1.

To recover Fitting height, we add the axiom:

(Nil) Nilpotent groups have complexity at most 1.

For solvability, we can have an analogous axiom:

(Solv) Solvable groups have complexity at most 1.

Since finite abelian groups are nilpotent, and nilpotent groups are solvable (e.g. [10, Ch. 9 & 10]), we have the following implications:

$$(\text{Solv}) \Rightarrow (\text{Nil}) \Rightarrow (\text{Com}).$$

Theorem 6.13. *There exist unique maximal complexity functions \mathbf{cx}_{Der} , \mathbf{cx}_{Fit} and $\mathbf{cx}_{\text{Solv}}$ on finite groups satisfying the complexity axioms together with the (Com), (Nil) and (Solv) axioms, respectively.*

Proof. The proof follows same reasoning as in Theorem 2.3 but considering only complexity functions also satisfying the added axiom. \square

Note that the normal subgroup property is *not* assumed, but will be found to hold nonetheless, as we shall see from alternative characterizations of these functions (just as for \mathbf{cx} and μ).

Starting from the notion of a span of gems (or mālā or necklace of simple groups), i.e., a finite group that is the direct product of simple groups, we have the following increasingly general concepts:

Definition 6.14. (1) A *derived necklace* is a finite group G that is the direct product of abelian groups and simple groups, i.e.,

$$G \in \text{Span}(\text{Abelian groups} \cup \text{SIMPLE}).$$

(2) A *Fitting necklace* is a finite group that is the direct product of nilpotent groups and simple groups, i.e.,

$$G \in \text{Span}(\text{Nilpotent groups} \cup \text{SIMPLE}).$$

(3) A *solvability necklace* is a finite group that is the direct product of solvable groups and simple groups,

$$G \in \text{Span}(\text{Solvable groups} \cup \text{SIMPLE}).$$

Note that in these definitions, zero or more simple factors are permitted, also the abelian, nilpotent or solvable groups, respectively, may be trivial.

Fact 6.15. (1) *Since the direct product of abelian groups is abelian, each derived necklace is the direct product of a single abelian group and (zero or more) simple non-abelian groups.*

- (2) *Similarly, since the direct product of nilpotent groups is nilpotent and simple abelian groups are nilpotent, each Fitting span of gems is the direct product of a single nilpotent group and simple non-abelian groups.*
- (3) *A solvability necklace is the direct product of a single solvable group and simple non-abelian groups, since solvable groups are closed under product and abelian simple groups are solvable.*

One has the following consequences of the SNAG rigidity lemma 4.1 and its corollary.

Lemma 6.16. *A normal subgroup of a derived necklace is a derived necklace. A quotient of a derived necklace is a derived necklace. A product of derived necklaces is a derived necklace. Similarly, Fitting necklaces are closed under normal subgroups, quotients and products. Solvability necklaces are closed under normal subgroups, quotients, and products.*

Proof. A derived necklace G is the product of an abelian group K and zero or more SNAGs H_i ($1 \leq i \leq \ell$, $\ell \geq 0$). Hence, by Lemma 4.1 and its corollary, $N \triangleleft G$, is a product of some of these SNAGs and $\pi(N)$ where $\pi : G \rightarrow K$ is the projection and $\pi(N)$ is normal in K . Since K is abelian so is $\pi(N)$. The assertion about product is follows from Fact 6.15 on products of necklaces. Similarly, replacing the word “abelian” by “nilpotent”, resp. “solvable” in this reasoning shows the analogous assertions about Fitting necklaces and solvability necklaces are also correct. \square

Just as for spans of gems, these types of necklaces are not closed under taking subgroups unless the necklaces have no SNAG factors, i.e., unless the necklaces are solvable groups.

Definition 6.17. Now we introduce three functions defined on each finite group G :

- (1) **Der**(G) is the length of a shortest subnormal series such that the quotients are products of an abelian group and zero or more finite simple non-abelian groups.
- (2) **Fit**(G) is the length of a shortest subnormal series such that the quotients are products of a nilpotent group and zero or more finite simple non-abelian groups.
- (3) **Solv**(G) is the length of a shortest subnormal series such that the quotients are products of a solvable group and zero or more finite simple non-abelian groups.

Theorem 6.18. *We have the following equalities of functions on finite groups:*

- (1) **Der** = **cx_{Der}**, *the unique maximal complexity function assigning 1 to all nontrivial abelian groups.*
- (2) **Fit** = **cx_{Fit}** *is the unique maximal complexity function assigning 1 to all nontrivial nilpotent groups.*
- (3) **Solv** = **cx_{Solv}** *is the unique maximal complexity function assigning 1 to all nontrivial solvable groups.*

Proof. The proof that each of **Der**, **Fit** and **Solv** is a complexity function satisfying (Com), (Nil) or (Solv), respectively, follows exactly as for

μ in Proposition 4.5, but uses Lemma 6.16 in place of the Span of Gems Lemma 4.3 and Fact 6.15 in place of the fact that spans of gems are closed under products. That each one is the unique maximal complexity function \mathbf{cDer} , \mathbf{cFit} or \mathbf{cSolv} satisfying (Com), (Nil) and (Solv), respectively, is then established in the manner of Theorem 4.6 showing their maximality among the appropriate class of complexity functions replacing spans of gems by necklaces of the appropriate type (derived necklaces, Fitting necklaces, resp. solvability necklaces). \square

The normal property for each of the three functions is shown just as in Theorem 4.8.

Theorem 6.19 (Normal Property). *The finite group complexity functions **Der**, **Fit** and **Solv** each satisfy the Normal property $\mathbf{c}(N) \triangleleft \mathbf{c}(G)$ for $N \triangleleft G$.*

Next we obtain a series of complexity lower bounds.

Theorem 6.20 (Complexity Lower Bounds). *Let G be a finite group, then*

$$\mathbf{Solv}(G) \leq \mathbf{Fit}(G) \leq \mathbf{Der}(G) \leq \mathbf{cx}(G)$$

Proof. This follows immediately from examining the quotients in a minimal subnormal series of the appropriate kind since a span of gems is a derived necklace, a derived necklace is a Fitting necklace, and a Fitting necklace is a solvability necklace. \square

Theorem 6.21. (1) *There exist unique maximal complexity functions defined on solvable groups satisfying $\mathbf{Solv}_{\mathbf{Solv}}$, $\mathbf{Fit}_{\mathbf{Solv}}$ and $\mathbf{Der}_{\mathbf{Solv}}$ solvability (Solv), nilpotent (Nil) and commutative axioms (Com), respectively.*

- (2) $\mathbf{Fit}_{\mathbf{Solv}}$ is the restriction of \mathbf{Fit} to solvable groups.
- (3) $\mathbf{Der}_{\mathbf{Solv}}$ is the restriction of \mathbf{Der} to solvable groups.
- (4) $\mathbf{Solv}_{\mathbf{Solv}}$ is the restriction of \mathbf{Solv} to solvable groups, and $\mathbf{Solv}_{\mathbf{Solv}} = \delta$ which assigns the value 1 to all non-trivial solvable groups.
- (5) $\mathbf{Der}_{\mathbf{Solv}}$, $\mathbf{Fit}_{\mathbf{Solv}}$ and $\mathbf{Solv}_{\mathbf{Solv}}$ each satisfy the subgroup axiom.

Proof. (1) That these maximal complexity functions on solvable groups exist again follows by the same reasoning as in Theorem 2.3 adding the appropriate axiom. Arguing just as in the proof that $\mathbf{cx}_{\mathbf{Solv}}(G) = \mathbf{cx}(G)$ (Theorem 6.12(1)), we have (2) and (3): For finite solvable groups, no simple non-abelian groups can occur as factors of a subnormal series, since otherwise they would occur as Jordan-Hölder factors, contradicting the solvability. Hence the definition of \mathbf{Fit} coincides with \mathbf{Fit} on solvable groups, and the definition of \mathbf{Der} coincides with \mathbf{Der} on solvable groups.

The assertion (4) is trivially true, since a solvable group is the product of itself and zero simple groups. (5) That these complexity measures each satisfy the subgroup axiom follows as in Theorem 6.12(2) replacing “direct product of abelian simple groups” in the proof by “abelian group”, “nilpotent group”, and “solvable group”, respectively. \square

Definition 6.22. The *Fitting subgroup* $F(G)$ of a group G is generated by all normal nilpotent subgroups of G . By Fitting’s Theorem (e.g. [17, Theorem 5.2.8]), the product of two normal nilpotent subgroups is nilpotent

and normal in G , so it follows $F(G)$ is the unique maximal nilpotent normal subgroup of G . Let $F_0 = \mathbf{1}$, and inductively define $F_{n+1} = \varphi_i^{-1}(F(G/F_n))$ where $\varphi_i : G \twoheadrightarrow G/F_i$ is the natural quotient map. It is easy to see this sequence terminates at G if and only if G is solvable. The *Fitting height* of a solvable group G is the least n such that $F_n = G$.

We use the following Lemma from [15, p. 17].

Lemma 6.23 (Rhodes). *The Fitting height of a solvable group G equals*

- (1) *the smallest integer n such that $G = N_n \triangleright N_{n-1} \triangleright \cdots \triangleright N_1 \triangleright N_0 = \mathbf{1}$ with N_{i+1}/N_i nilpotent ($0 \leq i < n$).*
- (2) *the least m such that G is a homomorphic image of a subgroup of $M_m \wr \cdots \wr M_1$, where the M_i are nilpotent ($0 \leq i < m$).*
- (3) *the least k such that there are surjective morphisms $G = G_k \twoheadrightarrow G_{k-1} \twoheadrightarrow \cdots \twoheadrightarrow G_1 \twoheadrightarrow G_0 = \mathbf{1}$ whose kernels are nilpotent.*

Theorem 6.24. *Let G be a finite solvable group. The derived length of group G is $\mathbf{Der}(G)$. The Fitting height of G is $\mathbf{Fit}(G)$.*

Proof. Since G is solvable, simple non-abelian factors cannot occur in the quotients of any subnormal series for G . Since the derived series of G is the shortest subnormal series with abelian quotients (e.g., [10, proof of Theorem 9.2.5]), its length coincides with $\mathbf{Der}(G)$ for solvable G by definition of \mathbf{Der} . Lemma 6.23(1) says exactly that $\mathbf{Fit}(G)$ is the Fitting height of G . \square

6.6. Wreath Product of Abelian Groups and Hierarchical Complexity of Iterated Wreath Products.

Fact 6.25. *The derived length of an n -fold wreath product of non-trivial abelian groups $W = H_n \wr \cdots \wr H_1$ is $\mathbf{Der}(W) = n$.*

Proof. One can show this by induction of n : $[W, W]$ has trivial coordinate at the top level, and each successive derived subgroup has only trivial coordinates at the next level, but non-trivial coordinates below that. Precisely, consider a wreath product $G = G_2 \wr G_1$. Let $w = (1, w_1)$, $n = (n_2, 1) \in G_2 \wr G_1$, with $w_1 \neq 1$. Consider the action of the commutator $[w, n] = w^{-1}n^{-1}wn$ on $(y, x) \in G_2 \times G_1$:¹⁰

$$\begin{aligned}
 (y, x)[w, n] &= (y, x)(1, w_1)^{-1}(n_2, 1)^{-1}(1, w_1)(n_2, 1) \\
 &= (y, xw_1^{-1})(n_2, 1)^{-1}(1, w_1)(n_2, 1) \\
 &= (y(n_2(xw_1^{-1}))^{-1}, xw_1^{-1})(1, w_1)(n_2, 1) \\
 &= (y(n_2(xw_1^{-1}))^{-1}, xw_1^{-1}w_1)(n_2, 1) \\
 &= (y(n_2(xw_1^{-1}))^{-1}n_2(x), x).
 \end{aligned}$$

This computation shows that $[w, n] \in G_2^{G_1} \rtimes \mathbf{1} \triangleleft G_2^{G_1} \rtimes G_1 = G_2 \wr G_1$. Since $1 \neq w_1 \in G_1$, for any $x_0 \in G_2$, let $n_2(x_0) = g_2$ but $n_2(x) = 1$ for $x \neq x_0$. This yields $[w, n] = (n', 1)$ with $n'(x_0) = g_2$, $n'(x_0w_1) = g_2^{-1}$ and $n'(x) = 1$ for $x \notin \{x_0, x_0w_1\}$. In particular, in $[G, G]$ we can obtain any $g_2 \in G_2$ in the x_0 position of the direct product $G_2^{G_1} \rtimes \mathbf{1} \cong G_2 \times \cdots \times G_2$. It follows

¹⁰Here we are using the fact that the inverse of $w = (w_2, w_1) \in W$ is $w^{-1} = (w_2, w_1)^{-1}$ mapping $(y, x) \in G_2 \times G_1$ to $(y(w_2(xw_1^{-1}))^{-1}, xw_1^{-1})$.

the $[G, G]$ projects onto G_2 . Hence $\mathbf{Der}([G, G]) \geq \mathbf{Der}(G_2)$ by the quotient axiom for \mathbf{Der} . Therefore

$$\mathbf{Der}(G_2 \wr G_1) = \mathbf{Der}([G_2 \wr G_1, G_2 \wr G_1]) + 1 \geq \mathbf{Der}(G_2) + 1.$$

We have $\mathbf{Der}(H_1) = 1$. Suppose the proposition holds for n -factors, then by induction hypothesis $G_2 = H_{n+1} \wr \cdots \wr H_2$ has $\mathbf{Der}(G_2) = n$ so $W = G_2 \wr H_1$ has $\mathbf{Der}(W) \geq \mathbf{Der}(G_2) + 1 = n + 1$. By the extension and product axioms for \mathbf{Der} , $\mathbf{Der}(W) = \mathbf{Der}(G_2^{H_1}) + \mathbf{Der}(H_1) = \mathbf{Der}(G_2) + 1 = n + 1$. The result now follows by induction. \square

Proposition 6.26 (Iterated Wreath Product of Complexity 1 Groups). *Let $W = H_n \wr \cdots \wr H_1$ with each H_i be a commutative group of complexity 1 ($\mathbf{cx}(H_i) = 1$) for $1 \leq i \leq n$. Then $\mathbf{cx}(W) = n$.*

Proof. Each H_i be a direct product of one or more simple groups for $1 \leq i \leq n$. Let $W = \wr_{i=1}^n H_i$ be the wreath product. Then by the extension axiom, $\mathbf{cx}(W) \leq \sum_{i=1}^n \mathbf{cx}(H_i)$, but $\mathbf{cx}(H_i) = 1$ by the product axiom since simple groups have complexity 1. Thus $\mathbf{cx}(W) \leq n$. Now suppose each of the simple groups involved in the construction of W is cyclic, so that H_i is commutative (and hence W is solvable). By Fact 6.25, it follows the derived length of W is n . So $n = \mathbf{Der}(W) \leq \mathbf{cx}(W) \leq n$. Therefore, $\mathbf{cx}(W) = n$. \square

Open Question 3. Let W be a iterated wreath product of n (non-trivial) spans of gems G_1, \dots, G_n , for $n \geq 2$. We know if all the G_i are solvable then $\mathbf{cx}(W) = n$ by Prop. 6.26. Also, by Prop. 4.10, $\mathbf{cx}(W) = 2$ for $n = 2$ even if some G_i are simple non-abelian groups. Does $\mathbf{cx}(W) = n$ in general?

7. MUTUAL INDEPENDENCE OF THE COMPLEXITY AXIOMS

7.1. Normal Subgroup Property is Independent from the Other Complexity Axioms. Here we construct complexity functions that do not have the normal subgroup property. Let S be a finite simple group. For each finite group, let

$$\mathbf{sur}_S(G) = \begin{cases} 1 & \text{there is surjective morphism } \varphi : G \twoheadrightarrow S \\ 0 & \text{otherwise.} \end{cases}$$

This is the characteristic function of groups mapping onto S .

Then $\mathbf{sur}_S(S) = 1$ but $\mathbf{sur}_S(K) = 0$ for K any other finite simple group. Hence by Lemma 1.3, $\mathbf{sur}_S(G)$ satisfies the constructability axiom.

$\mathbf{sur}_S(1) = 0$ so the initial condition holds.

The product axiom holds: Consider a product group $G \times H$. If $H \twoheadrightarrow S$, then pre-composing with the projection yields $G \times H \twoheadrightarrow H \twoheadrightarrow S$. Thus $\mathbf{sur}_S(H) = 1$ implies $\mathbf{sur}_S(G \times H) = 1$. Similarly, for the case $\mathbf{sur}_S(G) = 1$. This shows if $\max(\mathbf{sur}_S(G), \mathbf{sur}_S(H)) = 1$ then $\mathbf{sur}_S(G \times H) = 1$. Otherwise, $\max(\mathbf{sur}_S(G), \mathbf{sur}_S(H)) = 0$. Suppose there were a $\varphi : G \times H \twoheadrightarrow S$. Pre-composing with the injection

$$G \cong G \times 1 \rightarrow G \times H \twoheadrightarrow S.$$

The image of $G \cong G \times 1$ in S is a normal subgroup of S , since a surjective morphism maps normal subgroups to normal subgroups of its image, thus

$\varphi(G \times \mathbf{1})$ is either $\mathbf{1}$ or S . But the image cannot be S since $\mathbf{sur}_S(G) = 0$, i.e. G does not map onto S . Similarly, $\mathbf{1} \times H$ maps to $\mathbf{1}$ since $\mathbf{sur}_S(H) = 0$. This shows $G \times \mathbf{1}$ and $\mathbf{1} \times H$ are both in the kernel of φ , therefore $(G \times \mathbf{1})(\mathbf{1} \times H) = G \times H$ is in the kernel of φ , contradicting $\varphi(G \times H) = S$. This shows if $\max(\mathbf{sur}_S(G), \mathbf{sur}_S(H)) = 0$ then $\mathbf{sur}_S(G \times H) = 0$. Thus, the product axiom holds.

The quotient axiom holds: Suppose $G \twoheadrightarrow H$. If $\mathbf{sur}_S(G) = 1$, then we are done since $\mathbf{sur}_S(G) \geq \mathbf{sur}_S(H)$. Otherwise $\mathbf{sur}_S(G) = 0$. Suppose there were a surjective morphism $H \twoheadrightarrow K$. The precomposing with the map from G to H yields a surjective morphism $G \twoheadrightarrow H \twoheadrightarrow K$, contradicting $\mathbf{sur}_S(G) = 0$. Hence there can be no map $H \twoheadrightarrow K$ and $\mathbf{sur}_S(H) = 0$. In either case, $\mathbf{sur}_S(G) \geq \mathbf{sur}_S(H)$.

Extension axiom: Suppose $N \triangleleft G$. We must show $\mathbf{sur}_S(G) \leq \mathbf{sur}_S(N) + \mathbf{sur}_S(G/N)$. If $\mathbf{sur}_S(G) = 0$, we are done. If $\mathbf{sur}_S(G) = 1$, then we have a $\varphi : G \twoheadrightarrow S$. Let K be the kernel of φ . If $N \triangleleft K$, then $\varphi(N) = \mathbf{1}$. Otherwise, since surjective morphisms map normal subgroups to normal subgroups and S is simple, it follows that $\varphi(N) = S$, whence $\mathbf{sur}_S(G) = 1 \leq 1 + \mathbf{sur}_S(G/N) = \mathbf{sur}_S(N) + \mathbf{sur}_S(G/N)$ holds. We are left with the case $N \triangleleft K$. In this case, G/N maps homomorphically onto G/K , which maps onto S . That is, $\mathbf{sur}_S(G/N) = 1$ and extension holds.

Proposition 7.1. *The function \mathbf{sur}_S satisfies the initial condition, constructability axiom, product axiom, quotient axiom and the extension axiom, hence is a complexity function on finite groups. However, \mathbf{sur}_S does not have the normal subgroup property, that is there a group G and normal subgroup N with $\mathbf{sur}_S(N) > \mathbf{sur}_S(G)$.*

Proof. \mathbf{sur}_S is a complexity function since it satisfies the axioms listed as just shown. Suppose G is any group that is an extension by $N = S$ of any other simple group Q ($Q \not\cong S$), such that G is not a direct product. Then G maps onto Q but not S , so $\mathbf{sur}_S(G) = 0$ but $\mathbf{sur}_S(S) = 1$ even though $S \triangleleft G$. More generally, we can always take G to be the wreath product $S \wr Q$, which is a semidirect product of Q with N , a direct product of copies of S . The wreath product $S \wr Q$ maps onto Q but not S , and has N as a normal subgroup. By the product axiom, $\mathbf{sur}_S(N) = 1$, but $\mathbf{sur}_S(G) = 0$. \square

Counterexample 13 (Failure of Normal Subgroup Property). *The normal subgroup property need not hold for a complexity function.*

- (1) For the complexity function \mathbf{sur}_{A_5} , take $G = S_5$ the symmetric group on 5 objects, which is an extension of \mathbb{Z}_2 by the alternating simple group A_5 . We have $\mathbf{sur}_{A_5}(S_5) = 0$ but $\mathbf{sur}_{A_5}(A_5) = 1$ with the simple alternating group $A_5 \triangleleft S_5$.
- (2) For the complexity function $\mathbf{sur}_{\mathbb{Z}_2}$, consider the special linear group $SL(2, 5)$ of all 2×2 matrices over the field with 5 elements having determinant 1. $SL(2, 5)$ maps onto $PSL(2, 5) \cong A_5$ with kernel isomorphic to $\mathbb{Z}_2 = \pm I$, where I is the identity matrix. We have $\mathbf{sur}_{\mathbb{Z}_2}(SL(2, 5)) = 0$ but $\mathbb{Z}_2 \triangleleft SL(2, 5)$ with $\mathbf{sur}_{\mathbb{Z}_2}(\mathbb{Z}_2) = 1$.
- (3) (Smallest Counterexample.) The complexity function $\mathbf{sur}_{\mathbb{Z}_3}$ takes value 0 on S_3 , the symmetric group on 3 objects, which is an extension of \mathbb{Z}_2 by \mathbb{Z}_3 . But $\mathbb{Z}_3 \triangleleft S_3$ and $\mathbf{sur}_{\mathbb{Z}_3}(\mathbb{Z}_3) = 1$.

7.2. Independence of the Quotient Axiom.

Proposition 7.2. *Let \mathbf{V} be a class of finite groups closed under direct products and normal subgroups which is not the class of all finite groups and has a least one non-trivial member. Then the characteristic function of non-trivial members of \mathbf{V} defined by*

$$\mathbf{sub}_{\mathbf{V}}(G) = \begin{cases} 1 & \text{if } 1 \neq G \text{ lies in } \mathbf{V} \\ 0 & \text{otherwise.} \end{cases}$$

is a complexity function, but does not satisfy the quotient axiom.

Proof. It is immediate from the properties and \mathbf{V} and the definition of $\mathbf{sub}_{\mathbf{V}}$ that the initial, product and normal subgroup axioms hold. Constructability holds by Lemma 1.3.

Extension axiom: Suppose N is normal in a finite group G . If G is not in \mathbf{V} or $G = 1$, then $0 = \mathbf{sub}_{\mathbf{V}}(G) \leq \mathbf{sub}_{\mathbf{V}}(N) + \mathbf{sub}_{\mathbf{V}}(G/N)$, i.e., the extension property holds. Otherwise, $1 \neq G$ is in \mathbf{V} , then either $N = 1$ and so $\mathbf{sub}_{\mathbf{V}}(G) = 1 = 0 + 1 = \mathbf{sub}_{\mathbf{V}}(N) + \mathbf{sub}_{\mathbf{V}}(G/N)$, since $G/N \cong G$; or $N \neq 1$ and so $\mathbf{sub}_{\mathbf{V}}(G) = 1 \leq 1 + \mathbf{sub}_{\mathbf{V}}(G/N) = \mathbf{sub}_{\mathbf{V}}(N) + \mathbf{sub}_{\mathbf{V}}(G/N)$. Thus $\mathbf{sub}_{\mathbf{V}}$ satisfies the extension axioms. We conclude \mathbf{sub} is a complexity function.

Failure of quotient property: By the hypotheses, \mathbf{V} contains some non-trivial group V and there exists a least one group L not in \mathbf{V} . Observe that $V \times L$ is not in \mathbf{V} , since \mathbf{V} is closed under normal subgroups but its normal subgroup $L \cong 1 \times L$ is not in \mathbf{V} . Now V is a quotient of $V \times L$, but we have $\mathbf{sub}_V(V \times L) = 0$ and $\mathbf{sub}(V) = 1$. Hence, the quotient axiom does not hold for \mathbf{sub}_V . \square

Examples 2. Using Proposition 7.2 we may construct a plethora of complexity functions not satisfying the quotient axiom:

- (1) Consider $\text{Span}(\text{SJMPL}\mathcal{E})$ the collection of all finite products of finite simple groups. By Lemma 4.3, products, normal subgroups and quotients of members of $\text{Span}(\text{SJMPL}\mathcal{E})$ lie in $\text{Span}(\text{SJMPL}\mathcal{E})$. By Prop. 7.2, $\mathbf{sub}_{\text{Span}(\text{SJMPL}\mathcal{E})}$ satisfies all the complexity axioms except quotient axiom. For example, this function assigns the value 1 to the quotient $\mathbb{Z}_2 \cong S_5/A_5$ of the symmetric group S_5 but assigns 0 to S_5 .
- (2) Let $\mathbf{V} = \mathbf{Nil}$, all nilpotent groups, or $\mathbf{V} = \mathbf{Solv}$, all solvable groups, and applying Prop. 7.2, we obtain that the characteristic function $\mathbf{sub}_{\mathbf{Nil}}$ for non-trivial nilpotent groups and the characteristic function $\mathbf{sub}_{\mathbf{Solv}}$ for non-trivial groups solvable groups are both complexity functions not satisfying the quotient axiom.

7.3. Independence of Each Complexity Axiom from the Rest.

Theorem 7.3 (Independence). *Consider the set of five complexity axioms*

Axioms = {product axiom, extension axiom, initial, quotient, constructability},

(as they are stated in the Introduction) and also the adding the normal subgroup property, consider

$$\mathbf{Axioms}' = \mathbf{Axioms} \cup \{\text{normal subgroup property}\}.$$

Then for each $\phi \in \mathbf{Axioms}'$, there is a function on \mathbf{c} on finite groups satisfying all members of $\mathbf{Axioms}' \setminus \{\phi\}$, but not satisfying ϕ . Therefore, none of the axioms follows from a proper subset of \mathbf{Axioms}' .

Proof. Here in the table below we collect example functions satisfying all but one property ϕ and references to proofs that this is so. This proves the theorem.

Axiom / Property ϕ	Function	Proof
Product	JH	Theorem 2.2
Extension	sx	Corollary 5.15
Initial	$\mathbf{c} = 1$	(trivial check)
Quotient	sub_{Nil}	Prop. 7.2
Constructability	$\mathbf{c}(G) = 2$ for $G \neq 1$, $\mathbf{c}(1) = 0$	Lemma 1.3, plus trivial check.
Normal Subgroup	sur_S (S simple)	Prop. 7.1

Functions satisfying All Axioms/Properties except the one named.

□

Remarks 7.4. 1. While the subgroup axiom is not one of the complexity axioms, for completeness, we record here that **cx** is an example of a function that satisfies satisfies all the axioms and the normal subgroup property, but not the subgroup axiom (Prop. 3.1).

2. In addition, the following complexity functions on finite groups satisfy all the axioms including the normal subgroup property: z , δ , **cx**, χ_s , χ_s^* , **Der**, **Fit**, **Solv**.

3. On solvable groups, the following satisfy all the axioms including the subgroup axiom: z , δ , **cx** = **cx_{solv}**, χ_s , χ_s^* , \log_p (p prime), **Der** = **Der_{solv}**, **Fit** = **Fit_{solv}**, **Solv_{solv}** = δ .

REFERENCES

- [1] G. Baumslag. *Topics in Combinatorial Group Theory*. Lectures in Mathematics / ETH Zürich, Birkhäuser, 1993.
- [2] J. J. Cannon and D. F. Holt. Chief series, composition series and socles. *Journal of Symbolic Computation*, 24:285–301, 1997.
- [3] J. D. Dixon and B. Mortimer. *Permutation Groups*. Springer, 1996.
- [4] P. Dömösi and C. L. Nehaniv. *Algebraic Theory of Finite Automata Networks: An Introduction*, volume 11 of *SIAM Series on Discrete Mathematics and Applications*. Society for Industrial and Applied Mathematics, 2005.
- [5] A. Egri-Nagy, J. D. Mitchell, and C. L. Nehaniv. SgpDec: Cascade (de)compositions of finite transformation semigroups and permutation groups. In *International Congress on Mathematical Software*, volume 8592 of *Lecture Notes in Computer Science*, pages 75–82. Springer, 2014.
- [6] A. Egri-Nagy and C. L. Nehaniv. Cascade product of permutation groups. arXiv:1303.0091 [math.GR], <https://arxiv.org/abs/1303.0091>, 2013.
- [7] S. Eilenberg. *Automata, Languages and Machines*, volume B. Academic Press, 1976.
- [8] G. Frobenius. Über auflösbare Gruppen. *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin*, XXIII:337–345, 4 Mai 1893. <https://www.biodiversitylibrary.org/item/93363#page/367/mode/2up>.
- [9] I. Ganev. Groups of a square-free order. *Rose-Hulman Undergraduate Mathematics Journal*, 11(1):Article 7, 2010. <https://scholar.rose-hulman.edu/rhumj/vol11/iss1/7>.
- [10] M. Hall, Jr. *The Theory of Groups*. The Macmillan Company, 1959.

- [11] L. Kaloujnine and M. Krasner. Produit complet des groupes de permutations et le problème d'extension de groupes I, II, III. *Acta Scientiarum Mathematicarum (Szeged)*, 13:208–230, 1950; 14:33–66, 69–82, 1951.
- [12] K. Krohn and J. Rhodes. Results on finite semigroups derived from the algebraic theory of machines. *Proceedings of the National Academy of Sciences U.S.A.*, 53:499–501, 1965.
- [13] C. L. Nehaniv. Complexity of finite aperiodic semigroups and star-free languages. In J. Almeida, G. Gomes, and P. Silva, editors, *Semigroups, Automata, Languages*, pages 195–209. World Scientific, 1996.
- [14] C. L. Nehaniv and J. Rhodes. The evolution and understanding of hierarchical complexity in biology from an algebraic perspective. *Artificial Life*, 6(1):45–67, 2000.
- [15] J. Rhodes. *Applications of Automata Theory and Algebra via the Mathematical Theory of Complexity to Biology, Physics, Psychology, Philosophy, and Games*. Edited by C. L. Nehaniv, Foreword by M. W. Hirsch, World Scientific, 2010.
- [16] J. Rhodes and B. Steinberg. *The q -Theory of Finite Semigroups*. Springer, 2009.
- [17] D. J. Robinson. *A Course in the Theory of Groups*. Springer, 1993.
- [18] J. J. Rotman. *An Introduction to the Theory of Groups*. Springer, 4th edition, 1995.
- [19] B. Tilson. Complexity of semigroups and morphisms. In S. Eilenberg, editor, *Automata, Languages and Machines*, volume B, chapter XII. Academic Press, 1976.

UNIVERSITY OF WATERLOO, ONTARIO, CANADA
Email address: `chrystopher.nehaniv@uwaterloo.ca`