

Counting and countable sets

COUNTABLE SETS and counting schemes for infinite countable sets are the topics of the problem in this note.

Problem

Let $P = \{N \subset \mathbb{N} : N \text{ finite}\}$. Prove P is countable.

Let's revisit what it means for an infinite set to be countable: An infinite set M is countable if there is a bijection ¹ from M to \mathbb{N} .

Given this definition, the problem statement is quite remarkable: the set of all the finite subsets of \mathbb{N} is not "bigger" than \mathbb{N} .

Our strategy will be to start smaller and prove certain subsets of P are countable. We then expand it to P . We start by proving that the set of all subsets of \mathbb{N} of size two is countable. We actually will prove something stronger, namely the set of ordered pairs of natural numbers is countable.

Theorem 1.1. *The set of ordered pairs $\mathbb{N} \times \mathbb{N}$ is countable.*

Proof. We need a bijection from $\mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$. There are many ways to do this ².

The main idea we are going to use for our bijection is to order the pairs $(i, j) \in \mathbb{N} \times \mathbb{N}$ in rows, such that each pair in a row has the same value when summing the components of the pair. Figure 1.1 illustrates the idea. Row one has all pairs with components that sum up to two (in this case only one pair). Row two has all pairs with components that sum up to three, row three all pairs which sum to four, Notice also that in a row the pairs are sorted in increasing order of the first component.

We count the pairs from left to right in each row and go down the rows starting at the first row. For a given pair (i, j) , how many pairs come before it in our counting scheme? It is in row $i + j - 1$, so there are $k : 1 \leq k < i + j - 1$ rows before it. Each row k has k pairs in it. This means there are

$$\sum_{k=1}^{i+j-2} k = \frac{(i+j-2)(i+j-1)}{2}$$

pairs in rows before our pair (i, j) . There are $i - 1$ pairs before (i, j) in the same row. Therefore, our counting function is

$$f : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}, \quad f(i, j) = i + \frac{(i+j-2)(i+j-1)}{2}$$

¹ A bijection is a function that is one-to-one and onto.

Mention puzzle 136 (Catching a Spy) from Levitin: Algorithmic Puzzles

² A very elegant way is described at <http://www.math.upenn.edu/~wilf/website/recounting.pdf>




Figure 1.1: Counting all pairs.

Suppose we have two pairs $(i_1, j_1) \neq (i_2, j_2)$. We have two cases:

- $i_1 + j_1 = i_2 + j_2$, same row, then $i_1 \neq i_2$, so $f(i_1, j_1) \neq f(i_2, j_2)$
- $i_1 + j_1 \neq i_2 + j_2$, different rows, so $f(i_1, j_1) \neq f(i_2, j_2)$

This means, f is one-to-one.

To prove that f is onto, we consider an arbitrary $n \in \mathbb{N}$ and find a pair (i, j) with $f(i, j) = n$. Working backwards and assuming we have a pair (i, j) with $f(i, j) = n$, it would fall on a row $r = i + j - 1$. In each row k there are k pairs, n is on row r , so

$$\sum_{k=1}^{r-1} k = \frac{r(r-1)}{2} < n \leq \sum_{k=1}^r k = \frac{r(r+1)}{2}$$

Solving for r we have:³

$$r^2 - r - 2n < 0, \quad r^2 + r - 2n \geq 0, \quad r = \left\lceil \frac{-1 + \sqrt{1 + 8n}}{2} \right\rceil$$

And then

$$i = n - \frac{r(r-1)}{2}, \quad j = r - i + 1$$

This means that given an arbitrary n , there exists a pair (i, j) with $f(i, j) = n$, so f is onto.

It follows that f is a bijection and $\mathbb{N} \times \mathbb{N}$ is countable. \square

A corollary to Theorem 1.1 let's us expand the countable subsets of P even more.

Corollary. *Set of all finite sequences of length k , \mathbb{N}^k is countable.*⁴

Proof. Follows by induction on k : Assuming \mathbb{N}^{k-1} is countable, then

$$\mathbb{N}^k = \mathbb{N}^{k-1} \times \mathbb{N}$$

is also countable according to Theorem 1.1. \square

From the corollary we now know ⁵ that the set of all subsets of \mathbb{N} of size k is countable (it's a subset of \mathbb{N}^k). The problem in this section asks us to prove that P is countable, which means the union of all these countable sets is countable. The next theorem⁶ will prove just that.

Theorem 1.2. *Let A_n , $n \in \mathbb{N}$ be countable sets. Then*

$$\bigcup_{n=1}^{\infty} A_n$$

is countable.

³ Note that $\frac{1+\sqrt{1+8n}}{2} - \frac{-1+\sqrt{1+8n}}{2} = 1$

⁴ \mathbb{N}^k is the set of sequences of length k , or the cartesian product $\mathbb{N} \times \mathbb{N} \times \dots \times \mathbb{N}$. The set of pairs is $\mathbb{N}^2 = \mathbb{N} \times \mathbb{N}$.

⁵ We keep using the fact that the set of all finite subsets of \mathbb{N} of size k is a subset of the set of all sequences of size k . To see this impose an order on a set of size k and you get a sequence.

⁶ Exercise 1.5.3 on page 30 from Stephen Abbott. *Understanding Analysis*. Springer, 2 edition, 2015. ISBN 978-1-4939-2711-1.

Proof. A_n is countable, so there exists a bijection $f_n : \mathbb{N} \rightarrow A_n$. We already know that $\mathbb{N} \times \mathbb{N}$ is countable, so there exists a bijection $g : \mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$.

We define $F : \mathbb{N} \rightarrow \bigcup_{n=1}^{\infty} A_n$

$$F(n) = f_i(j), \text{ where } (i, j) = g(n)$$

We claim that F is a bijection.

Take $n_1 \neq n_2$. Then $g(n_1) \neq g(n_2)$ and $(i_1, j_1) \neq (i_2, j_2)$, so

$$f_{i_1}(j_1) \neq f_{i_2}(j_2)$$

It means $F(n_1) \neq F(n_2)$ and F is one-to-one.

Now pick an arbitrary $a \in \bigcup_{n=1}^{\infty} A_n$. Then there exists $i \in \mathbb{N}$ with $a \in A_i$.⁷ There also exists $j \in \mathbb{N}$ with $f_i(j) = a$. The pair (i, j) is in $\mathbb{N} \times \mathbb{N}$, so there exists $n \in \mathbb{N}$ with $g(n) = (i, j)$. It follows that $F(n) = a$ and F is onto. \square

⁷We assume here the A_i are disjoint, if not we make them disjoint and their union stays the same.

Bibliography

Stephen Abbott. *Understanding Analysis*. Springer, 2 edition, 2015.
ISBN 978-1-4939-2711-1.