



Triangulation


Triangulation



Triangulation



Triangulation



Triangulation

Given a set of (noisy) matched points

$$\{\mathbf{x}_i, \mathbf{x}'_i\}$$

and camera matrices

$$\mathbf{P}, \mathbf{P}'$$


Estimate the 3D point

$$\mathbf{X}$$

$$\mathbf{x} = \mathbf{P} \mathbf{X}$$

known known

*Can we compute **X** from a single
correspondence **x**?*



$$\mathbf{x} = \mathbf{P}\mathbf{X}$$

known known

*Can we compute **X** from two
correspondences **x** and **x'**?*

$$\underset{\text{known}}{\mathbf{x}} = \underset{\text{known}}{\mathbf{P}} \mathbf{X}$$

*Can we compute **X** from two
correspondences **x** and **x'**?*

yes if perfect measurements

$$\underset{\text{known}}{\mathbf{x}} = \underset{\text{known}}{\mathbf{P}} \mathbf{X}$$

Can we compute \mathbf{X} from two correspondences \mathbf{x} and \mathbf{x}' ?

yes if perfect measurements

There will not be a point that satisfies both constraints
because the measurements are usually noisy

$$\mathbf{x}' = \mathbf{P}' \mathbf{X} \quad \mathbf{x} = \mathbf{P} \mathbf{X}$$

Need to find the **best fit**

$$\mathbf{x} = \mathbf{P} \mathbf{X}$$

(homogeneous
coordinate)

Also, this is a similarity relation because it involves homogeneous coordinates

(inhomogeneous
coordinate)

Same ray direction but differs by a scale factor

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

How do we solve for unknowns in a similarity relation?

$$\mathbf{x} = \mathbf{P} \mathbf{X}$$

(homogeneous
coordinate)

Also, this is a similarity relation because it involves homogeneous coordinates

Same ray direction but differs by a scale factor

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

How do we solve for unknowns in a similarity relation?

Direct Linear Transform

Remove scale factor, convert to linear system and solve with 

$$\mathbf{x} = \mathbf{P} \mathbf{X}$$

(homogeneous
coordinate)

Also, this is a similarity relation because it involves homogeneous coordinates

(inhomogeneous
coordinate)

Same ray direction but differs by a scale factor

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

How do we solve for unknowns in a similarity relation?

Direct Linear Transform

Remove scale factor, convert to linear system and solve with SVD.

$$\mathbf{x} \times \mathbf{P}X = \mathbf{0}$$


Cross product of two vectors of same direction is zero
(this equality removes the scale factor)

Recall: Cross Product

Vector (cross) product

takes two vectors and returns a vector perpendicular to both

$$\mathbf{c} = \mathbf{a} \times \mathbf{b}$$



$$\mathbf{c} \cdot \mathbf{a} = 0$$

$$\mathbf{c} \cdot \mathbf{b} = 0$$

$$\mathbf{a} \times \mathbf{b} = \begin{bmatrix} a_2b_3 - a_3b_2 \\ a_3b_1 - a_1b_3 \\ a_1b_2 - a_2b_1 \end{bmatrix}$$

cross product of two vectors in the same direction is zero

$$\mathbf{a} \times \mathbf{a} = 0$$

remember this!!!

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \text{---} & \boldsymbol{p_1}^\top & \text{---} \\ \text{---} & \boldsymbol{p_2}^\top & \text{---} \\ \text{---} & \boldsymbol{p_3}^\top & \text{---} \end{bmatrix} \begin{bmatrix} | \\ \boldsymbol{X} \\ | \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \boldsymbol{p_1}^\top \boldsymbol{X} \\ \boldsymbol{p_2}^\top \boldsymbol{X} \\ \boldsymbol{p_3}^\top \boldsymbol{X} \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \text{---} & \mathbf{p_1^\top} & \text{---} \\ \text{---} & \mathbf{p_2^\top} & \text{---} \\ \text{---} & \mathbf{p_3^\top} & \text{---} \end{bmatrix} \begin{bmatrix} | \\ \mathbf{X} \\ | \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{p_1^\top X} \\ \mathbf{p_2^\top X} \\ \mathbf{p_3^\top X} \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \times \begin{bmatrix} \mathbf{p_1^\top X} \\ \mathbf{p_2^\top X} \\ \mathbf{p_3^\top X} \end{bmatrix} = \begin{bmatrix} y\mathbf{p_3^\top X} - \mathbf{p_2^\top X} \\ \mathbf{p_1^\top X} - x\mathbf{p_3^\top X} \\ x\mathbf{p_2^\top X} - y\mathbf{p_1^\top X} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Using the fact that the cross product should be zero

$$\mathbf{x} \times \mathbf{P}\mathbf{X} = \mathbf{0}$$

$$\begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \times \begin{bmatrix} \mathbf{p}_1^\top \mathbf{X} \\ \mathbf{p}_2^\top \mathbf{X} \\ \mathbf{p}_3^\top \mathbf{X} \end{bmatrix} = \begin{bmatrix} y\mathbf{p}_3^\top \mathbf{X} - \mathbf{p}_2^\top \mathbf{X} \\ \mathbf{p}_1^\top \mathbf{X} - x\mathbf{p}_3^\top \mathbf{X} \\ x\mathbf{p}_2^\top \mathbf{X} - y\mathbf{p}_1^\top \mathbf{X} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Third line is a linear combination of the first and second lines.
(x times the first line plus y times the second line)

One 2D to 3D point correspondence give you  equations

Using the fact that the cross product should be zero

$$\mathbf{x} \times \mathbf{P}\mathbf{X} = \mathbf{0}$$

$$\begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \times \begin{bmatrix} \mathbf{p}_1^\top \mathbf{X} \\ \mathbf{p}_2^\top \mathbf{X} \\ \mathbf{p}_3^\top \mathbf{X} \end{bmatrix} = \begin{bmatrix} y\mathbf{p}_3^\top \mathbf{X} - \mathbf{p}_2^\top \mathbf{X} \\ \mathbf{p}_1^\top \mathbf{X} - x\mathbf{p}_3^\top \mathbf{X} \\ x\mathbf{p}_2^\top \mathbf{X} - y\mathbf{p}_1^\top \mathbf{X} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Third line is a linear combination of the first and second lines.
(x times the first line plus y times the second line)

One 2D to 3D point correspondence give you 2 equations

$$\begin{bmatrix} yp_3^\top X - p_2^\top X \\ p_1^\top X - xp_3^\top X \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} yp_3^\top - p_2^\top \\ p_1^\top - xp_3^\top \end{bmatrix} X = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\mathbf{A}_i X = \mathbf{0}$$

Now we can make a system of linear equations
(two lines for each 2D point correspondence)

Concatenate the 2D points from both images

$$\begin{bmatrix} y\mathbf{p}_3^\top - \mathbf{p}_2^\top \\ \mathbf{p}_1^\top - x\mathbf{p}_3^\top \\ y'\mathbf{p}'_3{}^\top - \mathbf{p}'_2{}^\top \\ \mathbf{p}'_1{}^\top - x'\mathbf{p}'_3{}^\top \end{bmatrix} \mathbf{X} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

sanity check! dimensions?

$$\mathbf{A}\mathbf{X} = \mathbf{0}$$

How do we solve homogeneous linear system?

Concatenate the 2D points from both images

$$\begin{bmatrix} y\mathbf{p}_3^\top - \mathbf{p}_2^\top \\ \mathbf{p}_1^\top - x\mathbf{p}_3^\top \\ y'\mathbf{p}'_3{}^\top - \mathbf{p}'_2{}^\top \\ \mathbf{p}'_1{}^\top - x'\mathbf{p}'_3{}^\top \end{bmatrix} \mathbf{X} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\mathbf{A}\mathbf{X} = \mathbf{0}$$

How do we solve homogeneous linear system?

S

Concatenate the 2D points from both images

$$\begin{bmatrix} y\mathbf{p}_3^\top - \mathbf{p}_2^\top \\ \mathbf{p}_1^\top - x\mathbf{p}_3^\top \\ y'\mathbf{p}'_3{}^\top - \mathbf{p}'_2{}^\top \\ \mathbf{p}'_1{}^\top - x'\mathbf{p}'_3{}^\top \end{bmatrix} \mathbf{X} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\mathbf{A}\mathbf{X} = \mathbf{0}$$

How do we solve homogeneous linear system?

S V

Concatenate the 2D points from both images

$$\begin{bmatrix} y\mathbf{p}_3^\top - \mathbf{p}_2^\top \\ \mathbf{p}_1^\top - x\mathbf{p}_3^\top \\ y'\mathbf{p}'_3{}^\top - \mathbf{p}'_2{}^\top \\ \mathbf{p}'_1{}^\top - x'\mathbf{p}'_3{}^\top \end{bmatrix} \mathbf{X} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\mathbf{A}\mathbf{X} = \mathbf{0}$$

How do we solve homogeneous linear system?

S V D

Concatenate the 2D points from both images

$$\begin{bmatrix} y\mathbf{p}_3^\top - \mathbf{p}_2^\top \\ \mathbf{p}_1^\top - x\mathbf{p}_3^\top \\ y'\mathbf{p}'_3{}^\top - \mathbf{p}'_2{}^\top \\ \mathbf{p}'_1{}^\top - x'\mathbf{p}'_3{}^\top \end{bmatrix} \mathbf{X} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\mathbf{A}\mathbf{X} = \mathbf{0}$$

How do we solve homogeneous linear system?

S V D !

Recall: Total least squares

(**Warning:** change of notation. \mathbf{x} is a vector of parameters!)

$$\begin{aligned} E_{\text{TLS}} &= \sum_i (\mathbf{a}_i \mathbf{x})^2 \\ &= \|\mathbf{A}\mathbf{x}\|^2 && \text{(matrix form)} \\ \|\mathbf{x}\|^2 &= 1 && \text{constraint} \end{aligned}$$

$$\begin{array}{ll} \text{minimize} & \|\mathbf{A}\mathbf{x}\|^2 \\ \text{subject to} & \|\mathbf{x}\|^2 = 1 \end{array} \quad \rightarrow \quad \begin{array}{l} \text{minimize} \quad \frac{\|\mathbf{A}\mathbf{x}\|^2}{\|\mathbf{x}\|^2} \\ \text{(Rayleigh quotient)} \end{array}$$

Solution is the eigenvector
corresponding to smallest eigenvalue of

$$\mathbf{A}^\top \mathbf{A}$$

```

# 3. Compute the Essential Matrix
E, mask = cv2.findEssentialMat(pts1, pts2, K, method=cv2.RANSAC, prob=0.999, threshold=1.0)
print("Essential Matrix:\n", E)

# 4. Recover the relative camera pose (R, t)
_, R, t, mask_pose = cv2.recoverPose(E, pts1, pts2, K)
print("Rotation Matrix (R):\n", R)
print("Translation Vector (t):\n", t)

# 5. Triangulate 3D points
# Build the projection matrices for both cameras
P1 = np.dot(K, np.hstack((np.eye(3), np.zeros((3, 1))))) # First camera (reference)
P2 = np.dot(K, np.hstack((R, t))) # Second camera

# Perform triangulation in homogeneous coordinates
points_4d_h = cv2.triangulatePoints(P1, P2, pts1.T, pts2.T)

# Convert homogeneous coordinates to 3D
points_3d = (points_4d_h / points_4d_h[3])[:3].T

print("Triangulated 3D Points:\n", points_3d)

```

	Structure (scene geometry)	Motion (camera geometry)	Measurements
Pose Estimation	known	estimate	3D to 2D correspondences
Triangulation	estimate	known	2D to 2D coorespondences
Reconstruction	estimate	estimate	2D to 2D coorespondences