Vanya Vorobiov

Basic definitions

What is integrability in elementary functions

Liouville's Theorem (Differential algebra)

Vanya Vorobiov

Sber

January 14, 2025

Through the all of presentation we will suppose that all fields have 0 characteristic.

Through the all of presentation we will suppose that all fields have 0 characteristic.

Definition

Field F is differential if it's equipped with the unary function $^\prime$ such that:

- (a + b)' = a' + b'
- (ab)' = a'b + ab'

Through the all of presentation we will suppose that all fields have 0 characteristic.

Definition

Field F is differential if it's equipped with the unary function $^\prime$ such that:

- (a + b)' = a' + b'
- (ab)' = a'b + ab'

Definition

Subfield $K \subseteq F, K = \{a \in F \mid a' = 0\}$ is called subfield of constants.

Through the all of presentation we will suppose that all fields have 0 characteristic.

Definition

Field F is differential if it's equipped with the unary function ' such that:

- (a + b)' = a' + b'
- (ab)' = a'b + ab'

Definition

Subfield $K \subseteq F, K = \{a \in F \mid a' = 0\}$ is called subfield of constants.

Definition

Differential extension of the differential field F is field E such that $E\supseteq F$ and there is the same differentiation ' on E.

Through the all of presentation we will suppose that all fields have 0 characteristic.

Definition

Field F is differential if it's equipped with the unary function ' such that:

- (a + b)' = a' + b'
- (ab)' = a'b + ab'

Definition

Subfield $K \subseteq F, K = \{a \in F \mid a' = 0\}$ is called subfield of constants.

Definition

Differential extension of the differential field F is field E such that $E\supseteq F$ and there is the same differentiation ' on E.

Definition

Let F be the differential field. Then

- **b** is called the logarithm of a if $b' = \frac{a'}{a}$
- ▶ b is called the exponent of a if $a' = \frac{b'}{b}$

The extension E is called elementary if it can be presented as $E = F(t_1, ..., t_n)$ and for all i t_i is logarithm or exponent or algebraic over $F(t_1, ..., t_{i-1})$.