

Liouville's Theorem on integrability via elementary functions

Vanya Vorobiov

Sber

January 15, 2025

Introduction

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
integrability in
elementary
functions

Liouville's
Theorem
(statement)

The Main
Lemma

TODO

Basic definitions

Remark

Through the all of presentation we will suppose that all fields have 0 characteristic.

Remark

Through the all of presentation we will suppose that all fields have 0 characteristic.

Definition

Field F is differential if it's equipped with the unary function $'$ such that:

- ▶ $(a + b)' = a' + b'$
- ▶ $(ab)' = a'b + ab'$

Remark

Through the all of presentation we will suppose that all fields have 0 characteristic.

Definition

Field F is differential if it's equipped with the unary function $'$ such that:

- ▶ $(a + b)' = a' + b'$
- ▶ $(ab)' = a'b + ab'$

Definition

Subfield $K \subseteq F$, $K = \{a \in F \mid a' = 0\}$ is called subfield of constants.

Remark

Through the all of presentation we will suppose that all fields have 0 characteristic.

Definition

Field F is differential if it's equipped with the unary function $'$ such that:

- ▶ $(a + b)' = a' + b'$
- ▶ $(ab)' = a'b + ab'$

Definition

Subfield $K \subseteq F$, $K = \{a \in F \mid a' = 0\}$ is called subfield of constants.

Definition

Differential extension of the differential field F is field E such that $E \supseteq F$ and there is the same differentiation $'$ on E .

Remark

Through the all of presentation we will suppose that all fields have 0 characteristic.

Definition

Field F is differential if it's equipped with the unary function $'$ such that:

- ▶ $(a + b)' = a' + b'$
- ▶ $(ab)' = a'b + ab'$

Definition

Subfield $K \subseteq F$, $K = \{a \in F \mid a' = 0\}$ is called subfield of constants.

Definition

Differential extension of the differential field F is field E such that $E \supseteq F$ and there is the same differentiation $'$ on E .

Definition

Let F be the differential field. Then

- ▶ b is called the logarithm of a if $b' = \frac{a'}{a}$
- ▶ b is called the exponent of a if $a' = \frac{b'}{b}$

What is integrability in elementary functions

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
integrability in
elementary
functions

Liouville's
Theorem
(statement)

The Main
Lemma

Definition

The extension E of F is called elementary if it can be presented as $E = F(t_1, \dots, t_n)$ and for all i t_i is logarithm or exponent or algebraic over $F(t_1, \dots, t_{i-1})$.

What is integrability in elementary functions

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
integrability in
elementary
functions

Liouville's
Theorem
(statement)

The Main
Lemma

Definition

The extension E of F is called elementary if it can be presented as $E = F(t_1, \dots, t_n)$ and for all i t_i is logarithm or exponent or algebraic over $F(t_1, \dots, t_{i-1})$.

Remark

Common sense says us that some function $f : \mathbb{C} \rightarrow \mathbb{C}$ is elementary iff it can be constructed via finite number of radicals, sines, cosines, exponents, logarithms and hyperbolic functions. One can see that it's consistent with our approach.

Liouville's Theorem (statement)

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
integrability in
elementary
functions

Liouville's
Theorem
(statement)

The Main
Lemma

Theorem (Liouville, 1833-1841)

Let F be a differential field, and K is its subfield of constants. If for $\alpha \in F$ equation $x' = \alpha$ has the solution in some elementary extension of F , then

$$\alpha = \sum_{i=1}^m c_i \frac{u_i'}{u_i} + v'$$

for some $c_1, \dots, c_m \in K$, $u_1, \dots, u_m, v \in F$.

Lemma

Let F be a differential field. And t is transcendental over F , and t is a logarithm or an exponent of some element from F . And let $f \in F[x]$ be a polynomial, $\deg f = k \geq 1$

- ▶ If t is a logarithm then $(f(t))'$ has degree k if the leading coefficient of f is not a constant, and it has degree $k - 1$ if the leading coefficient is a constant.
- ▶ If t is an exponent then $(f(t))'$ has degree k and it's multiple of f if and only if f is a monomial.