

Liouville's Theorem on integrability via elementary functions

Vanya Vorobiov

Sber

January 16, 2025

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

From high school, we are familiar with the idea that some integrals cannot be expressed in terms of elementary functions. For instance:

$$\int e^{\pm x^2} dx, \quad \int \frac{dx}{\ln x}, \quad \int \frac{e^x}{x} dx, \quad \int \frac{\sin x}{x} dx, \quad \int \frac{\sinh x}{x} dx, \quad \int \log \log x dx$$

These examples highlight the limitations of elementary functions in representing certain integrals.

From high school, we are familiar with the idea that some integrals cannot be expressed in terms of elementary functions. For instance:

$$\int e^{\pm x^2} dx, \quad \int \frac{dx}{\ln x}, \quad \int \frac{e^x}{x} dx, \quad \int \frac{\sin x}{x} dx, \quad \int \frac{\sinh x}{x} dx, \quad \int \log \log x dx$$

These examples highlight the limitations of elementary functions in representing certain integrals.

In this presentation, we will:

- ▶ Introduce and prove a powerful tool: Liouville's theorem.
- ▶ Derive these integrals as a consequence.
- ▶ If time permits, discuss some special form of elliptic integrals, which also cannot be expressed in elementary terms.

$$\int \frac{dx}{\sqrt{P(x)}}$$

for $\deg P = 2, 3$ and P has no multiple roots.

Basic definitions

Remark

Through the all of presentation we will suppose that all fields have 0 characteristic.

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Remark

Through the all of presentation we will suppose that all fields have 0 characteristic.

Definition

Field F is differential if it's equipped with the unary function $'$ such that:

- ▶ $(a + b)' = a' + b'$
- ▶ $(ab)' = a'b + ab'$

Remark

Through the all of presentation we will suppose that all fields have 0 characteristic.

Definition

Field F is differential if it's equipped with the unary function $'$ such that:

- ▶ $(a + b)' = a' + b'$
- ▶ $(ab)' = a'b + ab'$

Definition

Subfield $K \subseteq F$, $K = \{a \in F \mid a' = 0\}$ is called subfield of constants.

Remark

Through the all of presentation we will suppose that all fields have 0 characteristic.

Definition

Field F is differential if it's equipped with the unary function $'$ such that:

- ▶ $(a + b)' = a' + b'$
- ▶ $(ab)' = a'b + ab'$

Definition

Subfield $K \subseteq F$, $K = \{a \in F \mid a' = 0\}$ is called subfield of constants.

Definition

Differential extension of the differential field F is field E such that $E \supseteq F$ and there is the same differentiation $'$ on E .

Basic definitions

Remark

Through the all of presentation we will suppose that all fields have 0 characteristic.

Definition

Field F is differential if it's equipped with the unary function $'$ such that:

- ▶ $(a + b)' = a' + b'$
- ▶ $(ab)' = a'b + ab'$

Definition

Subfield $K \subseteq F$, $K = \{a \in F \mid a' = 0\}$ is called subfield of constants.

Definition

Differential extension of the differential field F is field E such that $E \supseteq F$ and there is the same differentiation $'$ on E .

Definition

Let F be the differential field. Then

- ▶ b is called the logarithm of a if $b' = \frac{a'}{a}$
- ▶ b is called the exponent of a if $a' = \frac{b'}{b}$

What is expression in elementary terms

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Definition

The extension E of F is called elementary if it can be presented as $E = F(t_1, \dots, t_n)$ and for all i t_i is logarithm or exponent or algebraic over $F(t_1, \dots, t_{i-1})$.

What is expression in elementary terms

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Definition

The extension E of F is called elementary if it can be presented as $E = F(t_1, \dots, t_n)$ and for all i t_i is logarithm or exponent or algebraic over $F(t_1, \dots, t_{i-1})$.

Remark

Common sense says us that some function $f : \mathbb{C} \rightarrow \mathbb{C}$ is elementary iff it can be constructed via finite number of radicals, sines, cosines, exponents, logarithms and hyperbolic functions. One can see that it's consistent with our approach. Furthermore our definition on elementarity is more general.

Liouville's Theorem (statement)

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Theorem (Liouville, 1833-1841)

Let F be a differential field, and K is its subfield of constants. If for $\alpha \in F$ equation $x' = \alpha$ has the solution in some elementary extension of F , such that its subfield of constants is still K , then

$$\alpha = \sum_{i=1}^m c_i \frac{u_i'}{u_i} + v'$$

for some $c_1, \dots, c_m \in K$, $u_1, \dots, u_m, v \in F$.

The Main Lemma

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Lemma

Let F be a differential field, t is transcendental over F , and t is a logarithm or an exponent of some element from F . And let $f \in F[x]$ be a polynomial, $\deg f = k \geq 1$

- ▶ If t is a logarithm then the degree of $(f(t))'$ is k if the leading coefficient of f is not a constant, and it has degree $k - 1$ if the leading coefficient is a constant.
- ▶ If t is an exponent then the degree of $(f(t))'$ is k and it's multiple of f if and only if f is a monomial.

The Main Lemma

Lemma

Let F be a differential field, t is transcendental over F , and t is a logarithm or an exponent of some element from F . And let $f \in F[x]$ be a polynomial, $\deg f = k \geq 1$

- ▶ If t is a logarithm then the degree of $(f(t))'$ is k if the leading coefficient of f is not a constant, and it has degree $k - 1$ if the leading coefficient is a constant.
- ▶ If t is an exponent then the degree of $(f(t))'$ is k and it's multiple of k if and only if f is a monomial.

Proof.

It's a quite simple technical exercise. □

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Liouville's Theorem (proof)

Let x be the solution of differential equation mentioned above. And $x \in F(t_1, \dots, t_n)$.

We will use induction on n (we don't fix the field F).

For short we denote $t = t_1$.

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Liouville's Theorem (proof)

Liouville's Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is expression in elementary terms

Liouville's Theorem (statement)

The Main Lemma

Liouville's Theorem (proof)

t is a transcendental logarithm

t is a transcendental exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Let x be the solution of differential equation mentioned above. And $x \in F(t_1, \dots, t_n)$.

We will use induction on n (we don't fix the field F).

For short we denote $t = t_1$.

Using the inductive assumption, we get

$$\alpha = \sum_{i=1}^m c_i \frac{u_i'}{u_i} + v'$$

for some $c_1, \dots, c_m \in K$, $u_1, \dots, u_m, v \in F(t)$.

Here we use that the subfield of constants of $F(t)$ is K .

Liouville's Theorem (proof)

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Let x be the solution of differential equation mentioned above. And $x \in F(t_1, \dots, t_n)$.

We will use induction on n (we don't fix the field F).

For short we denote $t = t_1$.

Using the inductive assumption, we get

$$\alpha = \sum_{i=1}^m c_i \frac{u_i'}{u_i} + v'$$

for some $c_1, \dots, c_m \in K$, $u_1, \dots, u_m, v \in F(t)$.

Here we use that the subfield of constants of $F(t)$ is K .

Now we consider 3 cases

- ▶ t is transcendental over F and it is a logarithm;
- ▶ t is transcendental over F and it is an exponent;
- ▶ t is algebraic over F .

t is a transcendental logarithm

Firstly let us consider the basic properties of logarithmical derivatives:

$$\frac{(ab)'}{ab} = \frac{a'}{a} + \frac{b'}{b}, \quad \frac{(1/a)'}{1/a} = -\frac{a'}{a}$$

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

t is a transcendental logarithm

Firstly let us consider the basic properties of logarithmical derivatives:

$$\frac{(ab)'}{ab} = \frac{a'}{a} + \frac{b'}{b}, \quad \frac{(1/a)'}{1/a} = -\frac{a'}{a}$$

Then we can suppose that all of u_i are distinct monic irreducible polynomials.

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

t is a transcendental logarithm

Firstly let us consider the basic properties of logarithmical derivatives:

$$\frac{(ab)'}{ab} = \frac{a'}{a} + \frac{b'}{b}, \quad \frac{(1/a)'}{1/a} = -\frac{a'}{a}$$

Then we can suppose that all of u_i are distinct monic irreducible polynomials.

Now suppose that some $u_i \notin F$. It's clear that $\frac{u_i'}{u_i}$ is already in lowest terms (because $\deg u_i > \deg u_i'$).

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

t is a transcendental logarithm

Firstly let us consider the basic properties of logarithmical derivatives:

$$\frac{(ab)'}{ab} = \frac{a'}{a} + \frac{b'}{b}, \quad \frac{(1/a)'}{1/a} = -\frac{a'}{a}$$

Then we can suppose that all of u_i are distinct monic irreducible polynomials.

Now suppose that some $u_i \notin F$. It's clear that $\frac{u_i'}{u_i}$ is already in lowest terms (because $\deg u_i > \deg u_i'$).

- If there's not u_i in the denominator of v , then there's not u_i in the denominator of v' . But then $\alpha \notin F$.

t is a transcendental logarithm

Firstly let us consider the basic properties of logarithmical derivatives:

$$\frac{(ab)'}{ab} = \frac{a'}{a} + \frac{b'}{b}, \quad \frac{(1/a)'}{1/a} = -\frac{a'}{a}$$

Then we can suppose that all of u_i are distinct monic irreducible polynomials.

Now suppose that some $u_i \notin F$. It's clear that $\frac{u_i'}{u_i}$ is already in lowest terms (because $\deg u_i > \deg u_i'$).

- ▶ If there's not u_i in the denominator of v , then there's not u_i in the denominator of v' . But then $\alpha \notin F$.
- ▶ If there's u_i in the denominator of v , then the denominator of v' is divisible by u_i^2 and it still cannot be reduced in the general sum and $\alpha \notin F$.

t is a transcendental logarithm

Firstly let us consider the basic properties of logarithmical derivatives:

$$\frac{(ab)'}{ab} = \frac{a'}{a} + \frac{b'}{b}, \quad \frac{(1/a)'}{1/a} = -\frac{a'}{a}$$

Then we can suppose that all of u_i are distinct monic irreducible polynomials.

Now suppose that some $u_i \notin F$. It's clear that $\frac{u_i'}{u_i}$ is already in lowest terms (because $\deg u_i > \deg u_i'$).

- ▶ If there's not u_i in the denominator of v , then there's not u_i in the denominator of v' . But then $\alpha \notin F$.
- ▶ If there's u_i in the denominator of v , then the denominator of v' is divisible by u_i^2 and it still cannot be reduced in the general sum and $\alpha \notin F$.

Therefore $u_1, \dots, u_m \in F$ and $v' \in F$

t is a transcendental logarithm

Firstly let us consider the basic properties of logarithmical derivatives:

$$\frac{(ab)'}{ab} = \frac{a'}{a} + \frac{b'}{b}, \quad \frac{(1/a)'}{1/a} = -\frac{a'}{a}$$

Then we can suppose that all of u_i are distinct monic irreducible polynomials.

Now suppose that some $u_i \notin F$. It's clear that $\frac{u_i'}{u_i}$ is already in lowest terms (because $\deg u_i > \deg u_i'$).

- ▶ If there's not u_i in the denominator of v , then there's not u_i in the denominator of v' . But then $\alpha \notin F$.
- ▶ If there's u_i in the denominator of v , then the denominator of v' is divisible by u_i^2 and it still cannot be reduced in the general sum and $\alpha \notin F$.

Therefore $u_1, \dots, u_m \in F$ and $v' \in F$

Then

$$v' = (ct + s)' = ct' + s' = c \frac{z'}{z} + s'$$

$$\alpha = \sum c_i \frac{u_i'}{u_i} + c \frac{z'}{z} + s'$$

QED.

t is a transcendental exponent

Let u_1, \dots, u_m be distinct monic irreducible again.

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

t is a transcendental exponent

Let u_1, \dots, u_m be distinct monic irreducible again.

It's clear that $u_i = t$ is only u that can be not in F because t is an only irreducible monomial, and we would get the same contradiction otherwise.

Liouville's Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is expression in elementary terms

Liouville's Theorem (statement)

The Main Lemma

Liouville's Theorem (proof)

t is a transcendental logarithm

t is a transcendental exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

t is a transcendental exponent

Let u_1, \dots, u_m be distinct monic irreducible again.

It's clear that $u_i = t$ is only u that can be not in F because t is an only irreducible monomial, and we would get the same contradiction otherwise.

By the main lemma $v' \in F$ iff $v \in F$.

Liouville's Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is expression in elementary terms

Liouville's Theorem (statement)

The Main Lemma

Liouville's Theorem (proof)

t is a transcendental logarithm

t is a transcendental exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

t is a transcendental exponent

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Let u_1, \dots, u_m be distinct monic irreducible again.

It's clear that $u_i = t$ is only u that can be not in F because t is an only irreducible monomial, and we would get the same contradiction otherwise.

By the main lemma $v' \in F$ iff $v \in F$.

Then

$$\alpha = c_1 \frac{t'}{t} + \sum_{i=2}^m \frac{u_i'}{u_i} + v' = \sum_{i=2}^m \frac{u_i'}{u_i} + (v + c_1 z)'$$

t is algebraic

Consider all congruent elements of t :

$$\tau_1 = t, \tau_2, \dots, \tau_k$$

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

t is algebraic

Consider all congruent elements of t :

$$\tau_1 = t, \tau_2, \dots, \tau_k$$

Then it's clear that for all j :

$$\alpha = \sum_i \frac{u_i'(\tau_j)}{u_i(\tau_j)} + v'(\tau_j)$$

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

t is algebraic

Consider all congruent elements of t :

$$\tau_1 = t, \tau_2, \dots, \tau_k$$

Then it's clear that for all j :

$$\alpha = \sum_i \frac{u_i'(\tau_j)}{u_i(\tau_j)} + v'(\tau_j)$$

$$\alpha = \frac{1}{k} \left(\sum_i \sum_j \frac{u_i'(\tau_j)}{u_i(\tau_j)} + \sum_j v'(\tau_j) \right)$$

t is algebraic

Consider all congruent elements of t :

$$\tau_1 = t, \tau_2, \dots, \tau_k$$

Then it's clear that for all j :

$$\alpha = \sum_i \frac{u_i'(\tau_j)}{u_i(\tau_j)} + v'(\tau_j)$$

$$\alpha = \frac{1}{k} \left(\sum_i \sum_j \frac{u_i'(\tau_j)}{u_i(\tau_j)} + \sum_j v'(\tau_j) \right)$$

$$\alpha = \frac{1}{k} \left(\sum_i \frac{\left(\prod_j u_i(\tau_j) \right)'}{\prod_j u_i(\tau_j)} + \left(\sum_j v(\tau_j) \right)' \right)$$

Since all rational functions of polynomial's roots belongs to the basic field, proof is completed.

t is algebraic

Consider all congruent elements of t:

$$\tau_1 = t, \tau_2, \dots, \tau_k$$

Then it's clear that for all j:

$$\alpha = \sum_i \frac{u_i'(\tau_j)}{u_i(\tau_j)} + v'(\tau_j)$$

$$\alpha = \frac{1}{k} \left(\sum_i \sum_j \frac{u_i'(\tau_j)}{u_i(\tau_j)} + \sum_j v'(\tau_j) \right)$$

$$\alpha = \frac{1}{k} \left(\sum_i \frac{\left(\prod_j u_i(\tau_j) \right)'}{\prod_j u_i(\tau_j)} + \left(\sum_j v(\tau_j) \right)' \right)$$

Since all rational functions of polynomial's roots belongs to the basic field, proof is completed.

Remark

Here we used that algebraic extension can be equipped with the unique differentiation, but it's trivial for the case of meromorphic functions.

The main corollary

Corollary

Let $f, g \in \mathbb{C}(x)$, $f \neq 0$ and g is not constant. If the integral of $f(x)\exp(g(x))$ can be expressed in elementary terms, then there's $r \in \mathbb{C}(x)$ such that $f = r' + rg'$.

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Liouville's Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's Theorem (statement)

The Main Lemma

Liouville's Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

Some special cases

On elliptic integrals

Bibliography



Liouville's Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's Theorem (statement)

The Main Lemma

Liouville's Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

Some special cases

On elliptic integrals

Bibliography



The main corollary

Corollary

Let $f, g \in \mathbb{C}(x)$, $f \neq 0$ and g is not constant. If the integral of $f(x)\exp(g(x))$ can be expressed in elementary terms, then there's $r \in \mathbb{C}(x)$ such that $f = r' + rg'$.

Proof.

Denoting $F = \mathbb{C}(x)$, $t = \exp(g)$.

Using Liouville's theorem

$$ft = \sum c_i \frac{u_i'}{u_i} + v'$$

One can use the main lemma and get:

$$c_i \frac{u_i'}{u_i} \in F, \quad v = \sum_{i \in \mathbb{Z}} b_i t^i$$

The rest of the proof is trivial. □

Corollary

Functions

$$\int e^{\pm x^2} dx, \quad \int \frac{dx}{\ln x}, \quad \int \frac{e^x}{x} dx, \quad \int \log \log x dx$$

are not elementary.

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Corollary

Functions

$$\int e^{\pm x^2} dx, \quad \int \frac{dx}{\ln x}, \quad \int \frac{e^x}{x} dx, \quad \int \log \log x dx$$

are not elementary.

Proof.

All of these cases can be simply proven with the main corollary and some basic asymptotical investigations. \square

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Corollary

Functions

$$\int e^{\pm x^2} dx, \quad \int \frac{dx}{\ln x}, \quad \int \frac{e^x}{x} dx, \quad \int \log \log x dx$$

are not elementary.

Proof.

All of these cases can be simply proven with the main corollary and some basic asymptotical investigations. \square

Corollary

Functions

$$\int \frac{\sin x}{x} dx, \quad \int \frac{\sinh x}{x} dx$$

are not elementary.

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Corollary

Functions

$$\int e^{\pm x^2} dx, \quad \int \frac{dx}{\ln x}, \quad \int \frac{e^x}{x} dx, \quad \int \log \log x dx$$

are not elementary.

Proof.

All of these cases can be simply proven with the main corollary and some basic asymptotical investigations. \square

Corollary

Functions

$$\int \frac{\sin x}{x} dx, \quad \int \frac{\sinh x}{x} dx$$

are not elementary.

Proof.

This case is slightly more complicated.

Denoting $F = \mathbb{C}(x)$, $t = \exp(x)$ and using Liouville's theorem

$$\frac{t^2 - 1}{tz} = \sum c_i \frac{u_i'}{u_i} + v'$$

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

Corollary

Functions

$$\int e^{\pm x^2} dx, \quad \int \frac{dx}{\ln x}, \quad \int \frac{e^x}{x} dx, \quad \int \log \log x dx$$

are not elementary.

Proof.

All of these cases can be simply proven with the main corollary and some basic asymptotical investigations. \square

Corollary

Functions

$$\int \frac{\sin x}{x} dx, \quad \int \frac{\sinh x}{x} dx$$

are not elementary.

Proof.

This case is slightly more complicated.

Denoting $F = \mathbb{C}(x)$, $t = \exp(x)$ and using Liouville's theorem

$$\frac{t^2 - 1}{tz} = \sum c_i \frac{u_i'}{u_i} + v'$$

After some considirations using the main lemma we will deduce an impossible equation

$$\frac{1}{x} = a' + a$$

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

On elliptic integrals

Corollary

Let P be a monic polynomial with degree ≥ 3 and no repeated roots. Then $\int \frac{dx}{\sqrt{P(x)}}$ is not elementary.

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography

On elliptic integrals

Corollary

Let P be a monic polynomial with degree ≥ 3 and no repeated roots. Then $\int \frac{dx}{\sqrt{P(x)}}$ is not elementary.

Proof.

Here we will use a bit more advanced stuff on Riemann surfaces.
Liouville's theorem enable us to write

$$\frac{1}{\sqrt{P(x)}} = \sum c_i \frac{u_i'}{u_i} + v'$$

with $c_1, \dots, c_m \in \mathbb{C}$ and $u_1, \dots, u_m, v \in \mathbb{C}(x, \sqrt{P(x)})$.



On elliptic integrals

Corollary

Let P be a monic polynomial with degree ≥ 3 and no repeated roots. Then $\int \frac{dx}{\sqrt{P(x)}}$ is not elementary.

Proof.

Here we will use a bit more advanced stuff on Riemann surfaces.
Liouville's theorem enable us to write

$$\frac{1}{\sqrt{P(x)}} = \sum c_i \frac{u_i'}{u_i} + v'$$

with $c_1, \dots, c_m \in \mathbb{C}$ and $u_1, \dots, u_m, v \in \mathbb{C}(x, \sqrt{P(x)})$.

Now we deduce an equation on the compact Riemann surface C associated to the equation $y^2 = P(x)$

$$\frac{dx}{y} = \sum c_i \frac{du_i}{u_i} + dv$$



Liouville's Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's Theorem (statement)

The Main Lemma

Liouville's Theorem (proof)

t is a transcendental
logarithm

t is a transcendental exponent

t is algebraic

Corollaries

The main corollary

Some special cases

Bibliography

☐

Bibliography



Joseph Liouville, Sur la détermination des intégrales dont la valeur est algébrique, Journal de l'École Polytechnique, vol. 14, 1833, pp. 93–123.



Maxwell Rosenlicht, Integration in finite terms, American Mathematical Monthly, vol. 79, no. 9, 1972, pp. 963–972.



Keith Conrad, Impossibility theorems for elementary integration, American Mathematical Monthly, vol. 110, no. 5, 2003, pp. 459–462.

Liouville's
Theorem

Vanya Vorobiov

Introduction

Basic definitions

What is
expression in
elementary terms

Liouville's
Theorem
(statement)

The Main
Lemma

Liouville's
Theorem (proof)

t is a transcendental
logarithm

t is a transcendental
exponent

t is algebraic

Corollaries

The main corollary

Some special cases

On elliptic integrals

Bibliography