

Probability and Statistics. Week 3

1. Distribution of random variable ξ is given by $\xi \sim \begin{pmatrix} -2 & -1 & 0 & 1 & 2 \\ 0.1 & 0.3 & 0.2 & p & 0.2 \end{pmatrix}$. Determine the probability p . Draw the graph of a cumulative distribution function $F_\xi(t)$.

Answer: $p = 0.2$.

2. Let us consider the circle of radius R centered at O . Point M is chosen at random inside this circle. Random variable ξ is equal to the length of OM . Find the cumulative distribution and probability density functions.

Answer:
$$F_\xi(r) = \begin{cases} 0, & r < 0 \\ \frac{r^2}{R^2}, & 0 \leq r < R \\ 1, & r \geq R \end{cases}, \quad f_\xi(r) = \begin{cases} \frac{2r}{R^2}, & 0 \leq r < R \\ 0, & \text{otherwise} \end{cases}.$$

3. Let us consider the ball of radius R centered at O . Point M is chosen at random inside this ball. Random variable ξ is equal to the distance from point M to the sphere. Find the cumulative distribution and probability density functions.

Answer:
$$F_\xi(x) = \begin{cases} 0, & x < 0 \\ 1 - \frac{(R-x)^3}{R^3}, & 0 \leq x < R \\ 1, & x \geq R \end{cases}; \quad f_\xi(x) = \begin{cases} 3\frac{(R-x)^2}{R^3}, & 0 < x < R \\ 0, & \text{otherwise} \end{cases}$$

4. Find all values of C such that function $F(x) = \begin{cases} 0, & x < 1 \\ 1 - \frac{C}{x}, & x \geq 1 \end{cases}$ can be a cumulative distribution function for some random variable ζ . Find the probability density function.

Answer: $C = 1, f(x) = \begin{cases} \frac{1}{x^2}, & x > 1 \\ 0, & \text{otherwise} \end{cases}$

5. Is it possible that for some values of C the functions below are probability density functions of random variables? If it is so, find the values of C .

(a) $f(x) = \begin{cases} Ce^{-2x}, & x > 0 \\ 0, & x \leq 0 \end{cases}$

(b) $f(x) = Ce^{-|x|}, x \in \mathbb{R}$.

Answer: (a) $C = 2$ (b) $C = 0.5$.

6. Find such value of C that function $f(x) = \frac{C}{1+x^2}$ is probability density function of a random variable.

Answer: $C = \frac{1}{\pi}$

7. (Walpole 3.5) Determine the value c so that each of the following functions can serve as a probability distribution of the discrete random variable X .

(a) $f(x) = c(x^2 + 4), x \in \{0, 1, 2, 3\};$

(b) $f(x) = c\binom{2}{x}\binom{3}{3-x}, x \in \{0, 1, 2\}.$

Answer: (a) $\frac{1}{30}$, (b) $\frac{1}{10}$.

8. (Walpole 3.9) The proportion of people who respond to a certain mail-order solicitation is a continuous random variable X that has the density function $f(x) = \begin{cases} \frac{2(x+2)}{5}, & 0 < x < 1 \\ 0, & \text{elsewhere} \end{cases}$.

- (a) Show that $P(0 < X < 1) = 1$;
(b) Find the probability that more than $\frac{1}{4}$ but fewer than $\frac{1}{2}$ of the people contacted will respond to this type of solicitation.

Answer: (b) $\frac{19}{80}$.

9. (Walpole 3.35) Suppose it is known from large amounts of historical data that X , the number of cars that arrive at a specific intersection during a 20-second time period, is characterized by the following discrete probability function: $f(x) = e^{-6} \frac{6^x}{x!}$, for $x = 0, 1, 2, \dots$.

- (a) Find the probability that in a specific 20-second time period, more than 8 cars arrive at the intersection.
(b) Find the probability that only 2 cars arrive.

Hint: $e^6 \approx 400$.

Answer: (a) 0.1528 (b) 0.0446

10. (Walpole 3.42) Let X and Y denote the lengths of life, in years, of two components in an electronic system. If the joint density function of these variables is $f(x, y) = \begin{cases} e^{-(x+y)}, & x > 0, y > 0 \\ 0, & \text{elsewhere} \end{cases}$

find $P(0 < X < 1 | Y = 2)$.

Answer: $1 - \frac{1}{e} \approx 0.6321$

11. (Walpole 3.68) Consider the following joint probability density function of the random variables X and Y :

$$f(x, y) = \begin{cases} \frac{3x-y}{9}, & 1 < x < 3, 1 < y < 2 \\ 0, & \text{elsewhere} \end{cases}$$

- (a) Find the marginal density functions of X and Y ;
(b) Are X and Y independent?
(c) Find $P(X > 2)$.

Answer: (a) $g(x) = \frac{x}{3} - \frac{1}{6}$ for $1 < x < 3$ and $h(y) = \frac{4}{3} - \frac{2}{9}y$; (b) No; (c) $\frac{2}{3}$.

12. Consider a system of components in which there are 5 independent components, each of which possesses an operational probability of 0.92. The system does have a redundancy built in such that it does not fail if 3 out of the 5 components are operational.

- (a) Write down the probability mass function;
(b) What is the probability that the total system is operational?

Answer: (a) $f(x) = \begin{cases} \binom{5}{x} \cdot 0.92^x \cdot (1 - 0.92)^{5-x}, & x \in \{0, 1, 2, 3, 4, 5\} \\ 0, & \text{otherwise} \end{cases}$; (b) 0.9955