

AstroBytes, Data Analytics Report, NSSC 2021

Rohan Kumar^{1,*} and Ankush Kumar Singh^{2,†}

¹*IISER, Kolkata*

²*IIT(ISM), Dhanbad*

^{*}*Corresponding author: Rohan Kumar,
rohankumarprasad@yahoo.com*

[†]*Team Leader: Ankush Kumar Singh, 20je0155@fme.iitism.ac.in*

December 13, 2021

Problem 1

To begin the analysis we will first import the standard library *Lightcurve*.

```
1 import lightcurve as lk
```

We can now download the lightcurve data for *Kepler-17*¹ and plot it.

```
1 lc_KIC=lk.search_lightcurve("KIC 10619192")[38].download()  
2 lc_KIC.plot()
```




Figure 1: Kepler-17 Lightcurve

¹TIC 273874849, KIC 10619192

To make the lightcurve usable we will remove the outliers and NAN-values and plot the clean lightcurve.

```
1 lc_KIC=lc_KIC.remove_outliers()
2 lc_KIC=lc_KIC.remove_nans()
3 lc_KIC.plot()
```




Figure 2: Clean lightcurve

To remove the long term trends we will flatten² out the lightcurve, doing this will in a sense magnify the local trends in which we are interested in for exo-planets study.

```
1 flat_lc=lc_KIC.flatten()
2 flat_lc.plot()
```




Figure 3: Flat Lightcurve

²normalize

We can now convert the `flat_lc` to an astropy table and store the flux values in new variable and calculate the mean and standard deviation using *NumPy*. For CDPP³ we will use the `lightcurve.estimate_cdpp()`.

```
1 flux=flat_lc.to_table()['flux']
2 import numpy as np
3 std=np.std(flux)
4 mean=np.mean(flux)
5 cdpp=flat_lc.estimate_cdpp()
```

$$\sigma = 0.0045 \text{ } es^{-1}$$

$$\text{mean flux} = \bar{x} = 0.9989 \text{ } es^{-1}$$

$$\text{CDPP} = 1271.0180 \text{ ppm}$$

Problem 2

We will now calculate and plot the BLS⁴ periodogram of the `flat_lc`,

```
1 bls=flat_lc.to_periodogram(method='bls')
2 bls.plot()
```




Figure 4: BLS Periodogram

We know that BLS Periodogram is a power spectrum which shows the dominant periods in the time series data. So we can use this fact to extract the transit duration and time period of revolution.

```
1 planet_period = bls.period_at_max_power
2 planet_dur = bls.duration_at_max_power
```

$$\text{Transit Duration} = 0.1 \text{ d}$$

$$\text{Time Period} = 1.4857 \text{ d}$$

³Combined Differential Photometric Precision

⁴Box Least Square

To verify whether the time period obtained above is correct or not we will fold the light curve with stated period and plot the folded lightcurve,

```
1 folded_lc = flat_lc.fold(period=1.4856943)
2 folded_lc.plot()
```




Figure 5: Folded Lightcurve with Period=1.4857 d

We can see from the above figure a distinct dip around -0.4 which means we have correctly determined the period of the planet.

We can now make a model curve by using the BLS⁵ and giving it the same time period and transit time to get a model lightcurve.

```
1 planet_t0 = bls.transit_time_at_max_power
2 planet_model = bls.get_transit_model(period=planet_period,
3 transit_time=planet_t0,duration=planet_dur)
4 ax = flat_lc.fold(planet_period, planet_t0).scatter()
5 planet_model.fold(planet_period, planet_t0).plot(ax=ax, c='r', lw=1)
```




Figure 6: Model Lightcurve

⁵Box-fitting Least Squares

To calculate the relative flux dip($\frac{\Delta F}{F}$),

```
1 model_flux=planet_model.fold(planet_period, planet_t0)['flux']
2 min=np.min(model_flux)
3 max=np.max(model_flux)
4 rfd=(max-min)/max
```

$$\text{Relative Flux Dip} = \frac{\Delta F}{F} = 0.015807545862905243 \approx 0.0158$$

Problem 3

To start the analysis of the star *Kepler-21*⁶ in quarter 2,5,6,7 we will first download the short cadence lightcurve data and plot the lightcurve,

```
1 search_result = lk.search_lightcurve("Kepler-21", author="Kepler",
2   cadence="short", quarter=[2,5,6,7])
3 search_result
4 lc_collection=search_result.download_all()
5 for i in range(len(search_result)):
6     lc_collection[i].plot()
```






Figure 7: Quarter 2 Time[BKJD] 231-258


⁶KIC 3632418, TIC 121214185



(a)




(b)




(c)


Figure 8: Quarter 5 Time[BKJD] (a)443-475, (b)476-503, (c)504-538



(a)




(b)




(c)


Figure 9: Quarter 6 Time[BKJD] (a)539-566, (b)567-598, (c)599-629



(a)



(b)



(c)

Figure 10: Quarter 7 Time[BKJD] (a)630-660, (b)661-690, (c)691-719

Now we will stitch the lightcurve of different quarter to get the long term behaviour of the star over all the quarters.

```
1 lc_stitched = lc_collection.stitch()
2 lc_stitched.plot()
```




Figure 11: Stitched lightcurve over quarters 2,5,6,7

Now we will remove the outliers and the NAN-values from the composite lightcurve to make the result more accurate.

```
1 clean_lc=lc_stitched.remove_outliers()
2 clean_lc=lc_stitched.remove_nans()
3 clean_lc.plot()
```




Figure 12: Clean lightcurve

As we are interested in the short term details of this composite lightcurve⁷ we can flatten the lightcurve to remove the long term behaviour of the star and plot the flattened lightcurve.

⁷Here we are interested in the stellar oscillations which happens at very high frequencies compared to the long the term behaviour we can see here.

```

1 lc_stitched_flat=clean_lc.flatten()
2 lc_stitched_flat.plot()

```




Figure 13: Flattened lightcurve

We will now plot the periodogram to see the data in frequency domain.

```

1 pg=lc_stitched_flat.to_periodogram(normalization='psd')
2 pg.plot(scale='log')

```




Figure 14: Periodogram in frequency domain

From the given document⁸ we know that when we draw the periodogram of a solar-type oscillator star, we get a peak around $10^3 \mu\text{Hz}$. Therefore we zoomed into the region within the range of 700-1500 μHz and found a peak around $10^3 \mu\text{Hz}$ as mentioned above.

```
1 pg=lc_stitched_flat.to_periodogram(normalization='psd',
2   minimum_frequency=700,maximum_frequency=1500)
3 pg.plot()
```




Figure 15: Periodogram zoomed near power excess region around $10^3 \mu\text{Hz}$

As the above plot is very noisy thus to make it usable we will smooth it out using `scipy.signal.medfilt()`,

```
1 freq=pg.to_table()['frequency']
2 pw=pg.to_table()['power']
3 from scipy.signal import medfilt
4 smooth=medfilt(pw)
5 fig=plt.figure(figsize=(12.5,7),facecolor='white')
6 plt.plot(freq,smooth,'k')
7 plt.xlabel(r'Frequency [μHz]',fontsize=20)
8 plt.ylabel(r'Power [1/μHz]',fontsize=20)
9 plt.legend(['KIC 3632418'])
10 plt.show()
```

⁸Astroseismology




Figure 16: Smoothed out Periodogram

For the *Gaussian* envelope we have taken the distinct peaks with power greater than $4 \times 10^{-11} \text{Hz}^{-1}$ were selected and stored in an array which was then used for the *Gaussian* fitting to get the envelope, for the fitting process we used `lmfit.Model()`.

```

1 x=[]
2 y=[]
3 for i in range(len(freq)):
4     if (smooth[i]>4e-11):
5         x.append(freq[i])
6         y.append(smooth[i])
7
8 from lmfit import Model
9
10 def gaussian(x, amp, cen, wid):
11     "1-d gaussian: gaussian(x, amp, cen, wid)"
12     return (amp/(np.sqrt(2*np.pi)*wid)) * np.exp(-(x-cen)**2 / (2*
13         wid**2))
14
15 gmod = Model(gaussian)
16 result = gmod.fit(y, x=x, amp=9e-11, cen=1150, wid=1000)

```

We then use `print(result.fit_report())` to get the parameters for the *Gaussian* envelope,

```

1
2 [[Model]]
3     Model(gaussian)
4 [[Fit Statistics]]
5     # fitting method      = leastsq
6     # function evals      = 123
7     # data points         = 13
8     # variables            = 3
9     chi-square            = 2.5757e-21
10    reduced chi-square     = 2.5757e-22
11    Akaike info crit       = -643.650708
12    Bayesian info crit     = -641.955860
13 [[Variables]]
14    amp: 2.5869e-08 +/- 1.6467e-08 (63.65%) (init = 9e-11)
15    cen: 1125.47295 +/- 56.1484650 (4.99%) (init = 1150)
16    wid: 168.600608 +/- 116.586197 (69.15%) (init = 1000)
17 [[Correlations]] (unreported correlations are < 0.100)
18    C(amp, wid) = 0.993
19    C(amp, cen) = 0.680
20    C(cen, wid) = 0.674

```

We can now use the calculated parameters of the *Gaussian* envelope to plot it.

```

1 gauss=gaussian(freq,2.5869e-08,1125.47295,168.600608)
2 fig =plt.figure(figsize=(12.5,7),facecolor='white')
3 plt.plot(freq,smooth,'k')
4 plt.plot(freq,gauss,'r')
5 plt.xlabel(r'Frequency [ $\mu$ Hz]',fontsize=20)
6 plt.ylabel(r'Power [ $\frac{1}{\mu$ Hz]$',fontsize=20)
7 plt.legend(['KIC 3632418','Gaussian Fit'])
8 plt.show()

```




Figure 17: Periodogram with *Gaussian* envelope

From the *Gaussian* envelope parameter we can get the ν_{\max} and for the $\Delta\nu$ we can use the following code.

```
1 seismology = pg.flatten().to_seismology() #using seismology
   function of Lightkurve library
2 seismology.estimate_numax(method='acf2d')
3 seismology.estimate_deltanu()
```

$$\nu_{\max} = 1125.2730 \mu Hz$$

$$\Delta\nu = 60.72 \mu Hz$$

Now we will use above calculated values to get the radius(R) and mass(M) of *Kepler-21* using the following expressions,

$$\frac{M}{M_{\odot}} \simeq \left(\frac{\nu_{\max}}{\nu_{\max, \odot}} \right)^3 \left(\frac{\Delta\nu}{\Delta\nu_{\odot}} \right)^{-4} \left(\frac{T_{\text{eff}}}{T_{\text{eff}, \odot}} \right)^{1.5}$$

$$\frac{R}{R_{\odot}} \simeq \left(\frac{\nu_{\max}}{\nu_{\max, \odot}} \right) \left(\frac{\Delta\nu}{\Delta\nu_{\odot}} \right)^{-2} \left(\frac{T_{\text{eff}}}{T_{\text{eff}, \odot}} \right)^{0.5}$$

where $\nu_{\max, \odot} = 3090 \mu Hz$, $\Delta\nu_{\max, \odot} = 135.1 \mu Hz$, $T_{\text{eff}, \odot} = 5777.2 K$, temperature of the star $T_{\text{eff}} = 6200 K$,

```
1 M = ((1125.273/3090)**3)*((60.72/135.1)**(-4))*((6200/5777.2)
   **(3/2))
2 R = (1125.273/3090)*((60.72/135.1)**(-2))*((6200/5777.2)**(1/2))
```

$$R = 1.8676 R_{\odot}$$

$$M = 1.3158 M_{\odot}$$

Problem 4

We will first import the standard libraries,

```
1 import numpy as np
2 import astropy.constants as const
```

To calculate the radius of *Kepler-17 b* we will use,

$$\frac{\Delta F}{F} = \left(\frac{R_p}{R_s} \right)^2 \quad (1)$$

We are given the value of $R_s = 1.01R_\odot$ and we already calculated the value of $\frac{\Delta F}{F}$ here, we can now calculate R_p which is the radius of *Kepler-17 b*.⁹

```
1 R_sun = const.R_sun
2 R_s = 1.01*R_sun    #converting radius of star given in solar radius
                      to metre
3 delF_F = 0.015807545862905243
4 R_p = (R_s)*np.sqrt(delF_F)
```

$$R_p = 1.2357R_J = 13.8511R_\oplus = 88343703.6673 \text{ m}$$

Since the time period of the planet and mass of star is known, we can use Kepler's third law to compute the orbital radius of the planet.

$$T^2 = \frac{4\pi^2 a^3}{G(M_s + M_p)}$$

We can make the following assumptions $a = R_{\text{orbit}}$ ¹⁰ and $M_s + M_p \approx M_s$ ¹¹ we already have the value of $M_s = 1.04M_\odot$, these assumptions simplifies the above equation.

$$T^2 = \frac{4\pi^2 R_{\text{orbit}}^3}{G(M_s)}$$

```
1 M_s = 1.04*const.M_sun    #converting mass of star given in
                              solar mass to kg
2 T = 1.4856943*24*3600
3 R_cube = (const.G*M_s*(T**2))/(4*(np.pi**2))
4 R_orbit = R_cube**(1/3)
```

$$R_{\text{orbit}} = 0.0258 \text{ AU} = 3862103236.9458 \text{ m}$$

⁹Here \odot , \oplus and J stands for solar, earth and jupiter parameters respectively.

¹⁰It's given in the problem statement to assume that the orbit is circular.

¹¹Here we are talking about a star planet binary system which means that $M_s \gg M_p$.

Mass of *Kepler-17 b* can be found using conservation of angular momentum.

$$M_s V_s = M_p V_p$$

We already have the values of $M_s = 1.04 M_\odot$ and $V_s = 0.228$ km/s, V_p can be calculated by,

$$V_p = \frac{D}{T}$$

where D is the circumference of the orbit $2\pi R_{orbit}$ and T is the time period,

$$V_p = \frac{2\pi R_{orbit}}{T}$$

This means the expression for M_p becomes.

$$M_p = \frac{M_s V_s T}{2\pi R_{orbit}}$$

```

1 V_s = 0.228*1000 #conversion to m/s
2 M_p = (M_s*V_s*T)/(2*np.pi*R_orbit)
3 vol_planet = (4*np.pi*(R_p**3))/3
4 D = M_p/vol_planet

```

$$M_p = 1.3137 M_J = 417.6202 M_\oplus = 2.4941 \times 10^{27} kg$$

$$\rho = 863.5712 \text{ kg } m^{-3}$$

Now we will do the classification of *Kepler-17 b* according to the above calculated parameters.




Figure 18: Classification of Exo-Planets¹²

Coordinate of *Kepler-17 b* in this plot will be (M_p, R_p) and this comes out to be $(417.6202M_{\oplus}, 13.8511R_{\oplus})$, we can clearly see that it is in the *Jovian worlds* region. This means that *Kepler-17 b* is a gas giant. We can also say that it's a *Jupiter* like planet because it's mass is $1.3137M_J$.

Now to comment whether the planet is *Hot Jupiter* or *Cold Jupiter* we will use the surface temperature of the host star *Kepler-17* which is $5781 \pm 85K$ ¹³ which is very close to the surface temperature of the *Sun*. We also know that $R_{orbit} = 0.0258AU$ and for a *Sun* like star *Continued Habitable Zone* is around $0.95AU - 1.15AU$ and we can see that R_{orbit} is much less than $1AU$, this means that the planet is very close to the host star which implies that *Kepler-17 b* is a ***Hot Jupiter***.

¹²Plot used is taken from <https://arxiv.org/abs/1603.08614>

¹³This data is taken from <https://arxiv.org/abs/1110.5462>

Error Analysis

Kepler-17 b			
Parameter	Standard data	Calculated data	Error(percentage)
Orbital Period[d]	1.4857	1.4857	0.00
Transit Duration[h]	2.276	2.4	5.45
Orbital Semi-Major Axis[AU]	0.0268	0.0258	3.73
Radius [R_J]	1.33	1.2357	7.09
Mass [M_J]	2.47	1.3137	46.81
Density [g cm^{-3}]	1.3	0.8636	33.57
Kepler-21			
Mass [M_\odot]	1.408	1.3158	6.55
Radius [R_\odot]	1.902	1.8676	1.81

In the above table for the standard data column we have used the values from [here](#) for *Kepler-17 b* and from [here](#) for *Kepler-21*.

We can observe large error in the measurment of mass of *Kepler-17 b*, hence a large difference in its density, from the standard data. This *could* be due to incorrect measurement of stellar velocity.