

Pilot data analysis

valeriia beliaeva

11/8/2021

```
.libPaths('C:/Users/vbeliaev/Documents/r_packages')

knitr::opts_chunk$set(echo = TRUE)

rm(list = ls())

library(ggsignif)
library(ggplot2)
library(gridExtra)
library(lme4)
library(plyr)
library(dplyr)
library(boot)
library(data.table)

'%!in%' <- function(x,y)!('%in%'(x,y))
```

Data analysis Part II

vmPFC TI neuromodulation

Data analysis is separated into 2 parts:

- 1) we test whether the task is working correctly (Behavioral calibration)
- 2) we test the effect of TI stimulation on choice consistency / accuracy (vmPFC TI neuromodulation)

In this file the second part of the data analysis is presented.

Main dependent variable is choice consistency / accuracy

Correct answers (variable corr) correspond to trials, in which participant has chosen a picture, which was earlier rated higher during the rating task. For example, participant during the rating task estimated taste of banana for 0.8 and Twix for 0.9. Then in the choice task, when twix and banana are presented in the same trial, if participant chooses Twix, answer is correct, if they choose banana the answer is considered incorrect.

TI stimulation was delivered during the 1st part (112 trials) of the choice task (immediate effects) and switched off during the second task (after effects, last 64 trials).

While the vmPFC that was stimulated with the TI is more involved in value-based decisions like taste, we expect to find effect of stimulation on taste trials and not on size trials in the choice task.

For the model: dependent variable - vector, which contains correct (1) or incorrect (0) trials.

Regressors: (1) active stimulation = 1, sham stimulation = 0;

(2) difference in size or taste between two food items (the smaller the difference - the harder is the choice)

First, we will compare results across sessions, e.g., comparing results of 112 trials from active session and sham session. Then, we will compare results within session: 112 trials with last 64 trials from active session.

```
# loading preprocessed data

setwd('C:/Users/vbeliaev/Documents/TI_fMRI/pre_registration_paper/methods/pilot_scripts_paper')
tab1 = as.data.table(read.csv("Data_collected_together_preprocessed.csv"))

dataVal = tab1[Cue_Taste1_Size2 == 1]
dataVal$Taste_diff.abs = scale(abs(dataVal$Taste_diff))
dataVal$Size_diff.abs = scale(abs(dataVal$Size_diff))

dataPer = tab1[Cue_Taste1_Size2 == 2]
dataPer$Taste_diff.abs = scale(abs(dataPer$Taste_diff))
dataPer$Size_diff.abs = scale(abs(dataPer$Size_diff))

ns = length(unique(tab1$Participant))
```

RUN MODELS

Between session comparison (active vs sham)

Immediate effects of TI on taste trials

There is no significant effect on correct taste trials during the 1st part of the choice task, when the TI is on.

```
idx = which(dataVal$Trial_Nr<113)           #Table: first 112 trials, when TI is on
dataVal2 = dataVal[idx,]
dataVal2$Trial_Nr.z = scale(dataVal2$Trial_Nr)

m1.val.stim1.abs = glmer(corr ~ Stim*Taste_diff.abs + Trial_Nr.z + (1+Stim*Taste_diff.abs + Trial_Nr.z | Participant), dataVal2)

s1 = summary(m1.val.stim1.abs)
s1
```

```
## Generalized linear mixed model fit by maximum likelihood (Laplace
##   Approximation) [glmerMod]
##   Family: binomial ( logit )
##   Formula:
##   corr ~ Stim * Taste_diff.abs + Trial_Nr.z + (1 + Stim * Taste_diff.abs +
##     Trial_Nr.z | Participant)
##   Data: dataVal2
##
##           AIC          BIC    logLik deviance df.resid
##    2775.2    2892.2  -1367.6   2735.2     2552
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
## -4.9943 -0.9722  0.4104  0.6414  1.3351
##
## Random effects:
##   Groups      Name                Variance Std.Dev. Corr
##   Participant (Intercept)      0.2510122 0.50101
##                        Stim      0.0544141 0.23327  0.23
##                        Taste_diff.abs 0.1083354 0.32914  0.98 0.06
##                        Trial_Nr.z     0.0155520 0.12471 -0.03 0.65 -0.21
```

```
## Stim:Taste_diff.abs 0.0003053 0.01747 -0.88 -0.12 -0.92 0.42
## Number of obs: 2572, groups: Participant, 23
##
## Fixed effects:
## Estimate Std. Error z value Pr(>|z|)
## (Intercept) 1.20923 0.11754 10.288 < 2e-16 ***
## Stim -0.08338 0.07231 -1.153 0.2489
## Taste_diff.abs 0.67614 0.08713 7.760 8.47e-15 ***
## Trial_Nr.z -0.09096 0.05519 -1.648 0.0993 .
## Stim:Taste_diff.abs -0.08031 0.05361 -1.498 0.1341
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
## (Intr) Stim Tst_d. Trl_N.
## Stim 0.115
## Tast_dff.bs 0.796 -0.001
## Trial_Nr.z -0.027 0.217 -0.112
## Stm:Tst_df. -0.083 0.276 -0.086 0.040
## convergence code: 0
## Model failed to converge with max|grad| = 0.00994695 (tol = 0.002, component 1)
```

After effects of TI on taste trials

After TI stimulation amount of correct choices in taste trials increases.

```
idx = which(dataVal$Trial_Nr>=113) #Table: last 64 trials, when TI is off
dataVal2 = dataVal[idx,]
dataVal2$Trial_Nr.z = scale(dataVal2$Trial_Nr)

m1.val.stim2.abs = glmer(corr ~ Stim*Taste_diff.abs + Trial_Nr.z + (1+Stim*Taste_diff.abs + Trial_Nr.z
s2 = summary(m1.val.stim2.abs)
s2

## Generalized linear mixed model fit by maximum likelihood (Laplace
## Approximation) [glmerMod]
## Family: binomial ( logit )
## Formula:
## corr ~ Stim * Taste_diff.abs + Trial_Nr.z + (1 + Stim * Taste_diff.abs +
## Trial_Nr.z | Participant)
## Data: dataVal2
##
## AIC BIC logLik deviance df.resid
## 1673.0 1778.9 -816.5 1633.0 1452
##
## Scaled residuals:
## Min 1Q Median 3Q Max
## -4.1634 -0.9884 0.4595 0.6675 1.2066
##
## Random effects:
## Groups Name Variance Std.Dev. Corr
## Participant (Intercept) 0.182077 0.42670
## Stim 0.001734 0.04165 0.52
## Taste_diff.abs 0.098177 0.31333 0.90 0.26
```

```
##          Trial_Nr.z          0.013927 0.11801  0.20 0.83 0.16
##          Stim:Taste_diff.abs 0.063596 0.25218  0.59 0.98 0.41 0.87
## Number of obs: 1472, groups: Participant, 23
##
## Fixed effects:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)    1.09854    0.11274   9.744 < 2e-16 ***
## Stim           0.15889    0.06951   2.286  0.0223 *
## Taste_diff.abs  0.64196    0.09538   6.731 1.69e-11 ***
## Trial_Nr.z      -0.13584    0.06784  -2.002  0.0453 *
## Stim:Taste_diff.abs 0.06113    0.08694   0.703  0.4819
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##          (Intr) Stim   Tst_d. Trl_N.
## Stim      0.137
## Tast_dff.bs 0.648  0.112
## Trial_Nr.z  0.024  0.020 -0.007
## Stm:Tst_df. 0.342  0.356  0.237  0.196
## convergence code: 0
## Model failed to converge with max|grad| = 0.00941021 (tol = 0.002, component 1)
```

Immediate effects of TI on size trials

No significant effect of TI on size trials.

```
idx = which(dataPer$Trial_Nr<113) #Table: first 112 trials, when TI is on
dataPer2 = dataPer[idx,]
dataPer2$Trial_Nr.z = scale(dataPer2$Trial_Nr)
```

```
m1.per.stim1.abs = glmer(corr ~ Stim*Size_diff.abs + Trial_Nr.z + (1+Stim*Size_diff.abs + Trial_Nr.z |P
```

```
s3 = summary(m1.per.stim1.abs)
s3
```

```
## Generalized linear mixed model fit by maximum likelihood (Laplace
## Approximation) [glmerMod]
## Family: binomial ( logit )
## Formula:
## corr ~ Stim * Size_diff.abs + Trial_Nr.z + (1 + Stim * Size_diff.abs +
## Trial_Nr.z | Participant)
## Data: dataPer2
##
##      AIC      BIC    logLik deviance df.resid
##    2790    2907     -1375     2750     2548
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
## -4.5751 -0.9669  0.4178  0.6501  1.2204
##
## Random effects:
## Groups      Name                Variance Std.Dev. Corr
## Participant (Intercept)  0.210375 0.45867
## Stim           0.017050 0.13058  0.03
```

```
##           Size_diff.abs      0.052520 0.22917   0.96  0.30
##           Trial_Nr.z         0.006926 0.08322   0.68  0.45  0.77
##           Stim:Size_diff.abs 0.004371 0.06612  -0.96  0.10 -0.89 -0.44
## Number of obs: 2568, groups: Participant, 23
##
## Fixed effects:
##           Estimate Std. Error z value Pr(>|z|)
## (Intercept)      1.16884    0.10952  10.672 <2e-16 ***
## Stim            -0.09844    0.05968  -1.650  0.099 .
## Size_diff.abs     0.72889    0.07171  10.164 <2e-16 ***
## Trial_Nr.z        -0.02546    0.05103  -0.499  0.618
## Stim:Size_diff.abs -0.04706    0.05514  -0.854  0.393
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##           (Intr) Stim   Sz_df. Trl_N.
## Stim      -0.021
## Size_dff.bs 0.703  0.045
## Trial_Nr.z  0.206  0.060  0.168
## Stm:Sz_dff. -0.244  0.347 -0.196 -0.064
## convergence code: 0
## unable to evaluate scaled gradient
## Model failed to converge: degenerate Hessian with 1 negative eigenvalues
```

After effects of TI on size trials

No significant effect of TI on size trials.

```
idx = which(dataPer$Trial_Nr>=113) #Table: last 64 trials, when TI is off
dataPer2 = dataPer[idx,]
dataPer2$Trial_Nr.z = scale(dataPer2$Trial_Nr)
```

```
m1.per.stim2.abs = glmer(corr ~ Stim*Size_diff.abs + Trial_Nr.z + (1+Stim*Size_diff.abs+ Trial_Nr.z |Pa
```

```
s4 = summary(m1.per.stim2.abs)
s4
```

```
## Generalized linear mixed model fit by maximum likelihood (Laplace
## Approximation) [glmerMod]
## Family: binomial ( logit )
## Formula:
## corr ~ Stim * Size_diff.abs + Trial_Nr.z + (1 + Stim * Size_diff.abs +
## Trial_Nr.z | Participant)
## Data: dataPer2
##
##           AIC      BIC   logLik deviance df.resid
##    1612.2    1718.1   -786.1   1572.2     1448
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
## -4.2737 -0.9460  0.4007  0.6366  1.2274
##
## Random effects:
## Groups      Name                Variance Std.Dev. Corr
```

```
## Participant (Intercept)      0.14550  0.3815
## Stim                        0.05053  0.2248  1.00
## Size_diff.abs               0.04742  0.2178  0.87 0.87
## Trial_Nr.z                   0.02331  0.1527  0.27 0.27 0.59
## Stim:Size_diff.abs          0.02816  0.1678  0.48 0.48 0.68 0.95
## Number of obs: 1468, groups: Participant, 23
##
## Fixed effects:
##               Estimate Std. Error z value Pr(>|z|)
## (Intercept)    1.15968    0.10612  10.928 <2e-16 ***
## Stim           0.02772    0.08459   0.328  0.743
## Size_diff.abs  0.77826    0.08383   9.283 <2e-16 ***
## Trial_Nr.z      0.01996    0.07157   0.279  0.780
## Stim:Size_diff.abs -0.02044  0.07837  -0.261  0.794
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##      (Intr) Stim  Sz_df. Trl_N.
## Stim      0.461
## Size_dff.bs 0.571  0.294
## Trial_Nr.z  0.097  0.046 0.136
## Stm:Sz_dff. 0.191  0.406 0.199  0.178
## convergence code: 0
## boundary (singular) fit: see ?isSingular
```

Within session comparison (immediate vs after effects)

Taste trials: immediate and after effect in active stimulation condition

In the session when TI was applied there was no difference in accuracy during the first part of the choice task (immediate effects) and the second part (after effects) for taste trials.

```
idx = which(dataVal$Session_type==1) # Table: active sessions
dataVal2 = dataVal[idx,]

m2.val.stim1.abs = glmer(corr ~ Block*Taste_diff.abs + (1+Block*Taste_diff.abs|Participant), data=dataVal2)

summary(m2.val.stim1.abs)
```

```
## Generalized linear mixed model fit by maximum likelihood (Laplace
## Approximation) [glmerMod]
## Family: binomial ( logit )
## Formula: corr ~ Block * Taste_diff.abs + (1 + Block * Taste_diff.abs |
## Participant)
## Data: dataVal2
##
##      AIC      BIC   logLik deviance df.resid
##  2219.5   2298.1 -1095.8   2191.5     2009
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
## -5.1404 -1.0086  0.4328  0.6651  1.0750
##
## Random effects:
```

```
## Groups      Name                Variance Std.Dev. Corr
## Participant (Intercept)          0.289995 0.5385
##           Block                  0.009546 0.0977    0.75
##           Taste_diff.abs         0.118349 0.3440    0.93 0.57
##           Block:Taste_diff.abs    0.034139 0.1848   -0.31 0.38 -0.52
## Number of obs: 2023, groups: Participant, 23
##
## Fixed effects:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)      1.18664    0.12915   9.188 < 2e-16 ***
## Block            -0.06374    0.06671  -0.955   0.339
## Taste_diff.abs     0.64847    0.09585   6.765 1.33e-11 ***
## Block:Taste_diff.abs -0.05164    0.07412  -0.697   0.486
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##              (Intr) Block  Tst_d.
## Block          0.044
## Tast_dff.bs    0.738  0.025
## Blck:Tst_d.   -0.212  0.386 -0.387
## convergence code: 0
## Model failed to converge with max|grad| = 0.00348866 (tol = 0.002, component 1)
```

Taste trials: immediate and after effect in sham condition

In sham session participants performed better during the first part of the choice task, but then accuracy dropped for the taste trials, in particular.

This pattern was not present in condition when TI was applied.

```
idx = which(dataVal$Session_type==0) # Table: sham sessions
dataVal2 = dataVal[idx,]

m2.val.stim2.abs = glmer(corr ~ Block*Taste_diff.abs + (1+Block*Taste_diff.abs|Participant), data=dataVal2)

summary(m2.val.stim2.abs)
```

```
## Generalized linear mixed model fit by maximum likelihood (Laplace
## Approximation) [glmerMod]
## Family: binomial ( logit )
## Formula: corr ~ Block * Taste_diff.abs + (1 + Block * Taste_diff.abs |
## Participant)
## Data: dataVal2
##
##      AIC      BIC    logLik deviance df.resid
## 2216.2  2294.8 -1094.1  2188.2     2007
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
## -4.8359 -1.0082  0.4289  0.6754  1.2560
##
## Random effects:
## Groups      Name                Variance Std.Dev. Corr
## Participant (Intercept)          0.155162 0.39391
##           Block                  0.043687 0.20901    0.36
```

```
##           Taste_diff.abs          0.103180 0.32122   0.93  0.67
##           Block:Taste_diff.abs 0.000597 0.02443   0.69 -0.43  0.37
## Number of obs: 2021, groups:  Participant, 23
##
## Fixed effects:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)      1.11199    0.10147  10.959 < 2e-16 ***
## Block            0.17649    0.07384   2.390  0.0168 *
## Taste_diff.abs    0.65682    0.08992   7.305 2.78e-13 ***
## Block:Taste_diff.abs 0.09207    0.06019   1.529  0.1261
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##              (Intr) Block  Tst_d.
## Block          0.108
## Tast_dff.bs    0.695  0.312
## Blck:Tst_d.    0.063  0.253 -0.082
## convergence code: 0
## boundary (singular) fit: see ?isSingular
```

Size trials: immediate and after effect in active stimulation condition

No significant effect of stimulation on the size trials.

```
idx = which(dataPer$Session_type==1)          # Table: active sessions
dataPer2 = dataPer[idx,]

m2.per.stim1.abs = glmer(corr ~ Block*Size_diff.abs + (1+Block*Size_diff.abs|Participant), data=dataPer2)

summary(m2.per.stim1.abs)
```

```
## Generalized linear mixed model fit by maximum likelihood (Laplace
## Approximation) [glmerMod]
## Family: binomial ( logit )
## Formula:
## corr ~ Block * Size_diff.abs + (1 + Block * Size_diff.abs | Participant)
## Data: dataPer2
##
##      AIC      BIC    logLik deviance df.resid
##  2221.1   2299.6 -1096.6   2193.1     2004
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
## -4.0695 -0.9779  0.4139  0.6643  1.2048
##
## Random effects:
## Groups      Name                Variance Std.Dev. Corr
## Participant (Intercept)      0.33833  0.5817
## Block              0.01567  0.1252   -0.68
## Size_diff.abs        0.09491  0.3081    0.91 -0.32
## Block:Size_diff.abs 0.01100  0.1049   -0.87  0.23 -0.99
## Number of obs: 2018, groups:  Participant, 23
##
## Fixed effects:
```



```
##               Estimate Std. Error z value Pr(>|z|)
## (Intercept)      1.14842    0.13746   8.354 < 2e-16 ***
## Block            -0.06822    0.06867  -0.993   0.321
## Size_diff.abs     0.73725    0.09050   8.146 3.76e-16 ***
## Block:Size_diff.abs -0.04329    0.06652  -0.651   0.515
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##      (Intr) Block  Sz_df.
## Block      -0.387
## Size_diff.bs 0.711 -0.205
## Blck:Sz_df. -0.331 0.378 -0.451
## convergence code: 0
## Model failed to converge with max|grad| = 0.0204894 (tol = 0.002, component 1)
```

Size trials: immediate and after effect in sham condition

No significant effect of stimulation on the size trials.

```
idx = which(dataPer$Session_type==0) # Table: sham sessions
dataPer2 = dataPer[idx,]

m2.per.stim2.abs = glmer(corr ~ Block*Size_diff.abs + (1+Block*Size_diff.abs|Participant), data=dataPer)

summary(m2.per.stim2.abs)
```

```
## Generalized linear mixed model fit by maximum likelihood (Laplace
## Approximation) [glmerMod]
## Family: binomial ( logit )
## Formula:
## corr ~ Block * Size_diff.abs + (1 + Block * Size_diff.abs | Participant)
## Data: dataPer2
##
##      AIC      BIC    logLik deviance df.resid
##  2148.6   2227.2 -1060.3   2120.6     2004
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
## -4.5082 -0.9491  0.4017  0.6325  1.2548
##
## Random effects:
## Groups      Name                Variance Std.Dev. Corr
## Participant (Intercept)      0.086558 0.29421
## Block              0.031933 0.17870   1.00
## Size_diff.abs        0.066072 0.25705   1.00 1.00
## Block:Size_diff.abs 0.001059 0.03254  -0.19 -0.20 -0.15
## Number of obs: 2018, groups: Participant, 23
##
## Fixed effects:
##               Estimate Std. Error z value Pr(>|z|)
## (Intercept)      1.19631    0.08698  13.754 <2e-16 ***
## Block              0.06322    0.07186   0.880   0.379
## Size_diff.abs     0.78505    0.08227   9.542 <2e-16 ***
## Block:Size_diff.abs -0.01380    0.06252  -0.221   0.825
```

```
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##           (Intr) Block  Sz_df.
## Block           0.249
## Size_dff.bs     0.671  0.308
## Blck:Sz_df.    -0.046  0.320 -0.172
## convergence code: 0
## unable to evaluate scaled gradient
## Model failed to converge: degenerate Hessian with 1 negative eigenvalues
```

PLOTS

Plot 1. Plot betas from the first 4 models for between-session comparison.

```
plot_betas = as.data.table(matrix(c(s1$coefficients[2,1], s2$coefficients[2,1],
                                   s3$coefficients[2,1], s4$coefficients[2,2],
                                   s1$coefficients[2,2], s2$coefficients[2,2],
                                   s3$coefficients[2,2], s4$coefficients[2,2]), nrow = 4, ncol = 2))


colnames(plot_betas) = c('beta_mean', 'beta_std')
plot_betas$cue = rep(c("Value-based", "Perceptual"), each = 2)
plot_betas$cue = factor(plot_betas$cue, levels = c('Value-based', 'Perceptual'))

plot_betas$stim = rep(c("During stimulation", "After stimulation"), 2)
plot_betas$stim = factor(plot_betas$stim, levels = c('During stimulation', 'After stimulation'))

p_betas = ggplot(plot_betas, aes(x=stim, y=beta_mean, fill = cue)) +
  geom_bar(stat="identity", position=position_dodge(), alpha = 0.5, color = 'black') +
  geom_errorbar(aes(ymin=beta_mean-beta_std, ymax=beta_mean+beta_std), width=.1,
               position=position_dodge(.9)) +
  geom_point(position=position_dodge(.9), show_guide = FALSE) +
  theme_bw() +
  ylab('Beta estimates') +
  xlab(' ') +
  scale_fill_manual(values=c('darkred', 'grey48')) +
  #ylim(c(0, 1)) +
  ggtitle('Impact of stimulation condition on choice consistency') +
  #scale_y_continuous(limits = c(0.5,1), ) +
  coord_cartesian(ylim= c(-0.2,0.3)) +
  #geom_signif(comparisons = list(c("Immediat effects", "After effects")), annotation = c('*'))
  theme(text = element_text(size=16), axis.text.x = element_text(size=16, colour = 'black'),
        axis.text.y = element_text(size=16), legend.text=element_text(size=16)) +
  theme(legend.position = c(0.2, 0.87)) +
  guides(fill=guide_legend(title="Trials")) +
  theme(legend.background=element_blank()) +
  annotate('text', x = 0.78, y = -0.18, label = 'NS') +
  annotate('text', x = 1.22, y = -0.18, label = 'NS') +
  annotate('text', x = 1.77, y = 0.26, label = '*') +
  annotate('text', x = 2.22, y = 0.22, label = 'NS')

p_betas
```

Impact of stimulation condition on choice consistency



```
#plot_dir = 'C:/Users/vbeliaev/Documents/TI_fmRI/pre_registration_paper/methods/paper_scripts_v4'
#ggsave('p_betas_stim.png', path = plot_dir, dpi=300)
```

Plot 2. Plot raw data for between-session comparison.

```
##### plot during stimulation
```

```
idx = which(dataVal$Trial_Nr<113)
imm_data_val = as.data.table(dataVal[idx,])
imm_data_val$Trial_type = c('Taste')
```

```
idx = which(dataPer$Trial_Nr<113)
imm_data_per = as.data.table(dataPer[idx,])
imm_data_per$Trial_type = c('Size')
```

```
tmp_val = ddply(imm_data_val, .(Participant, Session_type), summarise, acc = mean(corr))
tmp_val = ddply(tmp_val, .(Session_type), summarise, acc2 = mean(acc), se=sd(acc)/sqrt(ns))
tmp_val$cued = "Value-based\ntrials"
colnames(tmp_val)[1] = "Stim"
```

```
tmp_per = ddply(imm_data_per, .(Participant, Session_type), summarise, acc = mean(corr))
tmp_per = ddply(tmp_per, .(Session_type), summarise, acc2 = mean(acc), se=sd(acc)/sqrt(ns))
tmp_per$cued = "Perceptual\ntrials"
colnames(tmp_per)[1] = "Stim"
```

```
tmp_val_per = rbind(tmp_val,tmp_per)
tmp_val_per$Stim = factor(tmp_val_per$Stim, levels = c(1,0), labels = c('Active','Sham'))
```

```

tmp_val_per$cued = factor(tmp_val_per$cued, labels = c('Perceptual\ntrials', 'Value-based\ntrials'))
tmp_val_per$cued = ordered(tmp_val_per$cued, levels = c('Value-based\ntrials', 'Perceptual\ntrials'))

limits = aes(ymax = acc2+se, ymin=acc2-se, colour=cued, group=cued)

plot_during = ggplot(tmp_val_per, aes(x=cued, y=acc2, fill = Stim)) +
  geom_bar(stat="identity", position=position_dodge(), alpha = 0.5, color = 'black') +
  geom_errorbar(aes(ymin=acc2-se, ymax=acc2+se), width=.1,
    position=position_dodge(.9)) +
  geom_point(position=position_dodge(.9), show_guide = FALSE) +
  theme_bw() +
  ylab('Choice consistency') +
  xlab(' ') +
  scale_fill_manual(values=c('darkred', 'grey48')) +
  #ylim(c(0, 1)) +
  ggtitle('During stimulation') +
  theme(legend.title = element_blank()) +
  #scale_y_continuous(limits = c(0.5,1), ) +
  coord_cartesian(ylim= c(0.6,0.81)) +
  #geom_signif(comparisons = list(c("Immidiate effects", "After effects")), annotation = c('*'))
  theme(legend.position = c(0.5, 0.93)) +
  theme(legend.background=element_blank()) +
  theme(text = element_text(size=16), axis.text.x = element_text(size=12, colour = 'black'),
    axis.text.y = element_text(size=16), legend.text=element_text(size=12)) +
  theme(axis.title.y = element_text(vjust=2)) +
  geom_signif(y_position = c(0.78, 0.78), xmin = c(0.8, 1.8), xmax = c(1.2, 2.2),
    annotation = c("NS", 'NS'), tip_length = 0.1)

#plot_during

##### plot after stimulation

idx = which(dataVal$Trial_Nr>=113)
after_data_val = as.data.table(dataVal[idx,])
after_data_val$Trial_type = c('Taste')

idx = which(dataPer$Trial_Nr>=113)
after_data_per = as.data.table(dataPer[idx,])
after_data_per$Trial_type = c('Size')

tmp_val = ddply(after_data_val, .(Participant, Session_type), summarise, acc = mean(corr))
tmp_val = ddply(tmp_val, .(Session_type), summarise, acc2 = mean(acc), se=sd(acc)/sqrt(ns))
tmp_val$cued = "Value-based\ntrials"
colnames(tmp_val)[1] = "Stim"

tmp_per = ddply(after_data_per, .(Participant, Session_type), summarise, acc = mean(corr))
tmp_per = ddply(tmp_per, .(Session_type), summarise, acc2 = mean(acc), se=sd(acc)/sqrt(ns))
tmp_per$cued = "Perceptual\ntrials"
colnames(tmp_per)[1] = "Stim"

tmp_val_per = rbind(tmp_val,tmp_per)
tmp_val_per$Stim = factor(tmp_val_per$Stim, levels = c(1,0), labels = c('Active','Sham'))
tmp_val_per$cued = factor(tmp_val_per$cued, labels = c('Perceptual\ntrials', 'Value-based\ntrials'))

```

```

tmp_val_per$cued = ordered(tmp_val_per$cued, levels = c('Value-based\ntrials', 'Perceptual\ntrials'))


limits = aes(ymax = acc2+se, ymin=acc2-se, colour=cued, group=cued)

plot_after = ggplot(tmp_val_per, aes(x=cued, y=acc2, fill = Stim)) +
  geom_bar(stat="identity", position=position_dodge(), alpha = 0.5, color = 'black') +
  geom_errorbar(aes(ymin=acc2-se, ymax=acc2+se), width=.1,
    position=position_dodge(.9)) +
  geom_point(position=position_dodge(.9), show_guide = FALSE) +
  theme_bw() +
  ylab(' ') +
  xlab(' ') +
  scale_fill_manual(values=c('darkred', 'grey48')) +
  #ylim(c(0, 1)) +
  ggtitle('After stimulation') +
  theme(legend.title = element_blank()) +
  #scale_y_continuous(limits = c(0.5,1), ) +
  coord_cartesian(ylim= c(0.6,0.81)) +
  #geom_signif(comparisons = list(c("Immediate effects", "After effects")), annotation = c('*'))
  theme(legend.position = 'none') +
  theme(text = element_text(size=16), axis.text.x = element_text(size=12, colour = 'black'),
    axis.text.y = element_text(size=16)) +
  theme(axis.title.y = element_text(vjust=2)) +
  geom_signif(y_position = c(0.78, 0.78), xmin = c(0.8, 1.8), xmax = c(1.2, 2.2),
    annotation = c("?", 'NS'), tip_length = 0.1)

#plot_after

grid.arrange(plot_during, plot_after, ncol=2, widths=c(1,1))

```



```
#g = arrangeGrob(plot_during, plot_after, ncol=2)
#ggsave('p_stim.png', g, path = plot_dir, dpi=300)
```